

AN INTRODUCTION TO
PROBABILITY AND
STATISTICAL INFERENCE



ROUSSAS



Introduction to Probability and Statistical Inference

This Page Intentionally Left Blank



Introduction to Probability and Statistical Inference

George Roussas

University of California, Davis



ACADEMIC PRESS

An imprint of Elsevier Science

Amsterdam Boston London New York Oxford Paris
San Diego San Francisco Singapore Sydney Tokyo

Senior Sponsoring Editor	Barbara Holland
Project Manager	Nancy Zachor
Editorial Coordinator	Tom Singer
Cover Design	Shawn Girsberger
Copyeditor	Mary Prescott
Composition	International Typesetting and Composition
Printer	Maple-Vail

This book is printed on acid-free paper. ∞

Copyright 2003, Elsevier Science (USA)

All rights reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Requests for permission to make copies of any part of the work should be mailed to: Permissions Department, Harcourt, Inc., 6277 Sea Harbor Drive, Orlando, Florida 32887-6777.

Academic Press
An imprint of Elsevier Science
525 B Street, Suite 1900, San Diego, California 92101-4495, USA
<http://www.academicpress.com>

Academic Press
An imprint of Elsevier Science
200 Wheeler Road, Burlington, Massachusetts 01803, USA
<http://www.academicpressbooks.com>

Academic Press
An imprint of Elsevier Science
84 Theobald's Road, London WC1X 8RR, UK
<http://www.academicpress.com>

Library of Congress Control Number: 2002110812
International Standard Book Number: 0-12-599020-0

PRINTED IN THE UNITED STATES OF AMERICA
02 03 04 05 06 9 8 7 6 5 4 3 2 1

*To my wife and sons,
and the unforgettable Beowulf*

This Page Intentionally Left Blank



Contents

<i>Preface</i>		xi
1	SOME MOTIVATING EXAMPLES AND SOME FUNDAMENTAL CONCEPTS	1
1.1	Some Motivating Examples	1
1.2	Some Fundamental Concepts	8
1.3	Random Variables	19
2	THE CONCEPT OF PROBABILITY AND BASIC RESULTS	23
2.1	Definition of Probability and Some Basic Results	24
2.2	Distribution of a Random Variable	33
2.3	Conditional Probability and Related Results	41
2.4	Independent Events and Related Results	51
2.5	Basic Concepts and Results in Counting	59
3	NUMERICAL CHARACTERISTICS OF A RANDOM VARIABLE, SOME SPECIAL RANDOM VARIABLES	68
3.1	Expectation, Variance, and Moment Generating Function of a Random Variable	68
3.2	Some Probability Inequalities	77
3.3	Some Special Random Variables	79
3.4	Median and Mode of a Random Variable	102
4	JOINT AND CONDITIONAL P.D.F.'S, CONDITIONAL EXPECTATION AND VARIANCE, MOMENT GENERATING FUNCTION, COVARIANCE, AND CORRELATION COEFFICIENT	109
4.1	Joint d.f. and Joint p.d.f. of Two Random Variables	110
4.2	Marginal and Conditional p.d.f.'s, Conditional Expectation and Variance	117
4.3	Expectation of a Function of Two r.v.'s, Joint and Marginal m.g.f.'s, Covariance, and Correlation Coefficient	126
4.4	Some Generalizations to k Random Variables	137
4.5	The Multinomial, the Bivariate Normal, and the Multivariate Normal Distributions	139

5	INDEPENDENCE OF RANDOM VARIABLES AND SOME APPLICATIONS	150
<hr/>		
5.1	Independence of Random Variables and Criteria of Independence	150
5.2	The Reproductive Property of Certain Distributions	159
6	TRANSFORMATION OF RANDOM VARIABLES	168
<hr/>		
6.1	Transforming a Single Random Variable	168
6.2	Transforming Two or More Random Variables	173
6.3	Linear Transformations	185
6.4	The Probability Integral Transform	192
6.5	Order Statistics	193
7	SOME MODES OF CONVERGENCE OF RANDOM VARIABLES, APPLICATIONS	202
<hr/>		
7.1	Convergence in Distribution or in Probability and Their Relationship	202
7.2	Some Applications of Convergence in Distribution: The Weak Law of Large Numbers and the Central Limit Theorem	208
7.3	Further Limit Theorems	222
8	AN OVERVIEW OF STATISTICAL INFERENCE	227
<hr/>		
8.1	The Basics of Point Estimation	228
8.2	The Basics of Interval Estimation	230
8.3	The Basics of Testing Hypotheses	231
8.4	The Basics of Regression Analysis	235
8.5	The Basics of Analysis of Variance	236
8.6	The Basics of Nonparametric Inference	238
9	POINT ESTIMATION	240
<hr/>		
9.1	Maximum Likelihood Estimation: Motivation and Examples	240
9.2	Some Properties of Maximum Likelihood Estimates	253
9.3	Uniformly Minimum Variance Unbiased Estimates	261
9.4	Decision-Theoretic Approach to Estimation	270
9.5	Other Methods of Estimation	277
10	CONFIDENCE INTERVALS AND CONFIDENCE REGIONS	281
<hr/>		
10.1	Confidence Intervals	282
10.2	Confidence Intervals in the Presence of Nuisance Parameters	289

10.3	A Confidence Region for (μ, σ^2) in the $N(\mu, \sigma^2)$ Distribution	292
10.4	Confidence Intervals with Approximate Confidence Coefficient	294
11	TESTING HYPOTHESES	299
<hr/>		
11.1	General Concepts, Formulation of Some Testing Hypotheses	300
11.2	Neyman–Pearson Fundamental Lemma, Exponential Type Families, Uniformly Most Powerful Tests for Some Composite Hypotheses	302
11.3	Some Applications of Theorems 2 and 3	315
11.4	Likelihood Ratio Tests	324
12	MORE ABOUT TESTING HYPOTHESES	343
<hr/>		
12.1	Likelihood Ratio Tests in the Multinomial Case and Contingency Tables	343
12.2	A Goodness-of-Fit Test	349
12.3	Decision-Theoretic Approach to Testing Hypotheses	353
12.4	Relationship Between Testing Hypotheses and Confidence Regions	360
13	A SIMPLE LINEAR REGRESSION MODEL	363
<hr/>		
13.1	Setting-up the Model — The Principle of Least Squares	364
13.2	The Least Squares Estimates of β_1 and β_2, and Some of Their Properties	366
13.3	Normally Distributed Errors: MLE's of β_1, β_2, and σ^2, Some Distributional Results	374
13.4	Confidence Intervals and Hypotheses Testing Problems	383
13.5	Some Prediction Problems	389
13.6	Proof of Theorem 5	393
13.7	Concluding Remarks	395
14	TWO MODELS OF ANALYSIS OF VARIANCE	397
<hr/>		
14.1	One-Way Layout with the Same Number of Observations per Cell	398
14.2	A Multicomparison Method	407
14.3	Two-Way Layout with One Observation per Cell	412
15	SOME TOPICS IN NONPARAMETRIC INFERENCE	428
<hr/>		
15.1	Some Confidence Intervals with Given Approximate Confidence Coefficient	429
15.2	Confidence Intervals for Quantiles of a Distribution Function	431

15.3	The Two-Sample Sign Test	433
15.4	The Rank Sum and the Wilcoxon–Mann–Whitney Two-Sample Tests	435
15.5	Nonparametric Curve Estimation	442
APPENDIX		450
SOME NOTATION AND ABBREVIATIONS		480
ANSWERS TO EVEN-NUMBERED EXERCISES		483
INDEX		515



Preface

Overview

This book is an introductory textbook in probability and statistical inference. No prior knowledge of either probability or statistics is required, although prior exposure to an elementary precalculus course would prove beneficial in the sense that the student would not see the basic concepts discussed here for the first time.

The mathematical prerequisite is a year of calculus and familiarity with the basic concepts and some results of linear algebra. Elementary differential and integral calculus will suffice for the majority of the book. In some parts, such as Chapters 4, 5, and 6, the concept of a multiple integral is used. Also, in Chapter 6, the student is expected to be at least vaguely familiar with the basic techniques of changing variables in a single or a multiple integral.

Chapter Descriptions

The material discussed in this book is enough for a one-year course in introductory probability and statistical inference. It consists of a total of 15 chapters. Chapters 1 through 7 are devoted to probability, distributional theory, and related topics. Chapters 9 through 14 discuss the standard topics of parametric statistical inference, namely point estimation, interval estimation, and testing hypotheses. This is done first in a general setting and then in the special models of linear regression and analysis of variance. Chapter 15 is devoted to discussing selected topics from nonparametric inference.

Features

This book has a number of features that differentiate it from existing books. First, the material is arranged in such a manner that Chapters 1 through 8 can be used independently for an introductory course in probability. The desirable duration for such a course would be a semester, although a quarter would

also be long enough if some of the proofs were omitted. Chapters 1 through 7 would suffice for this purpose. The centrally placed Chapter 8 plays a twofold role. First, it serves as a window into what statistical inference is all about for those taking only the probability part of the course. Second, it paints a fairly broad picture of the material discussed in considerable detail in the subsequent chapters. Accordingly and purposely, no specific results are stated, no examples are discussed, no exercises are included. All these things are done in the chapters following it. As already mentioned, the sole objective here is to take the reader through a brief orientation trip to statistical inference; to indicate why statistical inference is needed in the first place, how the relevant main problems are formulated, and how we go about resolving them.

The second differentiating feature of the book is the relatively large number of examples discussed in detail. There are more than 220 such examples, not including scores of numerical examples and applications. The first chapter alone is replete with 44 examples selected from a variety of applications. Their purpose is to impress upon the student the breadth of applications of probability and statistics, to draw attention to the wide range of applications where probabilistic and statistical questions are pertinent. At this stage, one could not possibly provide answers to the questions posed without the methodology developed in the subsequent chapters. Answers to these questions are given in the form of examples and exercises throughout the remaining chapters.

The book contains more than 560 exercises placed strategically at the ends of sections. The exercises are closely related to the material discussed in the respective sections, and they vary in the degree of difficulty. Detailed solutions to all of them are available in the form of a *Solutions Manual* for the instructors of the course, when this textbook is used. Brief answers to even-numbered exercises are provided at the end of the book. Also included in the textbook are approximately 60 figures that help illustrate some concepts and operations.

Still another desirable feature of this textbook is the effort made to minimize the so-called arm waving. This is done by providing a substantial number of proofs, without ever exceeding the mathematical prerequisites set. This also helps ameliorate the not so unusual phenomenon of insulting students' intelligence by holding them incapable of following basic reasoning.

Regardless of the effort made by the author of an introductory book in probability and statistics to cover the largest possible number of areas where probability and statistics apply, such a goal is unlikely to be attained. Consequently, no such textbook will ever satisfy students who focus exclusively on their own area of interest. It is also expected that this book will come as a disappointment to students who are oriented more toward vocational training rather than college or university education. This book is not meant to codify answers to questions in the form of framed formulas and prescription recipes. Rather, its purpose is to introduce the student to a thinking process and guide her or him toward the answer sought to a posed question. To paraphrase a Chinese saying, if you are taught how to fish, you eat all the time, whereas if you are given a fish, you eat only once.

On several occasions the reader is referred for proofs and more comprehensive treatment of some topics to the book *A Course in Mathematical Statistics, 2nd edition* (1997), Academic Press, by G.G. Roussas. This reference book was originally written for the same audience as that of the present book. However, circumstances dictated the adjustment of the level of the reference book to match the mathematical preparation of the anticipated audience.

On the practical side, a number of points of information are given here. Thus, $\log x$ (logarithm of x), whenever it occurs, is always the natural logarithm of x (the logarithm of x with base e), whether it is explicitly stated or not.

The rule followed in the use of decimal numbers is that we retain three decimal digits, the last of which is rounded up to the next higher number, if the fourth omitted decimal is greater or equal 5. An exemption to this rule is made when the division is exact, and also when the numbers are read out of tables. The book is supplied with an appendix consisting of excerpts of tables: *Binomial tables, Poisson tables, Normal tables, t-tables, Chi-Square tables, and F-tables*. The last table, Table 7, consists of a list of certain often-occurring distributions along with some of their characteristics. The appendix is followed by a list of some notation and abbreviations extensively used throughout the book, and the body of the book is concluded with brief answers to the even-numbered exercises.

In closing, a concerted effort has been made to minimize the number of inevitable misprints and oversights in the book. We have no illusion, however, that the book is free of them. This author would greatly appreciate being informed of any errors; such errors will be corrected in a subsequent printing of the book.

Acknowledgments and Credits

I would like to thank Subhash Bagui, University of West Florida; Matthew Carlton, Cal Polytechnic State University; Tapas K. Das, University of South Florida; Jay Devore, Cal Polytechnic State University; Charles Donaghey, University of Houston; Pat Goeters, Auburn University; Xuming He, University of Illinois, and Krzysztof M. Ostaszewski, Illinois State University, Champaign-Urbana for their many helpful comments.

Some of the examples discussed in this book have been taken and/or adapted from material included in the book *Statistics: Principles and Methods*, 2nd edition (1992), [ISBN: 0471548421], by R. A. Johnson, G. K. Bhattacharyya, Copyright ©1987, 1992, by John Wiley & Sons, Inc., and are reprinted by permission of John Wiley & Sons, Inc. They are Table 4 on page 74, Examples 8, 1, 2, 4, 12, 4, 2, 1, and 7 on pages 170, 295, 296, 353, 408, 439, 510, 544, and 562, respectively, and Exercises 4.18, 3.19, 4.21, 5.22, 5.34, 8.16, 4.14, 6.34, 3.16, 6.6 and 3.8 on pages 123, 199, 217, 222, 225, 265, 323, 340, 356, 462, and 525, respectively. The reprinting permission is kindly acknowledged herewith.

This Page Intentionally Left Blank



Chapter 1

Some Motivating Examples and Some Fundamental Concepts

This chapter consists of three sections. The first section is devoted to presenting a number of examples (25 to be precise), drawn from a broad spectrum of human activities. Their purpose is to demonstrate the wide applicability of probability and statistics. In the formulation of these examples, certain terms, such as at random, average, data fit by a line, event, probability (estimated probability, probability model), rate of success, sample, and sampling (sample size), are used. These terms are presently to be understood in their everyday sense, and will be defined precisely later on.

In the second section, some basic terminology and fundamental quantities are introduced and are illustrated by means of examples. In the closing section, the concept of a random variable is defined and is clarified through a number of examples.

1.1 Some Motivating Examples

EXAMPLE 1

In a certain state of the Union, n landfills are classified according to their concentration of three hazardous chemicals: arsenic, barium, and mercury. Suppose that the concentration of each one of the three chemicals is characterized as either high or low. Then some of the questions which can be posed are as follows: (i) If a landfill is chosen at random from among the n , what is the probability it is of a specific configuration? In particular, what is the probability that it has: (a) High concentration of barium? (b) High concentration of mercury and low concentration of both arsenic and barium? (c) High


concentration of any two of the chemicals and low concentration of the third?
 (d) High concentration of any one of the chemicals and low concentration of the other two? (ii) How can one check whether the proportions of the landfills falling into each one of the eight possible configurations (regarding the levels of concentration) agree with a priori stipulated numbers?

EXAMPLE 2

Suppose a disease is present in $100p_1\%$ ($0 < p_1 < 1$) of a population. A diagnostic test is available but is yet to be perfected. The test shows $100p_2\%$ false positives ($0 < p_2 < 1$) and $100p_3\%$ false negatives ($0 < p_3 < 1$). That is, for a patient not having the disease, the test shows positive (+) with probability p_2 and negative (−) with probability $1 - p_2$. For a patient having the disease, the test shows “−” with probability p_3 and “+” with probability $1 - p_3$. A person is chosen at random from the target population, and let D be the event that the person is diseased and N be the event that the person is not diseased. Then, it is clear that some important questions are as follows: In terms of p_1 , p_2 , and p_3 : (i) Determine the probabilities of the following configurations: D and +, D and −, N and +, N and −. (ii) Also, determine the probability that a person will test + or the probability the person will test −. (iii) If the person chosen tests +, what is the probability that he/she is diseased? What is the probability that he/she is diseased, if the person tests −?

EXAMPLE 3

In the circuit drawn below, suppose that switch $i = 1, \dots, 5$ turns on with probability p_i and independently of the remaining switches. What is the probability of having current transferred from point A to point B?

**EXAMPLE 4**

A travel insurance policy pays \$1,000 to a customer in case of a loss due to theft or damage on a 5-day trip. If the risk of such a loss is assessed to be 1 in 200, what is a fair premium for this policy?

EXAMPLE 5

Jones claims to have extrasensory perception (ESP). In order to test the claim, a psychologist shows Jones five cards that carry different pictures. Then Jones is blindfolded and the psychologist selects one card and asks Jones to identify

the picture. This process is repeated n times. Suppose, in reality, that Jones has no ESP but responds by sheer guesses.

- (i) Decide on a suitable probability model describing the number of correct responses.
- (ii) What is the probability that at most $n/5$ responses are correct?
- (iii) What is the probability that at least $n/2$ responses are correct?

EXAMPLE 6

A government agency wishes to assess the prevailing rate of unemployment in a particular county. It is felt that this assessment can be done quickly and effectively by sampling a small fraction n , say, of the labor force in the county. The obvious questions to be considered here are: (i) What is a suitable probability model describing the number of unemployed? (ii) What is an estimate of the rate of unemployment?

EXAMPLE 7

Suppose that, for a particular cancer, chemotherapy provides a 5-year survival rate of 80% if the disease could be detected at an early stage. Suppose further that n patients, diagnosed to have this form of cancer at an early stage, are just starting the chemotherapy. Finally, let X be the number of patients among the n who survive 5 years.

Then the following are some of the relevant questions which can be asked:

- (i) What are the possible values of X , and what are the probabilities that each one of these values is taken on?
- (ii) What is the probability that X takes values between two specified numbers a and b , say?
- (iii) What is the average number of patients to survive 5 years, and what is the variation around this average?

EXAMPLE 8

An advertisement manager for a radio station claims that over 100p% ($0 < p < 1$) of all young adults in the city listen to a weekend music program. To establish this conjecture, a random sample of size n is taken from among the target population and those who listen to the weekend music program are counted.

- (i) Decide on a suitable probability model describing the number of young adults who listen to the weekend music program.
- (ii) On the basis of the collected data, check whether the claim made is supported or not.
- (iii) How large a sample size n should be taken to ensure that the estimated average and the true proportion do not differ in absolute value by more than a specified number with prescribed (high) probability?

EXAMPLE 9

When the output of a production process is stable at an acceptable standard, it is said to be “in control.” Suppose that a production process has been in control for some time and that the proportion of defectives has been p . As a means of monitoring the process, the production staff will sample n items. Occurrence of k or more defectives will be considered strong evidence for “out of control.”

- (i) Decide on a suitable probability model describing the number X of defectives; what are the possible values of X , and what is the probability that each of

these values is taken on? (ii) On the basis of the data collected, check whether or not the process is out of control. (iii) How large a sample size n should be taken to ensure that the estimated proportion of defectives will not differ in absolute value from the true proportion of defectives by more than a specified quantity with prescribed (high) probability?

EXAMPLE 10

An electronic scanner is believed to be more efficient in determining flaws in a material than a mechanical testing method which detects $100p\%$ ($0 < p < 1$) of the flawed specimens. To determine its success rate, n specimens with flaws are tested by the electronic scanner.

(i) Decide on a suitable probability model describing the number X of the flawed specimens correctly detected by the electronic scanner; what are the possible values of X , and what is the probability that each one of these values is taken on? (ii) Suppose that the electronic scanner detects correctly k out of n flawed specimens. Check whether or not the rate of success of the electronic scanner is higher than that of the mechanical device.

EXAMPLE 11

At a given road intersection, suppose that X is the number of cars passing by until an observer spots a particular make of a car (e.g., a Mercedes).

Then some of the questions one may ask are as follows: (i) What are the possible values of X ? (ii) What is the probability that each one of these values is taken on? (iii) How many cars would the observer expect to observe until the first Mercedes appears?

EXAMPLE 12

A city health department wishes to determine whether the mean bacteria count per unit volume of water at a lake beach is within the safety level of 200. A researcher collected n water samples of unit volume and recorded the bacteria counts.

Relevant questions here are: (i) What is the appropriate probability model describing the number X of bacteria in a unit volume of water; what are the possible values of X , and what is the probability that each one of these values is taken on? (ii) Do the data collected indicate that there is no cause for concern?

EXAMPLE 13

Consider an aptitude test administered to aircraft pilot trainees, which requires a series of operations to be performed in quick succession.

Relevant questions here are: (i) What is the appropriate probability model for the time required to complete the test? (ii) What is the probability that the test is completed in no less than t_1 minutes, say? (iii) What is the percentage of candidates passing the test, if the test is to be completed within t_2 minutes, say?

EXAMPLE 14

Measurements of the acidity (pH) of rain samples were recorded at n sites in an industrial region.

(i) Decide on a suitable probability model describing the number X of the acidity of rain measured. (ii) On the basis of the measurements taken, provide an estimate of the average acidity of rain in that region.

EXAMPLE 15

To study the growth of pine trees at an early state, a nursery worker records n measurements of the heights of 1-year-old red pine seedlings.

(i) Decide on a suitable probability model describing the heights X of the pine seedlings. (ii) On the basis of the n measurements taken, determine average height of the pine seedlings. (iii) Also, check whether these measurements support the stipulation that the average height is a specified number.

EXAMPLE 16

It is claimed that a new treatment is more effective than the standard treatment for prolonging the lives of terminal cancer patients. The standard treatment has been in use for a long time, and from records in medical journals the mean survival period is known to have a certain numerical value (in years). The new treatment is administered to n patients, and their duration of survival is recorded.

(i) Decide on suitable probability models describing the survival times X and Y under the old and the new treatments, respectively. (ii) On the basis of the existing journal information and the data gathered, check whether or not the claim made is supported.

EXAMPLE 17

A medical researcher wishes to determine whether a pill has the undesirable side effect of reducing the blood pressure of the user. The study requires recording the initial blood pressures of n college-age women. After the use of the pill regularly for 6 months, their blood pressures are again recorded.

(i) Decide on suitable probability models describing the blood pressures, initially and after the 6-month period. (ii) Do the observed data support the claim that the use of the pill reduces blood pressure?

EXAMPLE 18

It is known that human blood is classified in four types denoted by A, B, AB, and O. Suppose that the blood of n persons who have volunteered to donate blood at a plasma center has been classified in these four categories. Then a number of questions can be posed; some of them are:

(i) What is the appropriate probability model to describe the distribution of the blood types of the n persons into the four types? (ii) What is the estimated probability that a person, chosen at random from among the n , has a specified blood type (e.g., O)? (iii) What are the proportions of the n persons falling into each one of the four categories? (iv) How can one check whether the observed proportions are in agreement with a priori stipulated numbers?

EXAMPLE 19

The following record shows a classification of 41,208 births in Wisconsin (courtesy of Professor Jerome Klotz). Set up a suitable probability model and check whether or not the births are uniformly distributed over all 12 months of the year.

<i>Jan.</i>	3,478	<i>July</i>	3,476
<i>Feb.</i>	3,333	<i>Aug.</i>	3,495
<i>March</i>	3,771	<i>Sept.</i>	3,490
<i>April</i>	3,542	<i>Oct.</i>	3,331
<i>May</i>	3,479	<i>Nov.</i>	3,188
<i>June</i>	3,304	<i>Dec.</i>	3,321
		<i>Total</i>	41,208

EXAMPLE 20

To compare the effectiveness of two diets *A* and *B*, 150 infants were included in a study. Diet *A* was given to 80 randomly selected infants and diet *B* was given to the other 70 infants. At a later time, the health of each infant was observed and classified into one of the three categories: “excellent,” “average,” and “poor.” The frequency counts are tabulated as follows:

HEALTH UNDER TWO DIFFERENT DIETS				
	Excellent	Average	Poor	Sample Size
Diet A	37	24	19	80
Diet B	17	33	20	70
Total	54	57	39	150

Set up a suitable probability model for this situation, and, on the basis of the observed data, compare the effectiveness of the two diets.

EXAMPLE 21

Osteoporosis (loss of bone minerals) is a common cause of broken bones in the elderly. A researcher on aging conjectures that bone mineral loss can be reduced by regular physical therapy or by certain kinds of physical activity. A study is conducted on n elderly subjects of approximately the same age divided into control, physical therapy, and physical activity groups. After a suitable period of time, the nature of change in bone mineral content is observed.

Set up a suitable probability model for the situation under consideration, and check whether or not the observed data indicate that the change in bone mineral varies for different groups.

CHANGE IN BONE MINERAL				
	Appreciable Loss	Little Change	Appreciable Increase	Total
Control	38	15	7	60
Therapy	22	32	16	70
Activity	15	30	25	70
Total	75	77	48	200

EXAMPLE 22

In the following table, the data x = undergraduate GPA and y = score in the Graduate Management Aptitude Test (GMAT) are recorded.

DATA OF UNDERGRADUATE GPA (x) AND GMAT SCORE (y)					
x	y	x	y	x	y
3.63	447	2.36	399	2.80	444
3.59	588	2.36	482	3.13	416
3.30	563	2.66	420	3.01	471
3.40	553	2.68	414	2.79	490
3.50	572	2.48	533	2.89	431
3.78	591	2.46	509	2.91	446
3.44	692	2.63	504	2.75	546
3.48	528	2.44	336	2.73	467
3.47	552	2.13	408	3.12	463
3.35	520	2.41	469	3.08	440
3.39	543	2.55	538	3.03	419
				3.00	509

(i) Draw a scatter plot of the pairs (x, y) . (ii) On the basis of part (i), set up a reasonable model for the representation of the pairs (x, y) . (iii) Indicate roughly how this model can be used to predict a GMAT score on the basis of the corresponding GPA score.

EXAMPLE 23

In an experiment designed to determine the relationship between the doses of a compost fertilizer x and the yield y of a crop, n values of x and y are observed. On the basis of prior experience, it is reasonable to assume that the pairs (x, y) are fitted by a straight line, which can be determined by certain summary values of the data. Later on, it will be seen how this is specifically done and also how this model can be used for various purposes, including that of predicting a value of y on the basis of a given value of x .

EXAMPLE 24

In an effort to improve the quality of recording tapes, the effects of four kinds of coatings A, B, C, and D on the reproducing quality of sound are compared. Twenty two measurements of sound distortions are given in the following table.

SOUND DISTORTIONS OBTAINED WITH FOUR TYPES OF COATINGS	
Coating	Observations
A	10, 15, 8, 12, 15
B	14, 18, 21, 15
C	17, 16, 14, 15, 17, 15, 18
D	12, 15, 17, 15, 16, 15

In connection with these data, several questions may be posed (and will be posed later on). The most immediate of them all is the question of whether or not the data support the existence of any significant difference among the average distortions obtained using the four coatings.

EXAMPLE 25

Charles Darwin performed an experiment to determine whether self-fertilized and cross-fertilized plants have different growth rates. Pairs of *Zea mays* plants, one self- and the other cross-fertilized, were planted in pots, and their heights were measured after a specified period of time. The data Darwin obtained were:

Pair	PLANT HEIGHT (IN 1/8 INCHES)				
	Cross-	Self-	Pair	Cross-	Self-
1	188	139	9	146	132
2	96	163	10	173	144
3	168	160	11	186	130
4	176	160	12	168	144
5	153	147	13	177	102
6	172	149	14	184	124
7	177	149	15	96	144
8	163	122			

Source: Darwin, C., "The Effects of Cross- and Self-Fertilization in the Vegetable Kingdom," D. Appleton and Co., New York, 1902.

These data lead to many questions, the most immediate being whether cross-fertilized plants have a higher growth rate than self-fertilized plants. This example will be revisited later on.


1.2 Some Fundamental Concepts

One of the most basic concepts in probability and statistics is that of a *random experiment*. Although a more precise definition is possible, we will restrict ourselves here to understanding a random experiment as a procedure which is carried out under a certain set of conditions; it can be repeated any number of times under the same set of conditions, and upon the completion of the procedure certain results are observed. The results obtained are denoted by s and are called *sample points*. The set of all possible sample points is denoted by S and is called a *sample space*. Subsets of S are called *events* and are denoted by capital letters A, B, C , etc. An event consisting of one sample point only, $\{s\}$, is called a *simple event* and *composite* otherwise. An event A *occurs* (or *happens*) if the outcome of the random experiment (that is, the sample point s) belongs in A , $s \in A$; A *does not occur* (or *does not happen*) if $s \notin A$. The event S always occurs and is called the *sure* or *certain* event. On the other hand, the event \emptyset never happens and is called the *impossible* event. Of course, the relation $A \subseteq B$ between two events A and B means that the event B *occurs whenever A does*, but not necessarily the opposite. (See Figure 1.1 for the *Venn diagram* depicting the relation $A \subseteq B$.) The events A and B are *equal* if both $A \subseteq B$ and $B \subseteq A$.

Some random experiments are given in the following along with corresponding sample spaces and some events.

Figure 1.1

$A \subseteq B$; in Fact,
 $A \subset B$, Because
 $s_2 \in B$, But $s_2 \notin A$

**EXAMPLE 26**

Tossing three distinct coins once.

Then, with H and T standing for “heads” and “tails,” respectively, a sample space is:

$$S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}.$$

The event $A =$ “no more than 1 H occurs” is given by:

$$A = \{TTT, HTT, THT, TTH\}.$$

EXAMPLE 27

Rolling once two distinct dice.

Then a sample space is:

$$S = \{(1, 1), (1, 2), \dots, (1, 6), \dots, (6, 1), (6, 2), \dots, (6, 6)\},$$

and the event $B =$ “the sum of numbers on the upper faces is ≤ 5 ” is:

$$B = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (4, 1)\}.$$

EXAMPLE 28

Drawing a card from a well-shuffled standard deck of 52 cards. Denoting by C , D , H , and S clubs, diamonds, hearts, and spades, respectively, by J , Q , K Jack, Queen, and King, and using 1 for aces, the sample space is given by:

$$S = \{1_C, \dots, 1_S, \dots, 10_C, \dots, 10_S, \dots, K_C, \dots, K_S\}.$$

An event A may be described by: $A =$ “red and face card,” so that

$$A = \{J_D, J_H, Q_D, Q_H, K_D, K_H\}.$$

EXAMPLE 29

Drawing (without replacement) two balls from an urn containing m numbered black balls and n numbered red balls.

Then, in obvious notation, a sample space here is:

$$\begin{aligned} S = \{ & b_1 b_2, \dots, b_1 b_m, \dots, b_m b_1, \dots, b_m b_{m-1}, \\ & b_1 r_1, \dots, b_1 r_n, \dots, b_m r_1, \dots, b_m r_n, \\ & r_1 b_1, \dots, r_1 b_m, \dots, r_n b_1, \dots, r_n b_m, \\ & r_1 r_2, \dots, r_1 r_n, \dots, r_n r_1, \dots, r_n r_{n-1} \}. \end{aligned}$$

An event A may be the following: $A =$ “the sum of the numbers on the balls does not exceed 4.” Then

$$\begin{aligned} A = \{ & b_1 b_2, b_1 b_3, b_2 b_1, b_3 b_1, b_1 r_1, b_1 r_2, b_1 r_3, \\ & b_2 r_1, b_2 r_2, b_3 r_1, r_1 b_1, r_1 b_2, r_1 b_3, r_2 b_1, \\ & r_2 b_2, r_3 b_1, r_1 r_2, r_1 r_3, r_2 r_1, r_3 r_1 \} \quad (\text{assuming that } m, n \geq 3). \end{aligned}$$

EXAMPLE 30

Recording the gender of children of two-children families.

With b and g standing for boy and girl, and with the first letter on the left denoting the older child, a sample space is: $S = \{bb, bg, gb, gg\}$. An event B may be: $B =$ “children of both genders.” Then $B = \{bg, gb\}$.

EXAMPLE 31

Ranking five horses in a horse race.

Then the suitable sample space S consists of 120 sample points, corresponding to the 120 permutations of the numbers 1, 2, 3, 4, 5. (We exclude ties.) The event $A =$ “horse #3 comes second” consists of the 24 sample points, where 3 always occurs in the second place.

EXAMPLE 32

Tossing a coin repeatedly until H appears for the first time.

The suitable sample space here is:

$$S = \{H, TH, TTH, \dots, TT \dots TH, \dots\}.$$

Then the event $A =$ “the 1st H does not occur before the 10th tossing” is given by:

$$A = \{ \underbrace{T \dots T}_9 H, \underbrace{T \dots T}_{10} H, \dots \}.$$

EXAMPLE 33

Recording the number of telephone calls served by a certain telephone exchange center within a specified period of time.

Clearly, the sample space here is: $S = \{0, 1, \dots, C\}$, where C is a suitably large number associated with the capacity of the center. For mathematical convenience, we often take S to consist of all nonnegative integers; that is, $S = \{0, 1, \dots\}$.

EXAMPLE 34

Recording the number of traffic accidents which occurred in a specified location within a certain period of time.

As in the previous example, $\mathcal{S} = \{0, 1, \dots, M\}$ for a suitable number M . If M is sufficiently large, then \mathcal{S} is taken to be: $\mathcal{S} = \{0, 1, \dots\}$.

EXAMPLE 35

Recording the number of particles emitted by a certain radioactive source within a specified period of time.

As in the previous two examples, \mathcal{S} is taken to be: $\mathcal{S} = \{0, 1, \dots, M\}$, where M is often a large number, and then as before \mathcal{S} is modified to be: $\mathcal{S} = \{0, 1, \dots\}$.

EXAMPLE 36

Recording the lifetime of an electronic device, or of an electrical appliance, etc.

Here \mathcal{S} is the interval $(0, T)$ for some reasonable value of T ; that is, $\mathcal{S} = (0, T)$. Sometimes, for justifiable reasons, we take, $\mathcal{S} = (0, \infty)$.

EXAMPLE 37

Recording the distance from the bull's eye of the point where a dart, aiming at the bull's eye, actually hits the plane. Here it is clear that $\mathcal{S} = (0, \infty)$.

EXAMPLE 38

Measuring the dosage of a certain medication, administered to a patient, until a positive reaction is observed.

Here $\mathcal{S} = (0, D)$ for some suitable D (not rendering the medication lethal!).

EXAMPLE 39

Recording the yearly income of a target population.

If the incomes are measured in \$ and cents, the outcomes are fractional numbers in an interval $[0, M]$ for some reasonable M . Again, for reasons similar to those cited in Example 36, \mathcal{S} is often taken to be $\mathcal{S} = [0, \infty)$.

EXAMPLE 40

Waiting until the time the Dow–Jones Industrial Average index reaches or surpasses a specified level.

Here, with reasonable qualifications, we may chose to take $\mathcal{S} = (0, \infty)$.

Examples 1–25, suitably interpreted, may also serve as further illustrations of random experiments. All examples described previously will be revisited on various occasions.

For instance, in Example 1 and in self-explanatory notation, a suitable sample space is:

$$\mathcal{S} = \{A_h B_h M_h, A_h B_h M_\ell, A_h B_\ell M_h, A_\ell B_h M_h, A_h B_\ell M_\ell, \\ A_\ell B_h M_\ell, A_\ell B_\ell M_h, A_\ell B_\ell M_\ell\}.$$

Then the events $A =$ “no chemical occurs at high level” and $B =$ “at least two chemicals occur at high levels” are given by:

$$A = \{A_\ell B_\ell M_\ell\}, \quad B = \{A_h B_h M_\ell, A_h B_\ell M_h, A_\ell B_h M_h, A_h B_h M_h\}.$$

In Example 2, a patient is classified according to the result of the test, giving rise to the following sample space:

$$S = \{D+, D-, N+, N-\},$$

where D and N stand for the events “patient has the disease” and “patient does not have the disease,” respectively. Then the event $A =$ “false diagnosis of test” is given by: $A = \{D-, N+\}$.

In Example 5, the suitable probability model is the so-called binomial model. The sample space S is the set of 2^n points, each point consisting of a sequence of n S 's and F 's, S standing for success (on behalf of Jones) and F standing for failure. Then the questions posed can be answered easily.

Examples 6 through 10 can be discussed in the same framework as that of Example 5 with obvious modifications in notation.

In Example 11, a suitable sample space is:

$$S = \{M, M^c M, M^c M^c M, \dots, M^c \dots M^c M, \dots\},$$

where M stands for the passing by of a Mercedes car. Then the events A and B , where $A =$ “Mercedes was the 5th car passed by” and $B =$ “Mercedes was spotted after the first 3 cars passed by” are given by:

$$A = \{M^c M^c M^c M^c M\} \quad \text{and} \quad B = \{M^c M^c M^c M, M^c M^c M^c M^c M, \dots\}.$$

In Example 12, a suitable sample space is: $S = \{0, 1, \dots, M\}$ for an appropriately large (integer) M ; for mathematical convenience, S is often taken to be: $S = \{0, 1, 2, \dots\}$.

In Example 13, a suitable sample space is: $S = (0, T)$ for some reasonable value of T . In such cases, if T is very large, mathematical convenience dictates replacement of the previous sample space by: $S = (0, \infty)$.

Examples 14 and 15 can be treated in the same framework as Example 13 with obvious modifications in notation.

In Example 18, a suitable sample space S is the set of 4^n points, each point consisting of a sequence of n symbols A , B , AB , and O . The underlying probability model is the so-called multinomial model, and the questions posed can be discussed by available methodology. Actually, there is no need even to refer to the sample space S . All one has to do is to consider the outcomes in the n trials and then classify the n outcomes into four categories A , B , AB , and O .

Example 19 fits into the same framework as that of Example 18. Here the suitable \mathcal{S} consists of $12^{41,208}$ points, each point being a sequence of symbols representing the 12 months. As in the previous example, there is no need, however, even to refer to this sample space. Example 20 is also of the same type.


In many cases, questions posed can be discussed without reference to any explicit sample space. This is the case, for instance, in Examples 16–17 and 21–25.

In the examples discussed previously, we have seen sample spaces consisting of finitely many sample points (Examples 26–31), sample spaces consisting of countably infinite many points (for example, as many as the positive integers) (Example 32 and also Examples 33–35 if we replace C and M by ∞ for mathematical convenience), and sample spaces consisting of as many sample points as there are in a nondegenerate finite or infinite interval in the real line, which interval may also be the entire real line (Examples 36–40). Sample spaces with countably many points (i.e., either finitely many or countably infinite many) are referred to as *discrete* sample spaces. Sample spaces with sample points as many as the numbers in a nondegenerate finite or infinite interval in the real line $\mathfrak{R} = (-\infty, \infty)$ are referred to as *continuous* sample spaces.

Returning now to events, when one is dealing with them, one may perform the same operations as those with sets. Thus, the *complement* of the event A , denoted by A^c , is the event defined by: $A^c = \{s \in \mathcal{S}; s \notin A\}$. The event A^c is presented by the Venn diagram in Figure 1.2. So A^c occurs whenever A does not, and vice versa.

Figure 1.2

A^c Is the Shaded Region




The *union* of the events A_1, \dots, A_n , denoted by $A_1 \cup \dots \cup A_n$ or $\bigcup_{j=1}^n A_j$, is the event defined by $\bigcup_{j=1}^n A_j = \{s \in \mathcal{S}; s \in A_j, \text{ for at least one } j = 1, \dots, n\}$. So the event $\bigcup_{j=1}^n A_j$ occurs whenever at least one of $A_j, j = 1, \dots, n$ occurs. For $n = 2$, $A_1 \cup A_2$ is presented in Figure 1.3. The definition extends to an infinite number of events. Thus, for countably infinite many events $A_j, j = 1, 2, \dots$, one has $\bigcup_{j=1}^{\infty} A_j = \{s \in \mathcal{S}; s \in A_j, \text{ for at least one } j = 1, 2, \dots\}$.


The *intersection* of the events $A_j, j = 1, \dots, n$ is the event denoted by $A_1 \cap \dots \cap A_n$ or $\bigcap_{j=1}^n A_j$ and is defined by $\bigcap_{j=1}^n A_j = \{s \in \mathcal{S}; s \in A_j, \text{ for all } j = 1, \dots, n\}$. Thus, $\bigcap_{j=1}^n A_j$ occurs whenever all $A_j, j = 1, \dots, n$

Figure 1.3

$A_1 \cup A_2$ Is the
Shaded Region

**Figure 1.4**

$A_1 \cap A_2$ Is the
Shaded Region




occur simultaneously. For $n = 2$, $A_1 \cap A_2$ is presented in Figure 1.4. This definition extends to an infinite number of events. Thus, for countably infinite many events A_j , $j = 1, 2, \dots$, one has $\bigcap_{j=1}^{\infty} A_j = \{s \in S; s \in A_j, \text{ for all } j = 1, 2, \dots\}$.

If $A_1 \cap A_2 = \emptyset$, the events A_1 and A_2 are called *disjoint* (see Figure 1.5). The events A_j , $j = 1, 2, \dots$, are said to be *mutually* or *pairwise disjoint*, if $A_i \cap A_j = \emptyset$ whenever $i \neq j$.

Figure 1.5

A_1 and A_2 Are
Disjoint; That Is
 $A_i \cap A_j = \emptyset$




The *differences* $A_1 - A_2$ and $A_2 - A_1$ are the events defined by $A_1 - A_2 = \{s \in S; s \in A_1, s \notin A_2\}$, $A_2 - A_1 = \{s \in S; s \in A_2, s \notin A_1\}$ (see Figure 1.6).

From the definition of the preceding operations, the following properties follow immediately, and they are listed here for reference.

1. $S^c = \emptyset$, $\emptyset^c = S$, $(A^c)^c = A$.
2. $S \cup A = S$, $\emptyset \cup A = A$, $A \cup A^c = S$, $A \cup A = A$.
3. $S \cap A = A$, $\emptyset \cap A = \emptyset$, $A \cap A^c = \emptyset$, $A \cap A = A$.

Figure 1.6

$A_1 - A_2$ is $////$,
 $A_2 - A_1$ is $\\\\\\$



The previous statements are all obvious, as is the following: $\emptyset \subseteq A$ for every event A in S . Also,

4.
$$\left. \begin{aligned} A_1 \cup (A_2 \cup A_3) &= (A_1 \cup A_2) \cup A_3 \\ A_1 \cap (A_2 \cap A_3) &= (A_1 \cap A_2) \cap A_3 \end{aligned} \right\} \text{ (associative laws)}$$
5.
$$\left. \begin{aligned} A_1 \cup A_2 &= A_2 \cup A_1 \\ A_1 \cap A_2 &= A_2 \cap A_1 \end{aligned} \right\} \text{ (commutative laws)}$$
6.
$$\left. \begin{aligned} A \cap (\cup_j A_j) &= \cup_j (A \cap A_j) \\ A \cup (\cap_j A_j) &= \cap_j (A \cup A_j) \end{aligned} \right\} \text{ (distributive laws)}$$

In the last relations, as well as elsewhere, when the range of the index j is not indicated explicitly, it is assumed to be a finite set, such as $\{1, \dots, n\}$, or a countably infinite set, such as $\{1, 2, \dots\}$.

For the purpose of demonstrating some of the set-theoretic operations just defined, let us consider some further concrete examples.

EXAMPLE 41

Consider the sample space $S = \{s_1, s_2, s_3, s_4, s_5, s_6, s_7, s_8\}$ and define the events $A_1, A_2,$ and A_3 as follows: $A_1 = \{s_1, s_2, s_3\}$, $A_2 = \{s_2, s_3, s_4, s_5\}$, $A_3 = \{s_3, s_4, s_5, s_8\}$. Then observe that:

$$\begin{aligned} A_1^c &= \{s_4, s_5, s_6, s_7, s_8\}, & A_2^c &= \{s_1, s_6, s_7, s_8\}, & A_3^c &= \{s_1, s_2, s_6, s_7\}; \\ A_1 \cup A_2 &= \{s_1, s_2, s_3, s_4, s_5\}, & A_1 \cup A_3 &= \{s_1, s_2, s_3, s_4, s_5, s_8\}, \\ A_2 \cup A_3 &= \{s_2, s_3, s_4, s_5, s_8\}, & A_1 \cup A_2 \cup A_3 &= \{s_1, s_2, s_3, s_4, s_5, s_8\}; \\ A_1 \cap A_2 &= \{s_2, s_3\}, & A_1 \cap A_3 &= \{s_3\}, & A_1 \cap A_2 \cap A_3 &= \{s_3\}; \\ A_1 - A_2 &= \{s_1\}, & A_2 - A_1 &= \{s_4, s_5\}, & A_1 - A_3 &= \{s_1, s_2\}, \\ A_3 - A_1 &= \{s_4, s_5, s_8\}, & A_2 - A_3 &= \{s_2\}, & A_3 - A_2 &= \{s_8\}; \\ (A_1^c)^c &= \{s_1, s_2, s_3\} (=A_1), & (A_2^c)^c &= \{s_2, s_3, s_4, s_5\} (=A_2), \\ (A_3^c)^c &= \{s_3, s_4, s_5, s_8\} (=A_3). \end{aligned}$$

An identity and DeMorgan's laws stated subsequently are of significant importance. Their justifications are left as exercises (see Exercises 2.14 and 2.15).

$$\text{An identity} \quad \cup_j A_j = A_1 \cup (A_1^c \cap A_2) \cup (A_1^c \cap A_2^c \cap A_3) \cup \dots$$

EXAMPLE 42

From Example 41, we have:

$$A_1 = \{s_1, s_2, s_3\}, \quad A_1^c \cap A_2 = \{s_4, s_5\}, \quad A_1^c \cap A_2^c \cap A_3 = \{s_8\},$$

Note that $A_1, A_1^c \cap A_2, A_1^c \cap A_2^c \cap A_3$ are pairwise disjoint. Now $A_1 \cup (A_1^c \cap A_2) \cup (A_1^c \cap A_2^c \cap A_3) = \{s_1, s_2, s_3, s_4, s_5, s_8\}$, which is equal to $A_1 \cup A_2 \cup A_3$; that is,

$$A_1 \cup A_2 \cup A_3 = A_1 \cup (A_1^c \cap A_2) \cup (A_1^c \cap A_2^c \cap A_3)$$

as the preceding identity states.

The significance of the identity is that the events on the right-hand side are pairwise disjoint, whereas the original events $A_j, j \geq 1$, need not be so.

$$\text{DeMorgan's laws} \quad (\cup_j A_j)^c = \cap_j A_j^c, \quad (\cap_j A_j)^c = \cup_j A_j^c.$$

EXAMPLE 43

Again from Example 41, one has:

$$\begin{aligned} (A_1 \cup A_2)^c &= \{s_6, s_7, s_8\}, & A_1^c \cap A_2^c &= \{s_6, s_7, s_8\}; \\ (A_1 \cup A_2 \cup A_3)^c &= \{s_6, s_7\}, & A_1^c \cap A_2^c \cap A_3^c &= \{s_6, s_7\}; \\ (A_1 \cap A_2)^c &= \{s_1, s_4, s_5, s_6, s_7, s_8\}, & A_1^c \cup A_2^c &= \{s_1, s_4, s_5, s_6, s_7, s_8\}; \\ (A_1 \cap A_2 \cap A_3)^c &= \{s_1, s_2, s_4, s_5, s_6, s_7, s_8\}, \\ A_1^c \cup A_2^c \cup A_3^c &= \{s_1, s_2, s_4, s_5, s_6, s_7, s_8\}, \end{aligned}$$

so that

$$\begin{aligned} (A_1 \cup A_2)^c &= A_1^c \cap A_2^c, & (A_1 \cup A_2 \cup A_3)^c &= A_1^c \cap A_2^c \cap A_3^c, & \text{as DeMorgan's} \\ (A_1 \cap A_2)^c &= A_1^c \cup A_2^c, & (A_1 \cap A_2 \cap A_3)^c &= A_1^c \cup A_2^c \cup A_3^c, & \text{laws state.} \end{aligned}$$

As a further demonstration of how complements, unions, and intersections of sets are used for the expression of new sets, consider the following example.

EXAMPLE 44

In terms of the events A_1, A_2 , and A_3 (in some sample space S) and, perhaps, their complements, unions, and intersections, express the following events:

$$D_i = \text{"}A_i \text{ does not occur,"} \quad i = 1, 2, 3, \text{ so that } D_1 = A_1^c, D_2 = A_2^c, D_3 = A_3^c;$$

$$E = \text{"all } A_1, A_2, A_3 \text{ occur,"} \text{ so that } E = A_1 \cap A_2 \cap A_3;$$

$$F = \text{"none of } A_1, A_2, A_3 \text{ occurs,"} \text{ so that } F = A_1^c \cap A_2^c \cap A_3^c;$$

$$G = \text{"at least one of } A_1, A_2, A_3 \text{ occurs,"} \text{ so that } G = A_1 \cup A_2 \cup A_3;$$

$$H = \text{"exactly two of } A_1, A_2, A_3 \text{ occur,"} \text{ so that } H = (A_1 \cap A_2 \cap A_3^c) \cup (A_1 \cap A_2^c \cap A_3) \cup (A_1^c \cap A_2 \cap A_3);$$

$$I = \text{"exactly one of } A_1, A_2, A_3 \text{ occurs,"} \text{ so that } I = (A_1 \cap A_2^c \cap A_3^c) \cup (A_1^c \cap A_2 \cap A_3^c) \cup (A_1^c \cap A_2^c \cap A_3).$$

It also follows that:

$$\begin{aligned} G &= \text{"exactly one of } A_1, A_2, A_3 \text{ occurs"} \cup \text{"exactly two of } A_1, A_2, A_3 \text{ occur"} \cup \\ &\quad \text{"all } A_1, A_2, A_3 \text{ occur"} \\ &= I \cup H \cup E. \end{aligned}$$

This section is concluded with the concept of a monotone sequence of events. Namely, the sequence of events $\{A_n\}, n \geq 1$, is said to be *monotone*, if either

$A_1 \subseteq A_2 \subseteq \dots$ (*increasing*) or $A_1 \supseteq A_2 \supseteq \dots$ (*decreasing*). In case of an increasing sequence, the union $\bigcup_{j=1}^{\infty} A_j$ is called the *limit* of the sequence, and in case of a decreasing sequence, the intersection $\bigcap_{j=1}^{\infty} A_j$ is called its *limit*.

The concept of the limit is also defined, under certain conditions, for non-monotone sequences of events, but we are not going to enter into it here. The interested reader is referred to Definition 1, page 5, of the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G. G. Roussas.

Exercises

2.1 An airport limousine departs from a certain airport with three passengers to be delivered in any one of three hotels denoted by H_1, H_2, H_3 . Let (x_1, x_2, x_3) denote the number of passengers left at hotels H_1, H_2 , and H_3 , respectively.

- (i) Write out the sample space S of all possible deliveries.
- (ii) Consider the events A, B, C , and D , defined as follows, and express them in terms of sample points.

A = "one passenger in each hotel,"

B = "all passengers in H_1 ,"

C = "all passengers in one hotel,"

D = "at least two passengers in H_1 ,"

E = "fewer passengers in H_1 than in any one of H_2 or H_3 ."

2.2 A machine dispenses balls which are either red or black or green. Suppose we operate the machine three successive times and record the color of the balls dispensed, to be denoted by r, b, and g for the respective colors.

- (i) Write out an appropriate sample space S for this experiment.
- (ii) Consider the events A, B , and C , defined as follows, and express them by means of sample points.

A = "all three colors appear,"

B = "only two colors appear,"

C = "at least two colors appear."

2.3 A university library has five copies of a textbook to be used in a certain class. Of these copies, numbers 1 through 3 are of the 1st edition, and numbers 4 and 5 are of the 2nd edition. Two of these copies are chosen at random to be placed on a 2-hour reserve.

- (i) Write out an appropriate sample space S .
- (ii) Consider the events A, B, C , and D , defined as follows, and express them in terms of sample points.

A = "both books are of the 1st edition,"

B = "both books are of the 2nd edition,"

C = "one book of each edition,"

D = "no book is of the 2nd edition."

2.4 A large automobile dealership sells three brands of American cars, denoted by a_1, a_2, a_3 ; two brands of Asian cars, denoted by b_1, b_2 ; and one brand of a European car, denoted by c . We observe the cars sold in two consecutive sales. Then:

- (i) Write out an appropriate sample space for this experiment.
 (ii) Express the events defined as follows in terms of sample points:

A = "American brands in both sales,"

B = "American brand in the first sale and Asian brand in the second sale,"

C = "American brand in one sale and Asian brand in the other sale,"

D = "European brand in one sale and Asian brand in the other sale."

2.5 Of two gas stations I and II located at a certain intersection, I has five gas pumps and II has six gas pumps. On a given time of a day, observe the numbers x and y of pumps in use in stations I and II, respectively.

- (i) Write out the sample space \mathcal{S} for this experiment.
 (ii) Consider the events A, B, C , and D , defined as follows, and express them in terms of sample points.

A = "three pumps are in use in station I,"

B = "the number of pumps in use in both stations is the same,"

C = "the number of pumps in use in station II is larger than that in station I,"

D = "the total number of pumps in use in both stations is not greater than 4."

2.6 At a certain busy airport, denote by A, B, C , and D the events defined as follows:

A = "at least 5 planes are waiting to land,"

B = "at most 3 planes are waiting to land,"

C = "at most 2 planes are waiting to land,"

D = "exactly 2 planes are waiting to land."

In terms of the events A, B, C , and D and, perhaps, their complements, express the following events:

E = "at most 4 planes are waiting to land,"

F = "at most 1 plane is waiting to land,"

G = "exactly 3 planes are waiting to land,"

H = "exactly 4 planes are waiting to land,"

I = "at least 4 planes are waiting to land."

2.7 Let $\mathcal{S} = \{(x, y) \in \mathbb{N}^2; -3 \leq x \leq 3, 0 \leq y \leq 4, x \text{ and } y \text{ integers}\}$, and define the events A, B, C , and D as follows:

$$A = \{(x, y) \in \mathcal{S}; x = y\}, \quad B = \{(x, y) \in \mathcal{S}; x = -y\},$$

$$C = \{(x, y) \in \mathcal{S}; x^2 = y^2\}, \quad D = \{(x, y) \in \mathcal{S}; x^2 + y^2 \leq 5\}.$$

List the members of the events just defined.

2.8 In terms of the events A_1, A_2, A_3 in a sample space \mathcal{S} and, perhaps, their complements, express the following events:

- (i) $B_0 = \{s \in \mathcal{S}; s \text{ belongs to none of } A_1, A_2, A_3\}$,
- (ii) $B_1 = \{s \in \mathcal{S}; s \text{ belongs to exactly one of } A_1, A_2, A_3\}$,
- (iii) $B_2 = \{s \in \mathcal{S}; s \text{ belongs to exactly two of } A_1, A_2, A_3\}$,
- (iv) $B_3 = \{s \in \mathcal{S}; s \text{ belongs to all of } A_1, A_2, A_3\}$,
- (v) $C = \{s \in \mathcal{S}; s \text{ belongs to at most two of } A_1, A_2, A_3\}$,
- (vi) $D = \{s \in \mathcal{S}; s \text{ belongs to at least one of } A_1, A_2, A_3\}$.

2.9 If for three events A, B , and C it happens that either $A \cup B \cup C = A$ or $A \cap B \cap C = A$, what conclusions can you draw?

2.10 Show that A is the impossible event (that is, $A = \emptyset$), if and only if $(A \cap B^c) \cup (A^c \cap B) = B$ for every event B .

2.11 Let A, B , and C be arbitrary events in \mathcal{S} . Determine whether each of the following statements is correct or incorrect.

- (i) $(A - B) \cup B = (A \cap B^c) \cup B = B$,
- (ii) $(A \cup B) - A = (A \cup B) \cap A^c = B$,
- (iii) $(A \cap B) \cap (A - B) = (A \cap B) \cap (A \cap B^c) = \emptyset$,
- (iv) $(A \cup B) \cap (B \cup C) \cap (C \cup A) = (A \cap B) \cup (B \cap C) \cup (C \cap A)$.

2.12 For any three events A, B , and C in a sample space \mathcal{S} show that the transitive property, $A \subseteq B$ and $B \subseteq C$, implies that $A \subseteq C$ holds.

2.13 Establish the distributive laws, namely $A \cap (\cup_j A_j) = \cup_j (A \cap A_j)$ and $A \cup (\cap_j A_j) = \cap_j (A \cup A_j)$.

2.14 Establish the identity:

$$\cup_j A_j = A_1 \cup (A_1^c \cap A_2) \cup (A_1^c \cap A_2^c \cap A_3) \cup \dots$$

2.15 Establish DeMorgan's laws, namely

$$(\cup_j A_j)^c = \cap_j A_j^c \quad \text{and} \quad (\cap_j A_j)^c = \cup_j A_j^c.$$

2.16 Let $\mathcal{S} = \Re$ and, for $n = 1, 2, \dots$, define the events A_n and B_n by:

$$A_n = \left\{ x \in \Re; -5 + \frac{1}{n} < x < 20 - \frac{1}{n} \right\}, \quad B_n = \left\{ x \in \Re; 0 < x < 7 + \frac{3}{n} \right\}.$$

- (i) Show that the sequence $\{A_n\}$ is increasing and the sequence $\{B_n\}$ is decreasing.
- (ii) Identify the limits, $\lim_{n \rightarrow \infty} A_n = \bigcup_{n=1}^{\infty} A_n$ and $\lim_{n \rightarrow \infty} B_n = \bigcap_{n=1}^{\infty} B_n$.

1.3 Random Variables

For every random experiment, there is at least one sample space appropriate for the random experiment under consideration. In many cases, however, much of the work can be done without reference to an explicit sample space. Instead, what are used extensively are random variables and their distributions.

Those quantities will be studied extensively in subsequent chapters. What is done in this section is the introduction of the concept of a random variable.

Formally, a *random variable*, to be shortened to r.v., is simply a function defined on a sample space \mathcal{S} and taking values in the real line $\mathfrak{R} = (-\infty, \infty)$. Random variables are denoted by capital letters, such as X, Y, Z , with or without subscripts. Thus, the value of the r.v. X at the sample point s is $X(s)$, and the set of all values of X , that is, the *range* of X , is usually denoted by $X(\mathcal{S})$. The only difference between a r.v. and a function in the usual calculus sense is that the domain of a r.v. is a sample space \mathcal{S} , which may be an abstract set, unlike the usual concept of a function, whose domain is a subset of \mathfrak{R} or of a Euclidean space of higher dimension. The usage of the term “random variable” employed here rather than that of a function may be explained by the fact that a r.v. is associated with the outcomes of a random experiment. Thus, one may argue that $X(s)$ is not known until the random experiment is actually carried out and s becomes available. Of course, on the same sample space, one may define many distinct r.v.’s.

In reference to Example 26, instead of the sample space \mathcal{S} exhibited there, one may be interested in the number of heads appearing each time the experiment is carried out. This leads to the definition of the r.v. X by: $X(s) = \#$ of H ’s in s . Thus, $X(HHH) = 3$, $X(HHT) = X(HTH) = X(THH) = 2$, $X(HTT) = X(THT) = X(TTH) = 1$, and $X(TTT) = 0$, so that $X(\mathcal{S}) = \{0, 1, 2, 3\}$. The notation $(X \leq 1)$ stands for the event $\{s \in \mathcal{S}; X(s) \leq 1\} = \{TTT, HTT, THT, TTH\}$. In the general case and for $B \subseteq \mathfrak{R}$, the notation $(X \in B)$ stands for the event A in the sample space \mathcal{S} defined by: $A = \{s \in \mathcal{S}; X(s) \in B\}$. It is also denoted by $X^{-1}(B)$.

In reference to Example 27, a r.v. X of interest may be defined by $X(s) =$ sum of the numbers in the pair s . Thus, $X((1, 1)) = 2$, $X((1, 2)) = X((2, 1)) = 3$, \dots , $X((6, 6)) = 12$, and $X(\mathcal{S}) = \{2, 3, \dots, 12\}$. Also, $X^{-1}(\{7\}) = \{s \in \mathcal{S}; X(s) = 7\} = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$. Similarly for Examples 28–31.

In reference to Example 32, a natural r.v. X is defined to denote the number of tosses needed until the first head occurs. Thus, $X(H) = 1$, $X(TH) = 2$, \dots , $X(\underbrace{T \dots T}_{n-1}H) = n$, \dots , so that $X(\mathcal{S}) = \{1, 2, \dots\}$. Also, $(X > 5) = (X \geq 6) = \{\underbrace{TTTTT}_{n-1}H, TTTTTTH, \dots\}$.

In reference to Example 33, an obvious r.v. X is: $X(s) = s$, $s = 0, 1, \dots$, and similarly for Examples 34–35.

In reference to Example 36, a r.v. X of interest is $X(s) = s$, $s \in \mathcal{S}$, and similarly for Examples 37–40.

Also, in reference to Example 5, an obvious r.v. X may be defined as follows: $X(s) = \#$ of S ’s in s . Then, clearly, $X(\mathcal{S}) = \{0, 1, \dots, n\}$. Similarly for Examples 6–10.

In reference to Example 11, a r.v. X may be defined thus: $X(s) =$ the position of M in s . Then, clearly, $X(\mathcal{S}) = \{1, 2, \dots\}$.

In reference to Example 18, the r.v.’s of obvious interests are: $X_A = \#$ of those persons, out of n , having blood type A , and similarly for X_B, X_{AB}, X_O . Similarly for Examples 19 and 20.

From the preceding examples, two kinds of r.v.'s emerge: random variables which take on countably many values, such as those defined in conjunction with Examples 26–31 and 32–35, and r.v.'s which take on all values in a nondegenerate (finite or not) interval in \mathfrak{R} . Such are r.v.'s defined in conjunction with Examples 36–40. Random variables of the former kind are called discrete r.v.'s (or r.v.'s of the discrete type), and r.v.'s of the latter type are called continuous r.v.'s (or r.v.'s of the continuous type).

More generally, a r.v. X is called *discrete* (or *of the discrete type*), if X takes on countably many values; i.e., either finitely many values such as x_1, \dots, x_n , or countably infinite many values such as x_0, x_1, \dots or x_1, x_2, \dots . On the other hand, X is called *continuous* (or *of the continuous type*) if X takes all values in a proper interval $I \subseteq \mathfrak{R}$. Although there are other kinds of r.v.'s, in this book we will restrict ourselves to discrete and continuous r.v.'s as just defined.

The study of r.v.'s is one of the main objectives of this book.

Exercises

- 3.1** In reference to Exercise 2.1, define the r.v.'s $X_i, i = 1, 2, 3$ as follows:
 $X_i = \#$ of passengers delivered to hotel H_i .
 Determine the values of each $X_i, i = 1, 2, 3$, and specify the values of the sum $X_1 + X_2 + X_3$.
- 3.2** In reference to Exercise 2.2, define the r.v.'s X and Y as follows: $X = \#$ of red balls dispensed, $Y = \#$ of balls other than red dispensed.
 Determine the values of X and Y , and specify the values of the sum $X + Y$.
- 3.3** In reference to Exercise 2.5, define the r.v.'s X and Y as follows: $X = \#$ of pumps in use in station I, $Y = \#$ of pumps in use in station II.
 Determine the values of X and Y , and also of the sum $X + Y$.
- 3.4** In reference to Exercise 2.7, define the r.v. X by: $X((x, y)) = x + y$.
 Determine the values of X , as well as the following events: $(X \leq 2)$, $(3 < X \leq 5)$, $(X > 6)$.
- 3.5** Consider a year with 365 days, which are numbered serially from 1 to 365. Ten of those numbers are chosen at random and without replacement, and let X be the r.v. denoting the largest number drawn.
 Determine the values of X .
- 3.6** A four-sided die has the numbers 1 through 4 written on its sides, one on each side. If the die is rolled twice:
 (i) Write out a suitable sample space \mathcal{S} .
 (ii) If X is the r.v. denoting the sum of numbers appearing, determine the values of X .
 (iii) Determine the events: $(X \leq 3)$, $(2 \leq X < 5)$, $(X > 8)$.
- 3.7** From a certain target population, n individuals are chosen at random and their blood types are determined. Let X_1, X_2, X_3 , and X_4 be the r.v.'s

denoting the number of individuals having blood types A, B, AB, and O, respectively.

Determine the values of each one of these r.v.'s, as well as the values of the sum $X_1 + X_2 + X_3 + X_4$.

3.8 A bus is expected to arrive at a specified bus stop any time between 8:00 and 8:15 a.m., and let X be the r.v. denoting the actual time of arrival of the bus.

- (i) Determine the suitable sample space \mathcal{S} for the experiment of observing the arrival of the bus.
- (ii) What are the values of the r.v. X ?
- (iii) Determine the event: "The bus arrives within 5 minutes before the expiration of the expected time of arrival."



Chapter 2

The Concept of Probability and Basic Results

This chapter consists of five sections. The first section is devoted to the definition of the concept of probability. We start with the simplest case, where complete symmetry occurs, proceed with the definition by means of relative frequency, and conclude with the axiomatic definition of probability. The defining properties of probability are illustrated by way of examples. Also, a number of basic properties, resulting from the definition, are stated and justified. Some of them are illustrated by means of examples. The section is concluded with two theorems, which are stated but not proved.

In the second section, the distribution of a r.v. is introduced. Also, the distribution function and the probability density function of a r.v. are defined, and we explain how they determine the distribution of the r.v.

The concept of the conditional probability of an event, given another event, is taken up in the following section. Its definition is given, and its significance is demonstrated through a number of examples. This section is concluded with three theorems, formulated in terms of conditional probabilities. Through these theorems, conditional probabilities greatly simplify calculation of otherwise complicated probabilities.

In the fourth section, the independence of two events is defined, and we also indicate how it carries over to any finite number of events. A result (Theorem 6) is stated which is often used by many authors without its use even being acknowledged. The section is concluded with an indication of how independence extends to random experiments. The definition of independence of r.v.'s is deferred to another chapter (Chapter 5).

In the final section of the chapter, the so-called fundamental principle of counting is discussed; combinations and permutations are then obtained as applications of this principle. Several illustrative examples are also provided.

2.1 Definition of Probability and Some Basic Results

When a random experiment is entertained, one of the first questions which arise is, what is the probability that a certain event occurs? For instance, in reference to Example 26 in Chapter 1, one may ask: What is the probability that exactly one head occurs; in other words, what is the probability of the event $B = \{HTT, THT, TTH\}$? The answer to this question is almost automatic and is $3/8$. The relevant reasoning goes like this: Assuming that the three coins are balanced, the probability of each one of the 8 outcomes, considered as simple events, must be $1/8$. Since the event B consists of 3 sample points, it can occur in 3 different ways, and hence its probability must be $3/8$.

This is exactly the intuitive reasoning employed in defining the concept of probability when two requirements are met: First, the sample space S has finitely many outcomes, $S = \{s_1, \dots, s_n\}$, say, and second, each one of these outcomes is “equally likely” to occur, has the same chance of appearing, whenever the relevant random experiment is carried out. This reasoning is based on the underlying symmetry. Thus, one is led to stipulating that each one of the (simple) events $\{s_i\}$, $i = 1, \dots, n$ has probability $1/n$. Then the next step, that of defining the probability of a composite event A , is simple; if A consists of m sample points, $A = \{s_{i_1}, \dots, s_{i_m}\}$, say ($1 \leq m \leq n$) (or none at all, in which case $m = 0$), then the probability of A must be m/n . The notation used is: $P(\{s_1\}) = \dots = P(\{s_n\}) = \frac{1}{n}$ and $P(A) = \frac{m}{n}$. Actually, this is the so-called *classical* definition of probability. That is,

CLASSICAL DEFINITION OF PROBABILITY Let S be a sample space, associated with a certain random experiment and consisting of finitely many sample points n , say, each of which is equally likely to occur whenever the random experiment is carried out. Then the probability of any event A , consisting of m sample points ($0 \leq m \leq n$), is given by $P(A) = \frac{m}{n}$.

In reference to Example 26 in Chapter 1, $P(A) = \frac{4}{8} = \frac{1}{2} = 0.5$. In Example 27 (when the two dice are unbiased), $P(X = 7) = \frac{6}{36} = \frac{1}{6} \simeq 0.167$, where the r.v. X and the event $(X = 7)$ are defined in Section 1.3. In Example 29, when the balls in the urn are thoroughly mixed, we may assume that all of the $(m+n)(m+n-1)$ pairs are equally likely to be selected. Then, since the event A occurs in 20 different ways, $P(A) = \frac{20}{(m+n)(m+n-1)}$. For $m = 3$ and $n = 5$, this probability is $P(A) = \frac{20}{56} = \frac{5}{14} \simeq 0.357$.

From the preceding (classical) definition of probability, the following simple properties are immediate: For any event A , $P(A) \geq 0$; $P(S) = 1$; if two events A_1 and A_2 are disjoint ($A_1 \cap A_2 = \emptyset$), then $P(A_1 \cup A_2) = P(A_1) + P(A_2)$. This is so because, if $A_1 = \{s_{i_1}, \dots, s_{i_k}\}$, $A_2 = \{s_{j_1}, \dots, s_{j_\ell}\}$, where all s_{i_1}, \dots, s_{i_k} are distinct from all $s_{j_1}, \dots, s_{j_\ell}$, then $A_1 \cup A_2 = \{s_{i_1}, \dots, s_{i_k}, s_{j_1}, \dots, s_{j_\ell}\}$ and $P(A_1 \cup A_2) = \frac{k+\ell}{n} = \frac{k}{n} + \frac{\ell}{n} = P(A_1) + P(A_2)$.

In many cases, the stipulations made in defining the probability as above are not met, either because S has not finitely many points (as is the case in Examples 32, 33–35 (by replacing C and M by ∞), and 36–40 in Chapter 1), or

because the (finitely many outcomes) are not equally likely. This happens, for instance, in Example 26 when the coins are not balanced and in Example 27 when the dice are biased. Strictly speaking, it also happens in Example 30. In situations like this, the way out is provided by the so-called *relative frequency* definition of probability. Specifically, suppose a random experiment is carried out a large number of times N , and let $N(A)$ be the *frequency* of an event A , the number of times A occurs (out of N). Then the *relative frequency* of A is $\frac{N(A)}{N}$. Next, suppose that, as $N \rightarrow \infty$, the relative frequencies $\frac{N(A)}{N}$ oscillate around some number (necessarily between 0 and 1). More precisely, suppose that $\frac{N(A)}{N}$ converges, as $N \rightarrow \infty$, to some number. Then this number is called the *probability* of A and is denoted by $P(A)$. That is, $P(A) = \lim_{N \rightarrow \infty} \frac{N(A)}{N}$. (It will be seen later in this book that the assumption of convergence of the relative frequencies $N(A)/N$ is justified subject to some qualifications.) To summarize,

RELATIVE FREQUENCY DEFINITION OF PROBABILITY Let $N(A)$ be the number of times an event A occurs in N repetitions of a random experiment, and assume that the relative frequency of A , $\frac{N(A)}{N}$, converges to a limit as $N \rightarrow \infty$. This limit is denoted by $P(A)$ and is called the probability of A .

At this point, it is to be observed that empirical data show that the relative frequency definition of probability and the classical definition of probability agree in the framework in which the classical definition applies.

From the relative frequency definition of probability and the usual properties of limits, it is immediate that: $P(A) \geq 0$ for every event A ; $P(S) = 1$; and for A_1, A_2 with $A_1 \cap A_2 = \emptyset$,

$$\begin{aligned} P(A_1 \cup A_2) &= \lim_{N \rightarrow \infty} \frac{N(A_1 \cup A_2)}{N} = \lim_{N \rightarrow \infty} \left(\frac{N(A_1)}{N} + \frac{N(A_2)}{N} \right) \\ &= \lim_{N \rightarrow \infty} \frac{N(A_1)}{N} + \lim_{N \rightarrow \infty} \frac{N(A_2)}{N} = P(A_1) + P(A_2); \end{aligned}$$

that is, $P(A_1 \cup A_2) = P(A_1) + P(A_2)$, provided $A_1 \cap A_2 = \emptyset$. These three properties were also seen to be true in the classical definition of probability. Furthermore, it is immediate that under either definition of probability, $P(A_1 \cup \dots \cup A_k) = P(A_1) + \dots + P(A_k)$, provided the events are pairwise disjoint; $A_i \cap A_j = \emptyset$, $i \neq j$.

The above two definitions of probability certainly give substance to the concept of probability in a way consonant with our intuition about what probability should be. However, for the purpose of cultivating the concept and deriving deep probabilistic results, one must define the concept of probability in terms of some basic properties, which would not contradict what we have seen so far. This line of thought leads to the so-called axiomatic definition of probability due to Kolmogorov.

AXIOMATIC DEFINITION OF PROBABILITY Probability is a function, denoted by P , defined for each event of a sample space S , taking on values in the real

line \mathfrak{N} , and satisfying the following three properties:

(P1) $P(A) \geq 0$ for every event A (nonnegativity of P).

(P2) $P(S) = 1$ (P is normed).

(P3) For countably infinite many pairwise disjoint events $A_i, i = 1, 2, \dots, A_i \cap A_j = \emptyset, i \neq j$, it holds

$$P(A_1 \cup A_2 \cup \dots) = P(A_1) + P(A_2) + \dots; \quad \text{or} \quad P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

(sigma-additivity (σ -additivity) of P).

COMMENTS ON THE AXIOMATIC DEFINITION

1) Properties (P1) and (P2) are the same as the ones we have seen earlier, whereas property (P3) is new. What we have seen above was its so-called *finitely-additive* version; that is, $P(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i)$, provided $A_i \cap A_j = \emptyset, i \neq j$. It will be seen below that finite-additivity is implied by σ -additivity but not the other way around. Thus, if we are to talk about the probability of the union of countably infinite many pairwise disjoint events, property (P3) must be stipulated. Furthermore, the need for such a union of events is illustrated as follows: In reference to Example 32, calculate the probability that the first head does not occur before the n th tossing. By setting $A_i = \{\underbrace{T \dots T}_i H\}, i = n, n+1, \dots$, what we are actually after here

is $P(A_n \cup A_{n+1} \cup \dots)$ with $A_i \cap A_j = \emptyset, i \neq j, i$ and $j \geq n$.

2) Property (P3) is superfluous (reduced to finite-additivity) when the sample space S is finite, which implies that the total number of events is finite.

3) Finite-additivity is implied by additivity for two events, $P(A_1 \cup A_2) = P(A_1) + P(A_2), A_1 \cap A_2 = \emptyset$, by way of induction.

Here are two examples in calculating probabilities.

EXAMPLE 1

In reference to Example 1 in Chapter 1, take $n = 58$, and suppose we have the following configuration:

Arsenic	BARIUM			
	HIGH		LOW	
	High	Low	High	Low
High	1	3	5	9
Low	4	8	10	18

Calculate the probabilities mentioned in (i) (a)–(d).

DISCUSSION For simplicity, denote by B_h the event that the site selected has a high barium concentration, and likewise for other events figuring below. Then:

- (i)(a) $B_h = (A_h \cap B_h \cap M_h) \cup (A_h \cap B_h \cap M_\ell) \cup (A_\ell \cap B_h \cap M_h) \cup (A_\ell \cap B_h \cap M_\ell)$ and the events on the right-hand side are pairwise disjoint. Therefore (by the following basic property 2 in Subsection 2.1.1):

$$\begin{aligned} P(B_h) &= P(A_h \cap B_h \cap M_h) + P(A_h \cap B_h \cap M_\ell) \\ &\quad + P(A_\ell \cap B_h \cap M_h) + P(A_\ell \cap B_h \cap M_\ell) \\ &= \frac{1}{58} + \frac{3}{58} + \frac{4}{58} + \frac{8}{58} = \frac{16}{58} = \frac{8}{29} \simeq 0.276. \end{aligned}$$

- (i)(b) Here $P(M_h \cap A_\ell \cap B_\ell) = P(A_\ell \cap B_\ell \cap M_h) = \frac{10}{58} = \frac{5}{29} \simeq 0.172$.
 (i)(c) Here the required probability is as in (a):

$$P(A_h \cap B_h \cap M_\ell) + P(A_h \cap B_\ell \cap M_h) + P(A_\ell \cap B_h \cap M_h) = \frac{12}{58} = \frac{6}{29} \simeq 0.207.$$

- (i)(d) As above,

$$P(A_h \cap B_\ell \cap M_\ell) + P(A_\ell \cap B_h \cap M_\ell) + P(A_\ell \cap B_\ell \cap M_h) = \frac{27}{58} \simeq 0.466.$$

EXAMPLE 2

In ranking five horses in a horse race (Example 31 in Chapter 1), calculate the probability that horse #3 terminates at least second.

DISCUSSION Let A_i be the event that horse #3 terminates in the i th position, $i = 1, \dots, 5$. Then the required event is $A_1 \cup A_2$, where A_1, A_2 are disjoint. Thus,

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) = \frac{24}{120} + \frac{24}{120} = \frac{2}{5} = 0.4.$$

EXAMPLE 3

In tossing a coin repeatedly until H appears for the first time (Example 32 in Chapter 1), suppose that $P\{\underbrace{T \dots T}_i H\} = P(A_i) = q^{i-1}p$ for some $0 < p < 1$ and $q = 1 - p$ (in anticipation of Definition 3 in Section 2.4). Then

$$P\left(\bigcup_{i=n}^{\infty} A_i\right) = \sum_{i=n}^{\infty} P(A_i) = \sum_{i=n}^{\infty} q^{i-1}p = p \sum_{i=n}^{\infty} q^{i-1} = p \frac{q^{n-1}}{1-q} = p \frac{q^{n-1}}{p} = q^{n-1}.$$

For instance, for $p = 1/2$ and $n = 3$, this probability is $\frac{1}{4} = 0.25$. That is, when tossing a fair coin, the probability that the first head does not appear either the first or the second time (and therefore it appears either the third time or the fourth time etc.) is 0.25. For $n = 10$, this probability is approximately $0.00195 \simeq 0.002$.

Next, we present some basic results following immediately from the defining properties of the probability. First, we proceed with their listing and then with their justification.

2.1.1 Some Basic Properties of a Probability Function

1. $P(\emptyset) = 0$.
2. For any pairwise disjoint events A_1, \dots, A_n , $P(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i)$.
3. For any event A , $P(A^c) = 1 - P(A)$.
4. $A_1 \subseteq A_2$ implies $P(A_1) \leq P(A_2)$ and $P(A_2 - A_1) = P(A_2) - P(A_1)$.
5. $0 \leq P(A) \leq 1$ for every event A .
6. (i) For any two events A_1 and A_2 :

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2).$$

- (ii) For any three events A_1, A_2 , and A_3 :

$$P(A_1 \cup A_2 \cup A_3) = P(A_1) + P(A_2) + P(A_3) - [P(A_1 \cap A_2) + P(A_1 \cap A_3) + P(A_2 \cap A_3)] + P(A_1 \cap A_2 \cap A_3).$$

7. For any events A_1, A_2, \dots , $P(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} P(A_i)$ (σ -sub-additivity), and $P(\bigcup_{i=1}^n A_i) \leq \sum_{i=1}^n P(A_i)$ (finite-sub-additivity).

2.1.2 Justification

1. From the obvious fact that $S = S \cup \emptyset \cup \emptyset \cup \dots$ and property (P3),

$$P(S) = P(S \cup \emptyset \cup \emptyset \cup \dots) = P(S) + P(\emptyset) + P(\emptyset) + \dots$$

or $P(\emptyset) + P(\emptyset) + \dots = 0$. By (P1), this can only happen when $P(\emptyset) = 0$. Of course, that the impossible event has probability 0 does not come as a surprise. Any reasonable definition of probability should imply it.

2. Take $A_i = \emptyset$ for $i \geq n+1$, consider the following obvious relation, and use (P3) and #1 to obtain:

$$P\left(\bigcup_{i=1}^n A_i\right) = P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i) = \sum_{i=1}^n P(A_i).$$

3. From (P2) and #2, $P(A \cup A^c) = P(S) = 1$ or $P(A) + P(A^c) = 1$, so that $P(A^c) = 1 - P(A)$.
4. The relation $A_1 \subseteq A_2$, clearly, implies $A_2 = A_1 \cup (A_2 - A_1)$, so that, by #2, $P(A_2) = P(A_1) + P(A_2 - A_1)$. Solving for $P(A_2 - A_1)$, we obtain $P(A_2 - A_1) = P(A_2) - P(A_1)$, so that, by (P1), $P(A_1) \leq P(A_2)$. At this point it must be pointed out that $P(A_2 - A_1)$ need not be $P(A_2) - P(A_1)$, if A_1 is not contained in A_2 .
5. Clearly, $\emptyset \subseteq A \subseteq S$ for any event A . Then (P1), #1 and #4 give: $0 = P(\emptyset) \leq P(A) \leq P(S) = 1$.
6. (i) It is clear (by means of a Venn diagram, for example) that

$$A_1 \cup A_2 = A_1 \cup (A_2 \cap A_1^c) = A_1 \cup (A_2 - A_1 \cap A_2).$$

Then, by means of #2 and #4:

$$P(A_1 \cup A_2) = P(A_1) + P(A_2 - A_1 \cap A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2).$$

(ii) Apply part (i) to obtain:

$$\begin{aligned}
 P(A_1 \cup A_2 \cup A_3) &= P[(A_1 \cup A_2) \cup A_3] = P(A_1 \cup A_2) + P(A_3) \\
 &\quad - P[(A_1 \cup A_2) \cap A_3] \\
 &= P(A_1) + P(A_2) - P(A_1 \cap A_2) + P(A_3) \\
 &\quad - P[(A_1 \cap A_3) \cup (A_2 \cap A_3)] \\
 &= P(A_1) + P(A_2) + P(A_3) - P(A_1 \cap A_2) \\
 &\quad - [P(A_1 \cap A_3) + P(A_2 \cap A_3) - P(A_1 \cap A_2 \cap A_3)] \\
 &= P(A_1) + P(A_2) + P(A_3) - P(A_1 \cap A_2) - P(A_1 \cap A_3) \\
 &\quad - P(A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_3).
 \end{aligned}$$

7. By the identity in Section 2 of Chapter 1 and (P3):

$$\begin{aligned}
 P\left(\bigcup_{i=1}^{\infty} A_i\right) &= P[A_1 \cup (A_1^c \cap A_2) \cup \dots \cup (A_1^c \cap \dots \cap A_{n-1}^c \cap A_n) \cup \dots] \\
 &= P(A_1) + P(A_1^c \cap A_2) + \dots + P(A_1^c \cap \dots \cap A_{n-1}^c \cap A_n) + \dots \\
 &\leq P(A_1) + P(A_2) + \dots + P(A_n) + \dots \text{ (by \#4)}.
 \end{aligned}$$

For the finite case:

$$\begin{aligned}
 P\left(\bigcup_{i=1}^n A_i\right) &= P[A_1 \cup (A_1^c \cap A_2) \cup \dots \cup (A_1^c \cap \dots \cap A_{n-1}^c \cap A_n)] \\
 &= P(A_1) + P(A_1^c \cap A_2) + \dots + P(A_1^c \cap \dots \cap A_{n-1}^c \cap A_n) \\
 &\leq P(A_1) + P(A_2) + \dots + P(A_n).
 \end{aligned}$$

Next, some examples are presented to illustrate some of the properties #1–#7.

EXAMPLE 4

- (i) For two events A and B , suppose that $P(A) = 0.3$, $P(B) = 0.5$, and $P(A \cup B) = 0.6$. Calculate $P(A \cap B)$.
- (ii) If $P(A) = 0.6$, $P(B) = 0.3$, $P(A \cap B^c) = 0.4$, and $B \subset C$, calculate $P(A \cup B^c \cup C^c)$.

DISCUSSION

- (i) From $P(A \cup B) = P(A) + P(B) - P(A \cap B)$, we get $P(A \cap B) = P(A) + P(B) - P(A \cup B) = 0.3 + 0.5 - 0.6 = 0.2$.
- (ii) The relation $B \subset C$ implies $C^c \subset B^c$ and hence $A \cup B^c \cup C^c = A \cup B^c$. Then $P(A \cup B^c \cup C^c) = P(A \cup B^c) = P(A) + P(B^c) - P(A \cap B^c) = 0.6 + (1 - 0.3) - 0.4 = 0.9$.

EXAMPLE 5

Let A and B be the respective events that two contracts I and II, say, are completed by certain deadlines, and suppose that: $P(\text{at least one contract is completed by its deadline}) = 0.9$ and $P(\text{both contracts are completed by$

their deadlines) = 0.5. Calculate the probability: $P(\text{exactly one contract is completed by its deadline})$.

DISCUSSION The assumptions made are translated as follows: $P(A \cup B) = 0.9$ and $P(A \cap B) = 0.5$. What we wish to calculate is: $P((A \cap B^c) \cup (A^c \cap B)) = P(A \cap B^c) + P(A^c \cap B)$. Clearly, $A = (A \cap B) \cup (A \cap B^c)$ and $B = (A \cap B) \cup (A^c \cap B)$, so that $P(A) = P(A \cap B) + P(A \cap B^c)$ and $P(B) = P(A \cap B) + P(A^c \cap B)$. Hence, $P(A \cap B^c) = P(A) - P(A \cap B)$ and $P(A^c \cap B) = P(B) - P(A \cap B)$. Then $P(A \cap B^c) + P(A^c \cap B) = P(A) + P(B) - 2P(A \cap B) = [P(A) + P(B) - P(A \cap B)] - P(A \cap B) = P(A \cup B) - P(A \cap B) = 0.9 - 0.5 = 0.4$.

EXAMPLE 6

- (i) For three events A , B , and C , suppose that $P(A \cap B) = P(A \cap C)$ and $P(B \cap C) = 0$. Then show that $P(A \cup B \cup C) = P(A) + P(B) + P(C) - 2P(A \cap B)$.
- (ii) For any two events A and B , show that $P(A^c \cap B^c) = 1 - P(A) - P(B) + P(A \cap B)$.

DISCUSSION

- (i) We have $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$. But $A \cap B \cap C \subset B \cap C$, so that $P(A \cap B \cap C) \leq P(B \cap C) = 0$, and therefore $P(A \cup B \cup C) = P(A) + P(B) + P(C) - 2P(A \cap B)$.
- (ii) Indeed, $P(A^c \cap B^c) = P((A \cup B)^c) = 1 - P(A \cup B) = 1 - P(A) - P(B) + P(A \cap B)$.

EXAMPLE 7

In ranking five horses in a horse race (Example 31 in Chapter 1), what is the probability that horse #3 will terminate either first or second or third?

DISCUSSION Denote by B the required event and let $A_i = \text{"horse \#}i\text{ terminates in the }i\text{th place, }i = 1, 2, 3$. Then the events A_1, A_2, A_3 are pairwise disjoint, and therefore

$$P(B) = P(A_1 \cup A_2 \cup A_3) = P(A_1) + P(A_2) + P(A_3).$$

But $P(A_1) = P(A_2) = P(A_3) = \frac{24}{120} = 0.2$, so that $P(B) = 0.6$.

EXAMPLE 8

Consider a well-shuffled deck of 52 cards (Example 28 in Chapter 1), and suppose we draw at random three cards. What is the probability that at least one is an ace?

DISCUSSION Let A be the required event, and let A_i be defined by: $A_i = \text{"exactly }i\text{ cards are aces, }i = 0, 1, 2, 3$. Then, clearly, $P(A) = P(A_1 \cup A_2 \cup A_3)$. Instead, we may choose to calculate $P(A)$ through $P(A^c) = 1 - P(A_0)$, where

$$P(A_0) = \frac{\binom{48}{3}}{\binom{52}{3}} = \frac{48 \times 47 \times 46}{52 \times 51 \times 50} = \frac{4,324}{5,525}, \quad \text{so that } P(A) = \frac{1,201}{5,525} \simeq 0.217.$$

EXAMPLE 9

Refer to Example 3 in Chapter 1 and let C_1, C_2, C_3 be defined by: C_1 = “both S_1 and S_2 work,” C_2 = “ S_5 works,” C_3 = “both S_3 and S_4 work,” and let C = “current is transferred from point A to point B .” Then $P(C) = P(C_1 \cup C_2 \cup C_3)$. At this point (in anticipation of Definition 3 in Section 2.4; see also Exercise 4.14 in this chapter), suppose that:

$$\begin{aligned} P(C_1) &= p_1 p_2, & P(C_2) &= p_5, & P(C_3) &= p_3 p_4, \\ P(C_1 \cap C_2) &= p_1 p_2 p_5, & P(C_1 \cap C_3) &= p_1 p_2 p_3 p_4, \\ P(C_2 \cap C_3) &= p_3 p_4 p_5, & P(C_1 \cap C_2 \cap C_3) &= p_1 p_2 p_3 p_4 p_5. \end{aligned}$$

Then:

$$P(C) = p_1 p_2 + p_5 + p_3 p_4 - p_1 p_2 p_5 - p_1 p_2 p_3 p_4 - p_3 p_4 p_5 + p_1 p_2 p_3 p_4 p_5.$$

For example, for $p_1 = p_2 = p_3 = 0.9$, we obtain

$$P(C) = 0.9 + 2(0.9)^2 - 2(0.9)^3 - (0.9)^4 + (0.9)^5 \simeq 0.996.$$

This section is concluded with two very useful results stated as theorems. The first is a generalization of property #6 to more than three events, and the second is akin to the concept of continuity of a function as it applies to a probability function.

THEOREM 1

The probability of the union of any n events, A_1, \dots, A_n , is given by:

$$\begin{aligned} P\left(\bigcup_{j=1}^n A_j\right) &= \sum_{j=1}^n P(A_j) - \sum_{1 \leq j_1 < j_2 \leq n} P(A_{j_1} \cap A_{j_2}) \\ &\quad + \sum_{1 \leq j_1 < j_2 < j_3 \leq n} P(A_{j_1} \cap A_{j_2} \cap A_{j_3}) - \dots \\ &\quad + (-1)^{n+1} P(A_1 \cap \dots \cap A_n). \end{aligned}$$

Although its proof (which is by induction) will not be presented, the pattern of the right-hand side above follows that of property #6(i) and it is clear. First, sum up the probabilities of the individual events, then subtract the probabilities of the intersections of the events, taken two at a time (in the ascending order of indices), then add the probabilities of the intersections of the events, taken three at a time as before, and continue like this until you add or subtract (depending on n) the probability of the intersection of all n events.

Recall that, if $A_1 \subseteq A_2 \subseteq \dots$, then $\lim_{n \rightarrow \infty} A_n = \bigcup_{n=1}^{\infty} A_n$, and if $A_1 \supseteq A_2 \supseteq \dots$, then $\lim_{n \rightarrow \infty} A_n = \bigcap_{n=1}^{\infty} A_n$.

THEOREM 2

For any monotone sequence of events $\{A_n\}$, $n \geq 1$, it holds $P(\lim_{n \rightarrow \infty} A_n) = \lim_{n \rightarrow \infty} P(A_n)$.

This theorem will be employed in many instances, and its use will be then pointed out.

Exercises

1.1 If $P(A) = 0.4$, $P(B) = 0.6$, and $P(A \cup B) = 0.7$, calculate $P(A \cap B)$.

1.2 If for two events A and B , it so happens that $P(A) = \frac{3}{4}$ and $P(B) = \frac{3}{8}$, show that:

$$P(A \cup B) \geq \frac{3}{4} \quad \text{and} \quad \frac{1}{8} \leq P(A \cap B) \leq \frac{3}{8}.$$

1.3 If for the events A , B , and C , it so happens that $P(A) = P(B) = P(C) = 1$, then show that:

$$P(A \cap B) = P(A \cap C) = P(B \cap C) = P(A \cap B \cap C) = 1.$$

1.4 If the events A , B , and C are related as follows: $A \subset B \subset C$ and $P(A) = \frac{1}{4}$, $P(B) = \frac{5}{12}$, and $P(C) = \frac{7}{12}$, compute the probabilities of the following events:

$$A^c \cap B, \quad A^c \cap C, \quad B^c \cap C, \quad A \cap B^c \cap C^c, \quad A^c \cap B^c \cap C^c.$$

1.5 Let S be the set of all outcomes when flipping a fair coin four times, so that all 16 outcomes are equally likely. Define the events A and B by:

$$A = \{s \in S; s \text{ contains more } Ts \text{ than } Hs\},$$

$$B = \{s \in S; \text{any } T \text{ in } s \text{ precedes every } H \text{ in } s\}.$$

Compute the probabilities $P(A)$, $P(B)$.

1.6 Let $S = \{x \text{ integer}; 1 \leq x \leq 200\}$, and define the events A , B , and C as follows:

$$A = \{x \in S; x \text{ is divisible by } 7\},$$

$$B = \{x \in S; x = 3n + 10, \text{ for some positive integer } n\},$$

$$C = \{x \in S; x^2 + 1 \leq 375\}.$$

Calculate the probabilities $P(A)$, $P(B)$, and $P(C)$.

1.7 If two fair dice are rolled once, what is the probability that the total number of spots shown is:

(i) Equal to 5?

(ii) Divisible by 3?

1.8 Students in a certain college subscribe to three news magazines A , B , and C according to the following proportions:

$$A : 20\%, \quad B : 15\%, \quad C : 10\%,$$

both A and B : 5%, both A and C : 4%, both B and C : 3%, all three A , B , and C : 2%.

If a student is chosen at random, what is the probability he/she subscribes to none of the news magazines?

1.9 A high school senior applies for admissions to two colleges A and B , and suppose that: $P(\text{admitted at } A) = p_1$, $P(\text{rejected by } B) = p_2$, and $P(\text{rejected by at least one, } A \text{ or } B) = p_3$.

- (i) Calculate the probability that the student is admitted by at least one college.
- (ii) Find the numerical value of the probability in part (i), if $p_1 = 0.6$, $p_2 = 0.2$, and $p_3 = 0.3$.

1.10 An airport limousine service has two vans, the smaller of which can carry 6 passengers and the larger 9 passengers. Let x and y be the respective numbers of passengers carried by the smaller and the larger van in a given trip, so that a suitable sample space \mathcal{S} is given by:

$$\mathcal{S} = \{(x, y); x = 0, \dots, 6 \text{ and } y = 0, 1, \dots, 9\}.$$

Also, suppose that, for all values of x and y , the probabilities $P(\{(x, y)\})$ are equal. Finally, define the events A , B , and C as follows:

A = “the two vans together carry either 4 or 6 or 10 passengers,”

B = “the larger van carries twice as many passengers as the smaller van,”

C = “the two vans carry different numbers of passengers.”

Calculate the probabilities: $P(A)$, $P(B)$, and $P(C)$.

1.11 In the sample space $\mathcal{S} = (0, \infty)$, consider the events $A_n = (0, 1 - \frac{2}{n})$, $n = 1, 2, \dots$, $A = (0, 1)$, and suppose that $P(A_n) = \frac{2n-1}{4n}$.

- (i) Show that the sequence $\{A_n\}$ is increasing and that $\lim_{n \rightarrow \infty} A_n = \bigcup_{n=1}^{\infty} A_n = A$.
- (ii) Use part (i) and the appropriate theorem (cite it!) in order to calculate the probability $P(A)$.

2.2 Distribution of a Random Variable

For a r.v. X , define the set function $P_X(B) = P(X \in B)$. Then P_X is a probability function because: $P_X(B) \geq 0$ for all B , $P_X(\mathfrak{N}) = P(X \in \mathfrak{N}) = 1$, and, if B_j , $j = 1, 2, \dots$ are pairwise disjoint then, clearly, $(X \in B_j)$, $j \geq 1$, are also pairwise disjoint and $X \in (\bigcup_{j=1}^{\infty} B_j) = \bigcup_{j=1}^{\infty} (X \in B_j)$. Therefore

$$\begin{aligned} P_X\left(\bigcup_{j=1}^{\infty} B_j\right) &= P\left[X \in \left(\bigcup_{j=1}^{\infty} B_j\right)\right] = P\left[\bigcup_{j=1}^{\infty} (X \in B_j)\right] \\ &= \sum_{j=1}^{\infty} P(X \in B_j) = \sum_{j=1}^{\infty} P_X(B_j). \end{aligned}$$

The probability function P_X is called the *probability distribution of the r.v. X* . Its significance is extremely important because it tells us the probability that X takes values in any given set B . Indeed, much of probability and statistics revolves around the distribution of r.v.'s in which we have an interest.

By selecting B to be $(-\infty, x]$, $x \in \mathfrak{R}$, we have $P_X(B) = P(X \in (-\infty, x]) = P(X \leq x)$. In effect, we define a point function which we denote by F_X ; that is, $F_X(x) = P(X \leq x)$, $x \in \mathfrak{R}$. The function F_X is called the *distribution function* (d.f.) of X . Clearly, if we know P_X , then we certainly know F_X . Somewhat unexpectedly, the converse is also true. Namely, if we know the (relatively “few”) probabilities $F_X(x)$, $x \in \mathfrak{R}$, then we can determine precisely all probabilities $P_X(B)$ for B subset of \mathfrak{R} . This converse is a deep theorem in probability that we cannot deal with here. It is, nevertheless, the reason for which it is the d.f. F_X we deal with, a familiar point function for which so many calculus results hold, rather than the unfamiliar set function P_X .

Clearly, the expressions $F_X(+\infty)$ and $F_X(-\infty)$ have no meaning because $+\infty$ and $-\infty$ are not real numbers. They are defined as follows:

$$F_X(+\infty) = \lim_{n \rightarrow \infty} F_X(x_n), x_n \uparrow \infty \quad \text{and} \quad F_X(-\infty) = \lim_{n \rightarrow \infty} F_X(y_n), y_n \downarrow -\infty.$$

These limits exist because $x < y$ implies $(-\infty, x] \subset (-\infty, y]$ and hence

$$P_X((-\infty, x]) = F_X(x) \leq F_X(y) = P_X((-\infty, y]).$$

The d.f. of a r.v. X has the following basic properties:

1. $0 \leq F_X(x) \leq 1$ for all $x \in \mathfrak{R}$;
2. F_X is a nondecreasing function;
3. F_X is continuous from the right;
4. $F_X(+\infty) = 1$, $F_X(-\infty) = 0$.

The first and the second properties are immediate from the definition of the d.f.; the third follows by Theorem 2, by taking $x_n \downarrow x$; so does the fourth, by taking $x_n \uparrow +\infty$, which implies $(-\infty, x_n] \uparrow \mathfrak{R}$, and $y_n \downarrow -\infty$, which implies $(-\infty, y_n] \downarrow \emptyset$. Figures 2.1 and 2.2 show the graphs of the d.f.'s of some typical cases.

Now, suppose that the r.v. X is discrete and takes on the values x_j , $j = 1, 2, \dots, n$. Take $b = \{x_j\}$ and on the set $\{x_1, x_2, \dots, x_n\}$ define the function f_X as follows: $f_X(x_j) = P_X(\{x_j\})$. Next, extend f_X over the entire \mathfrak{R} by setting

Figure 2.1

Examples of Graphs of
d.f.'s





Figure 2.2

Examples of Graphs of d.f.'s




$f_X(x) = 0$ for $x \neq x_j$, $j = 1, 2, \dots, n$. Then $f_X(x) \geq 0$ for all x , and it is clear that $P(X \in B) = \sum_{x_j \in B} f_X(x_j)$ for $B \subseteq \mathfrak{R}$. In particular, $\sum_{j=1}^n f_X(x_j) = \sum_{x_j \in \mathfrak{R}} f_X(x_j) = P(X \in \mathfrak{R}) = 1$. The function f_X just defined is called the *probability density function* (p.d.f.) of the r.v. X . By selecting $B = (-\infty, x]$ for some $x \in \mathfrak{R}$, we have $F_X(x) = \sum_{x_j \leq x} f_X(x_j)$. Furthermore, if we assume at this point that $x_1 < x_2 < \dots < x_n$, it is clear that

$$f_X(x_j) = F_X(x_j) - F_X(x_{j-1}), \quad j = 2, 3, \dots, n \quad \text{and} \quad f_X(x_1) = F_X(x_1);$$

we may also allow j to take the value 1 above by setting $F_X(x_0) = 0$. Likewise if X takes the values x_j , $j = 1, 2, \dots$. These two relations state that, in the case that X is a discrete r.v. as above, either one of the F_X or f_X specifies uniquely the other. Setting $F_X(x_j-)$ for the limit from the left (left-limit) of F_X at x_j , $F_X(x_j-) = \lim_{x \uparrow x_j} F_X(x)$, we see that $F_X(x_j) - F_X(x_{j-1}) = F_X(x_j) - F_X(x_j-)$, so that $f_X(x_j) = F_X(x_j) - F_X(x_j-)$. In other words, the value of f_X at x_j is the size of the jump of F_X at the point x_j . These points are illustrated quite clearly in Figure 2.3. For a numerical example (associated with Figure 2.3), let the r.v. X take on the values: $-14, -6, 5, 9,$ and 24 with respective probabilities: $0.17, 0.28, 0.22, 0.22,$ and 0.11 .

Figure 2.3



Several specific cases of discrete r.v.'s are discussed in Section 3 of the following chapter. Also, Examples 4–12, 18–20, 26, 27, and 32–35 in Chapter 1 lead to discrete r.v.'s.

Now, suppose that X is a continuous r.v., one which takes on all values in a proper interval I (finite or not) in the real line \mathfrak{R} , so that $I = (a, b)$ with $-\infty \leq a < b \leq \infty$. Suppose further that there exists a function $f : I \rightarrow [0, \infty)$ having the following property: $F_X(x) = \int_a^x f(t) dt$, $x \in I$. In particular,

$$\int_a^b f(t) dt = F_X(b) = P(X \leq b) = P(a \leq X \leq b).$$

If I is not all of \mathfrak{R} , extend f off I by setting $f(x) = 0$ for $x \notin I$. Thus, for all $x : f(x) \geq 0$ and $F_X(x) = \int_{-\infty}^x f(t) dt$. As has already been pointed out elsewhere, F_X uniquely determines P_X . The implication of it is that $P(X \in B) = P_X(B) = \int_B f(t) dt$, $B \subseteq \mathfrak{R}$, and, in particular,

$$\int_{\mathfrak{R}} f(t) dt = \int_{-\infty}^{\infty} f(t) dt = P(X \in \mathfrak{R}) = 1.$$

The function f with the properties: $f(x) \geq 0$ all x and $P(X \in B) = \int_B f(t) dt$, $B \subseteq \mathfrak{R}$, is the p.d.f. of the r.v. X . In order to emphasize its association with the r.v. X , we often write f_X .

Most of the continuous r.v.'s we are dealing with in this book do have p.d.f.'s. In Section 3 of the following chapter, a number of such r.v.'s will be presented explicitly.

Also, Examples 13–17, 21–25, and 36–40 in Chapter 1, under reasonable assumptions, lead to continuous r.v.'s, as will be seen on various occasions later. Continuous r.v.'s having p.d.f.'s, actually, form a subclass of all continuous r.v.'s and are referred to as *absolutely continuous* r.v.'s. In this book, the term continuous r.v. will be used in the sense of an absolutely continuous r.v.

It is to be observed that for a continuous r.v. X , $P(X = x) = 0$ for all $x \in \mathfrak{R}$. That is, the probability that X takes on any specific value x is 0; X takes on values with positive probabilities in a nondegenerate interval around x . That $P(X = x) = 0$ follows, of course, from the definition of the p.d.f. of a continuous r.v., as

$$P(X = x) = \int_{\{x\}} f(t) dt = 0.$$

For a case of a genuine (absolutely) continuous r.v., refer to Example 37 in Chapter 1 and let X and Y be r.v.'s denoting the cartesian coordinates of the point P of impact. Then the distance of P from the origin is the r.v. $R = \sqrt{X^2 + Y^2}$, which truly takes every value in $[0, \infty)$. As will be seen, it is reasonable to assume that X and Y are independently normally distributed with mean 0 and variance σ^2 . This leads to the fact that R^2 is a multiple of a chi-square distributed r.v., so that the p.d.f. of R is precisely determined. (See Exercise 2.14 in Chapter 5.)

If X is a continuous r.v. with p.d.f. f_X , then its d.f. F_X is given by $F_X(x) = \int_{-\infty}^x f_X(t) dt$, $x \in \mathfrak{R}$, so that f_X uniquely determines F_X . It is also true that $\frac{dF_X(x)}{dx} = f_X(x)$ (for continuity points x of f_X). Thus, F_X also determines f_X .

In summary then, with a r.v. X , we have associated three quantities:

- (i) The probability distribution of X , denoted by P_X , which is a set function and gives the probabilities $P(X \in B)$, $B \subseteq \mathfrak{R}$.
- (ii) The d.f. of X , denoted by F_X , which is a point function and gives the probabilities $P(X \in (-\infty, x]) = P(X \leq x) = F_X(x)$, $x \in \mathfrak{R}$.
- (iii) The p.d.f. f_X which is a nonnegative (point) function and gives all probabilities we may be interested in, either through a summation (for the discrete case) or through an integration (for the continuous case). Thus, for every $x \in \mathfrak{R}$:

$$F_X(x) = P(X \leq x) = \begin{cases} \sum_{x_i \leq x} f_X(x_i) & \text{for the discrete case} \\ \int_{-\infty}^x f_X(t) dt & \text{for the continuous case,} \end{cases}$$

or, more generally:

$$P_X(B) = P(X \in B) = \begin{cases} \sum_{x_i \in B} f_X(x_i) & \text{for the discrete case} \\ \int_B f_X(t) dt & \text{for the continuous case.} \end{cases}$$

In the discrete case, $f_X(x_i) = P(X = x_i) = P_X(\{x_i\})$, whereas in the continuous case, $f_X(x) = 0$ for every x . The p.d.f. f_X , clearly, determines the d.f. F_X , and the converse is also true. Of course, the p.d.f. f_X also determines the probability distribution P_X , but what is also true, although not obvious, is that the d.f. F_X determines the probability distribution P_X .

Given a r.v. X , we are primarily if not exclusively interested in its probability distribution P_X . Because of the above, it suffices to restrict ourselves either to the d.f. F_X , or even better, to the p.d.f. f_X , which is easier to work with.

The notation $X \sim F_X$ or $X \sim f_X$ stands for the statement that the r.v. X has the d.f. F_X or p.d.f. f_X , respectively.

This section is concluded with the following observation and some follow-up discussion. If Q is any probability distribution in \mathfrak{R} , then there is a r.v. X such that $P_X = Q$. To see this, let Y be a r.v. with p.d.f. $f_Y(y) = 1$ in $[0, 1]$ (and 0 outside this interval). Then its d.f. F_Y is given by:

$$F_Y(y) = 0, \quad y < 0; \quad F_Y(y) = y, \quad 0 \leq y \leq 1; \quad F_Y(y) = 1, \quad y > 1.$$

Next, let F be the d.f. determined by Q , and suppose it is strictly increasing, so that the inverse F^{-1} exists. Set $X = F^{-1}(Y)$. Then we assert that $F_X = F$. Indeed,

$$\begin{aligned} F_X(x) &= P(X \leq x) = P[F^{-1}(Y) \leq x] = P\{F[F^{-1}(Y)] \leq F(x)\} \\ &= P\{Y \leq F(x)\} = F(x), \end{aligned}$$

because $0 \leq F(x) \leq 1$. That is, $F_X(x) = F(x)$, $x \in \mathfrak{R}$. The same result follows even if F is not strictly increasing by a modified definition of the inverse F^{-1} .

Along the same line, it makes sense to ask whether a given function f is the p.d.f. of a r.v. X . The required conditions for this to be the case are: $f(x) \geq 0$

for all x , and either

$f(x_j) > 0$, $j = 1, 2, \dots$, with $\sum_j f(x_j) = 1$, and $f(x) = 0$ for all $x \neq x_j$, $j \geq 1$;

$$\text{or } \int_{-\infty}^{\infty} f(x) dx = 1.$$

This is so because in either case f defines a d.f. F and hence a r.v. X (discrete in the former case, and continuous in the latter case) with d.f. F .

Let us conclude this section with the following concrete examples.

EXAMPLE 10

The number of light switch turn-ons at which the first failure occurs is a r.v. X whose p.d.f. is given by: $f(x) = c(\frac{9}{10})^{x-1}$, $x = 1, 2, \dots$ (and 0 otherwise).

- (i) Determine the constant c .
- (ii) Calculate the probability that the first failure will not occur until after the 10th turn-on.
- (iii) Determine the corresponding d.f. F .

Hint: At this point, recall how we sum geometric series; namely, $\sum_{n=k}^{\infty} t^n = \frac{t^k}{1-t}$, $|t| < 1$, $k = 0, 1, \dots$

DISCUSSION

- (i) The constant c is determined through the relationship: $\sum_{x=1}^{\infty} f(x) = 1$ or $\sum_{x=1}^{\infty} c(\frac{9}{10})^{x-1} = 1$. However, $\sum_{x=1}^{\infty} c(\frac{9}{10})^{x-1} = c \sum_{x=1}^{\infty} (\frac{9}{10})^{x-1} = c[1 + (\frac{9}{10}) + (\frac{9}{10})^2 + \dots] = c \frac{1}{1-\frac{9}{10}} = 10c$, so that $c = \frac{1}{10}$.
- (ii) Here $P(X > 10) = P(X \geq 11) = c \sum_{x=11}^{\infty} (\frac{9}{10})^{x-1} = c[(\frac{9}{10})^{10} + (\frac{9}{10})^{11} + \dots] = c \frac{(\frac{9}{10})^{10}}{1-\frac{9}{10}} = c \cdot 10(\frac{9}{10})^{10} = \frac{1}{10} \cdot 10(\frac{9}{10})^{10} = (0.9)^{10} \simeq 0.349$.
- (iii) First, for $x < 1$, $F(x) = 0$. Next, for $x \geq 1$, $F(x) = \sum_{t=1}^x c(\frac{9}{10})^{t-1} = 1 - \sum_{t=x+1}^{\infty} c \cdot (\frac{9}{10})^{t-1} = 1 - c \sum_{t=x+1}^{\infty} (\frac{9}{10})^{t-1} = 1 - \frac{1}{10} \cdot \frac{(\frac{9}{10})^x}{1-\frac{9}{10}} = 1 - (\frac{9}{10})^x$. Thus, $F(x) = 0$ for $x < 1$, and $F(x) = 1 - (\frac{9}{10})^x$ for $x \geq 1$.

EXAMPLE 11

The recorded temperature in an engine is a r.v. X whose p.d.f. is given by: $f(x) = n(1-x)^{n-1}$, $0 \leq x \leq 1$ (and 0 otherwise), where $n \geq 1$ is a known integer.

- (i) Show that f is, indeed, a p.d.f.
- (ii) Determine the corresponding d.f. F .

DISCUSSION

- (i) Because $f(x) \geq 0$ for all x , we simply have to check that $\int_0^1 f(x) dx = 1$. To this end, $\int_0^1 f(x) dx = \int_0^1 n(1-x)^{n-1} dx = -n \frac{(1-x)^n}{n} \Big|_0^1 = -(1-x)^n \Big|_0^1 = 1$.
- (ii) First, $F(x) = 0$ for $x < 0$, whereas for $0 \leq x \leq 1$, $F(x) = \int_0^x n(1-t)^{n-1} dt = -(1-t)^n \Big|_0^x$ (from part (i)), and this is equal to: $-(1-x)^n + 1 = 1 - (1-x)^n$. Thus,

$$F(x) = \begin{cases} 0, & x < 0 \\ 1 - (1-x)^n, & 0 \leq x \leq 1 \\ 1, & x > 0. \end{cases}$$

Exercises

2.1 A sample space describing a three-children family is as follows: $S = \{bbb, bbg, bgb, gbb, bgg, gbg, ggb, ggg\}$, and assume that all eight outcomes are equally likely to occur. Next, let X be the r.v. denoting the number of girls in such a family. Then:

- (i) Determine the set of all possible values of X .
- (ii) Determine the p.d.f. of X .
- (iii) Calculate the probabilities: $P(X \geq 2)$, $P(X \leq 2)$.

2.2 A r.v. X has d.f. F given by:

$$F(x) = \begin{cases} 0, & x \leq 0 \\ 2c(x^2 - \frac{1}{3}x^3), & 0 < x \leq 2 \\ 1, & x > 2. \end{cases}$$

- (i) Determine the corresponding p.d.f. f .
- (ii) Determine the constant c .

2.3 The r.v. X has d.f. F given by:

$$F(x) = \begin{cases} 0, & x \leq 0 \\ x^3 - x^2 + x, & 0 < x \leq 1 \\ 1, & x > 1. \end{cases}$$

- (i) Determine the corresponding p.d.f. f .
- (ii) Calculate the probability $P(X > \frac{1}{2})$.

2.4 The r.v. X has d.f. F given by:

$$F(x) = \begin{cases} 0, & x < 4 \\ 0.1, & 4 \leq x < 5 \\ 0.4, & 5 \leq x < 6 \\ 0.7, & 6 \leq x < 8 \\ 0.9, & 8 \leq x < 9 \\ 1, & x \geq 9. \end{cases}$$

- (i) Draw the graph of F .
- (ii) Calculate the probabilities

$$P(X \leq 6.5), \quad P(X > 8.1), \quad P(5 < x < 8).$$

2.5 Let X be a r.v. with p.d.f. $f(x) = cx^{-(c+1)}$, for $x \geq 1$, where c is a positive constant.


- (i) Determine the constant c , so that f is, indeed, a p.d.f.
- (ii) Determine the corresponding d.f. F .

2.6 Let X be a r.v. with p.d.f. $f(x) = cx + d$, for $0 \leq x \leq 1$, and suppose that $P(X > \frac{1}{2}) = \frac{1}{3}$. Then:

- (i) Determine the constants c and d .
- (ii) Find the d.f. F of X .

- 2.7** Show that the function $f(x) = (\frac{1}{2})^x$, $x = 1, 2, \dots$ is a p.d.f.
- 2.8** For what value of c is the function $f(x) = c\alpha^x$, $x = 0, 1, \dots$ a p.d.f.? The quantity α is a number such that $0 < \alpha < 1$.
- 2.9** For what value of the positive constant c is the function $f(x) = c^x$, $x = 1, 2, \dots$ a p.d.f.?
- 2.10** The p.d.f. of a r.v. X is $f(x) = c(\frac{1}{3})^x$, for $x = 0, 1, \dots$, where c is a positive constant.
- Determine the value of c .
 - Calculate the probability $P(X \geq 3)$.
- 2.11** The r.v. X has p.d.f. f given by: $f(x) = c(1 - x^2)$, $-1 \leq x \leq 1$.
- Determine the constant c .
 - Calculate the probability $P(-0.9 < X < 0.9)$.
- 2.12** Let X be a r.v. denoting the lifetime of an electrical equipment, and suppose that the p.d.f. of X is: $f(x) = ce^{-cx}$, for $x > 0$ (for some constant $c > 0$).
- Determine the constant c .
 - Calculate the probability that X is at least equal to 10 (time units).
 - If the probability in part (ii) is 0.5, what is the value of c ?
- 2.13** The r.v. X has the so-called Pareto p.d.f. given by: $f(x) = \frac{1+\alpha}{x^{2+\alpha}}$, for $x > 1$, where α is a positive constant.
- Verify that f is, indeed, a p.d.f.
 - Calculate the probability $P(X > c)$, for some $c > 1$.
- 2.14** Suppose that the r.v. X takes on the values $0, 1, \dots$ with the respective probabilities $P(X = j) = f(j) = \frac{c}{3^j}$, $j = 0, 1, \dots$. Then:
- Determine the constant c .
Compute the probabilities:
 - $P(X \geq 3)$.
 - $P(X = 2k + 1, \quad k = 0, 1, \dots)$.
 - $P(X = 3k + 1, \quad k = 0, 1, \dots)$.
- 2.15** Let X be a r.v. with p.d.f. f whose graph is given below. Without calculating f and by using geometric arguments, compute the following probabilities:

$$P(X \leq 3), \quad P(1 \leq X \leq 2), \quad P(X > 2), \quad P(X > 5).$$



2.16 Let X be the r.v. denoting the number of a certain item sold by a merchant in a given day, and suppose that its p.d.f. is given by:

$$f(x) = \left(\frac{1}{2}\right)^{x+1}, \quad x = 0, 1, \dots$$

Calculate the following probabilities:

- (i) No items are sold.
- (ii) More than three items are sold.
- (iii) An odd number of items is sold.

2.17 Suppose a r.v. X has p.d.f. given by: $f(x) = \lambda e^{-\lambda x}$, $x > 0$, ($\lambda > 0$), and you are invited to bet whether the observed value x of X would be $\geq c$ or $< c$ for some positive constant c .

- (i) For what c would you bet in favor of $x \geq c$?
- (ii) What is the answer in part (i) if $\lambda = 4 \log 2$? (log is the natural logarithm.)

2.18 The lifetime in hours of electric tubes is a r.v. X with p.d.f. $f(x) = c^2 x e^{-cx}$, for $x \geq 0$, where c is a positive constant.

- (i) Determine the constant c for which f is, indeed, a p.d.f.
- (ii) Calculate the probability that the lifetime will be at least t hours.
- (iii) Find the numerical value in part (ii) for $c = 0.2$ and $t = 10$.

2.19 Let X be the r.v. denoting the number of forms required to be filled out by a contractor for participation in contract bids, where the values of X are 1, 2, 3, 4, and 5, and suppose that the respective probabilities are proportional to x ; that is, $P(X = x) = f(x) = cx$, $x = 1, \dots, 5$.

- (i) Determine the constant c .
- (ii) Calculate the probabilities:

$$P(X \leq 3), \quad P(2 \leq X \leq 4).$$

2.20 The recorded temperature in an engine is a r.v. X whose p.d.f. is given by: $f(x) = n(1-x)^{n-1}$, $0 < x < 1$ ($n \geq 1$, known integer). The engine is equipped with a thermostat which is activated when the temperature exceeds a specified level x_0 . If the probability of the thermostat being activated is $1/10^{2n}$, determine x_0 .

2.3 Conditional Probability and Related Results

Conditional probability is a probability in its own right, as will be seen, and it is an extremely useful tool in calculating probabilities. Essentially, it amounts to suitably modifying a sample space \mathcal{S} , associated with a random experiment, on the evidence that a certain event has occurred. Consider the following examples, by way of motivation, before a formal definition is given.

EXAMPLE 12

In tossing three distinct coins once (Example 26 in Chapter 1), consider the events $B =$ “exactly 2 heads occur” $= \{HHT, HTH, THH\}$, $A =$ “2 specified coins (e.g., coins #1 and #2) show heads” $= \{HHH, HHT\}$. Then $P(B) = \frac{3}{8}$ and $P(A) = \frac{2}{8} = \frac{1}{4}$. Now, suppose we are told that event B has occurred and we are asked to evaluate the probability of A on the basis of this evidence. Clearly, what really matters here is the event B , and, given that B has occurred, the event A occurs only if the sample point HHT appeared; that is, the event $\{HHT\} = A \cap B$ occurred. The required probability is then $\frac{1}{3} = \frac{1/8}{3/8} = \frac{P(A \cap B)}{P(B)}$, and the notation employed is $P(A|B)$ (probability of A , given that B has occurred or, just, given B). Thus, $P(A|B) = \frac{P(A \cap B)}{P(B)}$. Observe that $P(A|B) = \frac{1}{3} > \frac{1}{4} = P(A)$.

EXAMPLE 13

In rolling two distinct dice once (Example 27 in Chapter 1), consider the event B defined by: $B =$ “the sum of numbers on the upper face is 5”, so that $B = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (4, 1)\}$, and let $A =$ “the sum of numbers on the upper faces is ≥ 4 .” Then $A^c =$ “the sum of numbers on the upper faces is ≤ 3 ” $= \{(1, 1), (1, 2), (2, 1)\}$, so that $P(B) = \frac{10}{36} = \frac{5}{18}$ and $P(A) = 1 - P(A^c) = 1 - \frac{3}{36} = \frac{33}{36} = \frac{11}{12}$. Next, if we are told that B has occurred, then the only way that A occurs is if $A \cap B$ occurs, where $A \cap B =$ “the sum of numbers on the upper faces is both ≥ 4 and ≤ 5 (i.e., either 4 or 5)” $= \{(1, 3), (1, 4), (2, 2), (2, 3), (3, 1), (3, 2), (4, 1)\}$. Thus, $P(A|B) = \frac{7}{10} = \frac{7/36}{10/36} = \frac{P(A \cap B)}{P(B)}$, and observe that $P(A|B) = \frac{7}{10} < \frac{11}{12} = P(A)$.

EXAMPLE 14

In recording the gender of children in a two-children family (Example 30 in Chapter 1), let $B = \{bg, gb\}$ and let $A =$ “older child is a boy” $= \{bb, bg\}$, so that $A \cap B = \{bg\}$. Then $P(B) = \frac{1}{2} = P(A)$, $P(A|B) = \frac{1}{2}$.

These examples motivate the following definition of conditional probability.

DEFINITION 1

The *conditional* probability of an event A , given the event B with $P(B) > 0$, is denoted by $P(A|B)$ and is defined by: $P(A|B) = P(A \cap B)/P(B)$.

Replacing B by the entire sample space S , we are led back to the (*unconditional*) probability of A , as $\frac{P(A \cap S)}{P(S)} = \frac{P(A)}{1} = P(A)$. Thus, the conditional probability is a generalization of the concept of probability where S is restricted to an event B .

That the conditional probability is, indeed, a probability is seen formally as follows: $P(A|B) \geq 0$ for every A by definition;

$$P(S|B) = \frac{P(S \cap B)}{P(B)} = \frac{P(B)}{P(B)} = 1;$$

and if A_1, A_2, \dots are pairwise disjoint, then

$$\begin{aligned} P\left(\bigcup_{j=1}^{\infty} A_j \mid B\right) &= \frac{P\left[\left(\bigcup_{j=1}^{\infty} A_j\right) \cap B\right]}{P(B)} = \frac{P\left[\bigcup_{j=1}^{\infty} (A_j \cap B)\right]}{P(B)} \\ &= \frac{\sum_{j=1}^{\infty} P(A_j \cap B)}{P(B)} = \sum_{j=1}^{\infty} \frac{P(A_j \cap B)}{P(B)} = \sum_{j=1}^{\infty} P(A_j \mid B). \end{aligned}$$

It is to be noticed, furthermore, that the $P(\bar{A} \mid B)$ can be smaller or larger than the $P(\bar{A})$, or equal to the $P(\bar{A})$. The case that $P(A \mid B) = P(A)$ is of special interest and will be discussed more extensively in the next section. This point is made by Examples 12, 13, and 14.

Here are another three examples pertaining to conditional probabilities.

EXAMPLE 15

When we are recording the number of particles emitted by a certain radioactive source within a specified period of time (Example 35 in Chapter 1), we are going to see that, if X is the number of particles emitted, then X is a r.v. taking on the values $0, 1, \dots$ and that a suitable p.d.f. for it is $f_X(x) = e^{-\lambda} \frac{\lambda^x}{x!}$, $x = 0, 1, \dots$, for some constant $\lambda > 0$. Next, let B and A be the events defined by: $B = (X \geq 10)$, $A = (X \leq 11)$, so that $A \cap B = (10 \leq X \leq 11) = (X = 10 \text{ or } X = 11)$. Then

$$P(B) = \sum_{x=10}^{\infty} e^{-\lambda} \frac{\lambda^x}{x!} = e^{-\lambda} \sum_{x=10}^{\infty} \frac{\lambda^x}{x!},$$

$$P(A) = \sum_{x=0}^{11} e^{-\lambda} \frac{\lambda^x}{x!} = e^{-\lambda} \sum_{x=0}^{11} \frac{\lambda^x}{x!}, \quad \text{and}$$

$$P(A \mid B) = \left(e^{-\lambda} \frac{\lambda^{10}}{10!} + e^{-\lambda} \frac{\lambda^{11}}{11!} \right) / e^{-\lambda} \sum_{x=10}^{\infty} \frac{\lambda^x}{x!}.$$

Once again, $P(A \mid B) = \frac{P(A \cap B)}{P(B)}$. For a numerical example, take $\lambda = 10$. Then we have (by means of Poisson tables):

$$P(B) \simeq 0.5421, \quad P(A) \simeq 0.6968, \quad \text{and} \quad P(A \mid B) \simeq 0.441.$$

EXAMPLE 16

When recording the lifetime of an electronic device, or of an electrical appliance etc. (Example 36 in Chapter 1), if X is the lifetime under consideration, then X is a r.v. taking values in $[0, \infty)$, and a suitable p.d.f. for it is seen to be the function $f_X(x) = \lambda e^{-\lambda x}$, $x \geq 0$, for some constant $\lambda > 0$. Let B and A be the events: $B =$ "at the end of 5 time units, the equipment was still operating" = $(X \geq 5)$, $A =$ "the equipment lasts for no more than 2 additional time

units" = $(X \leq 7)$. Then $A \cap B = (5 \leq X \leq 7)$, and:

$$P(B) = \int_5^{\infty} \lambda e^{-\lambda x} dx = e^{-5\lambda}, \quad P(A) = \int_0^7 \lambda e^{-\lambda x} dx = 1 - e^{-7\lambda},$$

$$P(A \cap B) = \int_5^7 \lambda e^{-\lambda x} dx = e^{-5\lambda} - e^{-7\lambda}, \text{ so that}$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{e^{-5\lambda} - e^{-7\lambda}}{e^{-5\lambda}} = 1 - e^{-2\lambda}.$$

Take, for instance, $\lambda = \frac{1}{10}$. Then, given that $e^{-1} \simeq 0.36788$, the preceding probabilities are:

$$P(B) \simeq 0.607, \quad P(A) \simeq 0.503 \quad \text{and} \quad P(A|B) \simeq 0.181.$$

EXAMPLE 17

If for the events A and B , $P(A)P(B) > 0$, then show that: $P(A|B) > P(A)$ if and only if $P(B|A) > P(B)$. Likewise, $P(A|B) < P(A)$ if and only if $P(B|A) < P(B)$.

DISCUSSION Indeed, $P(A|B) > P(A)$ is equivalent to $\frac{P(A \cap B)}{P(B)} > P(A)$ or $\frac{P(A \cap B)}{P(A)} > P(B)$ or $P(B|A) > P(B)$. Likewise, $P(A|B) < P(A)$ is equivalent to $\frac{P(A \cap B)}{P(B)} < P(A)$ or $\frac{P(A \cap B)}{P(A)} < P(B)$ or $P(B|A) < P(B)$.

This section is concluded with three simple but very useful results. They are the so-called multiplicative theorem, the total probability theorem, and the Bayes formula.

THEOREM 3

(*Multiplicative Theorem*) For any n events A_1, \dots, A_n with $P(\bigcap_{j=1}^{n-1} A_j) > 0$, it holds:

$$P\left(\bigcap_{j=1}^n A_j\right) = P(A_n | A_1 \cap \dots \cap A_{n-1})P(A_{n-1} | A_1 \cap \dots \cap A_{n-2}) \\ \dots P(A_2 | A_1)P(A_1).$$

Its justification is simple, is done by induction, and is left as an exercise (see Exercise 3.8). Its significance is that we can calculate the probability of the intersection of n events, step by step, by means of conditional probabilities. The calculation of these conditional probabilities is far easier. Here is a simple example which amply illustrates the point.

EXAMPLE 18

An urn contains 10 identical balls of which 5 are black, 3 are red, and 2 are white. Four balls are drawn one at a time and without replacement. Find the probability that the first ball is black, the second red, the third white, and the fourth black.

DISCUSSION Denoting by B_1 the event that the first ball is black, and likewise for R_2 , W_3 , and B_4 , the required probability is:

$$P(B_1 \cap R_2 \cap W_3 \cap B_4) = P(B_4 | B_1 \cap R_2 \cap W_3)P(W_3 | B_1 \cap R_2)P(R_2 | B_1)P(B_1).$$

Assuming equally likely outcomes at each step, we have:

$$P(B_1) = \frac{5}{10}, \quad P(R_2 | B_1) = \frac{3}{9}, \quad P(W_3 | B_1 \cap R_2) = \frac{2}{8},$$

$$P(B_4 | B_1 \cap R_2 \cap W_3) = \frac{4}{7}.$$

Therefore,

$$P(B_1 \cap R_2 \cap W_3 \cap B_4) = \frac{4}{7} \times \frac{2}{8} \times \frac{3}{9} \times \frac{5}{10} = \frac{1}{42} \simeq 0.024.$$

For the formulation of the next result, the concept of a partition of S is required. The events $\{A_1, A_2, \dots, A_n\}$ form a *partition* of S , if these events are pairwise disjoint, $A_i \cap A_j = \emptyset$, $i \neq j$, and their union is S , $\bigcup_{j=1}^n A_j = S$. Then it is obvious that any event B in S may be expressed as follows, in terms of a partition of S ; namely, $B = \bigcup_{j=1}^n (A_j \cap B)$. Furthermore,

$$P(B) = \sum_{j=1}^n P(A_j \cap B) = \sum_{j=1}^n P(B | A_j)P(A_j), \quad \text{provided } P(A_j) > 0 \text{ for all } j.$$

The concept of partition is defined similarly for countably infinite many events, and the probability $P(B)$ is expressed likewise. In the sequel, by writing $j = 1, 2, \dots$ and \sum_j we mean to include both cases, finitely many indices and countably infinite many indices.

Thus, we have the following result.

THEOREM 4

(*Total Probability Theorem*) Let $\{A_1, A_2, \dots\}$ be a partition of S , and let $P(A_j) > 0$ for all j . Then, for any event B ,

$$P(B) = \sum_j P(B | A_j)P(A_j).$$

The significance of this result is that, if it happens that we know the probabilities of the partitioning events $P(A_j)$, as well as the conditional probabilities of B , given A_j , then these quantities may be combined, according to the preceding formula, to produce the probability $P(B)$. The probabilities $P(A_j)$, $j = 1, 2, \dots$ are referred to as *a priori* or *prior* probabilities. The following examples illustrate the theorem and also demonstrate its usefulness.

EXAMPLE 19

In reference to Example 2 in Chapter 1, calculate the probability $P(+)$.

DISCUSSION Without having to refer specifically to a sample space, it is clear that the events D and N form a partition. Then,

$$P(+)=P(+ \text{ and } D)+P(+ \text{ and } N)=P(+|D)P(D)+P(+|N)P(N).$$

Here the a priori probabilities are $P(D) = p_1$, $P(N) = 1 - p_1$, and

$$P(+ | D) = 1 - P(- | D) = 1 - p_3, \quad P(+ | N) = p_2.$$

Therefore, $P(+)= (1 - p_3)p_1 + p_2(1 - p_1)$. For a numerical application, take $p_1 = 0.02$ and $p_2 = p_3 = 0.01$. Then $P(+)= 0.0296$. So, on the basis of this testing procedure, about 2.96% of the population would test positive.

EXAMPLE 20

The proportion of motorists in a given gas station using regular unleaded gasoline, extra unleaded, and premium unleaded over a specified period of time are 40%, 35%, and 25%, respectively. The respective proportions of filling their tanks are 30%, 50%, and 60%. What is the probability that a motorist selected at random from among the patrons of the gas station under consideration and for the specified period of time will fill his/her tank?

DISCUSSION Denote by R , E , and P the events of a motorist using unleaded gasoline which is regular, extra unleaded, and premium, respectively, and by F the event of having the tank filled. Then the translation into terms of probabilities of the proportions given above is:

$$P(R) = 0.40, \quad P(E) = 0.35, \quad P(P) = 0.25,$$

$$P(F | R) = 0.30, \quad P(F | E) = 0.50, \quad P(F | P) = 0.60.$$

Then the required probability is:

$$\begin{aligned} P(F) &= P((F \cap R) \cup (F \cap E) \cup (F \cap P)) \\ &= P(F \cap R) + P(F \cap E) + P(F \cap P) \\ &= P(F | R)P(R) + P(F | E)P(E) + P(F | P)P(P) \\ &= 0.30 \times 0.40 + 0.50 \times 0.35 + 0.60 \times 0.25 \\ &= 0.445. \end{aligned}$$

In reference to Theorem 4, stipulating the prior probabilities $P(B | A_j)$, $j = 1, 2, \dots$, is often a precarious thing and guesswork. This being the case, the question then arises of whether experimentation may lead to reevaluation of the prior probabilities on the basis of new evidence. To put it more formally, is it possible to use $P(A_j)$ and $P(B | A_j)$, $j = 1, 2, \dots$ in order to calculate $P(A_j | B)$? The answer to this question is in the affirmative, is quite simple, and is the content of the next result.

THEOREM 5

(*Bayes' Formula*) Let $\{A_1, A_2, \dots\}$ and B be as in the previous theorem. Then, for any $j = 1, 2, \dots$:

$$P(A_j | B) = \frac{P(B | A_j)P(A_j)}{\sum_i P(B | A_i)P(A_i)}.$$

PROOF Indeed, $P(A_j | B) = P(A_j \cap B)/P(B) = P(B | A_j)P(A_j)/P(B)$, and then the previous theorem completes the proof. ▲

The probabilities $P(A_j | B)$, $j = 1, 2, \dots$, are referred to as *posterior* probabilities in that they are reevaluations of the respective prior $P(A_j)$ after the event B has occurred.

EXAMPLE 21

Referring to Example 19, a question of much importance is this: Given that the test shows positive, what is the probability that the patient actually has the disease? In terms of the notation adopted, this question becomes: $P(D | +) = ?$ Bayes' formula gives:

$$P(D | +) = \frac{P(+ | D)P(D)}{P(+ | D)P(D) + P(+ | N)P(N)} = \frac{p_1(1 - p_3)}{p_1(1 - p_3) + p_2(1 - p_1)}.$$

For the numerical values used above, we get:

$$P(D | +) = \frac{0.02 \times 0.99}{0.0296} = \frac{0.0198}{0.0296} = \frac{198}{296} \simeq 0.669.$$

So $P(D | +) \simeq 66.9\%$. This result is both reassuring and surprising. Reassuring, in that only 66.9% of those testing positive actually have the disease. Surprising, in that this proportion looks rather low, given that the test is quite good: it identifies correctly 99% of those having the disease. A reconciliation between these two seemingly contradictory aspects is as follows: The fact that $P(D) = 0.02$ means that, on the average, 2 out of 100 persons have the disease. So, in 100 persons, 2 will have the disease and 98 will not. When 100 such persons are tested, $2 \times 0.99 = 1.98$ will be correctly confirmed as positive (because 0.99 is the probability of a correct positive), and $98 \times 0.01 = 0.98$ will be incorrectly diagnosed as positive (because 0.01 is the probability of an incorrect positive). Thus, the proportion of correct positives is equal to:

$$\begin{aligned} & (\text{correct positives})/(\text{correct positives} + \text{incorrect positives}) \\ &= 1.98/(1.98 + 0.98) = 1.98/2.96 = 198/296 \simeq 0.669. \end{aligned}$$

REMARK 1 The fact that the probability $P(D | +)$ is less than 1 simply reflects the fact that the test, no matter how good, is imperfect. Should the test be perfect ($P(+ | D) = P(- | D^c) = 1$), then $P(D | +) = 1$, as follows from the preceding calculations, no matter what $P(D)$ is. The same, of course, is true for $P(D^c | -)$.

EXAMPLE 22

Refer to Example 20 and calculate the probabilities: $P(R | F)$, $P(E | F)$, and $P(P | F)$.

DISCUSSION By Bayes' formula and Example 20,

$$P(R | F) = \frac{P(R \cap F)}{P(F)} = \frac{P(F | R)P(R)}{P(F)} = \frac{0.30 \times 0.40}{0.445} \simeq 0.270,$$

and likewise,

$$P(E|F) = \frac{0.50 \times 0.35}{0.445} \simeq 0.393, \quad P(P|F) = \frac{0.60 \times 0.25}{0.445} \simeq 0.337.$$

Exercises

- 3.1** If $P(A|B) > P(A)$, then show that $P(B|A) > P(B)$, by assuming that both $P(A)$ and $P(B)$ are positive.
- 3.2** If $A \cap B = \emptyset$ and $P(A \cup B) > 0$, express the probabilities $P(A|A \cup B)$ and $P(B|A \cup B)$ in terms of $P(A)$ and $P(B)$.
- 3.3** A girls' club has in its membership rolls the names of 50 girls with the following descriptions:
 20 blondes, 15 with blue eyes and 5 with brown eyes;
 25 brunettes, 5 with blue eyes and 20 with brown eyes;
 5 redheads, 1 with blue eyes and 4 with green eyes.
 If one arranges a blind date with a club member, what is the probability that:
- The girl is blonde?
 - The girl is blonde, if it was revealed only that she has blue eyes?
- 3.4** Suppose that the probability that both of a pair of twins are boys is 0.30 and that the probability that they are both girls is 0.26. Given that the probability of the first child being a boy is 0.52, what is the probability that:
- The second twin is a boy, given that the first is a boy?
 - The second twin is a girl, given that the first is a girl?
 - The second twin is a boy?
 - The first is a boy and the second is a girl?
- 3.5** A shipment of 20 TV tubes contains 16 good tubes and 4 defective tubes. Three tubes are chosen successively and at random each time and are also tested successively. What is the probability that:
- The third tube is good if the first two were found to be good?
 - The third tube is defective if the first was found to be good and the second defective?
 - The third tube is defective if the first was found to be defective and the second was found to be good?
 - The third tube is defective if one of the other two was found to be good and the other was found to be defective?
- 3.6** For any three events A , B , and C with $P(A)P(B)P(C) > 0$, show that:
- $P(A^c|B) = 1 - P(A|B)$.
 - $P(A \cup B|C) = P(A|C) + P(B|C) - P(A \cap B|C)$.
- Also, by means of counterexamples, show that the following equations need not be true:

$$(iii) P(A | B^c) = 1 - P(A | B).$$

$$(iv) P(C | A \cup B) = P(C | A) + P(C | B), \text{ where } A \cap B = \emptyset.$$

3.7 If A , B , and C are any events in the sample space \mathcal{S} , show that $\{A, A^c \cap B, A^c \cap B^c \cap C, (A \cup B \cup C)^c\}$ is a partition of \mathcal{S} .

3.8 Use induction to prove Theorem 3.

3.9 Let $\{A_j, j = 1, \dots, 5\}$ be a partition of the sample space \mathcal{S} and suppose that:

$$P(A_j) = \frac{j}{15} \quad \text{and} \quad P(A | A_j) = \frac{5-j}{15}, \quad j = 1, \dots, 5.$$

Compute the probabilities $P(A_j | A), j = 1, \dots, 5$.

3.10 A box contains 15 identical balls except that 10 are red and 5 are black. Four balls are drawn successively and without replacement.

Calculate the probability that the first and the fourth balls are red.

3.11 A box contains $m + n$ identical balls except that m of them are red and n are black. A ball is drawn at random, its color is noticed, and then the ball is returned to the box along with r balls of the same color. Finally, a ball is drawn also at random.

- (i) What is the probability that the first ball is red?
- (ii) What is the probability that the second ball is red?
- (iii) Compare the probabilities in parts (i) and (ii) and comment on them.
- (iv) What is the probability that the first ball is black if the second is red?
- (v) Find the numerical values in parts (i), (ii), and (iv) if $m = 9, n = 6,$ and $r = 5$.

3.12 A test correctly identifies a disease D with probability 0.95 and wrongly diagnoses D with probability 0.01. From past experience, it is known that disease D occurs in a targeted population with frequency 0.2%. An individual is chosen at random from said population and is given the test. Calculate the probability that:

- (i) The test is +, $P(+)$.
- (ii) The individual actually suffers from disease D if the test turns out to be positive, $P(D | +)$.


3.13 Suppose that the probability of correct diagnosis (either positive or negative) of cervical cancer in the Pap test is 0.95 and that the proportion of women in a given population suffering from this disease is 0.01%. A woman is chosen at random from the target population and the test is administered.

What is the probability that:

- (i) The test is positive?
- (ii) The subject actually has the disease, given that the diagnosis is positive?

3.14 A signal S is sent from point A to point B and is received at B if both switches I and II are closed. It is assumed that the probabilities of I and

II being closed are 0.8 and 0.6, respectively, and that $P(\text{II is closed} | \text{I is closed}) = P(\text{II is closed})$.



Calculate the following probabilities:

- (i) The signal is received at B .
- (ii) The (conditional) probability that switch I was open, given that the signal was not received at B .
- (iii) The (conditional) probability that switch II was open, given that the signal was not received at B .

3.15 The student body in a certain college consists of 55% women and 45% men. Women and men smoke cigarettes in the proportions of 20% and 25%, respectively. If a student is chosen at random, calculate the probability that:

- (i) The student is a smoker.
- (ii) The student is a man, given that he/she is a smoker.

3.16 From a population consisting of 52% females and 48% males, an individual, drawn at random, is found to be color blind. If we assume that the proportions of color-blind females and males are 25% and 5%, respectively, what is the probability that the individual drawn is a male?

3.17 Drawers I and II contain black and red pencils as follows:

Drawer I: b_1 black pencils and r_1 red pencils,

Drawer II: b_2 black pencils and r_2 red pencils.

A drawer is chosen at random and then a pencil is also chosen at random from that drawer.

- (i) What is the probability that the pencil is black?
- (ii) If it is announced that the pencil is black, what is the probability it was chosen from drawer I?
- (iii) Give numerical values in parts (i) and (ii) for $b_1 = 36$, $r_1 = 12$, $b_2 = 60$, $r_2 = 24$.

3.18 Three machines I, II, and III manufacture 30%, 30%, and 40%, respectively, of the total output of certain items. Of them, 4%, 3%, and 2%, respectively, are defective. One item is drawn at random from the total output and is tested.

- (i) What is the probability that the item is defective?
- (ii) If it is found to be defective, what is the probability the item was produced by machine I?
- (iii) Same question as in part (ii) for each one of the machines II and III.

3.19 Suppose that a multiple choice test lists n alternative answers of which only one is correct. If a student has done the homework, he/she is certain to identify the correct answer; otherwise the student chooses an answer at random. Denote by A the event that the student does the homework, set $p = P(A)$, and let B be the event that he/she answers the question correctly.

- (i) Express the probability $P(A|B)$ in terms of p and n .
- (ii) If $0 < p < 1$ and fixed, show that the probability $P(A|B)$, as a function of n , is increasing.
- (iii) Does the result in part (ii) seem reasonable?

3.20 If the p.d.f. of the r.v. X is: $f(x) = \lambda e^{-\lambda x}$, for $x > 0$ ($\lambda > 0$), calculate:

- (i) $P(X > t)$ (for some $t > 0$).
- (ii) $P(X > s + t | X > s)$ (for some $s, t > 0$).
- (iii) Compare the probabilities in parts (i) and (ii), and draw your conclusion.

2.4 Independent Events and Related Results

In Example 14, it was seen that $P(A|B) = P(A)$. Thus, the fact that the event B occurred provides no information in reevaluating the probability of A . Under such a circumstance, it is only fitting to say that A is independent of B . For any two events A and B with $P(B) > 0$, we say that A is *independent* of B , if $P(A|B) = P(A)$. If, in addition, $P(A) > 0$, then B is also *independent* of A because

$$P(B|A) = \frac{P(B \cap A)}{P(A)} = \frac{P(A \cap B)}{P(A)} = \frac{P(A|B)P(B)}{P(A)} = \frac{P(A)P(B)}{P(A)} = P(B).$$

Because of this symmetry, we then say that A and B are independent. From the definition of either $P(A|B)$ or $P(B|A)$, it follows then that $P(A \cap B) = P(A)P(B)$. We further observe that this relation is true even if one or both of $P(A)$, $P(B)$ are equal to 0. We take this relation as the defining relation of independence.

DEFINITION 2

Two events A_1 and A_2 are said to be *independent* (statistically or stochastically or in the probability sense), if $P(A_1 \cap A_2) = P(A_1)P(A_2)$. When $P(A_1 \cap A_2) \neq P(A_1)P(A_2)$ they are said to be *dependent*.

REMARK 2 At this point, it should be emphasized that disjointness and independence of two events are two distinct concepts; the former does not even require the concept of probability. Nevertheless, they are related in that, if $A_1 \cap A_2 = \emptyset$, then they are independent if and only if at least one of $P(A_1)$, $P(A_2)$ is equal to 0. Thus (subject to $A_1 \cap A_2 = \emptyset$), $P(A_1)P(A_2) > 0$ implies that A_1 and A_2 are definitely dependent.

The definition of independence extends to three events A_1, A_2, A_3 , as well as to any number n of events A_1, \dots, A_n . Thus, three events A_1, A_2, A_3 for which $P(A_1 \cap A_2 \cap A_3) > 0$ are said to be independent, if all conditional probabilities coincide with the respective (unconditional) probabilities:

$$\left. \begin{aligned} P(A_1 | A_2) &= P(A_1 | A_3) = P(A_1 | A_2 \cap A_3) = P(A_1) \\ P(A_2 | A_1) &= P(A_2 | A_3) = P(A_2 | A_1 \cap A_3) = P(A_2) \\ P(A_3 | A_1) &= P(A_3 | A_2) = P(A_3 | A_1 \cap A_2) = P(A_3) \\ P(A_1 \cap A_2 | A_3) &= P(A_1 \cap A_2), P(A_1 \cap A_3 | A_2) \\ &= P(A_1 \cap A_3), P(A_2 \cap A_3 | A_1) = P(A_2 \cap A_3). \end{aligned} \right\} \quad (1)$$

From the definition of conditional probability, relations (1) are equivalent to:

$$\left. \begin{aligned} P(A_1 \cap A_2) &= P(A_1)P(A_2), P(A_1 \cap A_3) = P(A_1)P(A_3), \\ P(A_2 \cap A_3) &= P(A_2)P(A_3), P(A_1 \cap A_2 \cap A_3) = P(A_1)P(A_2)P(A_3). \end{aligned} \right\} \quad (2)$$

Furthermore, it is to be observed that relations (2) hold even if any of $P(A_1), P(A_2), P(A_3)$ are equal to 0. These relations are taken as defining relations of independence of three events A_1, A_2, A_3 .

As one would expect, all four relations (2) are needed for independence (that is, in order for them to imply relations (1)). That this is, indeed, the case is illustrated by the following examples.

EXAMPLE 23

Let $S = \{1, 2, 3, 4\}$ and let $P(\{1\}) = P(\{2\}) = P(\{3\}) = P(\{4\}) = 1/4$. Define the events A_1, A_2, A_3 by: $A_1 = \{1, 2\}$, $A_2 = \{1, 3\}$, $A_3 = \{1, 4\}$. Then it is easily verified that: $P(A_1 \cap A_2) = P(A_1)P(A_2)$, $P(A_1 \cap A_3) = P(A_1)P(A_3)$, $P(A_2 \cap A_3) = P(A_2)P(A_3)$. However, $P(A_1 \cap A_2 \cap A_3) \neq P(A_1)P(A_2)P(A_3)$.

EXAMPLE 24

Let $S = \{1, 2, 3, 4, 5\}$ and let $P(\{1\}) = \frac{2}{16}$, $P(\{2\}) = P(\{3\}) = P(\{4\}) = \frac{3}{16}$, $P(\{5\}) = \frac{5}{16}$. Define the events A_1, A_2, A_3 by: $A_1 = \{1, 2, 3\}$, $A_2 = \{1, 2, 4\}$, $A_3 = \{1, 3, 4\}$. Then it is easily verified that: $P(A_1 \cap A_2 \cap A_3) = P(A_1)P(A_2)P(A_3)$ but none of the other three relations in (2) is satisfied.

Relations (2) provide the pattern of the definition of independence of n events. Thus:

DEFINITION 3

The events A_1, \dots, A_n are said to be *independent (statistically or stochastically or in the probability sense)* if, for all possible choices of k out of n events ($2 \leq k \leq n$), the probability of their intersection equals the product of their probabilities. More formally, for any k with $2 \leq k \leq n$

and any integers j_1, \dots, j_k with $1 \leq j_1 < \dots < j_k \leq n$, we have:

$$P\left(\bigcap_{i=1}^k A_{j_i}\right) = \prod_{i=1}^k P(A_{j_i}). \quad (3)$$

If at least one of the relations in (3) is violated, the events are said to be *dependent*. The number of relations of the form (3) required to express independence of n events is:

$$\binom{n}{2} + \binom{n}{3} + \dots + \binom{n}{n} = 2^n - \binom{n}{1} - \binom{n}{0} = 2^n - n - 1.$$

For example, for $n = 2, 3$, these relations are: $2^2 - 2 - 1 = 1$ and $2^3 - 3 - 1 = 4$, respectively.

Typical cases where independent events occur are whenever we are sampling with replacement from finite populations, such as selecting successively and with replacement balls from an urn containing balls of several colors, pulling successively and with replacement playing cards out of a standard deck of such cards, and the like.

The following property of independence of events is often used without even being acknowledged; it is stated here as a theorem.

THEOREM 6

- (i) If the events A_1, A_2 are independent, then so are all three sets of events: $A_1, A_2^c; A_1^c, A_2; A_1^c, A_2^c$.
(ii) More generally, if the events A_1, \dots, A_n are independent, then so are the events A'_1, \dots, A'_n , where A'_i stands either for A_i or A_i^c , $i = 1, \dots, n$.
For illustrative purposes, we present the proof of part (i) only.

PROOF OF PART (i) Clearly, $A_1 \cap A_2^c = A_1 - A_1 \cap A_2$. Thus,

$$\begin{aligned} P(A_1 \cap A_2^c) &= P(A_1 - A_1 \cap A_2) = P(A_1) - P(A_1 \cap A_2) \text{ (since } A_1 \cap A_2 \subseteq A_1) \\ &= P(A_1) - P(A_1)P(A_2) \text{ (by independence of } A_1, A_2) \\ &= P(A_1)[1 - P(A_2)] = P(A_1)P(A_2^c). \end{aligned}$$

The proof of $P(A_1^c \cap A_2) = P(A_1^c)P(A_2)$ is entirely symmetric. Finally,

$$\begin{aligned} P(A_1^c \cap A_2^c) &= P((A_1 \cup A_2)^c) \text{ (by DeMorgan's laws)} \\ &= 1 - P(A_1 \cup A_2) \\ &= 1 - P(A_1) - P(A_2) + P(A_1 \cap A_2) \\ &= 1 - P(A_1) - P(A_2) + P(A_1)P(A_2) \text{ (by independence of } A_1, A_2) \\ &= [1 - P(A_1)] - P(A_2)[1 - P(A_1)] \\ &= P(A_1^c)P(A_2^c). \quad \blacktriangle \end{aligned}$$

The following examples will help illustrate concepts and results discussed in this section.

EXAMPLE 25

Suppose that $P(B)P(B^c) > 0$. Then the events A and B are independent if and only if $P(A|B) = P(A|B^c)$.

DISCUSSION First, if A and B are independent, then A and B^c are also independent, by Theorem 6. Thus, $P(A|B^c) = \frac{P(A \cap B^c)}{P(B^c)} = \frac{P(A)P(B^c)}{P(B^c)} = P(A)$. Since also $P(A|B) = P(A)$, the equality $P(A|B) = P(A|B^c)$ holds. Next, $P(A|B) = P(A|B^c)$ is equivalent to $\frac{P(A \cap B)}{P(B)} = \frac{P(A \cap B^c)}{P(B^c)}$ or $P(A \cap B)P(B^c) = P(A \cap B^c)P(B)$ or $P(A \cap B)[1 - P(B)] = P(A \cap B^c)P(B)$ or $P(A \cap B) - P(A \cap B)P(B) = P(A \cap B^c)P(B)$ or $P(A \cap B) = [P(A \cap B) + P(A \cap B^c)]P(B) = P(A)P(B)$, since $(A \cap B) \cup (A \cap B^c) = A$. Thus, A and B are independent.

REMARK 3 It is to be pointed out that the condition $P(A|B) = P(A|B^c)$ for independence of the events A and B is quite natural, intuitively. It says that the (conditional) probability of A remains the same no matter which one of B or B^c is given.

EXAMPLE 26

Let $P(C)P(C^c) > 0$. Then the inequalities $P(A|C) > P(B|C)$ and $P(A|C^c) > P(B|C^c)$ imply $P(A) > P(B)$.

DISCUSSION The inequalities $P(A|C) > P(B|C)$ and $P(A|C^c) > P(B|C^c)$ are equivalent to $P(A \cap C) > P(B \cap C)$ and $P(A \cap C^c) > P(B \cap C^c)$. Adding up these inequalities, we obtain $P(A \cap C) + P(A \cap C^c) > P(B \cap C) + P(B \cap C^c)$ or $P(A) > P(B)$, since $A = (A \cap C) \cup (A \cap C^c)$ and $B = (B \cap C) \cup (B \cap C^c)$.

REMARK 4 Once again, that the inequalities of the two conditional probabilities should imply the same inequality for the unconditional probabilities is quite obvious on intuitive grounds. The justification given above simply makes it rigorous.

EXAMPLE 27

If the events A , B , and C are independent, then $P(A \cup B \cup C) = 1 - [1 - P(A)][1 - P(B)][1 - P(C)]$.

DISCUSSION Clearly,

$$\begin{aligned} P(A \cup B \cup C) &= P[(A^c \cap B^c \cap C^c)^c] && \text{(by DeMorgan's laws)} \\ &= 1 - P(A^c \cap B^c \cap C^c) && \text{(by basic property (3))} \\ &= 1 - P(A^c)P(B^c)P(C^c) && \text{(by Theorem 6(ii)} \\ &&& \text{applied with } n = 3) \\ &= 1 - [1 - P(A)][1 - P(B)][1 - P(C)]. \end{aligned}$$

EXAMPLE 28

A mouse caught in a maze has to maneuver through three successive escape hatches in order to escape. If the hatches operate independently and the probabilities for the mouse to maneuver successfully through them are 0.6,

0.4, and 0.2, respectively, calculate the probabilities that the mouse: (i) will be able to escape, (ii) will not be able to escape.

DISCUSSION Denote by H_1 , H_2 , and H_3 the events that the mouse successfully maneuvers through the three hatches, and by E the event that the mouse is able to escape. We have that H_1 , H_2 , and H_3 are independent, $P(H_1) = 0.6$, $P(H_2) = 0.4$, and $P(H_3) = 0.2$, and $E = H_1 \cap H_2 \cap H_3$. Then: (i) $P(E) = P(H_1 \cap H_2 \cap H_3) = P(H_1)P(H_2)P(H_3) = 0.6 \times 0.4 \times 0.2 = 0.048$, and (ii) $P(E^c) = 1 - P(E) = 1 - 0.048 = 0.952$.

The concept of independence carries over to random experiments. Although a technical definition of independence of random experiments is available, we are not going to indulge in it. The concept of independence of random experiments will be taken in its intuitive sense, and somewhat more technically, in the sense that random experiments are independent if they give rise to independent events associated with them.

Finally, independence is also defined for r.v.'s. This topic will be taken up in Chapter 5 (see Definition 1 there). Actually, independence of r.v.'s is one of the founding blocks of most discussions taking place in this book.

Exercises

- 4.1** If $P(A) = 0.4$, $P(B) = 0.2$, and $P(C) = 0.3$, calculate the probability $P(A \cup B \cup C)$, if the events A , B , and C are:
- Pairwise disjoint.
 - Independent.
- 4.2** Show that the event A is independent of itself if and only if $P(A) = 0$ or $P(A) = 1$.
- 4.3** (i) For any two events A and B , show that $P(A \cap B) \geq P(A) + P(B) - 1$.
 (ii) If A and B are disjoint, then show that they are independent if and only if at least one of $P(A)$ and $P(B)$ is zero.
 (iii) If the events A , B , and C are pairwise disjoint, under what conditions are they independent?
- 4.4** Suppose that the events A_1 , A_2 , and B_1 are independent, the events A_1 , A_2 , and B_2 are independent, and that $B_1 \cap B_2 = \emptyset$. Then show that the events A_1 , A_2 , $B_1 \cup B_2$ are independent.
- 4.5** (i) If for the events A , B , and C , it so happens that $P(A) = P(B) = P(C) = \frac{1}{2}$, $P(A \cap B) = P(A \cap C) = P(B \cap C) = \frac{1}{4}$, and $P(A \cap B \cap C) = \frac{1}{6}$, determine whether or not these events are independent. Justify your answer.
 (ii) For the values given in part (i), calculate the probabilities: $P(A^c)$, $P(A \cup B)$, $P(A^c \cap B^c)$, $P(A \cup B \cup C)$, and $P(A^c \cap B^c \cap C^c)$.

4.6 For the events A, B, C and their complements, suppose that:

$$\begin{aligned} P(A \cap B \cap C) &= \frac{1}{16}, & P(A \cap B^c \cap C) &= \frac{5}{16}, & P(A \cap B \cap C^c) &= \frac{3}{16}, \\ P(A \cap B^c \cap C^c) &= \frac{2}{16}, & P(A^c \cap B \cap C) &= \frac{2}{16}, & P(A^c \cap B \cap C^c) &= \frac{1}{16}, \\ P(A^c \cap B^c \cap C) &= \frac{1}{16}, & \text{and } P(A^c \cap B^c \cap C^c) &= \frac{1}{16}. \end{aligned}$$

- (i) Calculate the probabilities: $P(A), P(B), P(C)$.
- (ii) Determine whether or not the events $A, B,$ and C are independent.
- (iii) Calculate the (conditional) probability $P(A|B)$.
- (iv) Determine whether or not the events A and B are independent.

4.7 If the events A_1, \dots, A_n are independent, show that

$$P\left(\bigcup_{j=1}^n A_j\right) = 1 - \prod_{j=1}^n P(A_j^c).$$

- 4.8** (i) Three coins, with probability of falling heads being p , are tossed once and you win, if all three coins show the same face (either all H or all T). What is the probability of winning?
- (ii) What are the numerical answers in part (i) for $p = 0.5$ and $p = 0.4$?
- 4.9** Suppose that men and women are distributed in the freshman and sophomore classes of a college according to the proportions listed in the following table.

Class\Gender	<i>M</i>	<i>W</i>	Totals
<i>F</i>	4	6	10
<i>S</i>	6	x	$6 + x$
Totals	10	$6 + x$	$16 + x$

A student is chosen at random and let $M, W, F,$ and S be the events, respectively, that the student is a man, a woman, a freshman, or a sophomore. Then, being a man or a woman and being a freshman or sophomore are independent, if:


$$\begin{aligned} P(M \cap F) &= P(M)P(F), & P(W \cap F) &= P(W)P(F), \\ P(M \cap S) &= P(M)P(S), & P(W \cap S) &= P(W)P(S). \end{aligned}$$

Determine the number x , so that the preceding independence relations hold.

4.10 The r.v. X has p.d.f. given by:

$$f(x) = \begin{cases} cx, & 0 \leq x < 5 \\ c(10 - x), & 5 \leq x < 10 \\ 0, & \text{elsewhere.} \end{cases}$$

- (i) Determine the constant c .
 (ii) Draw the graph of f .



Define the events A and B by: $A = (X > 5)$, $B = (5 < X < 7.5)$.

- (iii) Calculate the probabilities $P(A)$ and $P(B)$.
 (iv) Calculate the conditional probability $P(B | A)$.
 (v) Are the events A and B independent or not? Justify your answer.
- 4.11** Three players I , II , III throw simultaneously three coins with respective probabilities of falling heads (H) p_1 , p_2 , and p_3 . A sample space describing this experiment is:

$$S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}.$$

Define the events A_i , $i = 1, 2, 3$ and B by:

$$A_1 = \{HTT, THH\} \quad A_2 = \{THT, HTH\}, \quad A_3 = \{TTH, HHT\}$$

(i.e., the outcome for the i th player, $i = 1, 2, 3$, is different from those for the other two players),

$$B = \{HHH, TTT\}.$$

If any one of the events A_i , $i = 1, 2, 3$ occurs, the i th player wins and the game ends. If event B occurs, the game is repeated independently as many times as needed until one of the events A_1 , A_2 , A_3 occurs.

- (i) Calculate the probabilities: $P(A_i)$, $i = 1, 2, 3$.
 (ii) What do these probabilities become for $p_1 = p_2 = p_3 = p$?
 (iii) What is the numerical value in part (ii) if $p = 0.5$?

Hint: By symmetry, it suffices to calculate $P(A_1)$. Let A_{1j} = "event A_1 occurs the j th time," B_j = "event B occurs the j th time." Then (with slight abuse of notation)

$$A_1 = A_{11} \cup (B_1 \cap A_{12}) \cup (B_1 \cap B_2 \cap A_{13}) \cup \dots$$

At this point, also recall that: $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$, $|x| < 1$.

- 4.12** Jim takes the written and road driver's license tests repeatedly until he passes them. It is given that the probability that he passes the written test is 0.9, that he passes the road test is 0.6, and that the tests are independent of each other. Furthermore, it is assumed that the road test cannot be taken unless he passes the written test, and that once he passes the

written test, he does not have to take it again ever, no matter whether he passes or fails his road tests. Also, it is assumed that the written and the road test are distinct attempts.

- (i) What is the probability that he will pass the road test on his n th attempt?
- (ii) What is the numerical value in part (i) for $n = 5$?

Hint: Denote by W_i and R_j the events that Jim passes the written test and the road test the i th and j th time, respectively. Then the required event is expressed as follows:


$$(W_1 \cap R_1^c \cap \dots \cap R_{n-2}^c \cap R_{n-1}) \cup (W_1^c \cap W_2 \cap R_1^c \cap \dots \cap R_{n-3}^c \cap R_{n-2}) \\ \cup \dots \cup (W_1^c \cap \dots \cap W_{n-2}^c \cap W_{n-1} \cap R_n).$$

4.13 The probability that a missile fired against a target is not intercepted by an antimissile missile is $\frac{2}{3}$. If the missile is not intercepted, then the probability of a successful hit is $\frac{3}{4}$.

If four missiles are fired independently, what is the probability that:

- (i) All four will successfully hit the target?
- (ii) At least one will do so?
- (iii) What is the minimum number of missiles to be fired so that at least one is not intercepted with probability at least 0.95?
- (iv) What is the minimum number of missiles to be fired so that at least one hits the target with probability at least 0.99?

4.14 Electric current is transmitted from point A to point B, provided at least one of the circuits #1 through # n here is closed. It is assumed that the n circuits close independently of each other and with respective probabilities p_1, \dots, p_n .



Determine the following probabilities:

- (i) No circuit is closed.
- (ii) At least one circuit is closed.
- (iii) Exactly one circuit is closed.
- (iv) How do the expressions in parts (i)–(iii) simplify if $p_1 = \dots = p_n = p$?
- (v) What are the numerical values in part (iv) for $n = 5$ and $p = 0.6$?

4.15 Consider two urns U_1 and U_2 such that urn U_1 contains m_1 white balls and n_1 black balls, and urn U_2 contains m_2 white balls and n_2 black balls. All balls are identical except for color. One ball is drawn at random from each of the urns U_1 and U_2 and is placed into a third urn. Then a ball is drawn at random from the third urn. Compute the probability that the ball is:

- (i) Black; (ii) White.
- (iii) Give numerical answers to parts (i) and (ii) for: $m_1 = 10, n_1 = 15;$
 $m_2 = 35, n_2 = 25.$

2.5 Basic Concepts and Results in Counting

In this brief section, some basic concepts and results are discussed regarding the way of counting the total number of outcomes of an experiment or the total number of different ways we can carry out a task. Although many readers will, undoubtedly, be familiar with parts of or the entire material in this section, it would be advisable, nevertheless, to invest some time here in introducing and adopting some notation, establishing some basic results, and then using them in computing probabilities in the classical probability framework.

Problems of counting arise in a great number of different situations. Here are some of them. In each one of these situations, we are asked to compute the number of different ways that something or other can be done. Here are a few illustrative cases.

EXAMPLE 29

- (i) Attire yourself by selecting a T-shirt, a pair of trousers, a pair of shoes, and a cap out of n_1 T-shirts, n_2 pairs of trousers, n_3 pairs of shoes, and n_4 caps (e.g., $n_1 = 4, n_2 = 3, n_3 = n_4 = 2$).
- (ii) Form all k -digit numbers by selecting the k digits out of n available numbers (e.g., $k = 2, n = 4$ such as $\{1, 3, 5, 7\}$).
- (iii) Form all California automobile license plates by using one number, three letters and then three numbers in the prescribed order.
- (iv) Form all possible codes by using a given set of symbols (e.g., form all “words” of length 10 by using the digits 0 and 1).
- (v) Place k books on the shelf of a bookcase in all possible ways.
- (vi) Place the birthdays of k individuals in the 365 days of a year in all possible ways.
- (vii) Place k letters into k addressed envelopes (one letter to each envelope).
- (viii) Count all possible outcomes when tossing k distinct dice.
- (ix) Select k cards out of a standard deck of playing cards (e.g., for $k = 5$, each selection is a poker hand).
- (x) Form all possible k -member committees out of n available individuals.

The calculation of the numbers asked for in situations (i) through (x) just outlined is in actuality a simple application of the so-called *fundamental principle of counting*, stated next in the form of a theorem.

THEOREM 7

(*Fundamental Principle of Counting*) Suppose a task is completed in k stages by carrying out a number of subtasks in each one of the k stages. If the numbers of these subtasks are n_1, \dots, n_k for the k stages, respectively, then the total number of different ways the overall task is completed is: $n_1 \times \dots \times n_k$.

Thus, in (i) above the number of different attires is: $4 \times 3 \times 2 \times 2 = 48$.

In (ii), the number of all 2-digit numbers formed by using 1, 3, 5, 7 is: $4 \times 4 = 16$ (11, 13, 15, 17; 31, 33, 35, 37; 51, 53, 55, 57; 71, 73, 75, 77).

In (iii), the number of all possible license plates (by using indiscriminately all 10 digits from 0 through 9 and all 26 letters of the English alphabet, although this is not the case in practice) is: $10 \times (26 \times 26 \times 26) \times (10 \times 10 \times 10) = 175,760,000$.

In (iv), the number of all possible “words” is found by taking $k = 10$ and $n_1 = \dots = n_{10} = 2$ to obtain: $2^{10} = 1,024$.

In (v), all possible arrangements are obtained by taking $n_1 = k, n_2 = k - 1, \dots, n_k = k - (k - 1) = 1$ to get: $k(k - 1) \dots 1 = 1 \dots (k - 1)k$. For example, for $k = 10$, the number of arrangements is: 3,628,800.

In (vi), the required number is obtained by taking $n_1 = \dots = n_k = 365$ to get: 365^k . For example, for $k = 3$, we have $365^3 = 48,627,125$.

In (vii), the required number is: $k(k - 1) \dots 1 = 1 \dots (k - k)k$ obtained by taking $n_1 = k, n_2 = k - 1, \dots, n_k = k - (k - 1) = 1$.

In (viii), the required number is: 6^k obtained by taking $n_1 = \dots = n_k = 6$. For example, for $k = 3$, we have $6^3 = 216$, and for $k = 10$, we have $6^{10} = 60,466,176$.

In (ix), the number of poker hands is: $\frac{52 \times 51 \times 50 \times 49 \times 48}{120} = 2,598,960$. The numerator is obtained by taking $n_1 = 52, n_2 = 51, n_3 = 50, n_4 = 49, n_5 = 48$. The division by 120 accounts for elimination of hands consisting of the same cards but drawn in different order.

Finally, in (x), the required number is: $\frac{n(n-1)\dots(n-k+1)}{1 \times 2 \times \dots \times k}$, by arguing as in (ix). For example, for $n = 10$ and $k = 3$, we have: $\frac{10 \times 9 \times 8}{1 \times 2 \times 3} = 120$.

In all of the situations (i) through (x), the required numbers were calculated by the appropriate application of Theorem 7. Furthermore, in many cases, as clearly exemplified by cases (ii), (iii), (v), (vii), (ix), and (x), the task performed consisted of selecting and arranging a number of objects out of a set of available objects. In so doing, the order in which the objects appear in the arrangement may be of significance, as is, indeed, the case in situations (ii), (iii), (iv), (v), (vi), and (vii), or it may be just irrelevant, as happens, for example, in cases (ix) and (x). This observation leads us to the concepts of permutations and combinations. More precisely, we have

DEFINITION 4

An *ordered* arrangement of k objects taken from a set of n objects ($1 \leq k \leq n$) is a *permutation* of the n objects taken k at time. An *unordered*

arrangement of k objects taken from a set of n objects is a *combination* of the n objects taken k at a time.

The question then arises of how many permutations and how many combinations can be formed. The answer to this question is given next.

COROLLARY (to Theorem 7)

- (i) The number of ordered arrangements of a set of n objects taken k at a time ($1 \leq k \leq n$) is n^k when *repetitions are allowed*. When *no repetitions are allowed*, this number becomes the *permutations of n objects taken k at a time*, is denoted by $P_{n,k}$, and is given by:

$$P_{n,k} = n(n-1) \dots (n-k+1). \quad (4)$$

In particular, for $k = n$,

$$P_{n,n} = n(n-1) \dots 1 = 1 \dots (n-1)n = n!,$$

where the notation $n!$ is read “ n factorial.”

- (ii) The number of *combinations* (i.e., the number of unordered and without repetition arrangements) of n objects taken k at a time ($1 \leq k \leq n$) is denoted by $\binom{n}{k}$ and is given by:

$$\binom{n}{k} = \frac{P_{n,k}}{k!} = \frac{n!}{k!(n-k)!}. \quad (5)$$

REMARK 5 Whether permutations or combinations are appropriate in a given problem follows from the nature of the problem. For instance, in (ii), permutations rather than combinations are appropriate as, e.g., 13 and 31 are distinct entities. The same is true of cases (iii)–(viii), whereas combinations are appropriate for cases (ix) and (x).

As an example, in part (ii), $P_{4,2} = 4 \times 3 = 12$ (leave out the numbers with identical digits 11, 22, 33, and 44), and in part (ix), $\binom{52}{5} = \frac{52!}{5!47!} = 2,598,960$, after cancellations and by carrying out the arithmetic.

REMARK 6 In (5), set $k = n$. Then the left-hand side is, clearly, 1, and the right-hand side is $\frac{n!}{n!0!} = \frac{1}{0!}$. In order for this to be 1, we *define* $0! = 1$. From formula (5), it also follows that $\binom{n}{0} = 1$.

This section is concluded with the justification of Theorem 7 and its corollary and some applications of these results in calculating certain probabilities.

PROOF OF THEOREM 7 It is done by induction. For $k = 2$, all one has to do is to pair out each one of the n_1 ways of carrying out the subtask at stage 1 with each one of the n_2 ways of carrying out the subtask at stage 2 in order to obtain $n_1 \times n_2$ for the number of ways of completing the task. Next, make

the induction hypothesis that the conclusion is true for $k = m$ and establish it for $k = m + 1$. So, in the first m stages, the total number of ways of doing the job is: $n_1 \times \cdots \times n_m$, and there is still the final $(m + 1)$ st stage for completing the task. Clearly, all we have to do here is to combine each one of the $n_1 \times \cdots \times n_m$ ways of doing the job in the first m stages with each one of the n_{m+1} ways of carrying out the subtask in the $(m + 1)$ st stage to obtain the number $n_1 \times \cdots \times n_m \times n_{m+1}$ of completing the task. ▲

PROOF OF THE COROLLARY

- (i) Here, we are forming an ordered arrangement of objects in k stages by selecting one object at each stage from among the n available objects (because repetitions are allowed). Thus, the theorem applies with $n_1 = \cdots = n_k = n$ and gives the result n^k . When repetitions are not allowed, the only thing which changes from the case just considered is that: $n_1 = n, n_2 = n - 1, \dots, n_k = n - (k - 1) = n - k + 1$, and formula (4) follows.
- (ii) Let $\binom{n}{k}$ be the number of combinations (unordered without repetition arrangements) of the n objects taken k at a time. From each one of these unordered arrangements, we obtain $k!$ ordered arrangements by permutation of the k objects. Then $k! \times \binom{n}{k}$ is the total number of ordered arrangements of the n objects taken k at a time, which is $P_{n,k}$, by part (i). Solving for $\binom{n}{k}$, we obtain the first expression in (5). The second expression follows immediately by multiplying by $(n - k) \dots 1$ and dividing by $1 \dots (n - k) = (n - k)!$ ▲

There are many interesting variations and deeper results based on Theorem 7 and its corollary. Some of them may be found in Sections 2.4 and 2.6 of Chapter 2 of the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G.G. Roussas.

EXAMPLE 30

It happens that 4 hotels in a certain large city have the same name, e.g., Grand Hotel. Four persons make an appointment to meet at the Grand Hotel. If each one of the 4 persons chooses the hotel at random, calculate the following probabilities:

- (i) All 4 choose the same hotel.
 (ii) All 4 choose different hotels.

DISCUSSION

- (i) If $A =$ “all 4 choose the same hotel,” then $P(A) = \frac{n(A)}{n(S)}$, where $n(A)$ is the number of sample points in A . Here, $n(S) = 4 \times 4 \times 4 \times 4 = 4^4$, by Theorem 7 applied with $k = 4$ and $n_1 = n_2 = n_3 = n_4 = 4$, and $n(A) = 4$, by Theorem 7 again applied with $k = 1$ (the 4 people looked upon as a single unity) and $n_1 = 4$ (the 4 hotels they can choose). Thus, $P(A) = \frac{4}{4^4} = \frac{1}{4^3} = \frac{1}{64} = 0.015625 \simeq 0.016$.
- (ii) If $B =$ “all 4 choose different hotels,” then, by the first part of the corollary to Theorem 7, $n(B) = P_{4,4} = 4!$, so that $P(B) = \frac{4!}{4^4} = \frac{1 \times 2 \times 3}{4^3} = \frac{3}{32} = 0.09375 \simeq 0.094$.

EXAMPLE 31

Out of a set of 3 keys, only 1 opens a certain door. Someone tries the keys successively and let A_k be the event that the right key appears the k th time. Calculate the probability $P(A_k)$:

- (i) If the keys tried are not replaced, $k = 1, 2, 3$.
- (ii) If the keys tried are replaced, $k = 1, 2, \dots$

DISCUSSION

- (i) $P(A_1) = \frac{1}{3}$; $P(A_2) = \frac{2 \times 1}{3 \times 2} = \frac{1}{3}$; $P(A_3) = \frac{2 \times 1 \times 1}{3 \times 2 \times 1} = \frac{1}{3}$. So, $P(A_1) = P(A_2) = P(A_3) = \frac{1}{3} \simeq 0.333$.
- (ii) Clearly, $P(A_k) = P(W_1 \cap \dots \cap W_{k-1} \cap R_k) = \left(\frac{2}{3}\right)^{k-1} \times \frac{1}{3}$ for all $k = 1, 2, \dots$

REMARK 7 To calculate the probabilities in part (i) in terms of conditional probabilities, set: R_k = “the right key appears the k th time,” W_k = “a wrong key appears the k th time,” $k = 1, 2, 3$. Then: $P(A_1) = P(R_1) = \frac{1}{3}$, $P(A_2) = P(W_1 \cap R_2) = P(R_2 | W_1)P(W_1) = \frac{1}{2} \times \frac{2}{3} = \frac{1}{3}$, and $P(A_3) = P(W_1 \cap W_2 \cap R_3) = P(R_3 | W_1 \cap W_2)P(W_2 | W_1)P(W_1) = \frac{1}{1} \times \frac{1}{2} \times \frac{2}{3} = \frac{1}{3}$.

EXAMPLE 32

The faculty in an academic department in UC-Davis consists of 4 assistant professors, 6 associate professors, and 5 full professors. Also, it has 30 graduate students. An ad hoc committee of 5 is to be formed to study a certain curricular matter.

- (i) What is the number of all possible committees consisting of faculty alone?
- (ii) How many committees can be formed if 2 graduate students are to be included and all academic ranks are to be represented?
- (iii) If the committee is to be formed at random, what is the probability that the faculty will not be represented?

DISCUSSION It is clear that combinations are the appropriate tool here. Then we have:

- (i) This number is: $\binom{15}{5} = \frac{15!}{5!10!} = \frac{11 \times 12 \times 13 \times 14 \times 15}{1 \times 2 \times 3 \times 4 \times 5} = 3,003$.
- (ii) Here the number is: $\binom{30}{2} \binom{4}{1} \binom{6}{1} \binom{5}{1} = \frac{30!}{2!28!} \times 4 \times 6 \times 5 = \frac{29 \times 30}{2} \times 120 = 52,200$.
- (iii) The required probability is:

$$\frac{\binom{30}{5} \binom{15}{0}}{\binom{45}{5}} = \frac{\binom{30}{5}}{\binom{45}{5}} = \frac{30!/5!25!}{45!/5!40!} = \frac{26 \times 27 \times 28 \times 29 \times 30}{41 \times 42 \times 43 \times 44 \times 45} = \frac{2,262}{19,393} \simeq 0.117.$$

EXAMPLE 33

What is the probability that a poker hand contains 4 pictures, including at least 2 Jacks? It is recalled here that there are 12 pictures consisting of 4 Jacks, 4 Queens, and 4 Kings.

DISCUSSION A poker hand can be selected in $\binom{52}{5}$ ways. The event described, call it A , consists of the following number of sample points: $n(A) = n(J_2) + n(J_3) + n(J_4)$, where J_i = “the poker hand contains exactly i Jacks,”

$i = 2, 3, 4$. But

$$n(J_2) = \binom{4}{2} \binom{8}{2} \binom{40}{1}, \quad n(J_3) = \binom{4}{3} \binom{8}{1} \binom{40}{1}, \quad n(J_4) = \binom{4}{4} \binom{8}{0} \binom{40}{1},$$

so that

$$P(A) = \frac{[\binom{4}{2} \binom{8}{2} + \binom{4}{3} \binom{8}{1} + \binom{4}{4} \binom{8}{0}] \binom{40}{1}}{\binom{52}{5}} = \frac{8,040}{2,598,960} \simeq 0.003.$$

(For the calculation of $\binom{52}{5}$ see Example 29(ix).)

EXAMPLE 34

Each of the $2n$ members of a committee flips a fair coin in deciding whether or not to attend a meeting of the committee; a committee member attends the meeting if an H appears. What is the probability that a majority will show up for the meeting?

DISCUSSION There will be majority if there are at least $n + 1$ committee members present, which amounts to having at least $n + 1$ H 's in $2n$ independent throws of a fair coin. If X is the r.v. denoting the number of H 's in the $2n$ throws, then the required probability is: $P(X \geq n + 1) = \sum_{x=n+1}^{2n} P(X = x)$. However,

$$P(X = x) = \binom{2n}{x} \left(\frac{1}{2}\right)^x \left(\frac{1}{2}\right)^{2n-x} = \frac{1}{2^{2n}} \binom{2n}{x},$$

since there are $\binom{2n}{x}$ ways of having x H 's in $2n$ throws. Therefore

$$\begin{aligned} P(X \geq n + 1) &= \frac{1}{2^{2n}} \sum_{x=n+1}^{2n} \binom{2n}{x} = \frac{1}{2^{2n}} \left[\sum_{x=0}^{2n} \binom{2n}{x} - \sum_{x=0}^n \binom{2n}{x} \right] \\ &= \frac{1}{2^{2n}} \left[2^{2n} - \sum_{x=0}^n \binom{2n}{x} \right] = 1 - \frac{1}{2^{2n}} \sum_{x=0}^n \binom{2n}{x}. \end{aligned}$$

For example, for $2n = 10$, $P(X \geq 6) = 1 - 0.6230 = 0.377$ (from the binomial tables).

Exercises

- 5.1 Telephone numbers at UC-Davis consist of 7-digit numbers the first 3 of which are 752. It is estimated that about 15,000 different telephone numbers are needed to serve the university's needs. Are there enough telephone numbers available for this purpose? Justify your answer.
- 5.2 An experimenter is studying the effects of temperature, pressure, and a catalyst on the yield of a certain chemical reaction. Three different temperatures, four different pressures, and five different catalysts are under consideration.

-
- (i) If any particular experimental run involves the use of a single temperature, pressure, and catalyst, how many experimental runs are possible?
- (ii) How many experimental runs are there that involve use of the lowest temperature and two lowest pressures?
- (iii) How many experimental runs are possible if a specified catalyst is to be used?
- 5.3** (i) Given that a zip code consists of a 5-digit number, where the digits are selected from among the numbers 0, 1, . . . , 9, calculate the number of all different zip codes.
- (ii) If X is the r.v. defined by: $X(\text{zip code}) = \#$ of nonzero digits in the zip code, which are the possible values of X ?
- (iii) Give 3 zip codes and the respective values of X .
- 5.4** How many 5-digit numbers can be formed by using the numbers 1, 2, 3, 4, and 5, so that odd positions are occupied by odd numbers and even positions are occupied by even numbers, if:
- (i) Repetitions are allowed.
- (ii) Repetitions are not allowed.
- 5.5** Form three-digit numbers by using the numbers: 0, 1, 2, 3, 4, 5, 6, 7, 8, and 9, and satisfying one of the following requirements:
- (i) No restrictions are imposed.
- (ii) All three digits are distinct.
- (iii) All three-digit numbers start with 1 and end with 0.
If the three-digit numbers are formed at random, calculate the probability that such a number will be:
- (iv) As described in (ii).
- (v) As described in (iii).
- 5.6** On a straight line, there are n spots to be filled in by either a dot or a dash. What is the number of the distinct groups of resulting symbols? What is this number if $n = 5, 10, 15, 20,$ and 25 ?
- 5.7** A child's set of blocks consists of 2 red, 4 blue, and 5 yellow cubes. The blocks can be distinguished only by color. If the child lines up the blocks in a row at random, calculate the following probabilities:
- (i) Red blocks appear at both ends.
- (ii) All yellow blocks are adjacent.
- (iii) Blue blocks appear at both ends.
- 5.8** Suppose that the letters $C, E, F, F, I,$ and O are written on six chips and placed into a box. Then the six chips are mixed and drawn one by one without replacement. What is the probability that the word "OFFICE" is formed?
- 5.9** For any integers m and n with $0 \leq m \leq n$, show that $\binom{n}{m} = \binom{n}{n-m}$ either by calculation, or by using a suitable argument without writing out anything.
- 5.10** Show that $\binom{n+1}{m+1} / \binom{n}{m} = \frac{n+1}{m+1}$.

5.11 If M , N , and m are positive integers with $m \leq M$, show that:

$$\binom{M}{m} = \binom{M-1}{m} + \binom{M-1}{m-1},$$

by recalling that $\binom{k}{x} = 0$ for $x > k$.

5.12 Without any calculations and by recalling that $\binom{k}{x} = 0$ for $x > k$, show that:

$$\sum_{x=0}^r \binom{m}{x} \binom{n}{r-x} = \binom{m+n}{r}.$$

5.13 The binomial expansion formula states that, for any x and y real and n a positive integer:

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}.$$

Use this formula in order to show that:

$$\sum_{k=0}^n \binom{n}{k} = 2^n \quad \text{and} \quad \sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

5.14 In the plane, there are n points such that no three of them lie on a straight line. How many triangles can be formed? What is this number for $n = 10$?

5.15 Beethoven wrote 9 symphonies, Mozart wrote 27 piano concertos, and Schubert wrote 15 string quartets.

(i) If a university radio station announcer wishes to play first a Beethoven symphony, then a Mozart concerto, and then a Schubert string quartet, in how many ways can this be done?

(ii) What is the number in part (i) if all possible orderings are considered?

5.16 A course in English composition is taken by 10 freshmen, 15 sophomores, 30 juniors, and 5 seniors. If 10 students are chosen at random, calculate the probability that this group will consist of 2 freshman, 3 sophomores, 4 juniors, and 1 senior.

5.17 If n countries exchange ambassadors, how many ambassadors are involved? What is this number for $n = 10, 50, 100$?

5.18 From among n eligible draftees, m are to be drafted in such a way that all possible combinations are equally likely to occur. What is the probability that a specified man is not drafted?

5.19 From 10 positive and 6 negative numbers, 3 numbers are chosen at random and without repetitions. What is the probability that their product is a negative number?

5.20 Two people toss independently n times each a coin whose probability of falling heads is p . What is the probability that they have the same number of heads? What does this probability become for $p = \frac{1}{2}$ and any n ? Also, for $p = \frac{1}{2}$ and $n = 5$?

- 5.21** A shipment of 2,000 light bulbs contains 200 defective items and 1,800 good items. Five hundred bulbs are chosen at random and are tested, and the entire shipment is rejected if more than 25 bulbs from among those tested are found to be defective. What is the probability that the shipment will be accepted?
- 5.22** A student is given a test consisting of 30 questions. For each question, 5 different answers (of which only one is correct) are supplied. The student is required to answer correctly at least 25 questions in order to pass the test. If he/she knows the right answers to the first 20 questions and chooses an answer to the remaining questions at random and independently of each other, what is the probability that the student will pass the test?
- 5.23** Three cards are drawn at random and without replacement from a standard deck of 52 playing cards. Compute the probabilities $P(A_i)$, $i = 1, \dots, 4$, where the events A_i , $i = 1, \dots, 4$ are defined as follows:
 $A_1 =$ "all 3 cards are black," $A_2 =$ "exactly 1 card is an ace,"
 $A_3 =$ "1 card is a diamond, 1 card is a heart, and 1 card is a club."
 $A_4 =$ "at least 2 cards are red."
- 5.24** From an urn containing n_R red balls, n_B black balls, and n_W white balls (all identical except for color) 3 balls are drawn at random. Calculate the following probabilities:
 (i) All 3 balls are red.
 (ii) At least one ball is red.
 (iii) One ball is red, 1 is black, and 1 is white.
 Do this when the balls are drawn:
 (a) Successively and with replacement;
 (b) Without replacement.
- 5.25** A student committee of 12 people is to be formed from among 100 freshmen (40 male + 60 female), 80 sophomores (30 male and 50 female), 70 juniors (24 male and 46 female), and 40 seniors (12 male and 28 female). Calculate the following probabilities:
 (i) Seven students are female and 5 are male.
 (ii) The committee consists of the same number of students from each class.
 (iii) The committee consists of 2 female students and 1 male student from each class.
 (iv) The committee includes at least 1 senior (one of whom will serve as the chairperson of the committee).

The following tabular form of the data facilitates the calculations

Class\Gender	Male	Female	Totals
Freshman	40	60	100
Sophomore	30	50	80
Junior	24	46	70
Senior	12	28	40
Totals	106	184	290



Chapter 3

Numerical Characteristics of a Random Variable, Some Special Random Variables

In this chapter, we discuss the following material. In Section 3.1, the concepts of expectation and variance of a r.v. are introduced and interpretations are provided. Higher order moments are also defined and their significance is pointed out. Also, the moment generating function of a r.v. is defined, and its usefulness as a mathematical tool is commented upon. In Section 3.2, the Markov and Tchebichev inequalities are introduced and their role in estimating probabilities is explained. Section 3.3 is devoted to discussing some of the most commonly occurring distributions: They are the Binomial, Geometric, Poisson, Hypergeometric, Gamma (Negative Exponential and Chi-square), Normal, and Uniform distributions. In all cases, the mathematical expectation, variance, and the moment generating function involved are presented. The chapter is concluded with a discussion of the concepts of median and mode, which are illustrated by concrete examples.

3.1 Expectation, Variance, and Moment Generating Function of a Random Variable

The ideal situation in life would be to know with certainty what is going to happen next. This being almost never the case, the element of chance enters in all aspects of our life. A r.v. is a mathematical formulation of a random environment. Given that we have to deal with a r.v. X , the best thing to expect is to know the values of X and the probabilities with which these values are

taken on, for the case that X is discrete, or the probabilities with which X takes values in various subsets of the real line \Re when X is of the continuous type. That is, we would like to know the *probability distribution* of X . In real life, often, even this is not feasible. Instead, we are forced to settle for some numerical characteristics of the distribution of X . This line of arguments leads us to the concepts of the mathematical expectation and variance of a r.v., as well as to moments of higher order.

DEFINITION 1

Let X be a (discrete) r.v. taking on the values x_i with corresponding probabilities $f(x_i)$, $i = 1, \dots, n$. Then the *mathematical expectation* of X (or just *expectation* or *mean value* of X or just *mean* of X) is denoted by EX and is defined by:

$$EX = \sum_{i=1}^n x_i f(x_i). \quad (1)$$

If the r.v. X takes on (countably) infinite many values x_i with corresponding probabilities $f(x_i)$, $i = 1, 2, \dots$, then the expectation of X is defined by:

$$EX = \sum_{i=1}^{\infty} x_i f(x_i), \quad \text{provided } \sum_{i=1}^{\infty} |x_i| f(x_i) < \infty. \quad (2)$$

Finally, if the r.v. X is continuous with p.d.f. f , its expectation is defined by:

$$EX = \int_{-\infty}^{\infty} x f(x) dx, \quad \text{provided this integral exists.} \quad (3)$$

The alternative notations $\mu(X)$ or μ_X are also often used.

REMARK 1

- (i) The condition $\sum_{i=1}^{\infty} |x_i| f(x_i) < \infty$ is needed because, if it is violated, it is known that $\sum_{i=1}^{\infty} x_i f(x_i)$ may take on different values, depending on the order in which the terms involved are summed up. This, of course, would render the definition of EX meaningless.
- (ii) An example will be presented later on (see Exercise 1.16) where the integral $\int_{-\infty}^{\infty} x f(x) dx = \infty - \infty$, so that it does not exist.

The expectation has several interpretations, some of which are illustrated by the following Examples 1 and 2. One basic interpretation, however, is that of center of gravity. Namely, if one considers the material system where mass $f(x_i)$ is placed at the point x_i , $i = 1, \dots, n$, then EX is the *center of gravity* (point of equilibrium) of this system. In this sense, EX is referred to as a *measure of location* of the distribution of X . The same interpretation holds when X takes on (countably) infinite many values or is of the continuous type.

EXAMPLE 1

Suppose an insurance company pays the amount of \$1,000 for lost luggage on an airplane trip. From past experience, it is known that the company pays this amount in 1 out of 200 policies it sells. What premium should the company charge?

DISCUSSION Define the r.v. X as follows: $X = 0$ if no loss occurs, which happens with probability $1 - (1/200) = 0.995$, and $X = -1,000$ with probability $\frac{1}{200} = 0.005$. Then the expected loss to the company is: $EX = -1,000 \times 0.005 = -5$. Thus, the company must charge \$5 to break even. To this, it will normally add a reasonable amount for administrative expenses and a profit.

Even in this simple example, but most certainly so in more complicated cases, it is convenient to present the values of a (discrete) r.v. and the corresponding probabilities in a tabular form as follows.

x	0	-1,000	Total
$f(x)$	$\frac{199}{200}$	$\frac{1}{200}$	1

EXAMPLE 2

A roulette wheel consists of 18 black slots, 18 red slots, and 2 green slots. If a gambler bets \$10 on red, what is the gambler's expected gain or loss?

DISCUSSION Define the r.v. X by: $X = 10$ with probability $18/38$ and $X = -10$ with probability $20/38$, or in a tabular form

x	10	-10	Total
$f(x)$	$\frac{18}{38}$	$\frac{20}{38}$	1

Then $EX = 10 \times \frac{18}{38} - 10 \times \frac{20}{38} = -\frac{10}{19} \simeq -0.526$. Thus, the gambler's expected loss is about 53 cents.

From the definition of the expectation and familiar properties of summations or integrals, it follows that:

$$E(cX) = cEX, \quad E(cX + d) = cEX + d, \quad \text{where } c \text{ and } d \text{ are constants.} \quad (4)$$

Also (see Exercise 1.18),

$$X \geq c \text{ constant, implies } EX \geq c, \text{ and, in particular, } X \geq 0 \text{ implies } EX \geq 0. \quad (5)$$

Now if Y is a r.v. which is a function of X , $Y = g(X)$, then, in principle, one may be able to determine the p.d.f. of Y and proceed to defining its expectation by the appropriate version of formulas (1), (2), (3). It can be shown, however, that this is not necessary. Instead, the expectation of Y is defined by using the p.d.f. of X , namely:

$$EY = \sum_{i=1}^n g(x_i)f(x_i) \quad \text{or} \quad EY = \sum_{i=1}^{\infty} g(x_i)f(x_i) \quad \text{or} \quad EY = \int_{-\infty}^{\infty} g(x)f(x)dx, \quad (6)$$

under provisions similar to the ones mentioned in connection with (2) and (3). By taking $g(x) = x^k$, where k is a positive integer, we obtain the k th moment of X :

$$EX^k = \sum_{i=1}^n x_i^k f(x_i) \quad \text{or} \quad EX^k = \sum_{i=1}^{\infty} x_i^k f(x_i) \quad \text{or} \quad EX^k = \int_{-\infty}^{\infty} x^k f(x) dx. \quad (7)$$

For $k = 1$, we revert to the expectation of X , and for $k = 2$, we get its *second moment*. Moments are important, among other things, in that, in certain circumstances, a number of them completely determine the distribution of X . This will be illustrated by concrete cases in Section 3.3 (see also Remark 4).

The following simple example illustrates that the expectation, as a measure of location of the distribution, may reveal very little about the entire distribution. Indeed, let the r.v. X take on the values -1 , 1 , and 2 with corresponding probabilities $\frac{5}{8}$, $\frac{1}{8}$, and $\frac{2}{8}$, so that $EX = 0$. Also, let the r.v. Y take on the values -10 , 10 , and 20 with respective probabilities $\frac{5}{8}$, $\frac{1}{8}$, and $\frac{2}{8}$; then again $EY = 0$. The distribution of X is over an interval of length 3, whereas the distribution of Y is over an interval of length 10 times as large. Yet, they have the same center of location. This simple example, clearly, indicates that the expectation by itself is not an adequate measure of description of a distribution, and an additional measure is needed to be associated with the spread of a distribution. Such a measure exists and is the variance of a r.v. or of its distribution.

DEFINITION 2

The *variance* of a r.v. X is denoted by $\text{Var}(X)$ and is defined by:

$$\text{Var}(X) = E(X - EX)^2. \quad (8)$$

The explicit expression of the right-hand side in (8) is taken from (6) for $g(x) = (x - EX)^2$. The alternative notations $\sigma^2(X)$ and σ_X^2 are also often used for the $\text{Var}(X)$.

For the r.v.'s X and Y mentioned before, we have $\text{Var}(X) = 1.75$ and $\text{Var}(Y) = 175$. Thus, the variance does convey adequately the difference in size of the range of the distributions of the r.v.'s X and Y . More generally, for a r.v. X taking on finitely many values x_1, \dots, x_n with respective probabilities $f(x_1), \dots, f(x_n)$, the variance is: $\text{Var}(X) = \sum_{i=1}^n (x_i - EX)^2 f(x_i)$ and represents the sum of the weighted squared distances of the points x_i , $i = 1, \dots, n$ from the center of location of the distribution, EX . Thus, the further from EX the x_i 's are located, the larger the variance, and vice versa. The same interpretation holds for the case that X takes on (countably) infinite many values or is of the continuous type. Because of this characteristic property of the variance, the variance is referred to as a measure of *dispersion* of the underlying distribution. In mechanics, the variance is referred to as the moment of *inertia*.

The positive square root of the $\text{Var}(X)$ is called the *standard deviation* (s.d.) of X . Unlike the variance, the s.d. is measured in the same units

as X (and EX) and serves as a yardstick of measuring deviations of X from EX .

From (8), (6), and familiar properties of summations and integrals, one obtains:

$$\text{Var}(X) = EX^2 - (EX)^2. \quad (9)$$

This formula often facilitates the actual calculation of the variance. From (8), it also follows immediately that

$$\text{Var}(cX) = c^2 \text{Var}(X), \quad \text{Var}(cX + d) = c^2 \text{Var}(X),$$

where c and d are constants. (10)

For a r.v. Y which is a function of X , $Y = g(X)$, the calculation of the $\text{Var}[g(X)]$ reduces to calculating expectations as in (6) because, by means of (8) and (9):

$$\text{Var}[g(X)] = \text{Var}(Y) = E(Y - EY)^2 = EY^2 - (EY)^2 = Eg^2(X) - [Eg(X)]^2. \quad (11)$$

Formulas (8) and (9) are special cases of (11).

In reference to Examples 1 and 2, the variances and the s.d.'s of the r.v.'s. involved are: $\sigma^2(X) = 4,975$, $\sigma(X) \simeq 70.534$, and $\sigma^2(X) = \frac{36,000}{361} \simeq 99.723$, $\sigma(X) \simeq 9.986$, respectively.

EXAMPLE 3

Let X be a r.v. with p.d.f. $f(x) = 3x^2$, $0 < x < 1$. Then:

- (i) Calculate the quantities: EX , EX^2 , and $\text{Var}(X)$.
- (ii) If the r.v. Y is defined by: $Y = 3X - 2$, calculate the EY and the $\text{Var}(Y)$.

DISCUSSION

- (i) By (3), $EX = \int_0^1 x \cdot 3x^2 dx = \frac{3}{4}x^4 \Big|_0^1 = \frac{3}{4} = 0.75$, whereas by (7), applied with $k = 2$, $EX^2 = \int_0^1 x^2 \cdot 3x^2 dx = \frac{3}{5} = 0.60$, so that, by (9), $\text{Var}(X) = 0.60 - (0.75)^2 = 0.0375$.
- (ii) By (4) and (6), $EY = E(3X - 2) = 3EX - 2 = 3 \times 0.75 - 2 = 0.25$, whereas by (10), $\text{Var}(Y) = \text{Var}(3X - 2) = 9 \text{Var}(X) = 9 \times 0.0375 = 0.3375$.

In (6), the EY was defined for $Y = g(X)$, some function of X . In particular, we may take $Y = e^{tX}$ for an arbitrary but fixed $t \in \mathfrak{R}$. Assuming that there exist t 's in \mathfrak{R} for which Ee^{tX} is finite, then this expectation defines a function in t . This function is denoted by $M(t)$ and is called the moment generating function of X . That is,

DEFINITION 3

The function $M(t) = Ee^{tX}$, defined for all those t in \mathfrak{R} for which Ee^{tX} is finite, is called the *moment generating function* (m.g.f.) of X .

Sometimes the notation $M_X(t)$ is also used to emphasize the fact that the m.g.f. under discussion is that of the r.v. X . The m.g.f. of any r.v. always exists

for $t = 0$, since $Ee^{0X} = E1 = 1$; it may exist only for $t = 0$, or for t in a proper subset (interval) in \mathfrak{R} , or for every t in \mathfrak{R} . All these points will be demonstrated by concrete examples later on (see, for example, relations (20), (22), (24), (31), (33), (37), (44), and (46)). The following properties of $M(t)$ follow immediately from its definition.

$$M_{cX}(t) = M_X(ct), \quad M_{cX+d}(t) = e^{dt} M_X(ct), \quad \text{where } c \text{ and } d \text{ are constants.} \quad (12)$$

Indeed,

$$M_{cX}(t) = Ee^{t(cX)} = Ee^{(ct)X} = M_X(ct),$$

and

$$M_{cX+d}(t) = Ee^{t(cX+d)} = E[e^{dt} \cdot e^{(ct)X}] = e^{dt} Ee^{(ct)X} = e^{dt} M_X(ct).$$

Under certain conditions, it is also true that:

$$\left. \frac{d}{dt} M_X(t) \right|_{t=0} = EX \quad \text{and} \quad \left. \frac{d^n}{dt^n} M_X(t) \right|_{t=0} = EX^n, \quad n = 2, 3, \dots \quad (13)$$

For example, for the first property, we have:

$$\begin{aligned} \left. \frac{d}{dt} M_X(t) \right|_{t=0} &= \left. \left(\frac{d}{dt} Ee^{tX} \right) \right|_{t=0} = E \left(\left. \frac{d}{dt} e^{tX} \right|_{t=0} \right) \\ &= E(Xe^{tX}|_{t=0}) = EX. \end{aligned}$$

What is required for this derivation to be legitimate is that the order in which the operators $\frac{d}{dt}$ and E operate on e^{tX} can be interchanged. The justification of the property in (13) for $n \geq 2$ is quite similar. On account of property (13), $M_X(t)$ generates the moments of X through differentiation and evaluation of the derivatives at $t = 0$. It is from this property that the m.g.f. derives its name.

The m.g.f. is also a valuable mathematical tool in many other cases, some of which will be dealt with in subsequent chapters. Presently, it suffices only to state one fundamental property of the m.g.f. in the form of a proposition.

PROPOSITION 1 Under certain conditions, the m.g.f. M_X of a r.v. X uniquely determines the distribution of X .

This proposition is, actually, a rather deep probability result and it is referred to as the *inversion formula*.

Some forms of such a formula for characteristic functions, which are a version of a m.g.f. may be found, e.g., in pages 141–145 in *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G. G. Roussas.

Still another important result associated with m.g.f.'s is stated (but not proved) in the following proposition.

PROPOSITION 2 If for the r.v. X all moments EX^n , $n = 1, 2, \dots$ are finite, then, under certain conditions, these moments uniquely determine the m.g.f. M_X of X , and hence (by Proposition 1) the distribution of X .

Exercise 3.49 provides an example of an application of the proposition just stated.

For Examples 1 and 2, the m.g.f.'s of the r.v.'s involved are: $M_X(t) = 0.995 + 0.005e^{-1,000t}$, $t \in \Re$, and $M_X(t) = \frac{1}{19}(9e^{10t} + 10e^{-10t})$, $t \in \Re$. Then, by differentiation, we get: $\frac{d}{dt}M_X(t)|_{t=0} = -5 = EX$, $\frac{d^2}{dt^2}M_X(t)|_{t=0} = 5,000 = EX^2$, so that $\sigma^2(X) = 4,975$, and $\frac{d}{dt}M_X(t)|_{t=0} = -\frac{10}{19} = EX$, $\frac{d^2}{dt^2}M_X(t)|_{t=0} = 100 = EX^2$, so that $\sigma^2(X) = \frac{36,000}{361} \simeq 99.723$.

EXAMPLE 4

Let X be a r.v. with p.d.f. $f(x) = e^{-x}$, $x > 0$. Then:

- (i) Find the m.g.f. $M_X(t)$ for the t 's for which it is finite.
- (ii) Using M_X , obtain the quantities: EX , EX^2 , and $Var(X)$.
- (iii) If the r.v. Y is defined by: $Y = 2 - 3X$, determine $M_Y(t)$ for the t 's for which it is finite.

DISCUSSION

- (i) By Definition 3,

$$\begin{aligned} M_X(t) &= Ee^{tX} = \int_0^{\infty} e^{tx} \cdot e^{-x} dx = \int_0^{\infty} e^{-(1-t)x} dx \\ &= -\frac{1}{1-t} e^{-(1-t)x} \Big|_0^{\infty} \quad (\text{provided } t \neq 1) \\ &= -\frac{1}{1-t} (0 - 1) = \frac{1}{1-t} \quad (\text{provided } 1-t > 0 \text{ or } t < 1). \end{aligned}$$

Thus, $M_X(t) = \frac{1}{1-t}$, $t < 1$.

- (ii) By (13), $\frac{d}{dt}M_X(t)|_{t=0} = \frac{d}{dt}\left(\frac{1}{1-t}\right)|_{t=0} = \frac{1}{(1-t)^2}|_{t=0} = 1 = EX$, $\frac{d^2}{dt^2}M_X(t)|_{t=0} = \frac{d}{dt}\left(\frac{1}{(1-t)^2}\right)|_{t=0} = \frac{2}{(1-t)^3}|_{t=0} = 2 = EX^2$, so that, by (9), $Var(X) = 2 - 1^2 = 1$.
- (iii) By (12), $M_Y(t) = M_{2-3X}(t) = M_{-3X+2}(t) = e^{2t}M_X(-3t) = e^{2t}\frac{1}{1-(-3t)} = \frac{e^{2t}}{1+3t}$, provided $t > -\frac{1}{3}$.

Exercises

In several calculations required in solving some exercises in this section, the following formulas prove very useful.

First, in summing the infinite terms of a geometric series, we have:

$$\sum_{x=k}^{\infty} t^x = \frac{t^k}{1-t}, \quad k = 0, 1, \dots, \text{ and } |t| < 1.$$

Next,

$$\begin{aligned} \sum_{x=1}^{\infty} xt^x &= t \sum_{x=1}^{\infty} xt^{x-1} = t \sum_{x=1}^{\infty} \frac{d}{dt} t^x = t \frac{d}{dt} \sum_{x=1}^{\infty} t^x \\ &= t \frac{d}{dt} \left(\frac{t}{1-t} \right) = \frac{t}{(1-t)^2}. \end{aligned}$$

Also,

$$\begin{aligned}\sum_{x=2}^{\infty} x(x-1)t^x &= t^2 \sum_{x=2}^{\infty} x(x-1)t^{x-2} = t^2 \sum_{x=2}^{\infty} \frac{d^2}{dt^2} t^x \\ &= t^2 \frac{d^2}{dt^2} \sum_{x=2}^{\infty} t^x = t^2 \frac{d^2}{dt^2} \left(\frac{t^2}{1-t} \right) = \frac{2t^2}{(1-t)^3}.\end{aligned}$$

In the last two formulas, if there is a number instead of the variable t , the number is replaced by t , and in the final formulas t is replaced by the number.

- 1.1** Refer to Exercise 2.1 in Chapter 2 and calculate the quantities: EX , $Var(X)$, and the s.d. of X .
- 1.2** For the r.v. X for which $P(X = -c) = P(X = c) = 1/2$ (for some $c > 0$):
- Calculate the EX and the $Var(X)$.
 - Show that $P(|X - EX| \leq c) = Var(X)/c^2$.
- 1.3** A chemical company currently has in stock 100 lb of a certain chemical, which it sells to customers in 5 lb packages. Let X be the r.v. denoting the number of packages ordered by a randomly chosen customer, and suppose that the p.d.f. of X is given by: $f(1) = 0.2$, $f(2) = 0.4$, $f(3) = 0.3$, $f(4) = 0.1$.

x	1	2	3	4
$f(x)$	0.2	0.4	0.3	0.1

- Compute the following quantities: EX , EX^2 , and $Var(X)$.
 - Compute the expected number of pounds left after the order of the customer in question has been shipped, as well as the s.d. of the number of pounds around the expected value.
- 1.4** Let X be a r.v. denoting the damage incurred (in \$) in a certain type of accident during a given year, and suppose that the distribution of X is given by the following table:

x	0	1,000	5,000	10,000
$f(x)$	0.8	0.1	0.08	0.02

A particular company offers a \$500 deductible policy. If the company's expected profit on a given policy is \$100, what premium amount should it charge?

- 1.5** Let X be the r.v. denoting the number in the uppermost side of a fair die when rolled once.
- Determine the m.g.f. of X .
 - Use the m.g.f. to calculate: EX , EX^2 , $Var(X)$, and the s.d. of X .
- 1.6** For any r.v. X , for which the EX and the EX^2 are finite, show that:

$$Var(X) = EX^2 - (EX)^2 = E[X(X-1)] + EX - (EX)^2.$$

1.7 Suppose that for a r.v. X it is given that: $EX = 5$ and $E[X(X - 1)] = 27.5$.

Calculate:

- (i) EX^2 .
- (ii) $Var(X)$ and s.d. of X .

1.8 For the r.v. X with p.d.f. $f(x) = (1/2)^x$, $x = 1, 2, \dots$:

- (i) Calculate the EX and the $E[X(X - 1)]$.
- (ii) Use part (i) and Exercise 1.6 to compute the $Var(X)$.

1.9 The p.d.f. f of a r.v. X is given by: $f(x) = c(1/3)^x$, for $x = 0, 1, \dots$ (c some positive constant).

- (i) Calculate the EX .
- (ii) Determine the m.g.f. M_X of X and specify the range of its argument.
- (iii) Employ the m.g.f. in order to derive the EX .

1.10 For the r.v. X with p.d.f. $f(x) = 0.5x$, for $0 \leq x \leq 2$, calculate: EX , $Var(X)$, and the s.d. of X .

1.11 If the r.v. X has p.d.f. $f(x) = 3x^2 - 2x + 1$, for $0 < x < 1$, compute the expectation and variance of X .

1.12 If the r.v. X has p.d.f. f given by:

$$f(x) = \begin{cases} c_1x, & -2 < x < 0 \\ c_2x, & 0 \leq x < 1 \\ 0, & \text{otherwise,} \end{cases}$$

and if we suppose that $EX = \frac{1}{3}$, determine the constants c_1 and c_2 .

1.13 The lifetime in hours of electric tubes is a r.v. X with p.d.f. f given by: $f(x) = \lambda^2 x e^{-\lambda x}$, for $x > 0$ ($\lambda > 0$). Calculate the expected life of such tubes.

1.14 Let X be a r.v. whose $EX = \mu \in \mathfrak{R}$. Then:

- (i) For any constant c , show that:

$$E(X - c)^2 = E(X - \mu)^2 + (\mu - c)^2 = Var(X) + (\mu - c)^2.$$

- (ii) Use part (i) to show that $E(X - c)^2$, as a function of c , is minimized for $c = \mu$.

1.15 Let X be a r.v. with p.d.f. $f(x) = \frac{|x|}{c^2}$, for $-c < x < c$, $c > 0$. For any $n = 1, 2, \dots$, calculate the EX^n , and as a special case, derive the EX and the $Var(X)$.

1.16 Let X be a r.v. with p.d.f. given by: $f(x) = \frac{1}{\pi} \cdot \frac{1}{1+x^2}$, $x \in \mathfrak{R}$. Show that:

- (i) f is, indeed, a p.d.f. (called the *Cauchy* p.d.f.).
- (ii) $\int_{-\infty}^{\infty} x f(x) dx = \infty - \infty$, so that the EX does not exist.

1.17 If X is a r.v. for which all moments EX^n , $n = 0, 1, \dots$ are finite, show that

$$M_X(t) = \sum_{n=0}^{\infty} (EX^n) \frac{t^n}{n!}.$$

Hint: Use the expansion $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$.

Remark: The result in this exercise says that the moments of a r.v. determine (under certain conditions) the m.g.f. of the r.v., and hence its distribution.

1.18 Establish the inequalities stated in relation (5) for both the discrete and the continuous case.

3.2 Some Probability Inequalities

If the r.v. X has a known p.d.f. f , then, in principle, we can calculate probabilities $P(X \in B)$ for $B \subseteq \mathfrak{R}$. This, however, is easier said than done in practice. What one would be willing to settle for would be some suitable and computable bounds for such probabilities. This line of thought leads us to the inequalities discussed here.

THEOREM 1

- (i) For any nonnegative r.v. X and for any constant $c > 0$, it holds:

$$P(X \geq c) \leq EX/c.$$

- (ii) More generally, for any nonnegative function of any r.v. X , $g(X)$, and for any constant $c > 0$, it holds:

$$P[g(X) \geq c] \leq Eg(X)/c. \quad (14)$$

- (iii) By taking $g(X) = |X - EX|$ in part (ii), the inequality reduces to the *Markov* inequality, namely,

$$P(|X - EX| \geq c) = P(|X - EX|^r \geq c^r) \leq E|X - EX|^r / c^r, \quad r > 0. \quad (15)$$

- (iv) In particular, for $r = 2$ in (15), we get the *Tchebichev* inequality, namely,

$$P(|X - EX| \geq c) \leq \frac{E(X - EX)^2}{c^2} = \frac{\sigma^2}{c^2} \quad \text{or} \\ P(|X - EX| < c) \geq 1 - \frac{\sigma^2}{c^2}, \quad (16)$$

where σ^2 stands for the $Var(X)$. Furthermore, if $c = k\sigma$, where σ is the s.d. of X , then:

$$P(|X - EX| \geq k\sigma) \leq \frac{1}{k^2} \quad \text{or} \quad P(|X - EX| < k\sigma) \geq 1 - \frac{1}{k^2}. \quad (17)$$

REMARK 2 From the last expression, it follows that X lies within k s.d.'s from its mean with probability at least $1 - \frac{1}{k^2}$, regardless of the distribution of X . It is in this sense that the s.d. is used as a yardstick of deviations of X from its mean, as already mentioned elsewhere.

Thus, for example, for $k = 2, 3$, we obtain, respectively:

$$P(|X - EX| < 2\sigma) \geq 0.75, \quad P(|X - EX| < 3\sigma) \geq \frac{8}{9} \simeq 0.889. \quad (18)$$

PROOF OF THEOREM 1 Clearly, all one has to do is to justify (14) and this only for the case that X is continuous with p.d.f. f , because the discrete case is entirely analogous.

Indeed, let $A = \{x \in \mathfrak{R}; g(x) \geq c\}$, so that $A^c = \{x \in \mathfrak{R}; g(x) < c\}$. Then, clearly:

$$\begin{aligned} Eg(X) &= \int_{-\infty}^{\infty} g(x) f(x) dx = \int_A g(x) f(x) dx + \int_{A^c} g(x) f(x) dx \\ &\geq \int_A g(x) f(x) dx \quad (\text{since } g(x) \geq 0) \\ &\geq c \int_A f(x) dx \quad (\text{since } g(x) \geq c \text{ on } A) \\ &= cP(A) = cP[g(X) \geq c]. \end{aligned}$$

Solving for $P[g(X) \geq c]$, we obtain the desired result. \blacktriangle

EXAMPLE 5

Let the r.v. X take on the values $-2, -1/2, 1/2$, and 2 with respective probabilities $0.05, 0.45, 0.45$, and 0.05 . Then $EX = 0$ and $\sigma^2 = \text{Var}(X) = 0.625$, so that $2\sigma \simeq 1.582$. Then: $P(|X| < 2\sigma) = P(-1.582 < X < 1.582) = P(X = -\frac{1}{2}) + P(X = \frac{1}{2}) = 0.90$, compared with the lower bound of 0.75 .

EXAMPLE 6

Let the r.v. X take on the value x with probability $f(x) = e^{-\lambda} \frac{\lambda^x}{x!}$, $x = 0, 1, \dots$, some $\lambda > 0$. As will be seen later on (see relation (23)), this is a Poisson r.v. with parameter λ , and $EX = \text{Var}(X) = \lambda$. For selected values of λ , probabilities $P(X \leq k)$ are given by the Poisson tables. For illustrative purposes, let $\lambda = 9$. Then $\sigma = 3$ and therefore: $P(|X - 9| < 2 \times 3) = P(3 < X < 15) = 0.9373$, compared with 0.75 , and $P(|X - 9| < 3 \times 3) = P(0 < X < 18) = 0.9946$, compared with 0.889 .

Exercises

2.1 Suppose the distribution of the r.v. X is given by the following table:

x	-1	0	1
$f(x)$	$1/18$	$8/9$	$1/18$

- (i) Calculate the EX (call it μ), the $\text{Var}(X)$, and the s.d. of X (call it σ).
- (ii) Compute the probability: $P(|X - \mu| \geq k\sigma)$ for $k = 2, 3$.
- (iii) By the Tchebichev inequality: $P(|X - \mu| \geq k\sigma) \leq 1/k^2$. Compare the exact probabilities computed in part (ii) with the respective upper bounds.

2.2 If X is a r.v. with expectation μ and s.d. σ , use the Tchebichev inequality:

(i) To determine c in terms of σ and α , so that:

$$P(|X - \mu| < c) \geq \alpha \quad (0 < \alpha < 1).$$

(ii) Give the numerical value of c for $\sigma = 1$ and $\alpha = 0.95$.

2.3 Let X be a r.v. with p.d.f. $f(x) = c(1 - x^2)$, for $-1 \leq x \leq 1$. Refer to Exercise 2.11(i) in Chapter 2 for the determination of the constant c and then:

(i) Calculate the EX and $Var(X)$.

(ii) Use the Tchebichev inequality to find a lower bound for the probability $P(-0.9 < X < 0.9)$, and compare it with the exact probability calculated in Exercise 2.11(ii) in Chapter 2.

2.4 Let X be a r.v. with (finite) mean μ and variance 0. Then:

(i) Use the Tchebichev inequality to show that $P(|X - \mu| \geq c) = 0$ for all $c > 0$.

(ii) Use part (i) and Theorem 2 in Chapter 2 in order to conclude that $P(X = \mu) = 1$.

3.3 Some Special Random Variables

3.3.1 The Discrete Case

In this section, we discuss seven distributions—four discrete and three of the continuous type, which occur often. These are the Binomial, the Geometric, the Poisson, the Hypergeometric, the Gamma (which includes the Negative Exponential and the Chi-Square), the Normal, and the Uniform distributions. At this point, it should be mentioned that a p.d.f. is 0 for all the values of its argument not figuring in its definition.

Binomial Distribution We first introduced the concept of a *binomial experiment*, which is meant to be an experiment resulting in two possible outcomes, one termed a *success*, denoted by S and occurring with probability p , and the other termed a *failure*, denoted by F and occurring with probability $q = 1 - p$. A binomial experiment is performed n independent times (with p remaining the same), and let X be the r.v. denoting the number of successes. Then, clearly, X takes on the values $0, 1, \dots, n$, with the respective probabilities:

$$P(X = x) = f(x) = \binom{n}{x} p^x q^{n-x}, \quad x = 0, 1, \dots, n, \quad 0 < p < 1, \quad q = 1 - p. \quad (19)$$

The r.v. X is said to be *Binomially* distributed, its distribution is called *Binomial* with *parameters* n and p , and the fact that X is so distributed is denoted


by $X \sim B(n, p)$. The graph of f depends on n and p ; two typical cases, for $n = 12$, $p = \frac{1}{4}$, and $n = 10$, $p = \frac{1}{2}$ are given in Figures 3.1 and 3.2.

Values of the p.d.f. f of the $B(12, \frac{1}{4})$ distribution

$f(0) = 0.0317$	$f(7) = 0.0115$
$f(1) = 0.1267$	$f(8) = 0.0024$
$f(2) = 0.2323$	$f(9) = 0.0004$
$f(3) = 0.2581$	$f(10) = 0.0000$
$f(4) = 0.1936$	$f(11) = 0.0000$
$f(5) = 0.1032$	$f(12) = 0.0000$
$f(6) = 0.0401$	

Figure 3.1

Graph of the p.d.f. of the Binomial Distribution for $n = 12$, $p = \frac{1}{4}$




Values of the p.d.f. f of the $B(10, \frac{1}{2})$ distribution

$f(0) = 0.0010$	$f(6) = 0.2051$
$f(1) = 0.0097$	$f(7) = 0.1172$
$f(2) = 0.0440$	$f(8) = 0.0440$
$f(3) = 0.1172$	$f(9) = 0.0097$
$f(4) = 0.2051$	$f(10) = 0.0010$
$f(5) = 0.2460$	

For selected n and p , the d.f. $F(k) = \sum_{j=0}^k \binom{n}{j} p^j q^{n-j}$ is given by tables, the Binomial tables (see, however, Exercise 3.1). The individual probabilities $\binom{n}{j} p^j q^{n-j}$ may be found by subtraction. Alternatively, such probabilities can be calculated recursively (see Exercise 3.9).

Figure 3.2

Graph of the p.d.f. of the Binomial Distribution for $n = 10$, $p = \frac{1}{2}$



For $n = 1$, the corresponding r.v. is known as the *Bernoulli* r.v. It is then clear that a $B(n, p)$ r.v. X is the sum of $n B(1, p)$ r.v.'s. More precisely, in n independent binomial experiments, associate with the i th performance of the experiment the r.v. X_i defined by: $X_i = 1$ if the outcome is S (a success) and $X_i = 0$ otherwise, $i = 1, \dots, n$. Then, clearly, $\sum_{i=1}^n X_i$ is the number of 1's in the n trials, or, equivalently, the number of S 's, which is exactly what the r.v. X stands for. Thus, $X = \sum_{i=1}^n X_i$. Finally, it is mentioned here that, if $X \sim B(n, p)$, then:

$$EX = np, \quad \text{Var}(X) = npq, \quad \text{and} \quad M_X(t) = (pe^t + q)^n, \quad t \in \Re. \quad (20)$$

The relevant derivations are left as exercises (see Exercises 3.10 and 3.11). A brief justification of formula (19) is as follows: Think of the n outcomes of the n experiments as n points on a straight line segment, where an S or an F , is to be placed. By independence, the probability that there will be exactly x S 's in x specified positions (and therefore $n - x$ F 's in the remaining positions) is $p^x q^{n-x}$, and this probability is independent of the locations where the x S 's occur. Because there are $\binom{n}{x}$ ways of selected x points for the S 's, the conclusion follows.

Finally, for illustrative purposes, refer to Example 7 in Chapter 1. In that example, clearly $X \sim B(n, 0.8)$, and for the sake of specificity take $n = 25$, so that X takes on the values $0, 1, \dots, 25$. Next (see Exercise 3.1), $\binom{25}{x}(0.8)^x (0.2)^{25-x} = \binom{25}{y}(0.2)^y(0.8)^{25-y}$, where $y = 25 - x$. Therefore, for $a = 15$ and $b = 20$, for example, $P(15 \leq X \leq 25) = \sum_{y=5}^{10} \binom{25}{y}(0.2)^y(0.8)^{25-y} = 0.994 - 0.421 = 0.573$. Finally, $EX = 25 \times 0.8 = 20$, $\text{Var}(X) = 25 \times 0.8 \times 0.2 = 4$, so that $\sigma(X) = 2$. Examples 8–10 in Chapter 1 fit into the same framework.

Geometric Distribution This distribution arises in a binomial experiment situation when trials are carried out independently (with constant probability p of an S) until the *first* S occurs. The r.v. X denoting the number of required trials is a *Geometrically* distributed r.v. with *parameter* p and its distribution

is the *Geometric* distribution with *parameter* p . It is clear that X takes on the values $1, 2, \dots$ with the respective probabilities:

$$P(X = x) = f(x) = pq^{x-1}, \quad x = 1, 2, \dots, \quad 0 < p < 1, \quad q = 1 - p. \quad (21)$$


The justification of this formula is immediate because, if the first S is to appear in the x th position, the overall outcome is $\underbrace{FF \dots FS}_{x-1}$ whose probability (by independence) is $q^{x-1}p$.

The graph of f depends on p ; two typical cases for $p = \frac{1}{4}$ and $p = \frac{1}{2}$ are given in Figure 3.3.

Values of $f(x) = (0.25)(0.75)^{x-1}$, $x = 1, 2, \dots$	Values of $f(x) = (0.5)^x$, $x = 1, 2, \dots$
$f(1) = 0.2500$	$f(1) = 0.5000$
$f(2) = 0.1875$	$f(2) = 0.2500$
$f(3) = 0.1406$	$f(3) = 0.1250$
$f(4) = 0.1055$	$f(4) = 0.0625$
$f(5) = 0.0791$	$f(5) = 0.0313$
$f(6) = 0.0593$	$f(6) = 0.0156$
$f(7) = 0.0445$	$f(7) = 0.0078$
$f(8) = 0.0334$	
$f(9) = 0.0250$	
$f(10) = 0.0188$	

Figure 3.3

**Graphs of the p.d.f.'s
of the Geometric
Distribution with
 $p = \frac{1}{4}$ and $p = \frac{1}{2}$**



If the r.v. X is Geometrically distributed with parameter p , then:

$$EX = \frac{1}{p}, \quad \text{Var}(X) = \frac{q}{p^2}, \quad M_X(t) = \frac{pe^t}{1 - qe^t}, \quad t < -\log q. \quad (22)$$

REMARK 3 Sometimes the p.d.f. of X is given in the form: $f(x) = pq^x$, $x = 0, 1, \dots$; then $EX = \frac{q}{p}$, $Var(X) = \frac{p}{q^2}$ and $M_X(t) = \frac{p}{1-qe^t}$, $t < -\log q$.

In reference to Example 11 in Chapter 1, assume for mathematical convenience that the number of cars passing by may be infinite. Then the r.v. X described there has the Geometric distribution with some p . Here probabilities are easily calculated. For example, $P(X \geq 20) = \sum_{x=20}^{\infty} pq^{x-1} = pq^{19} \sum_{x=0}^{\infty} q^x = pq^{19} \frac{1}{1-q} = q^{19}$; i.e., $p(X \geq 20) = q^{19}$. For instance, if $p = 0.01$, then $q = 0.99$ and $P(X \geq 20) \simeq 0.826$.

Poisson Distribution A r.v. X taking on the values $0, 1, \dots$ with respective probabilities given in (23) is said to have the *Poisson distribution* with parameter λ ; its distribution is called the *Poisson distribution* with parameter λ . That X is Poisson distributed with parameter λ is denoted by $X \sim P(\lambda)$.


$$P(X = x) = f(x) = e^{-\lambda} \frac{\lambda^x}{x!}, \quad x = 0, 1, \dots, \lambda > 0. \quad (23)$$

The graph of f depends on λ ; for example, for $\lambda = 5$, it looks like that in Figure 3.4. That f is a p.d.f. follows from the formula $\sum_{x=0}^{\infty} \frac{\lambda^x}{x!} = e^\lambda$.

Values of the p.d.f. f of the $P(5)$ distribution

$f(0) = 0.0067$	$f(9) = 0.0363$
$f(1) = 0.0337$	$f(10) = 0.0181$
$f(2) = 0.0843$	$f(11) = 0.0082$
$f(3) = 0.1403$	$f(12) = 0.0035$
$f(4) = 0.1755$	$f(13) = 0.0013$
$f(5) = 0.1755$	$f(14) = 0.0005$
$f(6) = 0.1462$	$f(15) = 0.0001$
$f(7) = 0.1044$	
$f(8) = 0.0653$	$f(n)$ is negligible for $n \geq 16$.

Figure 3.4
Graph of the p.d.f. of the Poisson Distribution with $\lambda = 5$



For selected values of λ , the d.f. $F(k) = \sum_{j=0}^k e^{-\lambda} \frac{\lambda^j}{j!}$ is given by tables, the Poisson tables. The individual values $e^{-\lambda} \frac{\lambda^j}{j!}$ are found by subtraction. Alternatively, such probabilities can be calculated recursively (see Exercise 3.20). It is not hard to see (see Exercises 3.21 and 3.22) that, if $X \sim P(\lambda)$, then:

$$EX = \lambda, \quad \text{Var}(X) = \lambda, \quad \text{and} \quad M_X(t) = e^{\lambda e^t - \lambda}, \quad t \in \mathfrak{R}. \quad (24)$$

From these expressions, the parameter λ acquires a special meaning: it is both the mean and the variance of the r.v. X .

Example 12 in Chapter 1 may serve as an illustration of the usage of the Poisson distribution. Assuming, for mathematical convenience, that the number of bacteria may be infinite, then the Poisson distribution may be used to describe the actual distribution of bacteria (for a suitable value of λ) quite accurately. There is a host of similar cases for the description of which the Poisson distribution is appropriate. These include the number of telephone calls served by a certain telephone exchange center within a certain period of time, the number of particles emitted by a radioactive source within a certain period of time, the number of typographical errors in a book, etc.

There is an intimate relationship between the Poisson and the Binomial distributions: the former may be obtained as the limit of the latter, as explained in the following. Namely, it is seen (see Exercise 3.23) that in the Binomial, $B(n, p)$, situation, if n is large and p is small, then the Binomial probabilities $\binom{n}{x} p^x (1-p)^{n-x}$ are close to the Poisson probabilities $e^{-np} \frac{(np)^x}{x!}$. More precisely, $\binom{n}{x} p_n^x (1-p_n)^{n-x} \rightarrow e^{-\lambda} \frac{\lambda^x}{x!}$, provided $n \rightarrow \infty$ and $p_n \rightarrow 0$ so that $np_n \rightarrow \lambda \in (0, \infty)$. Here p_n is the probability of a success in the n th trial. Thus, for large values of n , $\binom{n}{x} p_n^x (1-p_n)^{n-x} \simeq e^{-\lambda} \frac{\lambda^x}{x!}$; or, upon replacing λ by np_n , we obtain the approximation mentioned before. A rough explanation of why Poisson probabilities are approximated by Binomial probabilities is given next. To this end, suppose an event A occurs once in a small time interval h with approximate probability proportional to h and coefficient of proportionality λ ; i.e., A occurs once in h with approximate probability λh . It occurs two or more times with probability approximately 0, so that it occurs zero times with probability approximately $1 - \lambda h$. Finally, occurrences in nonoverlapping intervals of length h are independent. Next, consider the unit interval $(0, 1]$ and divide it into a large number n of nonoverlapping subintervals of equal length h : $(t_{i-1}, t_i]$, $i = 1, \dots, n$, $t_0 = 0$, $t_n = 1$, $h = \frac{1}{n}$. With the i th interval $(t_{i-1}, t_i]$, associate the r.v. X_i defined by: $X_i = 1$ with approximate probability λh and 0 with approximate probability $1 - \lambda h$. Then the r.v. $X = \sum_{i=1}^n X_i$ denotes the number of occurrences of A over the unit $(0, 1]$ interval with approximate probabilities $\binom{n}{x} (\lambda h)^x (1 - \lambda h)^{n-x}$. The exact probabilities are found by letting $n \rightarrow \infty$ (which implies $h \rightarrow 0$). Because here $p_n = \lambda h$ and $np_n = n\lambda h = n\lambda \frac{1}{n} = \lambda$, we have that $\binom{n}{x} (\lambda h)^x (1 - \lambda h)^{n-x} \rightarrow e^{-\lambda} \frac{\lambda^x}{x!}$, as $n \rightarrow \infty$ (by Exercise 3.23), so that the exact probabilities are $e^{-\lambda} \frac{\lambda^x}{x!}$. So, the exact probability that A occurs x times in $(0, 1]$ is the Poisson probability $e^{-\lambda} \frac{\lambda^x}{x!}$, and the approximate probability

that A occurs the same number of times is the Binomial probability $\binom{n}{x}(\lambda h)^x(1 - \lambda h)^{n-x}$; these two probabilities are close to each other for large n .

The following example sheds some light on the approximation just discussed.

EXAMPLE 7

If X is a r.v. distributed as $B(25, \frac{1}{16})$, we find from the Binomial tables that $P(X = 2) = 0.2836$. Next, considering a r.v. Y distributed as $P(\lambda)$ with $\lambda = \frac{25}{16} = 1.5625$, we have that $P(Y = 2) = e^{-1.5625} \frac{(1.5625)^2}{2!} \simeq 0.2556$. Thus, the exact probability is underestimated by the amount 0.028. The error committed is of the order of 9.87%. Given the small value of $n = 25$, the approximate value is not bad at all.

Hypergeometric Distribution This distribution occurs quite often and is suitable in describing situations of the following type: m identical objects (e.g., balls) are thoroughly mixed with n identical objects (which again can be thought of as being balls) but distinct from the m objects. From these $m + n$ objects, r are drawn *without replacement*, and let X be the number among the r which come from the m objects. Then the r.v. X takes on the values $0, 1, \dots, \min(r, m)$ with respective probabilities given below. Actually, by defining $\binom{m}{x} = 0$ for $x > m$, we have:

$$P(X = x) = f(x) = \frac{\binom{m}{x} \binom{n}{r-x}}{\binom{m+n}{r}}, \quad x = 0, \dots, r; \quad (25)$$

m and n may be referred to as the *parameters* of the distribution. By assuming that the selections of r objects out of the $m + n$ are all equally likely, there are $\binom{m+n}{r}$ ways of selecting these r objects, whereas there are $\binom{m}{x}$ ways of selecting x out of the m objects, and $\binom{n}{r-x}$ ways of selecting the remaining $r - x$ objects out of n objects. Thus, the probability that $X = x$ is as given in the preceding formula. The simple justification that these probabilities actually sum to 1 follows from Exercise 5.12 in Chapter 2. For large values of any one of m , n , and r , actual calculation of the probabilities in (25) may be quite involved. A recursion formula (see Exercise 3.26) facilitates significantly these calculations. The calculation of the expectation and of the variance of X is based on the same ideas as those used in Exercise 3.10 in calculating the EX and $Var(X)$ when $X \sim B(n, p)$. We omit the details and give the relevant formulas, namely,

$$EX = \frac{mr}{m+n}, \quad Var(X) = \frac{mnr(m+n-r)}{(m+n)^2(m+n-1)}.$$

Finally, by utilizing ideas and arguments similar to those employed in Exercise 3.23, it is shown that as m and $n \rightarrow \infty$ so that $\frac{m}{m+n} \rightarrow p \in (0, 1)$, then $\binom{m}{x} \binom{n}{r-x} / \binom{m+n}{r}$ tends to $\binom{r}{x} p^x (1-p)^{r-x}$. Thus, for large values of m and n , the Hypergeometric probabilities $\binom{m}{x} \binom{n}{r-x} / \binom{m+n}{r}$ may be approximated by the simpler Binomial probabilities $\binom{r}{x} p_{m,n}^x (1-p_{m,n})^{r-x}$, where $p_{m,n} = \frac{m}{m+n}$.

EXAMPLE 8

As an application of formula (25) and the approximation discussed, take $m = 70$, $n = 90$, $r = 25$ and $x = 10$. Then:

$$f(10) = \frac{\binom{70}{10} \binom{90}{25-10}}{\binom{70+90}{25}} = \frac{\binom{70}{10} \binom{90}{15}}{\binom{160}{25}} \simeq 0.166,$$

after quite a few calculations. On the other hand, since $\frac{m}{m+n} = \frac{70}{160} = \frac{7}{16}$, the Binomial tables give for the $B(25, \frac{7}{16})$ distribution: $\binom{25}{10} (\frac{7}{16})^{10} (\frac{9}{16})^{15} = 0.15$. Therefore, the approximation overestimates the exact probability by the amount 0.016. The error committed is of the order of 10.7%.

3.3.2 The Continuous Case

Gamma Distribution For its introduction, a certain function, the so-called Gamma function, is to be defined first. It is shown that the integral $\int_0^{\infty} y^{\alpha-1} e^{-y} dy$ is finite for $\alpha > 0$ and is thus defining a function (in α), namely,

$$\Gamma(\alpha) = \int_0^{\infty} y^{\alpha-1} e^{-y} dy, \quad \alpha > 0. \quad (26)$$

This is the *Gamma function*. By means of the Gamma function, the *Gamma distribution* is defined as follows:

$$f(x) = \frac{1}{\Gamma(\alpha)\beta^{\alpha}} x^{\alpha-1} e^{-x/\beta}, \quad x > 0, \quad \alpha > 0, \quad \beta > 0; \quad (27)$$

α and β are the *parameters* of the distribution. That the function f integrates to 1 is an immediate consequence of the definition of $\Gamma(\alpha)$. A r.v. X taking on values in \mathfrak{R} and having p.d.f. f , given in (27), is said to be *Gamma distributed* with *parameters* α and β ; one may choose the notation $X \sim \Gamma(\alpha, \beta)$ to express this fact. The graph of f depends on α and β but is, of course, always concentrated on $(0, \infty)$. Typical cases for several values of the pair (α, β) are given in Figures 3.5 and 3.6.

The Gamma distribution is suitable for describing waiting times between successive occurrences of a random event and is also used for describing survival times. In both cases, it provides great flexibility through its two parameters α and β . For specific values of the pair (α, β) , we obtain the Negative

Figure 3.5

Graphs of the p.d.f. of the Gamma Distribution for Several Values of α, β





Figure 3.6

Graphs of the p.d.f. of the Gamma Distribution for Several Values of α, β



Exponential and Chi-Square distributions to be studied subsequently. By integration by parts, one may derive the following useful recursive relation for the Gamma function (see Exercise 3.27):

$$\Gamma(\alpha) = (\alpha - 1)\Gamma(\alpha - 1). \quad (28)$$

In particular, if α is an integer, repeated applications of recursive relation (28) produce

$$\Gamma(\alpha) = (\alpha - 1)(\alpha - 2) \dots \Gamma(1).$$

But $\Gamma(1) = \int_0^\infty e^{-y} dy = 1$, so that

$$\Gamma(\alpha) = (\alpha - 1)(\alpha - 2) \dots 1 = (\alpha - 1)! \quad (29)$$

For later reference, we mention here (see also Exercise 3.45) that, by integration, we obtain:

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}, \quad (30)$$

and then, by means of this and the recursive formula (28), we can calculate $\Gamma(\frac{3}{2})$, $\Gamma(\frac{5}{2})$, etc. Finally, by integration (see Exercises 3.28 and 3.29), it is seen that:

$$EX = \alpha\beta, \quad \text{Var}(X) = \alpha\beta^2 \quad \text{and} \quad M_X(t) = \frac{1}{(1 - \beta t)^\alpha}, \quad t < \frac{1}{\beta}. \quad (31)$$

EXAMPLE 9

The lifetime of certain equipment is described by a r.v. X whose distribution is Gamma with parameters $\alpha = 2$ and $\beta = \frac{1}{3}$, so that the corresponding p.d.f. is: $f(x) = 9xe^{-3x}$, for $x > 0$. Determine the expected lifetime, the variation around it, and the probability that the lifetime is at least 1 unit of time.

DISCUSSION Since $EX = \alpha\beta$ and $\text{Var}(X) = \alpha\beta^2$, we have here: $EX = \frac{2}{3}$ and $\text{Var}(X) = \frac{2}{9}$. Also,


$$P(X > 1) = \int_1^\infty 9xe^{-3x} dx = \frac{4}{e^3} \simeq 0.199.$$

Negative Exponential Distribution In (27), set $\alpha = 1$ and $\beta = \frac{1}{\lambda}$ ($\lambda > 0$) to obtain:

$$f(x) = \lambda e^{-\lambda x}, \quad x > 0, \lambda > 0. \quad (32)$$

This is the so-called *Negative Exponential* distribution with *parameter* λ . The graph of $f(x)$ depends on λ but, typically, looks as in Figure 3.7.

Figure 3.7
Graph of the
Negative
Exponential p.d.f.
with $\lambda = 1$



For a r.v. X having the *Negative Exponential* distribution with parameter λ , formulas (31) give:

$$EX = \frac{1}{\lambda}, \quad \text{Var}(X) = \frac{1}{\lambda^2}, \quad \text{and} \quad M_X(t) = \frac{\lambda}{\lambda - t}, \quad t < \lambda. \quad (33)$$

The expression $EX = \frac{1}{\lambda}$ provides special significance for the parameter λ : its inverse value is the mean of X . This fact also suggests the *reparameterization* of f ; namely, set $\frac{1}{\lambda} = \mu$, in which case:

$$f(x) = \frac{1}{\mu} e^{-\frac{x}{\mu}}, \quad x > 0, \quad EX = \mu, \quad \text{Var}(X) = \mu^2, \quad \text{and} \\ M_X(t) = \frac{1}{1 - \mu t}, \quad t < \frac{1}{\mu}. \quad (34)$$

From (32), one finds by a simple integration:

$$F(x) = 1 - e^{-\lambda x}, \quad x > 0, \quad \text{so that } P(X > x) = e^{-\lambda x}, \quad x > 0. \quad (35)$$

The *Negative Exponential* distribution is used routinely as a survival distribution; namely, as describing the lifetime of an equipment, etc., put in service at what may be termed as time zero. As such, it exhibits a *lack of memory* property, which may not be desirable in this context. Namely, if one poses the following question: What is the probability that an equipment will last for t additional units of time, given that it has already survived s units of time, the answer (by means of the *Negative Exponential* distribution) is, by (35):

$$P(X > s + t \mid X > s) = \frac{P(X > s + t, X > s)}{P(X > s)} = \frac{P(X > s + t)}{P(X > s)} = \frac{e^{-\lambda(s+t)}}{e^{-\lambda s}} \\ = e^{-\lambda t} = P(X > t);$$

i.e., $P(X > s + t \mid X > s) = P(X > t)$ independent of s ! Well, in real life, used pieces of equipment do not exactly behave as brand-new ones! Finally, it is to be mentioned that the Negative Exponential distribution is the waiting time distribution between the occurrence of any two successive events following the Poisson distribution (see also Exercise 3.20(ii) in Chapter 2).

EXAMPLE 10

The lifetime of an automobile battery is described by a r.v. X having the Negative Exponential distribution with parameter $\lambda = \frac{1}{3}$. Then:

- (i) Determine the expected lifetime of the battery and the variation around this mean.
- (ii) Calculate the probability that the lifetime will be between 2 and 4 time units.
- (iii) If the battery has lasted for 3 time units, what is the (conditional) probability that it will last for at least an additional time unit?

DISCUSSION

- (i) Since $EX = \frac{1}{\lambda}$ and $Var(X) = \frac{1}{\lambda^2}$, we have here: $EX = 3$, $Var(X) = 9$, and $s.d.(X) = 3$.
- (ii) Since, by (35), $F(x) = 1 - e^{-\frac{x}{3}}$ for $x > 0$, we have $P(2 < X < 4) = P(2 < X \leq 4) = P(X \leq 4) - P(X \leq 2) = F(4) - F(2) = (1 - e^{-\frac{4}{3}}) - (1 - e^{-\frac{2}{3}}) = e^{-\frac{2}{3}} - e^{-\frac{4}{3}} \simeq 0.252$.
- (iii) The required probability is: $P(X > 4 \mid X > 3) = P(X > 1)$, by the memoryless property of this distribution, and $P(X > 1) = 1 - P(X \leq 1) = 1 - F(1) = e^{-\frac{1}{3}} \simeq 0.716$.

Chi-Square Distribution In formula (27), set $\alpha = \frac{r}{2}$ for a positive integer r and $\beta = 2$ to obtain:

$$f(x) = \frac{1}{\Gamma(\frac{r}{2})2^{r/2}} x^{(r/2)-1} e^{-x/2}, \quad x > 0, \quad r > 0 \text{ integer.} \quad (36)$$

The resulting distribution is known as the *Chi-Square* distribution with r *degrees of freedom* (d.f.). This distribution is used in certain statistical inference problems involving confidence intervals for variances and testing hypotheses about variances. The notation used for a r.v. X having the Chi-Square distribution with r d.f. is $X \sim \chi_r^2$. For such a r.v., formulas (31) then become:

$$EX = r, \quad Var(X) = 2r \quad (\text{both easy to remember}) \text{ and}$$

$$M_X(t) = \frac{1}{(1 - 2t)^{r/2}}, \quad t < \frac{1}{2}. \quad (37)$$


The shape of the graph of f depends on r , and, typically, looks like that in Figure 3.8.

Later on (see Corollary to Theorem 5 in Chapter 5), it will be seen why r is referred to as the number of d.f. of the distribution.


Normal Distribution This is by far the most important distribution, in both probability and statistics. The reason for this is twofold: First, many observations do follow to a very satisfactory degree a Normal distribution

Figure 3.8

Graph of the p.d.f. of the Chi-Square Distribution for Several Values of r

**Figure 3.9**

Graph of the p.d.f. of the Normal Distribution with $\mu = 1.5$ and Several Values of σ



(see, for instance, Examples 13–17 in Chapter 1); and second, no matter what the underlying distribution of observations is, the sum of sufficiently many observations behaves pretty much as if it were normally distributed, under very mild conditions. This second property is referred to as *normal approximation* or as the *Central Limit Theorem* and will be revisited later on (see Section 7.2 in Chapter 7). The p.d.f. of a Normal distribution is given by:

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/\sigma^2}, \quad x \in \mathfrak{R}, \mu \in \mathfrak{R}, \sigma > 0; \quad (38)$$

μ and σ^2 (or σ) are referred to as the *parameters* of the distribution. The graph of f depends on μ and σ ; typical cases for $\mu = 1.5$ and various values of σ are given in Figure 3.9.

No matter what μ and σ are, the curve representing f attains its maximum at $x = \mu$ and this maximum is equal to $1/\sqrt{2\pi}\sigma$, is symmetric around μ (i.e., $f(\mu - x) = f(\mu + x)$), and $f(x)$ tends to 0 as $x \rightarrow \infty$ or $x \rightarrow -\infty$. All these observations follow immediately from formula (38). That the function $f(x)$ integrates to 1 is seen through a technique involving a double integral


and polar coordinates (see Exercise 3.44). For $\mu = 0$ and $\sigma = 1$, formula (38) is reduced to:

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}, \quad x \in \mathfrak{R}, \quad (39)$$

and this is referred to as the *standard Normal* distribution (see Figure 3.10 for its graph).

Figure 3.10

Graph of the p.d.f. of the standard Normal Distribution



The fact that a r.v. X is *Normally distributed* with parameters μ and σ^2 (or σ) is conveniently denoted by: $X \sim N(\mu, \sigma^2)$. In particular, $X \sim N(0, 1)$ for $\mu = 0$, $\sigma = 1$. We often use the notation Z for a $N(0, 1)$ distributed r.v.

The d.f. of the $N(0, 1)$ -distribution is usually denoted by Φ ; i.e., if $Z \sim N(0, 1)$, then:

$$P(Z \leq x) = \Phi(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt, \quad x \in \mathfrak{R}. \quad (40)$$

Calculations of probabilities of the form $P(a < X < b)$ for $-\infty \leq a \leq b < \infty$ are done through two steps: First, turn the r.v. $X \sim N(\mu, \sigma^2)$ into a $N(0, 1)$ -distributed r.v., or, as we say, *standardize* it, and then use available tables, the Normal tables (see Proposition 3). The standardization is based on the following simple result.

PROPOSITION 3 If $X \sim N(\mu, \sigma^2)$, then $Z = \frac{X - \mu}{\sigma}$ is $\sim N(0, 1)$.

PROOF Indeed, for $y \in \mathfrak{R}$,

$$\begin{aligned} F_Z(y) &= P(Z \leq y) = P\left(\frac{X - \mu}{\sigma} \leq y\right) \\ &= P(X \leq \mu + \sigma y) \\ &= \int_{-\infty}^{\mu + \sigma y} \frac{1}{\sqrt{2\pi}\sigma} e^{-(t-\mu)^2/2\sigma^2} dt. \end{aligned}$$

Set $\frac{t-\mu}{\sigma} = z$, so that $t = \mu + \sigma z$ with range from $-\infty$ to y , and $dt = \sigma dz$, to obtain:

$$\begin{aligned} F_Z(y) &= \int_{-\infty}^y \frac{1}{\sqrt{2\pi}\sigma} e^{-z^2/2} \sigma dz \\ &= \int_{-\infty}^y \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz, \quad \text{so that} \\ f_Z(y) &= \frac{d}{dy} F_Z(y) = \frac{1}{\sqrt{2\pi}} e^{-y^2/2}, \end{aligned}$$

which is the p.d.f. of the $N(0, 1)$ distribution. \blacktriangle

Thus, if $X \sim N(\mu, \sigma^2)$ and a, b are as above, then:

$$\begin{aligned} P(a < X < b) &= P\left(\frac{a-\mu}{\sigma} < \frac{X-\mu}{\sigma} < \frac{b-\mu}{\sigma}\right) = P\left(\frac{a-\mu}{\sigma} < Z < \frac{b-\mu}{\sigma}\right) \\ &= \Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right). \end{aligned}$$

That is,

$$P(a < X < b) = \Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right). \quad (41)$$

Any other probabilities (involving intervals) can be found by way of probability (40) by exploiting the symmetry (around 0) of the $N(0, 1)$ curve.

Now, if $Z \sim N(0, 1)$, it is clear that $EZ^{2n+1} = 0$ for $n = 0, 1, \dots$; by integration by parts, the following recursive relation is also easily established:

$$m_{2n} = (2n-1)m_{2n-2}, \quad \text{where } m_k = \int_{-\infty}^{\infty} x^k \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx, \quad (42)$$

from which it follows that $EZ = 0$ and $EZ^2 = 1$, so that $\text{Var}(Z) = 1$. (For details, see Exercise 3.48.)

If $X \sim N(\mu, \sigma^2)$, then $Z = \frac{X-\mu}{\sigma} \sim N(0, 1)$, so that (by properties (9) and (10)):

$$\begin{aligned} 0 = EZ &= \frac{EX}{\sigma} - \frac{\mu}{\sigma}, \quad 1 = \text{Var}(Z) = \frac{1}{\sigma^2} \text{Var}(X), \quad \text{or } EX = \mu \quad \text{and} \\ \text{Var}(X) &= \sigma^2. \end{aligned}$$

In other words:

$$\text{If } X \sim N(\mu, \sigma^2), \quad \text{then } EX = \mu \quad \text{and} \quad \text{Var}(X) = \sigma^2. \quad (43)$$

Thus, the parameters μ and σ^2 have specific interpretations: μ is the mean of X and σ^2 is its variance (so that σ is its s.d.).

If $Z \sim N(0, 1)$, it is seen from the Normal tables that:

$$\begin{aligned} P(-1 < Z < 1) &= 0.68269, & P(-2 < Z < 2) &= 0.95450, \\ P(-3 < Z < 3) &= 0.99730, \end{aligned}$$

so that almost all of the probability mass lies within 3 standard deviations from the mean. The same is true, by means of formula (41), applied with $a = \mu - k\sigma$

and $b = \mu + k\sigma$ with $k = 1, 2, 3$ in case $X \sim N(\mu, \sigma^2)$. That is:

$$\begin{aligned} P(\mu - \sigma < X < \mu + \sigma) &= 0.68269, & P(\mu - 2\sigma < X < \mu + 2\sigma) &= 0.95450, \\ P(\mu - 3\sigma < X < \mu + 3\sigma) &= 0.99730. \end{aligned}$$

Finally, simple integration produces the m.g.f. of X (see also Exercise 3.46), namely,

$$\begin{aligned} M_X(t) &= e^{\mu t + \sigma^2 t^2 / 2}, & t \in \Re, & \text{ for } X \sim N(\mu, \sigma^2); \\ M_Z(t) &= e^{t^2 / 2}, & t \in \Re, & \text{ for } Z \sim N(0, 1). \end{aligned} \quad (44)$$

As will be seen in subsequent chapters, the Normal distribution is widely used in problems of statistical inference, involving point estimation, interval estimation, and testing hypotheses. Some instances where the Normal distribution is assumed as an appropriate (approximate) underlying distribution are described in Examples 13–17 in Chapter 1, as mentioned already.

EXAMPLE 11

Suppose that numerical grades in a statistics class are values of a r.v. X which is (approximately) Normally distributed with mean $\mu = 65$ and s.d. $\sigma = 15$. Furthermore, suppose that letter grades are assigned according to the following rule: the student receives an A if $X \geq 85$; B if $70 \leq X < 85$; C if $55 \leq X < 70$; D if $45 \leq X < 55$; and F if $X < 45$.

- (i) If a student is chosen at random from that class, calculate the probability that the student will earn a given letter grade.
- (ii) Identify the expected proportions of letter grades to be assigned.

DISCUSSION

- (i) The student earns an A with probability $P(X \geq 85) = 1 - P(X < 85) = 1 - P\left(\frac{X - \mu}{\sigma} < \frac{85 - 65}{15}\right) \simeq 1 - P(Z < 1.34) \simeq 1 - \Phi(1.34) = 1 - 0.909877 = 0.090123 \simeq 0.09$. Likewise, the student earns a B with probability $P(70 \leq X < 85) = P\left(\frac{70 - 65}{15} \leq \frac{X - \mu}{\sigma} < \frac{85 - 65}{15}\right) \simeq P(0.34 \leq Z < 1.34) \simeq \Phi(1.34) - \Phi(0.34) = 0.909877 - 0.633072 = 0.276805 \simeq 0.277$. Similarly, the student earns a C with probability $P(55 \leq X < 70) \simeq \Phi(0.34) + \Phi(0.67) - 1 = 0.381643 \simeq 0.382$. The student earns a D with probability $P(45 \leq X < 55) \simeq \Phi(1.34) - \Phi(0.67) = 0.161306 \simeq 0.161$, and the student is assigned an F with probability $P(X < 45) \simeq \Phi(-1.34) = 1 - \Phi(1.34) = 0.09123 \simeq 0.091$.
- (ii) The respective expected proportions for A, B, C, D, and F are: 9%, 28%, 38%, 16%, and 9%.

Indeed, suppose there are n students, and let X_A be the number of those whose numerical grades are ≥ 85 . By assuming that the n events that the numerical grade of each one of the n students is ≥ 85 are independent, we have that $X_A \sim B(n, 0.09)$. Then, $\frac{X_A}{n}$ is the proportion of A grades, and $E\left(\frac{X_A}{n}\right) = \frac{1}{n}n \times 0.09 = 0.09 = 9\%$ is the expected proportion of A's. Likewise for the other grades.

This section is concluded with a simple distribution, the Uniform (or Rectangular) distribution.


Uniform (or Rectangular) Distribution Such a distribution is restricted to finite intervals between the *parameters* α and β with $-\infty < \alpha < \beta < \infty$, and its p.d.f. is given by:

$$f(x) = \frac{1}{\beta - \alpha}, \quad \alpha \leq x \leq \beta \quad (-\infty < \alpha < \beta < \infty). \quad (45)$$

Its graph is given in Figure 3.11, and it also justifies its name as rectangular.

Figure 3.11

Graph of the p.d.f. of the $U(\alpha, \beta)$ Distribution



The term “uniform” is justified by the fact that intervals of equal length in (α, β) are assigned the same probability regardless of their location. The notation used for such a distribution is $U(\alpha, \beta)$ (or $R(\alpha, \beta)$), and the fact that the r.v. X is distributed as such is denoted by $X \sim U(\alpha, \beta)$ (or $X \sim R(\alpha, \beta)$). Simple integrations give (see also Exercise 3.51):

$$EX = \frac{\alpha + \beta}{2}, \quad \text{Var}(X) = \frac{(\alpha - \beta)^2}{12}, \quad \text{and} \quad M_X(t) = \frac{e^{\beta t} - e^{\alpha t}}{(\beta - \alpha)t}, \quad t \in \Re. \quad (46)$$

EXAMPLE 12

A bus is supposed to arrive at a given bus stop at 10:00 a.m., but the actual time of arrival is a r.v. X which is uniformly distributed over the 16-minute interval from 9:52 to 10:08. If a passenger arrives at the bus stop at exactly 9:50, what is the probability that the passenger will board the bus no later than 10 minutes from the time of his/her arrival?

DISCUSSION The p.d.f. of X is $f(x) = 1/16$ for x ranging between 9:52 and 10:08, and 0 otherwise. The passenger will board the bus no later than 10 minutes from the time of his/her arrival at the bus stop if the bus arrives at the bus stop between 9:52 and 10:00 (as the passenger will necessarily have to wait for 2 minutes, between 9:50 and 9:52). The probability for the bus to arrive between 9:52 and 10:00 is $8/16 = 0.5$. This is the required probability.

REMARK 4 It has been stated (see comments after relation (7)) that sometimes a handful of moments of a r.v. X completely determine the distribution of X . Actually, this has been the case in all seven distributions examined in this section. In the Binomial distribution, knowledge of the mean amounts

to knowledge of p , and hence of f . The same is true in the Geometric distribution, as well as in the Poisson distribution. In the Hypergeometric distribution, knowledge of the first two moments, or equivalently, of the mean and variance of X (see expressions for the expectation and variance), determine m and n and hence the distribution itself. The same is true of the Gamma distribution, as well as the Normal and Uniform distributions.

Exercises

3.1 If $X \sim B(n, p)$ with $p > 0.5$, the Binomial tables (in this book) cannot be used, even if n is suitable. This problem is resolved by the following result.

(i) If $X \sim B(n, p)$, show that $P(X = x) = P(Y = n - x)$, where $Y \sim B(n, q)$ ($q = 1 - p$).

(ii) Apply part (i) for $n = 20$, $p = 0.625$, and $x = 8$.

3.2 Let X be a r.v. distributed as $B(n, p)$, and recall that $P(X = x) = f(x) = \binom{n}{x} p^x q^{n-x}$, $x = 0, 1, \dots, n$ ($q = 1 - p$). Set $B(n, p; x) = f(x)$.

(i) By using the relationship: $\binom{m+1}{y} = \binom{m}{y} + \binom{m}{y-1}$ (see Exercise 5.11 in Chapter 2), show that:

$$B(n+1, p; x) = pB(n, p; x-1) + qB(n, p; x).$$

(ii) By using this recursive relation of $B(n+1, p; \cdot)$, calculate the probabilities $B(n, p; x)$ for $n = 26$, $p = 0.25$, and $x = 10$.

3.3 Someone buys one ticket in each one of 50 lotteries, and suppose that each ticket has probability $1/100$ of winning a prize. Compute the probability that the person in question will win a prize:

(i) Exactly once.

(ii) At least once.

3.4 Suppose that 15 people, chosen at random from a target population, are asked if they favor a certain proposal. If 43.75% of the target population favor the proposal, calculate the probability that:

(i) At least 5 of the 15 polled favor the proposal.

(ii) A majority of those polled favor the proposal.

3.5 A fair die is tossed independently 18 times, and the appearance of a 6 is called a success. Find the probability that:

(i) The number of successes is greater than the number of failures.

(ii) The number of successes is twice as large as the number of failures.

(iii) The number of failures is 3 times the number of successes.

3.6 Suppose you are throwing darts at a target and that you hit the bull's eye with probability p . It is assumed that the trials are independent and that p remains constant throughout.

(i) If you throw darts 100 times, what is the probability that you hit the bull's eye at least 40 times?

- (ii) What does this expression become for $p = 0.25$?
 (iii) What is the expected number of hits, and what is the s.d. around this expected number?

3.7 If $X \sim B(100, 1/4)$, use the Tchebichev inequality to determine a lower bound for the probability: $P(|X - 25| < 10)$.

3.8 A manufacturing process produces defective items at the constant (but unknown to us) proportion p . Suppose that n items are sampled independently and let X be the r.v. denoting the number of defective items among the n , so that $X \sim B(n, p)$. Use the Tchebichev inequality in order to determine the smallest value of the sample size n , so that: $P(|\frac{X}{n} - p| < 0.05\sqrt{pq}) \geq 0.95$ ($q = 1 - p$).

3.9 If $X \sim B(n, p)$ show that $f(x+1) = \frac{p}{q} \cdot \frac{n-x}{x+1} f(x)$, $x = 0, 1, \dots, n-1$, ($q = 1 - p$).

3.10 If $X \sim B(n, p)$:

- (i) Calculate the EX and the $E[X(X-1)]$.
 (ii) Use part (i) and Exercise 1.6 to calculate the $Var(X)$.

Hint: For part (i), observe that:

$$\begin{aligned} EX &= \sum_{x=1}^n x \frac{n(n-1)!}{x(x-1)!(n-x)!} p^x q^{n-x} = np \sum_{x=1}^n \binom{n-1}{x-1} p^{x-1} q^{(n-1)-x} \\ &= np \sum_{y=0}^{n-1} \binom{n-1}{y} p^y q^{(n-1)-y} = np, \end{aligned}$$

$$\begin{aligned} \text{and } E[X(X-1)] &= \sum_{x=2}^n x(x-1) \frac{n(n-1)(n-2)!}{x(x-1)(x-2)!(n-x)!} p^x q^{n-x} \\ &= n(n-1)p^2 \sum_{x=2}^n \binom{n-2}{x-2} p^{x-2} q^{(n-2)-(x-2)} \\ &= n(n-1)p^2 \sum_{y=0}^{n-2} \binom{n-2}{y} p^y q^{(n-2)-y} = n(n-1)p^2. \end{aligned}$$

3.11 If $X \sim B(n, p)$:

- (i) Show that $M_X(t) = (pe^t + q)^n$, $t \in \Re$ ($q = 1 - p$).
 (ii) Use part (i) to rederive the EX and the $Var(X)$.

3.12 Let the r.v. X have the Geometric p.d.f. $f(x) = pq^{x-1}$, $x = 1, 2, \dots$ ($q = 1 - p$).

- (i) What is the probability that the first success will occur by the 10th trial?
 (ii) What is the numerical value of this probability for $p = 0.2$?

3.13 A manufacturing process produces defective items at the rate of 1%. Let X be the r.v. denoting the number of trials required until the first defective

item is produced. Then calculate the probability that X is not larger than 10.

3.14 A fair die is tossed repeatedly until a six appears for the first time. Calculate the probability that:

- (i) This happens on the 3rd toss.
- (ii) At least 5 tosses will be needed.

3.15 A coin with probability p of falling heads is tossed repeatedly and independently until the first head appears.

- (i) Determine the smallest number of tosses, n , required to have the first head appearing by the n th time with prescribed probability α .
- (ii) Determine the value of n for $\alpha = 0.95$, and $p = 0.25$ ($q = 0.75$) and $p = 0.50$ ($=q$).

3.16 If X has the Geometric distribution; i.e., $f(x) = pq^{x-1}$, for $x = 1, 2, \dots$ ($q = 1 - p$):

- (i) Calculate the EX and the $E[X(X - 1)]$.
- (ii) Use part (i) and Exercise 1.6 to calculate the $Var(X)$.

Hint: Refer to the comments made just before the Exercises in Section 3.1 (of this chapter).

3.17 If X has the Geometric distribution, then:

- (i) Derive the m.g.f. of X and specify the range of its argument.
- (ii) Employ the m.g.f. in order to derive: EX , EX^2 , and $Var(X)$.

3.18 Suppose that r.v. X is distributed as $P(\lambda)$; i.e., $f(x) = e^{-\lambda} \frac{\lambda^x}{x!}$, for $x = 0, 1, \dots$, and that $f(2) = 2f(0)$. Determine the value of the parameter λ .

3.19 Let X be a Poisson distributed r.v. with parameter λ , and suppose that $P(X = 0) = 0.1$. Calculate the probability $P(X = 5)$.

3.20 If $X \sim P(\lambda)$, show that: $f(x + 1) = \frac{\lambda}{x+1} f(x)$, $x = 0, 1, \dots$

3.21 If $X \sim P(\lambda)$:

- (i) Calculate the EX and the $E[X(X - 1)]$.
- (ii) Use part (i) and Exercise 1.6 to calculate the $Var(X)$.

Hint: For part (i), observe that:

$$EX = e^{-\lambda} \sum_{x=1}^{\infty} x \frac{\lambda \cdot \lambda^{x-1}}{x(x-1)!} = \lambda e^{-\lambda} \sum_{y=0}^{\infty} \frac{\lambda^y}{y!} = \lambda e^{-\lambda} e^{\lambda} = \lambda, \text{ and}$$

$$E[X(X - 1)] = \lambda^2 e^{-\lambda} \sum_{x=2}^{\infty} x(x-1) \frac{\lambda^{x-2}}{x(x-1)(x-2)!} = \lambda^2 e^{-\lambda} \sum_{y=0}^{\infty} \frac{\lambda^y}{y!} = \lambda^2.$$

3.22 If $X \sim P(\lambda)$:

- (i) Show that $M_X(t) = e^{\lambda(e^t - 1)}$, $t \in \mathfrak{R}$.
- (ii) Use the m.g.f. to rederive the EX and the $Var(X)$.

3.23 For $n = 1, 2, \dots$, let the r.v. $X_n \sim B(n, p_n)$ where, as $n \rightarrow \infty$, $0 < p_n \rightarrow 0$, and $np_n \rightarrow \lambda \in (0, \infty)$. Then show that:

$$\binom{n}{x} p_n^x q_n^{n-x} \xrightarrow{n \rightarrow \infty} e^{-\lambda} \frac{\lambda^x}{x!} \quad (q_n = 1 - p_n).$$

Hint: Write $\binom{n}{x}$ as $n(n-1)\dots(n-x+1)/x!$, set $np_n = \lambda_n$, so that $p_n = \frac{\lambda_n}{n} \xrightarrow{n \rightarrow \infty} 0$ and $q_n = 1 - p_n = 1 - \frac{\lambda_n}{n} \xrightarrow{n \rightarrow \infty} 1$. Group terms suitably, take the limit as $n \rightarrow \infty$, and use the calculus fact that $(1 + \frac{x_n}{n})^n \rightarrow e^x$ when $x_n \rightarrow x$ as $n \rightarrow \infty$.

3.24 In an undergraduate statistics class of 80, 10 of the students are, actually, graduate students. If 5 students are chosen at random from the class, what is the probability that:

- (i) No graduate students are included?
- (ii) At least 3 undergraduate students are included?

3.25 Suppose a geologist has collected 15 specimens of a certain rock, call it R_1 , and 10 specimens of another rock, call it R_2 . A laboratory assistant selects randomly 15 specimens for analysis, and let X be the r.v. denoting the number of specimens of rock R_1 selected for analysis.

- (i) Specify the p.d.f. of the r.v. X .
- (ii) What is the probability that at least 10 specimens of the rock R_1 are included in the analysis?
- (iii) What is the probability that all specimens come from the rock R_2 ?

3.26 If the r.v. X has the Hypergeometric distribution; i.e., $P(X = x) = f(x) = \frac{\binom{m}{x} \binom{n-x}{r-x}}{\binom{m+n}{r}}$, $x = 0, 1, \dots, r$, then show that:

$$f(x+1) = \frac{(m-x)(r-x)}{(n-r+x+1)(x+1)} f(x).$$

Hint: Start with $f(x+1)$ and write the numerator in terms of factorials. Then modify suitably some terms, and regroup them to arrive at the expression on the right-hand side.

3.27 By using the definition of $\Gamma(\alpha)$ by (26) and integrating by parts, show that: $\Gamma(\alpha) = (\alpha - 1)\Gamma(\alpha - 1)$, $\alpha > 1$.

3.28 Let the r.v. X have the Gamma distribution with parameters α and β . Then:

- (i) Show that: $EX = \alpha\beta$, $Var(X) = \alpha\beta^2$.
- (ii) As a special case of part (i), show that: If X has the Negative Exponential distribution with parameter λ , then $EX = \frac{1}{\lambda}$, $Var(X) = \frac{1}{\lambda^2}$.
- (iii) If $X \sim \chi_r^2$, then $EX = r$, $Var(X) = 2r$.

3.29 If the r.v. X is distributed as Gamma with parameters α and β , then:

- (i) Show that $M_X(t) = 1/(1 - \beta t)^\alpha$, provided $t < 1/\beta$.
- (ii) Use the m.g.f. to rederive the EX and the $Var(X)$.

3.30 Let X be a r.v. denoting the lifetime of a certain component of a system, and suppose that X has the Negative Exponential distribution with parameter λ . Also, let $g(x)$ be the cost of operating this equipment to time $X = x$.

- (i) Compute the expected cost of operation over the lifetime of the component under consideration, when:
- $g(x) = cx$, where c is a positive constant,
 - $g(x) = c(1 - 0.5e^{-\alpha x})$, where α is a positive constant.
- (ii) Specify the numerical values in part (i) when $\lambda = 1/5$, $c = 2$, and $\alpha = 0.2$.

3.31 If the r.v. X has the Negative Exponential p.d.f. with parameter λ , calculate the *failure rate* $r(x)$ defined by: $r(x) = \frac{f(x)}{1-F(x)}$, for $x > 0$, where F is the d.f. of X .

3.32 Suppose that certain events occur in a time interval λt according to the Poisson distribution with parameter λt . Then show that the waiting time between any two such successive events is a r.v. T which has the Negative Exponential distribution with parameter λ , by showing that $P(T > t) = e^{-\lambda t}$, $t > 0$.

3.33 Let X be the r.v. denoting the number of particles arriving independently at a detector at the average rate of 3 per second, and let Y be the r.v. denoting the waiting time between two successive arrivals. Refer to Exercise 3.32 in order to calculate:

- The probability that the first particle will arrive within 1 second.
- Given that we have waited for 1 second since the arrival of the last particle without a new arrival, what is the probability that we have to wait for at least another second?

3.34 Let X be a r.v. with p.d.f. $f(x) = \alpha\beta x^{\beta-1}e^{-\alpha x^\beta}$, for $x > 0$ (where the parameters α and β are > 0). This is the so-called *Weibull* distribution employed in describing the lifetime of living organisms or of mechanical systems.

- Show that f is, indeed, a p.d.f.
- For what values of the parameters does f become a Negative Exponential p.d.f.?
- Calculate the quantities: EX , EX^2 , and $Var(X)$.

Hint: For part (i), observe that: $\int_0^\infty \alpha\beta x^{\beta-1}e^{-\alpha x^\beta} dx = \int_0^\infty e^{-\alpha x^\beta} \times (\alpha\beta x^{\beta-1})dx = -\int_0^\infty de^{-\alpha x^\beta} = -e^{-\alpha x^\beta} \Big|_0^\infty = 1$.

For part (iii), set $\alpha x^\beta = t$, so that $x = t^{1/\beta}/\alpha^{1/\beta}$, $dx = (t^{\frac{1}{\beta}-1}/\beta\alpha^{1/\beta})dt$ and $0 < t < \infty$. Then:

$$EX^n = \frac{1}{\alpha^{n/\beta}} \int_0^\infty t^{(\frac{n}{\beta}+1)-1} e^{-t} dt.$$

Then multiply and divide by the constant $\Gamma(\frac{n}{\beta} + 1)$ and observe that $\frac{1}{\Gamma(\frac{n}{\beta}+1)} t^{(\frac{n}{\beta}+1)-1} e^{-t}$ ($t > 0$) is a Gamma p.d.f. with parameters $\frac{n}{\beta} + 1$ and 1.

3.35 In reference to Exercise 3.34, calculate:

- The *failure rate* $r(x) = \frac{f(x)}{1-F(x)}$, $x > 0$, where F is the d.f. of the r.v. X .
- The conditional probability $P(X > s + t \mid X > t)$, $s > 0$, $t > 0$.

(iii) Compare the results in parts (i) and (ii) with the respective results in Exercise 3.31 here, and Exercise 3.20(ii) in Chapter 2.

3.36 If Φ is the d.f. of the r.v. $Z \sim N(0, 1)$, show that:

- (i) For $0 \leq a < b$, $P(a < Z < b) = \Phi(b) - \Phi(a)$.
- (ii) For $a \leq 0 < b$, $P(a < Z < b) = \Phi(-a) + \Phi(b) - 1$.
- (iii) For $a \leq b < 0$, $P(a < Z < b) = \Phi(-a) - \Phi(-b)$.
- (iv) For $c > 0$, $P(-c < Z < c) = 2\Phi(c) - 1$.

3.37 If the r.v. $Z \sim N(0, 1)$, use the Normal tables in the appendix to verify that:

- (i) $P(-1 < Z < 1) = 0.68269$.
- (ii) $P(-2 < Z < 2) = 0.9545$.
- (iii) $P(-3 < Z < 3) = 0.9973$.

3.38 (i) If the r.v. X is distributed as $N(\mu, \sigma^2)$, identify the constant c , in terms of μ and σ , for which:

$$P(X < c) = 2 - 9P(X > c).$$

(ii) What is the numerical value of c for $\mu = 5$ and $\sigma = 2$?

3.39 For any r.v. X with expectation μ and variance σ^2 (both finite), use the Tchebichev inequality to determine a lower bound for the probabilities: $P(|X - \mu| < k\sigma)$, for $k = 1, 2, 3$. Compare these bounds with the respective probabilities when $X \sim N(\mu, \sigma^2)$ (see Exercise 3.37).

3.40 The distribution of I.Q.'s of the people in a given group is approximated well by the Normal distribution with $\mu = 105$ and $\sigma = 20$. What proportion of the individuals in the group in question has an I.Q. :

- (i) At least 50?
- (ii) At most 80?
- (iii) Between 95 and 125?

3.41 A certain manufacturing process produces light bulbs whose life length (in hours) is a r.v. X distributed as Normal with $\mu = 2,000$ and $\sigma = 200$. A light bulb is supposed to be defective if its lifetime is less than 1,800. If 25 light bulbs are tested, what is the probability that at most 15 of them are defective? (Use the required independence.)

3.42 A manufacturing process produces 1/2-inch ball bearings, which are assumed to be satisfactory if their diameter lies in the interval 0.5 ± 0.0006 and defective otherwise. A day's production is examined, and it is found that the distribution of the actual diameters of the ball bearings is approximately Normal with $\mu = 0.5007$ inch and $\sigma = 0.0005$ inch. What would you expect the proportion of defective ball bearings to be equal to?

3.43 Let f be the p.d.f. of the $N(\mu, \sigma^2)$ distribution. Then show that:

- (i) f is symmetric about μ .
- (ii) $\max_{x \in \mathcal{N}} f(x) = 1/\sqrt{2\pi}\sigma$.

- 3.44** (i) Show that $f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$, $x \in \mathfrak{R}$, is a p.d.f.
 (ii) Use part (i) in order to show that $f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$, $x \in \mathfrak{R}$ ($\mu \in \mathfrak{R}$, $\sigma > 0$) is also a p.d.f.

Hint: Set $I = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{x^2}{2}} dx$ and show that $I^2 = 1$, by writing I^2 as a product of two integrals and then as a double integral; at this point, use polar coordinates: $x = r \cos \theta$, $y = r \sin \theta$, $0 < r < \infty$, $0 \leq \theta < 2\pi$. Part (ii) is reduced to part (i) by letting $\frac{x-\mu}{\sigma} = y$.

- 3.45** Refer to the definition of $\Gamma(\alpha)$ by (26) and show that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$.
- 3.46** (i) If $X \sim N(0, 1)$, show that $M_X(t) = e^{t^2/2}$, $t \in \mathfrak{R}$.
 (ii) If $X \sim N(\mu, \sigma^2)$, use part (i) to show that $M_X(t) = e^{\mu t + \frac{\sigma^2 t^2}{2}}$, $t \in \mathfrak{R}$.
 (iii) Employ the m.g.f. in part (ii) in order to show that $EX = \mu$ and $Var(X) = \sigma^2$.
- 3.47** If the r.v. X has m.g.f. $M_X(t) = e^{\alpha t + \beta t^2}$, where $\alpha \in \mathfrak{R}$ and $\beta > 0$, identify the distribution of X .
- 3.48** If $X \sim N(0, 1)$, show that:
 (i) $EX^{2n+1} = 0$ and $EX^{2n} = \frac{(2n)!}{2^n(n!)}$, $n = 0, 1, \dots$
 (ii) From part (i), derive that $EX = 0$ and $Var(X) = 1$.
 (iii) Employ part (ii) in order to show that, if $X \sim N(\mu, \sigma^2)$, then $EX = \mu$ and $Var(X) = \sigma^2$.

Hint: For part (i), that $EX^{2n+1} = 0$ follows by the fact that the integrand is an odd function. For EX^{2n} , establish a recursive relation, integrating by parts, and then multiply out the resulting recursive relations to find an expression for EX^{2n} . The final form follows by simple manipulations. For part (iii), recall that $X \sim N(\mu, \sigma^2)$ implies $\frac{X-\mu}{\sigma} \sim N(0, 1)$.

- 3.49** Let X be a r.v. with moments given by:

$$EX^{2n+1} = 0, \quad EX^{2n} = \frac{(2n)!}{2^n(n!)}, \quad n = 0, 1, \dots$$

- (i) Use Exercise 1.17 in order to express the m.g.f. of X in terms of the moments given.
 (ii) From part (i) and Exercise 3.46(i), conclude that $X \sim N(0, 1)$.
- 3.50** If the r.v. X is distributed as $U(-\alpha, \alpha)$ ($\alpha > 0$), determine the parameter α , so that each of the following equalities holds:
 (i) $P(-1 < X < 2) = 0.75$.
 (ii) $P(|X| < 1) = P(|X| > 2)$.

- 3.51** If $X \sim U(\alpha, \beta)$, show that $EX = \frac{\alpha+\beta}{2}$, $Var(X) = \frac{(\alpha-\beta)^2}{12}$.

- 3.52** If the r.v. X is distributed as $U(0, 1)$, compute the expectations:
 (i) $E(3X^2 - 7X + 2)$.
 (ii) $E(2e^X)$.

3.4 Median and Mode of a Random Variable

Although the mean of a r.v. X does specify the center of location of the distribution of X , sometimes this is not what we actually wish to know. A case in point is the distribution of yearly income in a community (e.g., in a state or in a country). For the sake of illustration, consider the following (rather) extreme example. A community consisting of 100 households comprises 10 households with yearly income \$400,000 each and 90 households with yearly income \$12,000 each. Defining the r.v. X to take the values 400,000 and 12,000 with respective probabilities 0.10 and 0.90, we obtain: $EX = 50,800$. Thus, the average yearly income in this community would be \$50,800, significantly above the national average yearly income, which would indicate a rather prosperous community. The reality, however, is that this community is highly stratified, and the expectation does not reveal this characteristic. What is more appropriate for cases like this are numerical characteristics of a distribution known as median or, more generally, percentiles or quantiles.

The median of the distribution of a r.v. X is usually defined as a point, denoted by $x_{0.50}$, for which

$$P(X \leq x_{0.50}) \geq 0.50 \quad \text{and} \quad P(X \geq x_{0.50}) \geq 0.50, \quad (47)$$

or, equivalently,

$$P(X < x_{0.50}) \leq 0.50 \quad \text{and} \quad P(X \leq x_{0.50}) \geq 0.50. \quad (48)$$

If the underlying distribution is continuous, the median is (essentially) unique and may be simply defined by:

$$P(X \leq x_{0.50}) = P(X \geq x_{0.50}) = 0.50 \quad (49)$$

However, in the discrete case, relation (47) (or (48)) may not define the median in a unique manner, as the following example shows.

EXAMPLE 13

Examine the median of the r.v. X distributed as follows.

x	1	2	3	4	5	6	7	8	9	10
$f(x)$	2/32	1/32	5/32	3/32	4/32	1/32	2/32	6/32	2/32	6/32

DISCUSSION We have $P(X \leq 6) = 16/32 = 0.50 \geq 0.50$ and $P(X \geq 6) = 17/32 > 0.05 \geq 0.50$, so that (47) is satisfied. Also,

$$P(X \leq 7) = 18/32 > 0.50 \geq 0.50 \quad \text{and} \quad P(X \geq 7) = 16/32 = 0.50 \geq 0.50,$$

so that (47) is satisfied again. However, if we define the median as the point $(6 + 7)/2 = 6.5$, then $P(X \leq 6.5) = P(X \geq 6.5) = 0.50$, as (47) requires, and the median is uniquely defined.

Relations (47)–(49) and Example 13 suggest the following definition of the median.

DEFINITION 4

The *median* of the distribution of a continuous r.v. X is the (essentially) unique point $x_{0.50}$ defined by (49). For the discrete case, consider two cases: Let x_k be the value for which $P(X \leq x_k) = 0.50$, if such a value exists. Then the unique *median* is defined to be the midpoint between x_k and x_{k+1} : i.e., $x_{0.50} = (x_k + x_{k+1})/2$. If there is no such value, the unique *median* is defined by the relations: $P(X < x_{0.50}) < 0.50$ and $P(X \leq x_{0.50}) > 0.50$ (or $P(X \leq x_{0.50}) > 0.50$ and $P(X \geq x_{0.50}) > 0.50$).

Thus, in Example 14, $x_{0.50} = 6$, because $P(X < 6) = P(X \leq 5) = 15/32 < 0.50$ and $P(X \leq 6) = 17/32 > 0.50$.

EXAMPLE 14

Determine the median of the r.v. X distributed as follows.

x	1	2	3	4	5	6	7	8	9	10
$f(x)$	2/32	1/32	2/32	6/32	4/32	2/32	1/32	7/32	1/32	6/32

More generally, the p th quartile is defined as follows.

DEFINITION 5

For any p with $0 < p < 1$, the p th *quartile* of the distribution of a r.v. X , denoted by x_p , is defined as follows: If X is continuous, then the (essentially) unique x_p is defined by:

$$P(X \leq x_p) = p \quad \text{and} \quad P(X \geq x_p) = 1 - p.$$

For the discrete case, consider two cases: Let x_k be the value for which $P(X \leq x_k) = p$, if such a value exists. Then the unique p th *quartile* is defined to be the midpoint between x_k and x_{k+1} ; i.e., $x_p = (x_k + x_{k+1})/2$. If there is no such value, the unique p th *quartile* is defined by the relation: $P(X < x_p) < p$ and $P(X \leq x_p) > p$ (or $P(X \leq x_p) > p$ and $P(X \geq x_p) > 1 - p$).

Thus, the p th quartile is a point x_p , which divides the distribution of X into two parts, and $(-\infty, x_p]$ contains exactly $100p\%$ (or at least $100p\%$) of the distribution, and $[x_p, \infty)$ contains exactly $100(1 - p)\%$ (or at least $100(1 - p)\%$) of the distribution of X . For $p = 0.50$, we obtain the median. These concepts are illustrated further by the following examples.

EXAMPLE 15

Refer to Figure 3.1 ($B(12, 1/4)$) and determine $x_{0.25}$, $x_{0.50}$, and $x_{0.75}$.

DISCUSSION Here $x_{0.25} = 2$ since $P(X < 2) = P(X = 0) + P(X = 1) = 0.1584 \leq 0.25$ and $P(X \leq 2) = 0.1584 + P(X = 2) = 0.3907 \geq 0.25$. Likewise, $x_{0.50} = 3$ since $P(X < 3) = 0.3907 \leq 0.50$ and $P(X \leq 3) = 0.6488 \geq 0.50$. Finally, $x_{0.75} = 4$, since $P(X < 4) = 0.6488 \leq 0.75$ and $P(X \leq 4) = 0.8424 > 0.75$.

EXAMPLE 16

Refer to Figure 3.4 ($P(5)$) and determine $x_{0.25}$, $x_{0.50}$, and $x_{0.75}$.

As in the previous example, $x_{0.25} = 2$, $x_{0.50} = 4$, and $x_{0.75} = 6$.

EXAMPLE 17

If $X \sim U(0, 1)$, take $p = 0.10, 0.20, 0.30, 0.40, 0.50, 0.60, 0.70, 0.80$, and 0.90 and determine the corresponding x_p .

Here $F(x) = \int_0^x dt = x$, $0 \leq x \leq 1$. Therefore $F(x_p) = p$ gives $x_p = p$.

EXAMPLE 18

If $X \sim N(0, 1)$, take p as in the previous example and determine the corresponding x_p .

From the Normal tables, we obtain: $x_{0.10} = -x_{0.90} = -1.282$, $x_{0.20} = -x_{0.80} = -0.842$, $x_{0.30} = -x_{0.70} = -0.524$, $x_{0.40} = -x_{0.60} = -0.253$, and $x_{0.50} = 0$.

Another numerical characteristic which helps shed some light on the distribution of a r.v. X is the so-called mode.

DEFINITION 6

A *mode* of the distribution of a r.v. X is any point, if such points exist, which maximizes the p.d.f. of X , f .

A mode, being defined as a maximizing point, is subject to all shortcomings of maximization: It may not exist at all; it may exist but is not obtainable in closed form; there may be more than one mode (the distribution is a *multimodal* one). It may also happen that there is a unique mode (*unimodal* distribution). Clearly, if a mode exists, it will be of particular importance for discrete distributions, as the modes provide the values of the r.v. X which occur with the largest probability. With this in mind, we restrict ourselves to two of the most popular discrete distributions: the Binomial and the Poisson distribution.

THEOREM 2

Let X be $B(n, p)$; that is,

$$f(x) = \binom{n}{x} p^x q^{n-x}, \quad 0 < p < 1, \quad q = 1 - p, \quad x = 0, 1, \dots, n.$$

Consider the number $(n+1)p$ and set $m = [(n+1)p]$, where $[y]$ denotes the largest integer which is $\leq y$. Then, if $(n+1)p$ is *not* an integer, $f(x)$ has a unique mode at $x = m$. If $(n+1)p$ is an integer, then $f(x)$ has two modes obtained for $x = m$ and $x = m - 1$.

PROOF For $x \geq 1$, we have

$$\begin{aligned} \frac{f(x)}{f(x-1)} &= \frac{\binom{n}{x} p^x q^{n-x}}{\binom{n}{x-1} p^{x-1} q^{n-x+1}} \\ &= \frac{\frac{n!}{x!(n-x)!} p^x q^{n-x}}{\frac{n!}{(x-1)!(n-x+1)!} p^{x-1} q^{n-x+1}} = \frac{n-x+1}{x} \cdot \frac{p}{q}. \end{aligned}$$

That is,

$$\frac{f(x)}{f(x-1)} = \frac{n-x+1}{x} \cdot \frac{p}{q}.$$

Hence $f(x) > f(x-1)$ (f is increasing) if and only if

$$(n-x+1)p > x(1-p), \quad \text{or } np - xp + p > x - xp, \quad \text{or } (n+1)p > x, \quad \text{and}$$

$$f(x) = f(x-1) \quad \text{if and only if } x = (n+1)p \text{ in case } (n+1)p \text{ is an integer.}$$

Thus, if $(n+1)p$ is *not* an integer, we have:

$$0 < 1 < \cdots < m < (n+1)p < m+1 < \cdots < n,$$

and, by a slight abuse of notation,

$$f(0) < f(1) < \cdots < f(m) > f(m+1) > \cdots > f(n),$$

so that there is a unique mode at m . If $(n+1)p$ is an integer ($=m$), then we have:

$$0 < 1 < \cdots < m-1 < m < m+1 < \cdots < n,$$

and, as above,

$$f(0) < f(1) < \cdots < f(m-1) = f(m) > f(m+1) > \cdots > f(n),$$

so that there are two modes at m and $m-1$. ▲

THEOREM 3

Let X be $P(\lambda)$; that is,

$$f(x) = e^{-\lambda} \frac{\lambda^x}{x!}, \quad x = 0, 1, 2, \dots, \quad \lambda > 0.$$

Then, if λ is *not* an integer, $f(x)$ has a unique mode at $x = [\lambda]$. If λ is an integer, then $f(x)$ has two modes obtained for $x = \lambda$ and $x = \lambda - 1$.

PROOF For $x \geq 1$, we have

$$\frac{f(x)}{f(x-1)} = \frac{e^{-\lambda} \lambda^x / x!}{e^{-\lambda} \lambda^{x-1} / (x-1)!} = \frac{\lambda}{x}.$$

Hence $f(x) > f(x-1)$ if and only if $\lambda > x$, and $f(x) = f(x-1)$ if and only if $x = \lambda$ in case λ is an integer. Thus, if λ is *not* an integer, $f(x)$ keeps increasing for $x \leq [\lambda]$ and then decreases. Thus the maximum of $f(x)$ occurs at $x = [\lambda]$. If λ is an integer, then the maximum occurs at $x = \lambda$. But in this case $f(x) = f(x-1)$, which implies that $x = \lambda - 1$ is a second point which gives the maximum value to the p.d.f. ▲

EXAMPLE 19

Let $X \sim B(n, p)$ with $n = 20$ and $p = \frac{1}{4}$. Then $(n+1)p = \frac{21}{4}$ is not an integer and therefore there is a unique mode. Since $\frac{21}{4} = 5.25$, the mode is $[5.25] = 5$. The maximum probability is $\binom{20}{5} (0.25)^5 (0.75)^{15} = 0.2024$. If $n = 15$ and $p = \frac{1}{4}$,

then $(n+1)p = \frac{16}{4} = 4$ and therefore there are two modes; they are 4 and 3. The respective maximum probability is $\binom{15}{4}(0.25)^4(0.75)^{11} = 0.2252$.

EXAMPLE 20

Let $X \sim P(\lambda)$ and let $\lambda = 4.5$. Then there is a unique mode which is $[4.5] = 4$. The respective maximum probability is 0.1898. If, on the other hand, $\lambda = 7$, then there are two modes 7 and 6. The respective maximum probability is 0.149.

Exercises

- 4.1** Let X be a r.v. with p.d.f. $f(x) = 3x^2$, for $0 \leq x \leq 1$.
- Calculate the EX and the median of X and compare them.
 - Determine the 0.125-quantile of X .
- 4.2** Let X be a r.v. with p.d.f. $f(x) = x^n$, for $0 \leq x \leq c$ (n positive integer), and let $0 < p < 1$. Determine:
- The p th quantile x_p of X in terms of n and p .
 - The median $x_{0.50}$ for $n = 3$.
- 4.3** (i) If the r.v. X has p.d.f. $f(x) = \lambda e^{-\lambda x}$, for $x > 0$ ($\lambda > 0$), determine the p th quantile x_p in terms of λ and p .
- (ii) What is the numerical value of x_p for $\lambda = \frac{1}{10}$ and $p = 0.25$?
- 4.4** Let X be a r.v. with p.d.f. f given by:
- $$f(x) = \begin{cases} c_1 x^2, & -1 \leq x \leq 0 \\ c_2(1 - x^2), & 0 < x \leq 1 \\ 0, & \text{otherwise.} \end{cases}$$
- If it is also given that $EX = 0$, determine the constants c_1 and c_2 .
 - Determine the $\frac{1}{3}$ -quantile of the distribution.
- 4.5** Let X be a r.v. with d.f. F given in Exercise 2.2 of Chapter 2:
- Determine the mode of the respective p.d.f. f .
 - Show that $\frac{1}{2}$ is the $\frac{5}{32} = 0.15625$ -quantile of the distribution.
- 4.6** Two fair and distinct dice are rolled once, and let X be the r.v. denoting the sum of the numbers shown, so that the possible values of X are: 2, 3, ..., 12.
- Derive the p.d.f. f of the r.v. X .
 - Compute the EX .
 - Find the median of f , as well as its mode.
- 4.7** Determine the modes of the following p.d.f.'s:
- $f(x) = (\frac{1}{2})^x$, $x = 1, 2, \dots$
 - $f(x) = (1 - \alpha)^x$, $x = 1, 2, \dots$ ($0 < \alpha < 1$). Also, what is the value of α ?
 - $f(x) = \frac{2}{3^{x+1}}$, $x = 0, 1, \dots$

4.8 Let $X \sim B(100, 1/4)$ and suppose you were to bet on the observed value of X . On which value would you bet?

4.9 In reference to Exercise 3.33, which number(s) of particles arrives within 1 second with the maximum probability?

4.10 Let X be a r.v. (of the continuous type) with p.d.f. f symmetric about a constant c (i.e., $f(c - x) = f(c + x)$ for all x ; in particular, if $c = 0$, then $f(-x) = f(x)$ for all x). Then show that c is the median of X . (As a by-product of it, we have, for example, that the mean μ in the $N(\mu, \sigma^2)$ is also the median.)

Hint: Start with $P(X \leq c) = \int_{-\infty}^c f(x)dx$ and, by making a change of the variable x , show that this last integral equals $\int_0^{\infty} f(c - y)dy$. Likewise, $P(X \geq c) = \int_c^{\infty} f(x)dx$ and a change of the variable x leads to the integral $\int_0^{\infty} f(c + y)dy$. Then the use of symmetry completes the proof.

4.11 Let X be a r.v. of the continuous type with p.d.f. f , with finite expectation, and median m , and let c be any constant. Then:

(i) Show that:

$$E|X - c| = E|X - m| + 2 \int_m^c (c - x)f(x) dx.$$

(ii) Use part (i) to conclude that the constant c which minimizes the $E|X - c|$ is $c = m$.

Hint: For $m < c$, show that:

$$|x - c| - |x - m| = \begin{cases} c - m, & x < m \\ c + m - 2x, & m \leq x \leq c \\ m - c, & x > c. \end{cases}$$

Then

$$\begin{aligned} E|X - c| - E|X - m| &= \int_{-\infty}^m (c - m)f(x) dx + \int_m^c (c + m - 2x)f(x) dx \\ &\quad + \int_c^{\infty} (m - c)f(x) dx \\ &= \frac{c - m}{2} + (c + m) \int_m^c f(x) dx - 2 \int_m^c xf(x) dx \\ &\quad + (m - c) \int_m^{\infty} f(x) dx - (m - c) \int_m^c f(x) dx \\ &= \frac{c - m}{2} + \frac{m - c}{2} + 2c \int_m^c f(x) dx - 2 \int_m^c xf(x) dx \\ &= 2 \int_m^c (c - x)f(x) dx. \end{aligned}$$

For $m \geq c$, show that:

$$|x - c| - |x - m| = \begin{cases} c - m, & x < c \\ -c - m + 2x, & c \leq x \leq m \\ m - c, & x > m. \end{cases}$$

Then

$$\begin{aligned} E|X - c| - E|X - m| &= \int_{-\infty}^c (c - m)f(x) dx + \int_c^m (-c - m + 2x)f(x) dx \\ &\quad + \int_m^{\infty} (m - c)f(x) dx \\ &= (c - m) \int_{-\infty}^m f(x) dx - (c - m) \int_c^m f(x) dx \\ &\quad - (c + m) \int_c^m f(x) dx + 2 \int_c^m xf(x) dx \\ &\quad + (m - c) \int_m^{\infty} f(x) dx \\ &= \frac{c - m}{2} + \frac{m - c}{2} - 2c \int_c^m f(x) dx + 2 \int_c^m xf(x) dx \\ &= -2 \int_c^m (c - x)f(x) dx = 2 \int_m^c (c - x)f(x) dx. \end{aligned}$$

Combining the two results, we get

$$E|X - c| = E|X - m| + 2 \int_m^c (c - x)f(x) dx.$$

4.12 Let X be a continuous r.v. with p th quantile x_p , and let $Y = g(X)$, where g is a strictly increasing function, so that the inverse g^{-1} exists (and is also strictly increasing). Let y_p be the p th quantile of the r.v. Y .

- (i) Show that $y_p = g(x_p)$.
- (ii) If X has the Negative Exponential distribution with $\lambda = 1$, calculate x_p .
- (iii) Use parts (i) and (ii) to determine y_p without calculations, where $Y = e^X$.
- (iv) What do parts (ii) and (iii) become for $p = 0.5$?

Joint and Conditional p.d.f.'s, Conditional Expectation and Variance, Moment Generating Function, Covariance, and Correlation Coefficient

A brief description of the material discussed in this chapter is as follows. In the first section, two r.v.'s are considered and the concepts of their joint probability distribution, joint d.f., and joint p.d.f. are defined. The basic properties of the joint d.f. are given, and a number of illustrative examples are provided. On the basis of a joint d.f., marginal d.f.'s are defined. Also, through a joint p.d.f., marginal and conditional p.d.f.'s are defined, and illustrative examples are supplied. By means of conditional p.d.f.'s, conditional expectations and conditional variances are defined and are applied to some examples. These things are done in the second section of the chapter.

In the following section, the expectation is defined for a function of two r.v.'s and some basic properties are listed. As a special case, one obtains the joint m.g.f. of the r.v.'s involved, and from this, marginal m.g.f.'s are derived. Also, as a special case, one obtains the covariance and the correlation coefficient of two r.v.'s. Their significance is explained, and a basic inequality is established regarding the range of their values. Finally, a formula is provided for the calculation of the variance of the sum of two r.v.'s.

In the fourth section of the chapter, many of the concepts, defined for two r.v.'s in the previous sections, are generalized to k r.v.'s. In the final section, three specific multidimensional distributions are introduced, the Multinomial, the Bivariate (or 2-dimensional) Normal, and the Multivariate Normal. The derivation of marginal and conditional p.d.f.'s of the Multinomial and Bivariate Normal distributions is also presented. This section is concluded with a brief discussion of the Multivariate Normal distribution.

4.1 Joint d.f. and Joint p.d.f. of Two Random Variables

In carrying out a random experiment, we are often interested simultaneously in two outcomes rather than one. Then with each one of these outcomes a r.v. is associated, and thus we are furnished with two r.v.'s or a *2-dimensional random vector*. Let us denote by (X, Y) the two relevant r.v.'s or the 2-dimensional random vector. Here are some examples where two r.v.'s arise in a natural way. The pair of r.v.'s (X, Y) denote, respectively: the SAT and GPA scores of a student chosen at random from a specified student population; the number of customers waiting for service in two lines in your local favorite bank; the days of a given year that the Dow Jones Averages closed with a gain and the corresponding gains; the number of hours a student spends daily for studying and for other activities; the weight and the height of an individual chosen at random from a targeted population; the amount of fertilizer used and the yield of a certain agricultural commodity; the lifetimes of two components used in an electronic system; the dosage of a drug used for treating a certain allergy and the number of days a patient enjoys relief.


We are going to restrict ourselves to the case where both X and Y are either discrete or of the continuous type. The concepts of probability distribution, distribution function, and probability density function are defined by a straightforward generalization of the definition of these concepts in Section 2.2 of Chapter 2. Thus, the *joint probability distribution* of (X, Y) , to be denoted by $P_{X,Y}$, is defined by: $P_{X,Y}(B) = P[(X, Y) \in B]$, $B \subseteq \mathfrak{R}^2 = \mathfrak{R} \times \mathfrak{R}$, the 2-dimensional Euclidean space, the plane. In particular, by taking $B = (-\infty, x] \times (-\infty, y]$, we obtain the *joint d.f.* of X, Y , to be denoted by $F_{X,Y}$; namely, $F_{X,Y}(x, y) = P(X \leq x, Y \leq y)$, $x, y \in \mathfrak{R}$. The d.f. $F_{X,Y}$ has properties similar to the ones mentioned in the case of a single r.v., namely:

1. $0 \leq F_{X,Y}(x, y) \leq 1$ for all $x, y \in \mathfrak{R}$.
Whereas it is, clearly, still true that $x_1 \leq x_2$ and $y_1 \leq y_2$ imply $F_{X,Y}(x_1, y_1) \leq F_{X,Y}(x_2, y_2)$, property #2 in the case of a single r.v. may be restated as follows: $x_1 < x_2$ implies $F_X(x_2) - F_X(x_1) \geq 0$. This property is replaced here by:
2. The *variation* of $F_{X,Y}$ over rectangles with sides parallel to the axes, given in Figure 4.1, is ≥ 0 .
3. $F_{X,Y}$ is continuous from the right (right-continuous); i.e., if $x_n \downarrow x$ and $y_n \downarrow y$, then $F_{X,Y}(x_n, y_n) \rightarrow F_{X,Y}(x, y)$ as $n \rightarrow \infty$.

4. $F_{X,Y}(+\infty, +\infty) = 1$ and $F_{X,Y}(-\infty, -\infty) = F_{X,Y}(-\infty, y) = F_{X,Y}(x, -\infty) = 0$ for any $x, y \in \mathfrak{R}$, where, of course, $F_{X,Y}(+\infty, +\infty)$ is defined to be the $\lim_{n \rightarrow \infty} F_{X,Y}(x_n, y_n)$ as $x_n \uparrow \infty$ and $y_n \uparrow \infty$, and similarly for the remaining cases.

Figure 4.1

The Variation V of $F_{X,Y}$ over the Rectangle Is:
 $F_{X,Y}(x_1, y_1) +$
 $F_{X,Y}(x_2, y_2) -$
 $F_{X,Y}(x_1, y_2) -$
 $F_{X,Y}(x_2, y_1)$



Property #1 is immediate, and property #2 follows by the fact that the variation of $F_{X,Y}$ as described is simply the probability that the pair (X, Y) lies in the rectangle of Figure 4.1, or, more precisely, the probability $P(x_1 < X \leq x_2, y_1 < Y \leq y_2)$, which, of course, is ≥ 0 ; the justification of properties #3 and #4 is based on Theorem 2 in Chapter 2.

Now, suppose that the r.v.'s X and Y are discrete and take on the values x_j and y_j , $j \geq 1$, respectively. Then the *joint p.d.f.* of X and Y , to be denoted by $f_{X,Y}$, is defined by: $f_{X,Y}(x_j, y_j) = P(X = x_j, Y = y_j)$ and $f_{X,Y}(x, y) = 0$ when $(x, y) \neq (x_j, y_j)$ (i.e., at least one of x or y is not equal to x_j or y_j , respectively). It is then immediate that for $B \subseteq \mathfrak{R}^2$, $P[(X, Y) \in B] = \sum_{(x_j, y_j) \in B} f_{X,Y}(x_j, y_j)$, and, in particular, $\sum_{(x_j, y_j) \in \mathfrak{R}^2} f_{X,Y}(x_j, y_j) = 1$, and $F_{X,Y}(x, y) = \sum_{x_j \leq x, y_j \leq y} f_{X,Y}(x_j, y_j)$. In the last relation, $F_{X,Y}$ is expressed in terms of $f_{X,Y}$. The converse is also possible (as was done in the case of a single r.v.), but we do not intend to indulge in it. A simple illustrative example, however, may be in order.

EXAMPLE 1

Each one of the r.v.'s X and Y takes on four values only, 0, 1, 2, 3, with joint probabilities expressed best in a matrix form as in Table 4.1.

Table 4.1

$y \backslash x$	0	1	2	3	Totals
0	0.05	0.21	0	0	0.26
1	0.20	0.26	0.08	0	0.54
2	0	0.06	0.07	0.02	0.15
3	0	0	0.03	0.02	0.05
Totals	0.25	0.53	0.18	0.04	1

DISCUSSION The r.v.'s X and Y may represent, for instance, the number of customers waiting for service in two lines in a bank. Then, for example,

for (x, y) with $x = 2$ and $y = 1$, we have $F_{X,Y}(x, y) = F_{X,Y}(2, 1) = \sum_{u \leq 2, v \leq 1} f_{X,Y}(u, v) = f_{X,Y}(0, 0) + f_{X,Y}(0, 1) + f_{X,Y}(1, 0) + f_{X,Y}(1, 1) + f_{X,Y}(2, 0) + f_{X,Y}(2, 1) = 0.05 + 0.20 + 0.21 + 0.26 + 0 + 0.08 = 0.80$; also, $P(2 \leq X \leq 3, 0 \leq Y \leq 2) = f_{X,Y}(2, 0) + f_{X,Y}(2, 1) + f_{X,Y}(2, 2) + f_{X,Y}(3, 0) + f_{X,Y}(3, 1) + f_{X,Y}(3, 2) = 0 + 0.08 + 0.07 + 0 + 0 + 0.02 = 0.17$.

Now, suppose that both X and Y are of the continuous type, and, indeed, a little bit more; namely, there exists a nonnegative function $f_{X,Y}$ defined on \mathfrak{R}^2 such that, for all x and y in \mathfrak{R} : $F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(s, t) ds dt$. Then for $B \subseteq \mathfrak{R}^2$ (interpret B as a familiar geometric figure in \mathfrak{R}^2): $P[(X, Y) \in B] = \int_B \int f_{X,Y}(x, y) dx dy$, and, in particular, $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy = 1$. The function $f_{X,Y}$ is called the *joint p.d.f.* of X and Y . Analogously to the case of a single r.v., the relationship $\frac{\partial^2}{\partial x \partial y} F_{X,Y}(x, y) = f_{X,Y}(x, y)$ holds true (for continuity points (x, y) of $f_{X,Y}$), so that not only does the joint p.d.f. determine the joint d.f. through an integration process, but the converse is also true; i.e., the joint d.f. determines the joint p.d.f. through differentiation. Again, as in the case of a single r.v., $P(X = x, Y = y) = 0$ for all $x, y \in \mathfrak{R}$; also, if a nonnegative function f , defined on \mathfrak{R}^2 , integrates to 1, then there exist two r.v.'s X and Y for which f is their joint p.d.f.

This section is concluded with a reference to Example 37 in Chapter 1 where two continuous r.v.'s X and Y arise in a natural manner. Later on (see Subsection 4.5.2), it may be stipulated that the joint distribution of X and Y is the Bivariate Normal. For the sake of a simpler illustration, consider the following example.

EXAMPLE 2

Let the r.v.'s X and Y have the joint p.d.f. $f_{X,Y}(x, y) = \lambda_1 \lambda_2 e^{-\lambda_1 x - \lambda_2 y}$, $x, y > 0$, $\lambda_1, \lambda_2 > 0$. For example, X and Y may represent the lifetimes of two components in an electronic system. Derive the joint d.f. $F_{X,Y}$.

DISCUSSION The corresponding joint d.f. is: $F_{X,Y}(x, y) = \int_0^y \int_0^x \lambda_1 \lambda_2 \times e^{-\lambda_1 s - \lambda_2 t} ds dt = \int_0^y \lambda_2 e^{-\lambda_2 t} (\int_0^x \lambda_1 e^{-\lambda_1 s} ds) dt = \int_0^y \lambda_2 e^{-\lambda_2 t} (1 - e^{-\lambda_1 x}) dt = (1 - e^{-\lambda_1 x})(1 - e^{-\lambda_2 y})$ for $x > 0, y > 0$, and 0 otherwise. That is,

$$F_{X,Y}(x, y) = (1 - e^{-\lambda_1 x})(1 - e^{-\lambda_2 y}), \quad x > 0, y > 0,$$

$$\text{and } F_{X,Y}(x, y) = 0 \text{ otherwise.} \quad (1)$$

By letting x and $y \rightarrow \infty$, we obtain $F_{X,Y}(\infty, \infty) = 1$, which also shows that $f_{X,Y}$, as given above, is, indeed, a p.d.f., since $F_{X,Y}(\infty, \infty) = \int_0^{\infty} \int_0^{\infty} \lambda_1 \lambda_2 \times e^{-\lambda_1 s - \lambda_2 t} ds dt$.

EXAMPLE 3

It is claimed that the function $F_{X,Y}$ given by: $F_{X,Y} = \frac{1}{16}xy(x+y)$, $0 \leq x \leq 2$, $0 \leq y \leq 2$, is the joint d.f. of the r.v.'s X and Y . Then:

- (i) Verify that $F_{X,Y}$ is, indeed, a d.f.
- (ii) Determine the corresponding joint p.d.f. $f_{X,Y}$.
- (iii) Verify that $f_{X,Y}$ found in part (ii) is, indeed, a p.d.f.
- (iv) Calculate the probability: $P(0 \leq X \leq 1, 1 \leq Y \leq 2)$.

DISCUSSION

(i) We have to verify the validity of the defining relations 1–4. Clearly, $F_{X,Y}(x, y)$ attains its maximum for $x = y = 2$, which is 1. Since also $F_{X,Y}(x, y) \geq 0$, the first property holds. Next, for any rectangle as in Figure 4.1, we have:

$$\begin{aligned}
 & 16[F_{X,Y}(x_1, y_1) + F_{X,Y}(x_2, y_2) - F_{X,Y}(x_1, y_2) - F_{X,Y}(x_2, y_1)] \\
 &= x_1 y_1 (x_1 + y_1) + x_2 y_2 (x_2 + y_2) - x_1 y_2 (x_1 + y_2) - x_2 y_1 (x_2 + y_1) \\
 &= x_1^2 y_1 + x_1 y_1^2 + x_2^2 y_2 + x_2 y_2^2 - x_1^2 y_2 - x_1 y_2^2 - x_2^2 y_1 - x_2 y_1^2 \\
 &= -x_1^2 (y_2 - y_1) + x_2^2 (y_2 - y_1) - y_1^2 (x_2 - x_1) + y_2^2 (x_2 - x_1) \\
 &= (x_2^2 - x_1^2)(y_2 - y_1) + (x_2 - x_1)(y_2^2 - y_1^2) \\
 &= (x_1 + x_2)(x_2 - x_1)(y_2 - y_1) + (x_2 - x_1)(y_2 + y_1)(y_2 - y_1) \\
 &\geq 0,
 \end{aligned}$$

because $x_1 \leq y_1$ and $x_2 \leq y_2$, so that the second property also holds. The third property holds because $F_{X,Y}$ is continuous, and hence right-continuous. Finally, the fourth property holds because as either $x \rightarrow -\infty$ or $y \rightarrow -\infty$ (in fact, if either one of them is 0), then $F_{X,Y}$ is 0, and if $x \rightarrow \infty$ and $y \rightarrow \infty$ (in fact, if $x = y = 1$), then $F_{X,Y}$ is 1.

(ii) For $0 \leq x \leq 2$ and $0 \leq y \leq 2$, $f_{X,Y}(x, y) = \frac{\partial^2}{\partial x \partial y} (\frac{1}{16} xy(x+y)) = \frac{1}{16} \frac{\partial^2}{\partial x \partial y} (x^2 y + xy^2) = \frac{1}{16} \frac{\partial}{\partial y} \frac{\partial}{\partial x} (x^2 y + xy^2) = \frac{1}{16} \frac{\partial}{\partial y} (2xy + y^2) = \frac{1}{16} (2x + 2y) = \frac{1}{8} (x + y)$; i.e., $f_{X,Y}(x, y) = \frac{1}{8} (x + y)$, $0 \leq x \leq 2$, $0 \leq y \leq 2$. For (x, y) outside the rectangle $[0, 2] \times [0, 2]$, $f_{X,Y}$ is 0, since $F_{X,Y}$ is constantly either 0 or 1.

(iii) Since $f_{X,Y}$ is nonnegative, all we have to show is that it integrates to 1. In fact,

$$\begin{aligned}
 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy &= \int_0^2 \int_0^2 \frac{1}{8} (x + y) dx dy \\
 &= \frac{1}{8} \left(\int_0^2 \int_0^2 x dx dy + \int_0^2 \int_0^2 y dx dy \right) \\
 &= \frac{1}{8} (2 \times 2 + 2 \times 2) \\
 &= 1.
 \end{aligned}$$

(iv) Here, $P(0 \leq X \leq 1, 1 \leq Y \leq 2) = \int_1^2 \int_0^1 \frac{1}{8} (x + y) dx dy = \frac{1}{8} [\int_1^2 (\int_0^1 x dx) dy + \int_1^2 (y \int_0^1 dx) dy] = \frac{1}{8} (\frac{1}{2} \times 1 + 1 \times \frac{3}{2}) = \frac{1}{4}$.


EXAMPLE 4

If the function $f_{X,Y}$ is given by: $f_{X,Y}(x, y) = cx^2y$ for $0 < x^2 < y < 1$ (and 0 otherwise):

- (i) Determine the constant c , so that $f_{X,Y}$ is a p.d.f.
- (ii) Calculate the probability: $P(0 < X < \frac{3}{4}, \frac{1}{4} \leq Y < 1)$.

Figure 4.2

**Range of the Pair
(x, y)**



DISCUSSION

(i) Clearly, for the function to be nonnegative, c must be > 0 . The actual value of c will be determined through the relationship:


$$\iint_{\{(x,y); 0 < x^2 \leq y < 1\}} cx^2 y dx dy = 1.$$

The region over which the p.d.f. is positive is the shaded region in Figure 4.2, determined by a branch of the parabola $y = x^2$, the y -axis, and the line segment connecting the points $(0, 1)$ and $(1, 1)$. Since for each fixed x with $0 < x < 1$, y ranges from x^2 to 1, we have: $\iint_{\{x^2 \leq y < 1\}} cx^2 y dx dy = c \int_0^1 (x^2 \int_{x^2}^1 y dy) dx = \frac{c}{2} \int_0^1 x^2 (1 - x^4) dx = \frac{c}{2} (\frac{1}{3} - \frac{1}{7}) = \frac{2c}{21} = 1$ and $c = \frac{21}{2}$.

(ii) Since $y = x^2 = \frac{1}{4}$ for $x = \frac{1}{2}$, it follows that, for each x with $0 < x \leq \frac{1}{2}$, the range of y is from $\frac{1}{4}$ to 1; on the other hand, for each x with $\frac{1}{2} < x \leq \frac{3}{4}$, the range of y is from x^2 to 1 (see Figure 4.3).

Figure 4.3

**Diagram
Facilitating
Integration**



Thus,

$$\begin{aligned}
 P\left(0 < X \leq \frac{3}{4}, \frac{1}{4} \leq Y < 1\right) &= c \int_0^{\frac{1}{2}} \int_{\frac{1}{4}}^1 x^2 y dy dx + c \int_{\frac{1}{2}}^{\frac{3}{4}} \int_{x^2}^1 x^2 y dy dx \\
 &= c \int_0^{\frac{1}{2}} \left(x^2 \int_{\frac{1}{4}}^1 y dy\right) dx + c \int_{\frac{1}{2}}^{\frac{3}{4}} \left(x^2 \int_{x^2}^1 y dy\right) dx \\
 &= \frac{c}{2} \int_0^{\frac{1}{2}} x^2 \left(1 - \frac{1}{16}\right) dx + \frac{c}{2} \int_{\frac{1}{2}}^{\frac{3}{4}} x^2 (1 - x^4) dx \\
 &= \frac{15c}{3 \times 2^8} + \frac{38c}{3 \times 2^8} - \frac{2,059c}{7 \times 2^{15}} = c \times \frac{41,311}{21 \times 2^{15}} \\
 &= \frac{21}{2} \times \frac{41,311}{21 \times 2^{15}} = \frac{41,311}{2^{16}} = \frac{41,311}{65,536} \simeq 0.63.
 \end{aligned}$$

Exercises

1.1 Let X and Y be r.v.'s denoting, respectively, the number of cars and buses lined up at a stoplight at a given point in time, and suppose their joint p.d.f. is given by the following table:

$y \setminus x$	0	1	2	3	4	5
0	0.025	0.050	0.125	0.150	0.100	0.050
1	0.015	0.030	0.075	0.090	0.060	0.030
2	0.010	0.020	0.050	0.060	0.040	0.020

Calculate the following probabilities:

- There are exactly 4 cars and no buses.
- There are exactly 5 cars.
- There is exactly 1 bus.
- There are at most 3 cars and at least 1 bus.

1.2 In a sociological project, families with 0, 1, and 2 children are studied. Suppose that the numbers of children occur with the following frequencies:

0 children: 30%; 1 child: 40%; 2 children: 30%.

A family is chosen at random from the target population, and let X and Y be the r.v.'s denoting the number of children in the family and the number of boys among those children, respectively. Finally, suppose that $P(\text{observing a boy}) = P(\text{observing a girl}) = 0.5$.

Calculate the joint p.d.f. $f_{X,Y}(x, y) = P(X = x, Y = y)$, $0 \leq y \leq x$, $x = 0, 1, 2$.

Hint: Tabulate the joint probabilities as indicated below by utilizing the formula:

$$P(X = x, Y = y) = P(Y = y | X = x)P(X = x).$$

$y \backslash x$	0	1	2
0			
1			
2			

1.3 If the r.v.'s X and Y have the joint p.d.f. given by:

$$f_{X,Y}(x, y) = x + y, \quad 0 < x < 1, \quad 0 < y < 1,$$

calculate the probability $P(X < Y)$.

1.4 The r.v.'s X and Y have the joint p.d.f. $f_{X,Y}$ given by:

$$f_{X,Y}(x, y) = \frac{6}{7} \left(x^2 + \frac{xy}{2} \right), \quad 0 < x \leq 1, \quad 0 < y \leq 2.$$

(i) Show that $f_{X,Y}$ is, indeed, a p.d.f.

(ii) Calculate the probability $P(X > Y)$.

1.5 The r.v.'s X and Y have the joint p.d.f. $f_{X,Y}(x, y) = e^{-x-y}$, $x > 0$, $y > 0$.

(i) Calculate the probability $P(X \leq Y \leq c)$ for some $c > 0$.

(ii) Find the numerical value in part (i) for $c = \log 2$, where log is the natural logarithm.

1.6 If the r.v.'s X and Y have the joint p.d.f. $f_{X,Y}(x, y) = e^{-x-y}$, for $x > 0$ and $y > 0$, compute the following probabilities:

(i) $P(X \leq x)$; (ii) $P(Y \leq y)$; (iii) $P(X < Y)$; (iv) $P(X + Y \leq 3)$.

1.7 Let X and Y be r.v.'s jointly distributed with p.d.f. $f_{X,Y}(x, y) = 2/c^2$, for $0 < x \leq y < c$.

Determine the constant c .

1.8 The r.v.'s X and Y have the joint p.d.f. $f_{X,Y}$ given by:

$$f_{X,Y}(x, y) = cy e^{-xy/2}, \quad 0 < y < x.$$

Determine the constant c .

1.9 The joint p.d.f. of the r.v.'s X and Y is given by:

$$f_{X,Y}(x, y) = xy^2, \quad 0 < x \leq c_1, \quad 0 < y \leq c_2.$$

Determine the condition that c_1 and c_2 must satisfy so that $f_{X,Y}$ is, indeed, a p.d.f.


1.10 The joint p.d.f. of the r.v.'s X and Y is given by:

$$f_{X,Y}(x, y) = cx, \quad x > 0, \quad y > 0, \quad 1 \leq x + y < 2 \quad (c > 0).$$

Determine the constant c .

Hint: The following diagram should facilitate the calculations.

The range of the pair (x, y) is the shadowed area.



1.11 The r.v.'s X and Y have joint p.d.f. $f_{X,Y}$ given by:

$$f_{X,Y}(x, y) = c(y^2 - x^2)e^{-y}, \quad -y < x < y, \quad 0 < y < \infty.$$

Determine the constant c .

4.2 Marginal and Conditional p.d.f.'s, Conditional Expectation and Variance

In the case of two r.v.'s with joint d.f. $F_{X,Y}$ and joint p.d.f. $f_{X,Y}$, we may define quantities which were not available in the case of a single r.v. These quantities are marginal d.f.'s and p.d.f.'s, conditional p.d.f.'s, and conditional expectations and variances. To this end, consider the joint d.f. $F_{X,Y}(x, y) = P(X \leq x, Y \leq y)$, and let $y \rightarrow \infty$. Then we obtain $F_{X,Y}(x, \infty) = P(X \leq x, Y < \infty) = P(X \leq x) = F_X(x)$; thus, $F_X(x) = F_{X,Y}(x, \infty)$, and likewise, $F_Y(y) = F_{X,Y}(\infty, y)$. That is, the d.f.'s of the r.v.'s X and Y are obtained from their joint d.f. by eliminating one of the variables x or y through a limiting process. The d.f.'s F_X and F_Y are referred to as *marginal* d.f.'s. If the r.v.'s X and Y are discrete with joint p.d.f. $f_{X,Y}$, then $P(X = x_i) = P(X = x_i, -\infty < Y < \infty) = \sum_{y_j \in \mathfrak{R}} f_{X,Y}(x_i, y_j)$; i.e., $f_X(x_i) = \sum_{y_j \in \mathfrak{R}} f_{X,Y}(x_i, y_j)$, and likewise, $f_Y(y_j) = \sum_{x_i \in \mathfrak{R}} f_{X,Y}(x_i, y_j)$. Because of this marginalization process, the p.d.f.'s of the r.v.'s X and Y , f_X and f_Y , are referred to as *marginal* p.d.f.'s. In the continuous case, f_X and f_Y are obtained by integrating out the "superfluous" variables; i.e., $f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$ and $f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx$. The marginal f_X is, indeed, the p.d.f. of X because $P(X \leq x) = P(X \leq x, -\infty < Y < \infty) =$

$\int_{-\infty}^x \int_{-\infty}^{\infty} f_{X,Y}(s, t) dt ds = \int_{-\infty}^x [\int_{-\infty}^{\infty} f_{X,Y}(s, t) dt] ds = \int_{-\infty}^x f_X(s) ds$; i.e., $F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(s) ds$, so that $\frac{d}{dx} F_X(x) = f_X(x)$, and likewise, $\frac{d}{dy} F_Y(y) = f_Y(y)$ (for continuity points x and y of f_X and f_Y , respectively).

In terms of the joint and the marginal p.d.f.'s, one may define formally the functions:

$$f_{X|Y}(x|y) = f_{X,Y}(x, y)/f_Y(y) \quad \text{for fixed } y \text{ with } f_Y(y) > 0,$$

and

$$f_{Y|X}(y|x) = f_{X,Y}(x, y)/f_X(x) \quad \text{for fixed } x \text{ with } f_X(x) > 0.$$

These nonnegative functions are, actually, p.d.f.'s. For example, for the continuous case:

$$\int_{-\infty}^{\infty} f_{X|Y}(x|y) dx = \frac{1}{f_Y(y)} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx = \frac{f_Y(y)}{f_Y(y)} = 1,$$

and similarly for $f_{Y|X}(y|x)$; in the discrete case, integrals are replaced by summation signs. The p.d.f. $f_{X|Y}(\cdot|y)$ is called the *conditional* p.d.f. of X , given $Y = y$, and $f_{Y|X}(\cdot|x)$ is the *conditional* p.d.f. of Y , given $X = x$. The motivation for this terminology is as follows: For the discrete case, $f_{X|Y}(x|y) = \frac{f_{X,Y}(x, y)}{f_Y(y)} = \frac{P(X=x, Y=y)}{P(Y=y)} = P(X = x | Y = y)$; i.e., $f_{X|Y}(x|y)$ does, indeed, stand for the conditional probability that $X = x$, given that $Y = y$. Likewise for $f_{Y|X}(\cdot|x)$. In the continuous case, the points x and y are to be replaced by "small" intervals around them.

The concepts introduced so far are now illustrated by means of Examples.

EXAMPLE 5

Refer to Example 1 and derive the marginal and conditional p.d.f.'s involved.

DISCUSSION From Table 4.1, we have: $f_X(0) = 0.25$, $f_X(1) = 0.53$, $f_X(2) = 0.18$, and $f_X(3) = 0.04$; also, $f_Y(0) = 0.26$, $f_Y(1) = 0.54$, $f_Y(2) = 0.15$, and $f_Y(3) = 0.05$. Thus, the probability that there are 2 people in line one, for instance, regardless of how many people are in the other line, is: $P(X = 2) = f_X(2) = 0.18$. Next, $f_{X|Y}(0|0) = \frac{0.05}{0.26} = \frac{5}{26} \approx 0.192$, $f_{X|Y}(1|0) = \frac{0.21}{0.26} = \frac{21}{26} \approx 0.808$, $f_{X|Y}(2|0) = 0$, $f_{X|Y}(3|0) = 0$; $f_{X|Y}(0|1) = \frac{0.20}{0.54} = \frac{20}{54} \approx 0.37$, $f_{X|Y}(1|1) = \frac{0.26}{0.54} = \frac{26}{54} \approx 0.481$, $f_{X|Y}(2|1) = \frac{0.08}{0.54} = \frac{8}{54} \approx 0.148$, $f_{X|Y}(3|1) = 0$; $f_{X|Y}(0|2) = 0$, $f_{X|Y}(1|2) = \frac{0.06}{0.15} = \frac{6}{15} = 0.40$, $f_{X|Y}(2|2) = \frac{0.07}{0.15} = \frac{7}{15} \approx 0.467$, $f_{X|Y}(3|2) = \frac{0.02}{0.15} = \frac{2}{15} \approx 0.133$; $f_{X|Y}(0|3) = 0$, $f_{X|Y}(1|3) = 0$, $f_{X|Y}(2|3) = \frac{0.03}{0.05} = \frac{3}{5} = 0.60$, $f_{X|Y}(3|3) = \frac{0.02}{0.05} = \frac{2}{5} = 0.40$. Likewise for $f_{Y|X}(\cdot|\cdot)$. Thus, $f_{Y|X}(0|0) = 0.2$, $f_{Y|X}(1|0) = 0.8$, $f_{Y|X}(2|0) = f_{Y|X}(3|0) = 0$; $f_{Y|X}(0|1) = \frac{21}{53} \approx 0.396$, $f_{Y|X}(1|1) = \frac{26}{53} \approx 0.491$, $f_{Y|X}(2|1) = \frac{6}{53} \approx 0.113$, $f_{Y|X}(3|1) = 0$; $f_{Y|X}(0|2) = 0$, $f_{Y|X}(1|2) = \frac{8}{18} \approx 0.444$, $f_{Y|X}(2|2) = \frac{7}{18} \approx 0.389$, $f_{Y|X}(3|2) = \frac{3}{18} \approx 0.167$; $f_{Y|X}(0|3) = f_{Y|X}(1|3) = 0$, $f_{Y|X}(2|3) = f_{Y|X}(3|3) = 0.5$.

EXAMPLE 6

Refer to Example 2 and derive the marginal d.f.'s and p.d.f.'s, as well as the conditional p.d.f.'s, involved.

DISCUSSION In (1), let $y \rightarrow \infty$ to obtain $F_X(x) = 1 - e^{-\lambda_1 x}$, $x > 0$, and likewise $F_Y(y) = 1 - e^{-\lambda_2 y}$, $y > 0$, by letting $x \rightarrow \infty$. Next, by differentiation, $f_X(x) = \lambda_1 e^{-\lambda_1 x}$, $x > 0$, and $f_Y(y) = \lambda_2 e^{-\lambda_2 y}$, $y > 0$, so that the r.v.'s X and Y have the Negative Exponential distribution with parameters λ_1 and λ_2 , respectively. Finally, for $x > 0$ and $y > 0$:

$$f_{X|Y}(x|y) = \frac{\lambda_1 \lambda_2 e^{-\lambda_1 x - \lambda_2 y}}{\lambda_2 e^{-\lambda_2 y}} = \lambda_1 e^{-\lambda_1 x} = f_X(x), \text{ and likewise}$$

$$f_{Y|X}(y|x) = f_Y(y).$$

EXAMPLE 7

Refer to Example 4 and determine the marginal and conditional p.d.f.'s f_X , f_Y , $f_{X|Y}$, and $f_{Y|X}$.

DISCUSSION We have:

$$f_X(x) = \int_{x^2}^1 cx^2 y dy = cx^2 \int_{x^2}^1 y dy = \frac{21}{4} x^2 (1 - x^4), \quad 0 < x < 1,$$

$$f_Y(y) = \int_0^{\sqrt{y}} cx^2 y dx = cy \int_0^{\sqrt{y}} x^2 dx = \frac{21}{6} y^2 \sqrt{y}, \quad 0 < y < 1,$$

and therefore

$$f_{X|Y}(x|y) = \frac{\frac{21}{2} x^2 y}{\frac{21}{6} y^2 \sqrt{y}} = \frac{3x^2}{y\sqrt{y}}, \quad 0 < x \leq \sqrt{y}, \quad 0 < y < 1,$$

$$f_{Y|X}(y|x) = \frac{\frac{21}{2} x^2 y}{\frac{21}{4} x^2 (1 - x^4)} = \frac{2y}{1 - x^4}, \quad x^2 \leq y < 1, \quad 0 < x < 1.$$

EXAMPLE 8

Consider the function $f_{X,Y}$ defined by:

$$f_{X,Y}(x, y) = 8xy, \quad 0 < x \leq y < 1.$$

- (i) Verify that $f_{X,Y}$ is, indeed, a p.d.f.
- (ii) Determine the marginal and conditional p.d.f.'s.
- (iii) Calculate the quantities: EX , EX^2 , $Var(X)$, EY , EY^2 , $Var(Y)$, and $E(XY)$.

DISCUSSION

(i) Since $f_{X,Y}$ is nonnegative, all we have to check is that it integrates to 1. In fact,

$$\int_0^1 \int_0^y 8xy dx dy = 8 \int_0^1 \left(y \int_0^y x dx \right) dy = 4 \int_0^1 y^3 dy = 1.$$

(ii)

$$f_X(x) = \int_x^1 8xy dy = 8x \int_x^1 y dy = 4x(1 - x^2), \quad 0 < x < 1,$$

$$f_Y(y) = \int_0^y 8xy dx = 8y \int_0^y x dx = 4y^3, \quad 0 < y < 1,$$

and therefore

$$f_{X|Y}(x|y) = \frac{8xy}{4y^3} = \frac{2x}{y^2}, \quad 0 < x \leq y < 1,$$

$$f_{Y|X}(y|x) = \frac{8xy}{4x(1-x^2)} = \frac{2y}{1-x^2}, \quad 0 < x \leq y < 1.$$

(iii)

$$EX = \int_0^1 x \cdot 4x(1-x^2)dx = 4 \int_0^1 x^2(1-x^2)dx = 4 \left(\int_0^1 x^2 dx - \int_0^1 x^4 dx \right) = \frac{8}{15},$$

$$EX^2 = \int_0^1 x^2 \cdot 4x(1-x^2)dx = 4 \left(\int_0^1 x^3 dx - \int_0^1 x^5 dx \right) = \frac{1}{3}, \quad \text{so that}$$

$$\text{Var}(X) = EX^2 - (EX)^2 = \frac{1}{3} - \frac{64}{225} = \frac{11}{225};$$

$$EY = \int_0^1 y \cdot 4y^3 dy = 4 \int_0^1 y^4 dy = \frac{4}{5},$$

$$EY^2 = \int_0^1 y^2 \cdot 4y^3 dy = 4 \int_0^1 y^5 dy = \frac{2}{3}, \quad \text{so that}$$

$$\text{Var}(Y) = EY^2 - (EY)^2 = \frac{2}{3} - \frac{16}{25} = \frac{2}{75} = \frac{6}{225}.$$

Finally,

$$\begin{aligned} E(XY) &= \int_0^1 \int_0^y xy \cdot 8xy dx dy = 8 \int_0^1 y^2 \left(\int_0^y x^2 dx \right) dy \\ &= \frac{8}{3} \int_0^1 y^5 dy = \frac{4}{9}. \end{aligned}$$

Once a conditional p.d.f. is at hand, an expectation can be defined as done in relations (1), (2), and (3) of Chapter 3. However, a modified notation will be needed to reveal the fact that the expectation is calculated with respect to a *conditional* p.d.f. The resulting expectation is the *conditional* expectation of one r.v., given the other r.v., as specified below.

$$E(X|Y = y_j) = \sum_{x_i \in \mathfrak{N}} x_i f_{X|Y}(x_i | y_j) \quad \text{or} \quad E(X|Y = y) = \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx, \quad (2)$$

for the discrete and continuous case, respectively; similarly:

$$E(Y|X = x_i) = \sum_{y_j \in \mathfrak{N}} y_j f_{Y|X}(y_j | x_i) \quad \text{or} \quad E(Y|X = x) = \int_{-\infty}^{\infty} y f_{Y|X}(y|x) dy. \quad (3)$$

Of course, it is understood that the preceding expectations exist as explained right after relations (2) and (3) in Chapter 3 were defined. However, unlike the results in (1)–(3) in Chapter 3 which are numbers, in relations (2) and (3)

above the outcomes depend on y_j or y , and x_i or x , respectively, which reflect the values that the “conditioning” r.v.’s assume. For illustrative purposes, let us calculate some conditional expectations.

EXAMPLE 9

In reference to Example 1, calculate: $E(X | Y = 0)$ and $E(Y | X = 2)$.

DISCUSSION In Example 5, we have calculated the conditional p.d.f.’s $f_{X|Y}(\cdot | 0)$ and $f_{Y|X}(\cdot | 2)$. Therefore:

$$E(X | Y = 0) = 0 \times \frac{5}{26} + 1 \times \frac{21}{26} + 2 \times 0 + 3 \times 0 = \frac{21}{26} \simeq 0.808, \quad \text{and}$$

$$E(Y | X = 2) = 0 \times 0 + 1 \times \frac{8}{18} + 2 \times \frac{7}{18} + 3 \times \frac{3}{18} = \frac{31}{18} \simeq 1.722.$$

So, if in the y -line there are no customers waiting, the expected number of those waiting in the x -line will be about 0.81; likewise, if there are 2 customers waiting in the x -line, the expected number of those waiting in the y -line will be about 1.72.

EXAMPLE 10

In reference to Example 2, calculate: $E(X | Y = y)$ and $E(Y | X = x)$.

DISCUSSION In Example 6, we have found that $f_{X|Y}(x | y) = f_X(x) = \lambda_1 e^{-\lambda_1 x}$ ($x > 0$), and $f_{Y|X}(y | x) = f_Y(y) = \lambda_2 e^{-\lambda_2 y}$ ($y > 0$), so that: $E(X | Y = y) = \int_0^\infty x \lambda_1 e^{-\lambda_1 x} dx = 1/\lambda_1$, and $E(Y | X = x) = \int_0^\infty y \lambda_2 e^{-\lambda_2 y} dy = 1/\lambda_2$, by integration by parts, or simply by utilizing known results.

EXAMPLE 11

In reference to Example 4, calculate: $E(X | Y = y)$ and $E(Y | X = x)$.

DISCUSSION In Example 7, we have found that $f_{X|Y}(x | y) = \frac{3x^2}{y\sqrt{y}}$, $0 < x \leq \sqrt{y}$, so that

$$E(X | Y = y) = \int_0^{\sqrt{y}} x \cdot \frac{3x^2}{y\sqrt{y}} dx = \frac{3}{y\sqrt{y}} \int_0^{\sqrt{y}} x^3 dx = \frac{3\sqrt{y}}{4}, \quad 0 < y < 1.$$

Also, $f_{Y|X}(y | x) = \frac{2y}{1-x^4}$, $x^2 \leq y < 1$, so that

$$E(Y | X = x) = \int_{x^2}^1 y \cdot \frac{2y}{1-x^4} dy = \frac{2}{1-x^4} \int_{x^2}^1 y^2 dy = \frac{2(1-x^6)}{3(1-x^4)}, \quad 0 < x < 1.$$

EXAMPLE 12

In reference to Example 8, calculate: $E(X | Y = y)$ and $E(Y | X = x)$.

DISCUSSION In Example 8(ii), we have found that $f_{X|Y}(x | y) = \frac{2x}{y^2}$, $0 < x \leq y < 1$, and $f_{Y|X}(y | x) = \frac{2y}{1-x^2}$, $0 < x \leq y < 1$, so that

$$E(X | Y = y) = \int_0^y x \cdot \frac{2x}{y^2} dx = \frac{2}{y^2} \int_0^y x^2 dx = \frac{2y}{3}, \quad 0 < y < 1,$$

and

$$E(Y|X=x) = \int_x^1 y \cdot \frac{2y}{1-x^2} dy = \frac{2}{1-x^2} \int_x^1 y^2 dy = \frac{2(1-x^3)}{3(1-x^2)}, \quad 0 < x < 1.$$

Now, for the discrete case, set $g(y_j) = E(X|Y = y_j)$ and proceed to replace y_j by the r.v. Y . We obtain the r.v. $g(Y) = E(X|Y)$, and then it makes sense to talk about its expectation $Eg(Y) = E[E(X|Y)]$. Although the $E(X|Y = y_j)$ depends on the particular values of Y , it turns out that its average does not, and, indeed, is the same as the EX . More precisely, it holds:

$$E[E(X|Y)] = EX \quad \text{and} \quad E[E(Y|X)] = EY. \quad (4)$$

That is, the expectation of the conditional expectation of X is equal to its expectation, and likewise for Y . Relation (4) is true both for the discrete and the continuous case. Its justification for the continuous case, for instance, is as follows:

We have $g(Y) = E(X|Y)$ and therefore

$$\begin{aligned} Eg(Y) &= \int_{-\infty}^{\infty} g(y) f_Y(y) dy = \int_{-\infty}^{\infty} E(X|y) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx \right] f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [x f_{X|Y}(x|y) f_Y(y) dx] dy = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f_{X,Y}(x,y) dx dy \\ &= \int_{-\infty}^{\infty} x \left[\int_{-\infty}^{\infty} f_{X,Y}(x,y) dy \right] dx = \int_{-\infty}^{\infty} x f_X(x) dx = EX; \text{ i.e.,} \\ Eg(Y) &= E[E(X|Y)] = EX. \end{aligned}$$

REMARK 1 However, $\text{Var}[E(X|Y)] \leq \text{Var}(X)$ with equality holding, if and only if Y is a function of X (with probability 1). A proof of this fact may be found in Section 5.3.1 in the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G. G. Roussas.

EXAMPLE 13

Verify the first relation $E[E(X|Y)] = EX$, in (4) for Examples 4 and 8.

DISCUSSION By Example 7, $f_X(x) = \frac{21}{4}x^2(1-x^4)$, $0 < x < 1$, so that

$$EX = \int_0^1 x \cdot \frac{21}{4}x^2(1-x^4) dx = \frac{21}{4} \left(\int_0^1 x^3 dx - \int_0^1 x^7 dx \right) = \frac{21}{32}.$$

From Example 11, $E(X|Y) = \frac{3\sqrt{Y}}{4}$, $0 < Y < 1$, whereas, from Example 7, $f_Y(y) = \frac{21}{6}y^2\sqrt{y}$, $0 < y < 1$, so that

$$E[E(X|Y)] = \int_0^1 \frac{3\sqrt{y}}{4} \cdot \frac{21}{6}y^2\sqrt{y} dy = \frac{21}{8} \int_0^1 y^3 dy = \frac{21}{32} = EX.$$

(However, $\text{Var}[E(X | Y)] = \text{Var}(\frac{3\sqrt{Y}}{4}) = \frac{9}{16} \text{Var}(\sqrt{Y}) = \frac{9}{16} [EY - (E\sqrt{Y})^2] = \frac{9}{16} (\frac{4}{5} - \frac{49}{64}) = \frac{99}{5,120} < \frac{2}{75} = \text{Var}(Y)$.)

Also, from Examples 12 and 8(ii), $E(X | Y) = \frac{2Y}{3}$, $0 < Y < 1$, and $f_Y(y) = 4y^3$, $0 < y < 1$, $EX = \frac{8}{15}$ by Example 8, so that

$$E[E(X | Y)] = \int_0^1 \frac{2y}{3} \cdot 4y^3 dy = \frac{8}{3} \int_0^1 y^4 dy = \frac{8}{15} = EX.$$

(However, $\text{Var}[E(X | Y)] = \text{Var}(\frac{2Y}{3}) = \frac{4}{9} \text{Var}(Y) < \text{Var}(Y)$.)

In addition to the conditional expectation of X , given Y , one may define the *conditional variance* of X , given Y , by utilizing the conditional p.d.f. and formula (8) in Chapter 3; the notation to be used is $\text{Var}(X | Y = y_j)$ or $\text{Var}(X | Y = y)$ for the discrete and continuous case, respectively. Thus:

$$\text{Var}(X | Y = y_j) = \sum_{x_i \in \mathfrak{R}} [x_i - E(X | Y = y_j)]^2 f_{X|Y}(x_i | y_j), \quad (5)$$

and

$$\text{Var}(X | Y = y) = \int_{-\infty}^{\infty} [x - E(X | Y = y)]^2 f_{X|Y}(x | y) dx, \quad (6)$$

for the discrete and the continuous case, respectively. The conditional variances depend on the values of the conditioning r.v., as was the case for the conditional expectations. From formulas (5) and (6), it is not hard to see (see also Exercise 2.20) that:

$$\text{Var}(X | Y = y_j) = E(X^2 | Y = y_j) - [E(X | Y = y_j)]^2 \quad \text{or} \quad (7)$$

$$\text{Var}(X | Y = y) = E(X^2 | Y = y) - [E(X | Y = y)]^2,$$

for the discrete and the continuous case, respectively.

EXAMPLE 14

In reference to Example 8, determine $\text{Var}(X | Y = y)$ by using the second formula in (7).

DISCUSSION By (7),

$$\begin{aligned} \text{Var}(X | Y = y) &= E(X^2 | Y = y) - [E(X | Y = y)]^2 \\ &= \int_0^y x^2 \cdot \frac{2x}{y^2} dx - \left(\frac{2y}{3}\right)^2 \quad (\text{by Examples 8(ii) and 12}) \\ &= \frac{2}{y^2} \int_0^y x^3 dx - \frac{4y^2}{9} = \frac{y^2}{2} - \frac{4y^2}{9} = \frac{y^2}{18}, \quad 0 < y < 1. \end{aligned}$$

Exercises

2.1 Refer to Exercise 1.1 and calculate the marginal p.d.f.'s f_X and f_Y .

2.2 Refer to Exercise 1.2 and calculate the marginal p.d.f.'s f_X and f_Y .

2.3 If the joint p.d.f. of the r.v.'s X and Y is given by the following table, determine the marginal p.d.f.'s f_X and f_Y .

$y \setminus x$	-4	-2	2	4
-2	0	0.25	0	0
-1	0	0	0	0.25
1	0.25	0	0	0
2	0	0	0.25	0

2.4 The r.v.'s X and Y take on the values 1, 2, and 3, as indicated in the following table:

$y \setminus x$	1	2	3
1	2/36	2/36	3/36
2	1/36	10/36	3/36
3	4/36	5/36	6/36

- (i) Determine the marginal p.d.f.'s f_X and f_Y .
(ii) Determine the conditional p.d.f.'s $f_{X|Y}(\cdot | y)$ and $f_{Y|X}(\cdot | x)$.

2.5 The r.v.'s X and Y have joint p.d.f. $f_{X,Y}$ given by the entries of the following table:

$y \setminus x$	0	1	2	3
1	1/8	1/16	3/16	1/8
2	1/16	1/16	1/8	1/4

- (i) Determine the marginal p.d.f.'s f_X and f_Y , and the conditional p.d.f. $f_{X|Y}(\cdot | y)$, $y = 1, 2$.
(ii) Calculate: EX , EY , $E(X | Y = y)$, $y = 1, 2$, and $E[E(X | Y)]$.
(iii) Compare EX and $E[E(X | Y)]$.
(iv) Calculate: $Var(X)$ and $Var(Y)$.

2.6 Let the r.v.'s X and Y have the joint p.d.f.:

$$f_{X,Y}(x, y) = \frac{2}{n(n+1)}, \quad y = 1, \dots, x; x = 1, \dots, n.$$

Then compute:

- (i) The marginal p.d.f.'s f_X and f_Y .
(ii) The conditional p.d.f.'s $f_{X|Y}(\cdot | y)$ and $f_{Y|X}(\cdot | x)$.
(iii) The conditional expectations $E(X | Y = y)$ and $E(Y | X = x)$.

Hint: Recall that: $\sum_{t=1}^n t = \frac{n(n+1)}{2}$.

2.7 In reference to Exercise 1.3, calculate the marginal p.d.f.'s f_X and f_Y .

2.8 Determine the marginal p.d.f.'s of the r.v.'s X and Y whose joint p.d.f. is given by:

$$f_{X,Y}(x, y) = \frac{6}{5}(x + y^2), \quad 0 \leq x \leq 1, \quad 0 \leq y \leq 1.$$

2.9 Let X and Y be two r.v.'s with joint p.d.f. given by:

$$f_{X,Y}(x, y) = ye^{-x}, \quad 0 < y \leq x < \infty.$$

- (i) Determine the marginal p.d.f.'s f_X and f_Y , and specify the range of the arguments involved.
- (ii) Determine the conditional p.d.f.'s $f_{X|Y}(\cdot | y)$ and $f_{Y|X}(\cdot | x)$, and specify the range of the arguments involved.
- (iii) Calculate the (conditional) probability $P(X > 2 \log 2 | Y = \log 2)$, where always \log stands for the natural logarithm.

2.10 The joint p.d.f. of the r.v.'s X and Y is given by:

$$f_{X,Y}(x, y) = xe^{-(x+y)}, \quad x > 0, \quad y > 0.$$

- (i) Determine the marginal p.d.f.'s f_X and f_Y .
- (ii) Determine the conditional p.d.f. $f_{Y|X}(\cdot | x)$.
- (iii) Calculate the probability $P(X > \log 4)$, where always \log stands for the natural logarithm.

2.11 The joint p.d.f. of the r.v.'s X and Y is given by:

$$f_{X,Y}(x, y) = \frac{1}{2}ye^{-xy}, \quad 0 < x < \infty, \quad 0 < y < 2.$$

- (i) Determine the marginal p.d.f. f_Y .
- (ii) Find the conditional p.d.f. $f_{X|Y}(\cdot | y)$, and evaluate it at $y = 1/2$.
- (iii) Compute the conditional expectation $E(X | Y = y)$, and evaluate it at $y = 1/2$.

2.12 In reference to Exercise 1.4, calculate:

- (i) The marginal p.d.f.'s f_X , f_Y , and the conditional p.d.f. $f_{Y|X}(\cdot | x)$; in all cases, specify the range of the variables involved.
- (ii) EY and $E(Y | X = x)$.
- (iii) $E[E(Y | X)]$ and observe that it is equal to EY .
- (iv) The probability $P(Y > \frac{1}{2} | X < \frac{1}{2})$.

2.13 In reference to Exercise 1.7, calculate:

- (i) The marginal p.d.f.'s f_X and f_Y .
- (ii) The conditional p.d.f.'s $f_{X|Y}(\cdot | y)$ and $f_{Y|X}(\cdot | x)$.
- (iii) The probability $P(X \leq 1)$.

2.14 In reference to Exercise 1.8, determine the marginal p.d.f. f_Y and the conditional p.d.f. $f_{X|Y}(\cdot | y)$.

2.15 In reference to Exercise 1.9:

- (i) Determine the marginal p.d.f.'s f_X and f_Y .
- (ii) Determine the conditional p.d.f. $f_{X|Y}(\cdot | y)$.
- (iii) Calculate the EX and $E(X | Y = y)$.
- (iv) Show that $E[E(X | Y)] = EX$.

2.16 In reference to Exercise 1.10, determine:

- (i) The marginal p.d.f. f_X .
- (ii) The conditional p.d.f. $f_{Y|X}(\cdot | x)$.

2.17 In reference to Exercise 1.11, determine:

- (i) The marginal p.d.f. f_Y .
- (ii) The conditional p.d.f. $f_{X|Y}(\cdot | y)$.
- (iii) The marginal p.d.f. f_X .

2.18 (i) For a fixed $y > 0$, consider the function $f(x, y) = e^{-y \frac{y^x}{x!}}$, $x = 0, 1, \dots$ and show that it is the conditional p.d.f. of a r.v., given that another r.v. $Y = y$.

- (ii) Now, suppose that the marginal p.d.f. of Y is Negative Exponential with parameter $\lambda = 1$. Determine the joint p.d.f. of the r.v.'s X and Y .
- (iii) Show that the marginal p.d.f. f_X is given by:

$$f_X(x) = \left(\frac{1}{2}\right)^{x+1}, \quad x = 0, 1, \dots$$

2.19 Suppose the r.v. Y is distributed as $P(\lambda)$ and that the conditional p.d.f. of a r.v. X , given $Y = y$, is $B(y, p)$. Then show that:

- (i) The marginal p.d.f. f_X is Poisson with parameter λp .
- (ii) The conditional p.d.f. $f_{Y|X}(\cdot | x)$ is Poisson with parameter λq (with $q = 1 - p$) over the set: $x, x + 1, \dots$

2.20 (i) Let X and Y be two discrete r.v.'s with joint p.d.f. $f_{X,Y}$. Then show that the conditional variance of X , given Y , satisfies the following relation:

$$\text{Var}(X | Y = y_j) = E(X^2 | Y = y_j) - [E(X | Y = y_j)]^2.$$

- (ii) Establish the same relation, if the r.v.'s X and Y are of the continuous type.

4.3 Expectation of a Function of Two r.v.'s, Joint and Marginal m.g.f.'s, Covariance, and Correlation Coefficient

In this section, a function of the r.v.'s X and Y is considered and its expectation and variance are defined. As a special case, one obtains the joint m.g.f. of X and Y , the covariance of X and Y , and their correlation coefficient. To this end, let g be a real-valued function defined on \mathfrak{R}^2 , so that $g(X, Y)$ is a r.v. Then the expectation of $g(X, Y)$ is defined as in (6) in Chapter 3 except that the joint p.d.f. of X and Y is to be used. Thus:

$$Eg(X, Y) = \sum_{x_i \in \mathfrak{R}, y_j \in \mathfrak{R}} g(x_i, y_j) f_{X,Y}(x_i, y_j) \quad \text{or} \quad \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f_{X,Y}(x, y) dx dy, \quad (8)$$

for the discrete and the continuous case, respectively, provided, of course, the quantities defined exist. Properties analogous to those in (4) in Chapter 3 apply here, too. Namely, for c and d constants:

$$E[cg(X, Y)] = cEg(X, Y), \quad E[cg(X, Y) + d] = cEg(X, Y) + d. \quad (9)$$

Also, if h is another real-valued function, then (see also Exercise 3.17):

$$g(X, Y) \leq h(X, Y) \quad \text{implies} \quad Eg(X, Y) \leq Eh(X, Y), \quad (10)$$

and, in particular,

$$g(X) \leq h(X) \quad \text{implies} \quad Eg(X) \leq Eh(X). \quad (11)$$

For the special choice of the function $g(x, y) = e^{t_1x+t_2y}$, t_1, t_2 reals, the expectation $E \exp(t_1X + t_2Y)$ defines a function in t_1, t_2 for those t_1, t_2 for which this expectation is finite. That is:

$$M_{X,Y}(t_1, t_2) = Ee^{t_1X+t_2Y}, \quad (t_1, t_2) \in C \subseteq \mathfrak{N}^2. \quad (12)$$

Thus, for the discrete and the continuous case, we have, respectively,

$$M_{X,Y}(t_1, t_2) = \sum_{x_i \in \mathfrak{N}, y_j \in \mathfrak{N}} e^{t_1x_i+t_2y_j} f_{X,Y}(x_i, y_j), \quad (13)$$

and

$$M_{X,Y}(t_1, t_2) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{t_1x+t_2y} f_{X,Y}(x, y) dx dy. \quad (14)$$

The function $M_{X,Y}(\cdot, \cdot)$ so defined is called the *joint m.g.f.* of the r.v.'s X and Y . Clearly, $M_{X,Y}(0, 0) = 1$ for any X and Y , and it may happen that $C = \{(0, 0)\}$ or $C \subset \mathfrak{N}^2$ or $C = \mathfrak{N}^2$. Here are two examples of joint m.g.f.'s.

EXAMPLE 15

Refer to Example 1 and calculate the joint m.g.f. of the r.v.'s involved.

DISCUSSION For any $t_1, t_2 \in \mathfrak{N}$, we have, by means of (13):

$$\begin{aligned} M_{X,Y}(t_1, t_2) &= \sum_{x=0}^3 \sum_{y=0}^3 e^{t_1x+t_2y} f_{X,Y}(x, y) \\ &= 0.05 + 0.20e^{t_2} + 0.21e^{t_1} + 0.26e^{t_1+t_2} + 0.06e^{t_1+2t_2} + 0.08e^{2t_1+t_2} \\ &\quad + 0.07e^{2t_1+2t_2} + 0.03e^{2t_1+3t_2} + 0.02e^{3t_1+2t_2} + 0.02e^{3t_1+3t_2}. \end{aligned} \quad (15)$$

EXAMPLE 16

Refer to Example 2 and calculate the joint m.g.f. of the r.v.'s involved.

DISCUSSION By means of (14), we have here:

$$\begin{aligned} M_{X,Y}(t_1, t_2) &= \int_0^{\infty} \int_0^{\infty} e^{t_1x+t_2y} \lambda_1 \lambda_2 e^{-\lambda_1x-\lambda_2y} dx dy \\ &= \int_0^{\infty} \lambda_1 e^{-(\lambda_1-t_1)x} dx \cdot \int_0^{\infty} \lambda_2 e^{-(\lambda_2-t_2)y} dy. \end{aligned}$$

But $\int_0^\infty \lambda_1 e^{-(\lambda_1-t_1)x} dx = -\frac{\lambda_1}{\lambda_1-t_1} e^{-(\lambda_1-t_1)x} \Big|_0^\infty = \frac{\lambda_1}{\lambda_1-t_1}$, provided $t_1 < \lambda_1$, and likewise $\int_0^\infty \lambda_2 e^{-(\lambda_2-t_2)y} dy = \frac{\lambda_2}{\lambda_2-t_2}$ for $t_2 < \lambda_2$. (We arrive at the same results without integration by recalling (Example 6) that the r.v.'s X and Y have the Negative Exponential distributions with parameters λ_1 and λ_2 , respectively.) Thus,

$$M_{X,Y}(t_1, t_2) = \frac{\lambda_1}{\lambda_1 - t_1} \times \frac{\lambda_2}{\lambda_2 - t_2}, \quad t_1 < \lambda_1, \quad t_2 < \lambda_2. \quad (16)$$

In (12), by setting successively $t_2 = 0$ and $t_1 = 0$, we obtain:

$$M_{X,Y}(t_1, 0) = Ee^{t_1 X} = M_X(t_1), \quad M_{X,Y}(0, t_2) = Ee^{t_2 Y} = M_Y(t_2). \quad (17)$$

Thus, the m.g.f.'s of the individual r.v.'s X and Y are taken as marginals from the joint m.g.f. of X and Y , and they are referred to as *marginal m.g.f.'s*. For example, in reference to (15) and (16), we obtain:

$$M_X(t_1) = 0.25 + 0.53 e^{t_1} + 0.18 e^{2t_1} + 0.04 e^{3t_1}, \quad t_1 \in \mathfrak{R}, \quad (18)$$

$$M_Y(t_2) = 0.26 + 0.54 e^{t_2} + 0.15 e^{2t_2} + 0.05 e^{3t_2}, \quad t_2 \in \mathfrak{R}, \quad (19)$$

and

$$M_X(t_1) = \frac{\lambda_1}{\lambda_1 - t_1}, \quad t_1 < \lambda_1, \quad M_Y(t_2) = \frac{\lambda_2}{\lambda_2 - t_2}, \quad t_2 < \lambda_2. \quad (20)$$

The joint m.g.f., as defined in (12), has properties analogous to the ones stated in (12) of Chapter 3. Namely, for c_1, c_2 and d_1, d_2 constants:

$$M_{c_1 X + d_1, c_2 Y + d_2}(t_1, t_2) = e^{d_1 t_1 + d_2 t_2} M_{X,Y}(c_1 t_1, c_2 t_2). \quad (21)$$

Its simple justification is left as an exercise (see Exercise 3.2).

In the present context, a version of the properties stated in (13) of Chapter 3, is the following:

$$\frac{\partial}{\partial t_1} M_{X,Y}(t_1, t_2) \Big|_{t_1=t_2=0} = EX, \quad \frac{\partial}{\partial t_2} M_{X,Y}(t_1, t_2) \Big|_{t_1=t_2=0} = EY, \quad (22)$$

and

$$\frac{\partial^2}{\partial t_1 \partial t_2} M_{X,Y}(t_1, t_2) \Big|_{t_1=t_2=0} = E(XY), \quad (23)$$

provided one may interchange the order of differentiating and taking expectations. For example, for (23), we have:

$$\begin{aligned} \frac{\partial^2}{\partial t_1 \partial t_2} M_{X,Y}(t_1, t_2) \Big|_{t_1=t_2=0} &= \frac{\partial^2}{\partial t_1 \partial t_2} Ee^{t_1 X + t_2 Y} \Big|_{t_1=t_2=0} \\ &= E \left(\frac{\partial^2}{\partial t_1 \partial t_2} e^{t_1 X + t_2 Y} \Big|_{t_1=t_2=0} \right) \\ &= E(XY e^{t_1 X + t_2 Y} \Big|_{t_1=t_2=0}) = E(XY). \end{aligned}$$

REMARK 2 Although properties (21) and (22) allow us to obtain moments by means of the m.g.f.'s of the r.v.'s X and Y , the most significant property of the m.g.f. is that it allows (under certain conditions) to retrieve the distribution of the r.v.'s X and Y . This is done through the so-called *inversion formula*.

Now, select the function g as follows: $g(x, y) = cx + dy$, where c and d are constants. Then, for the continuous case:

$$\begin{aligned} Eg(X, Y) &= E(cX + dY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (cx + dy)f_{X,Y}(x, y)dx dy \\ &= c \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xf_{X,Y}(x, y)dx dy + d \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} yf_{X,Y}(x, y)dx dy \\ &= c \int_{-\infty}^{\infty} \left[x \int_{-\infty}^{\infty} f_{X,Y}(x, y)dy \right] dx + d \int_{-\infty}^{\infty} \left[y \int_{-\infty}^{\infty} f_{X,Y}(x, y)dx \right] dy \\ &= c \int_{-\infty}^{\infty} xf_X(x)dx + d \int_{-\infty}^{\infty} yf_Y(y)dy = cEX + dEY; \text{ i.e.,} \end{aligned}$$

assuming the expectations involved exist:

$$E(cX + dY) = cEX + dEY, \quad \text{where } c \text{ and } d \text{ are constants.} \quad (24)$$

In the discrete case, integrals are replaced by summation signs. On account of the usual properties of integrals and summations, property (24) applies to a more general situation. Thus, for two functions g_1 and g_2 , we have:

$$E[g_1(X, Y) + g_2(X, Y)] = Eg_1(X, Y) + Eg_2(X, Y), \quad (25)$$

provided the expectations involved exist.

Next, suppose the r.v.'s X and Y have finite expectations and take $g(x, y) = (x - EX)(y - EY)$. Then the $Eg(X, Y) = E[(X - EX)(Y - EY)]$ is called the *covariance* of the r.v.'s X and Y and is denoted by $Cov(X, Y)$. Thus:

$$Cov(X, Y) = E[(X - EX)(Y - EY)] = E(XY) - (EX)(EY). \quad (26)$$

The second equality in (26) follows by multiplying out $(X - EX)(Y - EY)$ and applying property (25).

The variance of a single r.v. has been looked upon as a measure of dispersion of the distribution of the r.v. Some motivation will be given subsequently to the effect that the $Cov(X, Y)$ may be thought of as a measure of the degree to which X and Y tend to increase or decrease simultaneously when $Cov(X, Y) > 0$ or to move toward opposite directions when $Cov(X, Y) < 0$. This point is sufficiently made by the following simple example.

EXAMPLE 17

Consider the events A and B with $P(A)P(B) > 0$ and set $X = I_A$ and $Y = I_B$ for the *indicator* functions, where $I_A(s) = 1$ if $s \in A$ and $I_A(s) = 0$ if $s \in A^c$. Then, clearly, $EX = P(A)$, $EY = P(B)$, and $XY = I_{A \cap B}$, so that $E(XY) = P(A \cap B)$.

It follows that $Cov(X, Y) = P(A \cap B) - P(A)P(B)$. Next,

$$\begin{aligned} P(A)[P(Y = 1 | X = 1) - P(Y = 1)] &= P(A \cap B) - P(A)P(B) \\ &= Cov(X, Y), \end{aligned} \quad (27)$$

$$\begin{aligned} P(A^c)[P(Y = 0 | X = 0) - P(Y = 0)] &= P(A^c \cap B^c) - P(A^c)P(B^c) \\ &= P(A \cap B) - P(A)P(B) = Cov(X, Y), \end{aligned} \quad (28)$$

$$\begin{aligned} P(A^c)[P(Y = 1 | X = 0) - P(Y = 1)] &= P(A^c \cap B) - P(A^c)P(B) \\ &= -[P(A \cap B) - P(A)P(B)] = -Cov(X, Y), \end{aligned} \quad (29)$$

$$\begin{aligned} P(A)[P(Y = 0 | X = 1) - P(Y = 0)] &= P(A \cap B^c) - P(A)P(B^c) \\ &= -[P(A \cap B) - P(A)P(B)] = -Cov(X, Y), \end{aligned} \quad (30)$$

(see also Exercise 3.3).

From (27) and (28), it follows that $Cov(X, Y) > 0$ if and only if $P(Y = 1 | X = 1) > P(Y = 1)$, or $P(Y = 0 | X = 0) > P(Y = 0)$. That is, $Cov(X, Y) > 0$ if and only if, given that X has taken a “large” value (namely, 1), it is more likely that Y does so as well than it otherwise would; also, given that X has taken a “small” value (namely, 0), it is more likely that Y does so too than it otherwise would. On the other hand, from relations (29) and (30), we see that $Cov(X, Y) < 0$ if and only if $P(Y = 1 | X = 0) > P(Y = 1)$, or $P(Y = 0 | X = 1) > P(Y = 0)$. That is, $Cov(X, Y) < 0$ if and only if, given that X has taken a “small” value, it is more likely for Y to take a “large” value than it otherwise would, and given that X has taken a “large” value, it is more likely for Y to take a “small” value than it otherwise would.

As a further illustration of the significance of the covariance we proceed to calculate the $Cov(X, Y)$ for the r.v.'s of Example 1.

EXAMPLE 18

Refer to Example 1 and calculate the $Cov(X, Y)$.

DISCUSSION In Example 5, the (marginal) p.d.f.'s f_X and f_Y were calculated. Then: $EX = 1.01$ and $EY = 0.99$. Next, the r.v. XY is distributed as follows, on the basis of Table 4.1.

xy	0	1	2	3	4	6	9
$f_{X,Y}$	0.46	0.26	0.14	0	0.07	0.05	0.02

Therefore $E(XY) = 1.3$ and then, by formula (26), $Cov(X, Y) = 1.3 - 1.01 \times 0.99 = 0.3001$.

Here the covariance is positive, and by comparing the values of the conditional probabilities in Example 5 with the appropriate unconditional probabilities, we see that this is consonant with the observation just made that X and Y tend to take simultaneously either “large” values or “small” values. (See also Example 19 later.)

The result obtained next provides the range of values of the covariance of two r.v.'s; it is also referred to as a version of the *Cauchy–Schwarz inequality*.

THEOREM 1

- (i) Consider the r.v.'s X and Y with $EX = EY = 0$ and $\text{Var}(X) = \text{Var}(Y) = 1$. Then always $-1 \leq E(XY) \leq 1$, and $E(XY) = 1$ if and only if $P(X = Y) = 1$, and $E(XY) = -1$ if and only if $P(X = -Y) = 1$.
- (ii) For any r.v.'s X and Y with finite expectations and positive variances σ_X^2 and σ_Y^2 , it always holds:

$$-\sigma_X\sigma_Y \leq \text{Cov}(X, Y) \leq \sigma_X\sigma_Y, \quad (31)$$

and $\text{Cov}(X, Y) = \sigma_X\sigma_Y$ if and only if $P[Y = EY + \frac{\sigma_Y}{\sigma_X}(X - EX)] = 1$,
 $\text{Cov}(X, Y) = -\sigma_X\sigma_Y$ if and only if $P[Y = EY - \frac{\sigma_Y}{\sigma_X}(X - EX)] = 1$.

PROOF

(i) Clearly, $0 \leq E(X - Y)^2 = EX^2 + EY^2 - 2E(XY) = 2 - 2E(XY)$, so that $E(XY) \leq 1$; also, $0 \leq E(X + Y)^2 = EX^2 + EY^2 + 2E(XY) = 2 + 2E(XY)$, so that $-1 \leq E(XY)$. Combining these results, we obtain $-1 \leq E(XY) \leq 1$. As for equalities, observe that, if $P(X = Y) = 1$, then $E(XY) = EX^2 = 1$, and if $P(X = -Y) = 1$, then $E(XY) = -EX^2 = -1$. Next, $E(XY) = 1$ implies $E(X - Y)^2 = 0$ or $\text{Var}(X - Y) = 0$. But then $P(X - Y = 0) = 1$ or $P(X = Y) = 1$ (see Exercise 2.4 in Chapter 3). Also, $E(XY) = -1$ implies $E(X + Y)^2 = 0$ or $\text{Var}(X + Y) = 0$, so that $P(X = -Y) = 1$ (by the exercise just cited).

(ii) Replace the r.v.'s X and Y by the r.v.'s $X^* = \frac{X - EX}{\sigma_X}$ and $Y^* = \frac{Y - EY}{\sigma_Y}$, for which $EX^* = EY^* = 0$ and $\text{Var}(X^*) = \text{Var}(Y^*) = 1$. Then the inequalities $-1 \leq E(X^*Y^*) \leq 1$ become

$$-1 \leq E \left[\left(\frac{X - EX}{\sigma_X} \right) \left(\frac{Y - EY}{\sigma_Y} \right) \right] \leq 1 \quad (32)$$

from which (31) follows. Also, $E(X^*Y^*) = 1$ if and only if $P(X^* = Y^*) = 1$ becomes $E[(X - EX)(Y - EY)] = \sigma_X\sigma_Y$ if and only if $P[Y = EY + \frac{\sigma_Y}{\sigma_X}(X - EX)] = 1$, and $E(X^*Y^*) = -1$ if and only if $P(X^* = -Y^*) = 1$ becomes $E[(X - EX)(Y - EY)] = -\sigma_X\sigma_Y$ if and only if $P[Y = EY - \frac{\sigma_Y}{\sigma_X}(X - EX)] = 1$. A restatement of the last two conclusions is: $\text{Cov}(X, Y) = \sigma_X\sigma_Y$ if and only if $P[Y = EY + \frac{\sigma_Y}{\sigma_X}(X - EX)] = 1$, and $\text{Cov}(X, Y) = -\sigma_X\sigma_Y$ if and only if $P[Y = EY - \frac{\sigma_Y}{\sigma_X}(X - EX)] = 1$. ▲

From the definition of the $\text{Cov}(X, Y)$ in (26), it follows that if X is measured in units, call them a , and Y is measured in units, call them b , then $\text{Cov}(X, Y)$ is measured in units ab . Furthermore, because the variance of a r.v. ranges from 0 to ∞ , it follows from (31) that $\text{Cov}(X, Y)$ may vary from $-\infty$ to ∞ . These two characteristics of a covariance are rather undesirable and are both eliminated through the standardization process of replacing X and Y by $\frac{X - EX}{\sigma_X}$ and $\frac{Y - EY}{\sigma_Y}$. By (32), the range of the covariance of these standardized r.v.'s is the interval

$[-1, 1]$. This covariance is called the *correlation coefficient* of the r.v.'s X and Y and is denoted by $\rho(X, Y)$. Thus:

$$\begin{aligned}\rho(X, Y) &= E\left[\left(\frac{X - EX}{\sigma_X}\right)\left(\frac{Y - EY}{\sigma_Y}\right)\right] \\ &= \frac{\text{Cov}(X, Y)}{\sigma_X\sigma_Y} = \frac{E(XY) - (EX)(EY)}{\sigma_X\sigma_Y}.\end{aligned}\quad (33)$$

Furthermore, by (32):

$$-1 \leq \rho(X, Y) \leq 1, \quad (34)$$

and, by part (ii) of Theorem 1:


$$\rho(X, Y) = 1 \text{ if and only if } P\left[Y = EY + \frac{\sigma_Y}{\sigma_X}(X - EX)\right] = 1, \quad (35)$$

$$\rho(X, Y) = -1 \text{ if and only if } P\left[Y = EY - \frac{\sigma_Y}{\sigma_X}(X - EX)\right] = 1. \quad (36)$$

The straight lines represented by $y = EY + \frac{\sigma_Y}{\sigma_X}(x - EX)$ and $y = EY - \frac{\sigma_Y}{\sigma_X}(x - EX)$ are depicted in Figure 4.4.

Figure 4.4

**Lines of Perfect
Linear Relation of x
and y**



From relation (35), we have that $\rho(X, Y) = 1$ if and only if (X, Y) are linearly related (with probability 1). On the other hand, from Example 17, we have that $\text{Cov}(X, Y) > 0$ if and only if X and Y tend to take simultaneously either “large” values or “small” values. Since $\text{Cov}(X, Y)$ and $\rho(X, Y)$ have the same sign, the same statement can be made about $\rho(X, Y)$, being positive if and only if X and Y tend to take simultaneously either “large” values or “small” values. The same arguments apply for the case that $\text{Cov}(X, Y) < 0$ (equivalently, $\rho(X, Y) < 0$). This reasoning indicates that $\rho(X, Y)$ may be looked upon as a measure of *linear* dependence between X and Y . The pair (X, Y) lies on the line $y = EY + \frac{\sigma_Y}{\sigma_X}(x - EX)$ if $\rho(X, Y) = 1$; pairs identical to (X, Y) tend to be arranged along this line, if $(0 <) \rho(X, Y) < 1$, and they tend to move further and further away from this line as $\rho(X, Y)$ gets closer to 0; the pairs bear no sign of linear tendency whatever, if $\rho(X, Y) = 0$. Rough arguments also hold for the reverse assertions. For $0 < \rho(X, Y) \leq 1$, the r.v.'s X and Y are said to be *positively correlated*, and *uncorrelated* if $\rho(X, Y) = 0$. Likewise, the pair (X, Y) lies on the line $y = EY - \frac{\sigma_Y}{\sigma_X}(x - EX)$ if $\rho(X, Y) = -1$; pairs identical


to (X, Y) tend to be arranged along this line if $-1 < \rho(X, Y) < 0$. Again, rough arguments can also be made for the reverse assertions. For $-1 \leq \rho(X, Y) < 0$, the r.v.'s X and Y are said to be *negatively correlated*.

Actually, a more precise argument to this effect can be made by considering the distance D of the (random) point (X, Y) from the lines $y = EY \pm \frac{\sigma_Y}{\sigma_X}(x - EX)$. It can be seen that:

$$ED^2 = \frac{2\sigma_X^2\sigma_Y^2}{\sigma_X^2 + \sigma_Y^2}(1 - |\rho(X, Y)|). \quad (37)$$

Then one may use the interpretation of the expectation as an average and exploit (37) in order to arrive at the same reasoning but in a more rigorous way.

Figure 4.5



As an illustration, let us calculate the $\rho(X, Y)$ for Examples 1 and 8.

EXAMPLE 19

In reference to Example 1, calculate the $Cov(X, Y)$ and the $\rho(X, Y)$.

DISCUSSION From Table 4.1, we find $EX^2 = 1.61$, $EY^2 = 1.59$. By Example 18, $EX = 1.01$, $EY = 0.99$, so that $Var(X) = EX^2 - (EX)^2 = 0.5899$,

$Var(Y) = EY^2 - (EY)^2 = 0.6099$. Since $Cov(X, Y) = 0.3001$ (by Example 18), we have: $\rho(X, Y) = \frac{Cov(X, Y)}{\sqrt{Var(X)Var(Y)}} = \frac{0.3001}{\sqrt{0.5899 \times 0.6099}} \simeq 0.5$.

EXAMPLE 20

In reference to Example 8, calculate the $Cov(X, Y)$ and the $\rho(X, Y)$.

DISCUSSION By Example 8(iii), $Cov(X, Y) = E(XY) - (EX)(EY) = \frac{4}{9} - \frac{8}{15} \cdot \frac{4}{5} = \frac{4}{225}$, $\sqrt{Var(X)} = \frac{\sqrt{11}}{15}$, $\sqrt{Var(Y)} = \frac{\sqrt{6}}{15}$, so that

$$\rho(X, Y) = \frac{Cov(X, Y)}{\sqrt{Var(X)}\sqrt{Var(Y)}} = \frac{\frac{4}{225}}{\frac{\sqrt{11}}{15} \cdot \frac{\sqrt{6}}{15}} = \frac{4}{\sqrt{66}} \simeq 0.492.$$

EXAMPLE 21

Let X and Y be two r.v.'s with finite expectations and equal (finite) variances, and set $U = X + Y$ and $V = X - Y$. Then the r.v.'s U and V are uncorrelated.

DISCUSSION Indeed,

$$\begin{aligned} E(UV) &= E[(X + Y)(X - Y)] = E(X^2 - Y^2) = EX^2 - EY^2, \\ (EU)(EV) &= [E(X + Y)][E(X - Y)] \\ &= (EX + EY)(EX - EY) = (EX)^2 - (EY)^2, \end{aligned}$$

so that

$$\begin{aligned} Cov(U, V) &= E(UV) - (EU)(EV) = [EX^2 - EY^2] - [EX^2 - EY^2] \\ &= Var(X) - Var(Y) = 0. \end{aligned}$$

Figure 4.5 illustrates the behavior of the correlation coefficient $\rho(X, Y)$ of the r.v.'s X and Y . In (a), $\rho(X, Y) = 1$, the r.v.'s X and Y are perfectly positively linearly related. In (b), $\rho(X, Y) = -1$, the r.v.'s X and Y are perfectly negatively linearly related. In (c), $0 < \rho(X, Y) < 1$, the r.v.'s X and Y are positively correlated. In (d), $-1 < \rho(X, Y) < 0$, the r.v.'s X and Y are negatively correlated. In (e), $\rho(X, Y) = 0$, the r.v.'s X and Y are uncorrelated.

The following result presents an interesting property of the correlation coefficient.

THEOREM 2

Let X and Y be r.v.'s with finite first and second moments and positive variances, and let c_1, c_2, d_1, d_2 be constants with $c_1c_2 \neq 0$. Then:

$$\rho(c_1X + d_1, c_2Y + d_2) = \pm \rho(X, Y), \text{ with } + \text{ if } c_1c_2 > 0 \text{ and } - \text{ if } c_1c_2 < 0.$$

(38)

PROOF Indeed, $Var(c_1X + d_1) = c_1^2Var(X)$, $Var(c_2Y + d_2) = c_2^2Var(Y)$, and $Cov(c_1X + d_1, c_2Y + d_2) = E\{[(c_1X + d_1) - E(c_1X + d_1)][(c_2Y + d_2) - E(c_2Y + d_2)]\} = E\{c_1(X - EX) \cdot c_2(Y - EY)\} = c_1c_2E[(X - EX)(Y - EY)] = c_1c_2Cov(X, Y)$. Therefore $\rho(c_1X + d_1, c_2Y + d_2) = \frac{c_1c_2Cov(X, Y)}{|c_1c_2|\sqrt{Var(X)Var(Y)}}$, and the conclusion follows. ▲

EXAMPLE 22

Let X and Y be temperatures in two localities measured in the Celsius scale, and let U and V be the same temperatures measured in the Fahrenheit scale. Then $\rho(X, Y) = \rho(U, V)$, as it should be. This is so because $U = \frac{9}{5}X + 32$ and $V = \frac{9}{5}Y + 32$, so that (38) applies with the + sign.

This section is concluded with the following result and an example.

THEOREM 3

For two r.v.'s X and Y with finite first and second moments, and (positive) standard deviations σ_X and σ_Y , it holds:

$$\text{Var}(X + Y) = \sigma_X^2 + \sigma_Y^2 + 2\text{Cov}(X, Y) = \sigma_X^2 + \sigma_Y^2 + 2\sigma_X\sigma_Y\rho(X, Y), \quad (39)$$

and

$$\text{Var}(X + Y) = \sigma_X^2 + \sigma_Y^2 \quad \text{if } X \text{ and } Y \text{ are uncorrelated.} \quad (40)$$

PROOF Since (40) follows immediately from (39), and $\text{Cov}(X, Y) = \sigma_X\sigma_Y \times \rho(X, Y)$, it suffices to establish only the first equality in (39). Indeed,

$$\begin{aligned} \text{Var}(X + Y) &= E[(X + Y) - E(X + Y)]^2 = E[(X - EX) + E(Y - EY)]^2 \\ &= E(X - EX)^2 + E(Y - EY)^2 + 2E[(X - EX)(Y - EY)] \\ &= \sigma_X^2 + \sigma_Y^2 + 2\text{Cov}(X, Y). \quad \blacktriangle \end{aligned}$$

EXAMPLE 23

In reference to Examples 1 and 8 and by means of results obtained in Examples 19, 8(iii), and 20, respectively, calculate $\text{Var}(X + Y)$.

DISCUSSION By (39),

$$\begin{aligned} \text{Var}(X + Y) &= \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y) \\ &= 0.5899 + 0.6099 + 2 \times 0.3001 = 1.8 \quad \text{for Example 1, and} \\ &= \frac{11}{225} + \frac{2}{75} + 2 \times \frac{4}{225} = \frac{1}{9} \quad \text{for Example 8.} \end{aligned}$$

Exercises

3.1 Let X and Y be the r.v.'s denoting the number of sixes when two fair dice are rolled independently 15 times each. Determine the $E(X + Y)$.

3.2 Show that the joint m.g.f. of two r.v.'s X and Y satisfies the following property, where c_1, c_2, d_1 , and d_2 are constants.

$$M_{c_1X+d_1, c_2Y+d_2}(t_1, t_2) = e^{d_1t_1+d_2t_2} M_{X,Y}(c_1t_1, c_2t_2).$$

3.3 Provide a justification of relations (28)–(30). That is:

- (i) $P(A^c \cap B^c) - P(A^c)P(B^c) = P(A \cap B) - P(A)P(B)$.
- (ii) $P(A^c \cap B) - P(A^c)P(B) = -P(A \cap B) + P(A)P(B)$.
- (iii) $P(A \cap B^c) - P(A)P(B^c) = -P(A \cap B) + P(A)P(B)$.

3.4 Let X and Y be two r.v.'s with $EX = EY = 0$. Then, if $\text{Var}(X - Y) = 0$, it follows that $P(X = Y) = 1$, and if $\text{Var}(X + Y) = 0$, then $P(X = -Y) = 1$.

Hint: Use Exercise 2.4 in Chapter 3.

3.5 In reference to Exercise 2.1 (see also Exercise 1.1), calculate:

- (i) EX , EY , $\text{Var}(X)$, and $\text{Var}(Y)$.
- (ii) $\text{Cov}(X, Y)$ and $\rho(X, Y)$.
- (iii) Decide on the kind of correlation of the r.v.'s X and Y .

3.6 Refer to Exercises 1.2 and 2.2 and calculate:

- (i) EX , EY , $\text{Var}(X)$, $\text{Var}(Y)$.
- (ii) $E(XY)$, $\text{Cov}(X, Y)$.
- (iii) $\rho(X, Y)$.
- (iv) What kind of correlation, if any, do the r.v.'s X and Y exhibit?

3.7 In reference to Exercise 2.3:

- (i) Calculate EX , EY , $\text{Var}(X)$, and $\text{Var}(Y)$.
- (ii) Calculate $\text{Cov}(X, Y)$ and $\rho(X, Y)$.
- (iii) Plot the points $(-4, 1)$, $(-2, 2)$, $(2, 2)$, and $(4, -1)$, and reconcile this graph with the value of $\rho(X, Y)$ found in part (ii).

3.8 In reference to Exercise 2.4, calculate the following quantities:

- (i) EX , EY , $\text{Var}(X)$, and $\text{Var}(Y)$.
- (ii) $\text{Cov}(X, Y)$ and $\rho(X, Y)$.

3.9 Refer to Exercise 2.5, and calculate the $\text{Cov}(X, Y)$ and the $\rho(X, Y)$.

3.10 Let X be a r.v. taking on the values $-2, -1, 1, 2$, each with probability $1/4$, and define the r.v. Y by: $Y = X^2$. Then calculate the quantities: EX , $\text{Var}(X)$, EY , $\text{Var}(Y)$, $E(XY)$, $\text{Cov}(X, Y)$, and $\rho(X, Y)$. Are you surprised by the value of $\rho(X, Y)$? Explain.

3.11 Refer to Example 8 and compute the covariance $\text{Cov}(X, Y)$ and the correlation coefficient $\rho(X, Y)$. Decide on the kind of correlation of the r.v.'s X and Y .

3.12 In reference to Exercise 2.7 (see also Exercise 1.3), calculate:

- (i) The expectations EX and EY .
- (ii) The variances $\text{Var}(X)$ and $\text{Var}(Y)$.
- (iii) The covariance $\text{Cov}(X, Y)$ and the correlation coefficient $\rho(X, Y)$.
- (iv) On the basis of part (iii), decide on the kind of correlation of the r.v.'s X and Y .

3.13 In reference to Exercise 2.8, calculate:

- (i) The expectations EX and EY .
- (ii) The variances $\text{Var}(X)$ and $\text{Var}(Y)$.

- (iii) The covariance $Cov(X, Y)$ and the correlation coefficient $\rho(X, Y)$.
- (iv) On the basis of part (iii), decide on the kind of correlation of the r.v.'s X and Y .

3.14 Let X be a r.v. with finite expectation and finite and positive variance, and set $Y = aX + b$, where a and b are constants and $a \neq 0$. Then show that $|\rho(X, Y)| = 1$ and, indeed $\rho(X, Y) = 1$ if and only if $a > 0$, and $\rho(X, Y) = -1$ if and only if $a < 0$.

3.15 For any two r.v.'s X and Y , set $U = X + Y$ and $V = X - Y$. Then show that:

- (i) $P(UV < 0) = P(|X| < |Y|)$.
- (ii) If $EX^2 = EY^2 < \infty$, then $E(UV) = 0$.
- (iii) If $EX^2 < \infty$, $EY^2 < \infty$ and $Var(X) = Var(Y)$, then the r.v.'s U and V are uncorrelated.

3.16 Let X and Y be r.v.'s with finite second moments EX^2 , EY^2 , and $Var(X) > 0$. Suppose we know X and we wish to *predict* Y in terms of X through the *linear* relationship $Y = \alpha X + \beta$, where α and β are (unknown) constants. Further, suppose there exist values $\hat{\alpha}$ and $\hat{\beta}$ of α and β , respectively, for which the expectation of the square difference $[Y - (\hat{\alpha}X + \hat{\beta})]^2$ is minimum. Then $\hat{Y} = \hat{\alpha}X + \hat{\beta}$ is called the *best linear predictor* of Y in terms of X (when the criterion of optimality is that of minimizing $E[Y - (\alpha X + \beta)]^2$ over all α and β). Then show that $\hat{\alpha}$ and $\hat{\beta}$ are given as follows:

$$\hat{\alpha} = \frac{\sigma_Y}{\sigma_X} \rho(X, Y), \quad \hat{\beta} = EY - \hat{\alpha}EX,$$

where σ_X and σ_Y are the s.d.'s of the r.v.'s X and Y , respectively.

3.17 Justify the statement made in relation (10), for both the discrete and the continuous case.

4.4 Some Generalizations to k Random Variables

If instead of two r.v.'s X and Y we have k r.v.'s X_1, \dots, X_k , most of the concepts defined and results obtained in the previous sections are carried over to the k -dimensional case in a straightforward way. Thus, the *joint probability distribution* of (X_1, \dots, X_k) , to be denoted by P_{X_1, \dots, X_k} , is defined by: $P_{X_1, \dots, X_k}(B) = P[(X_1, \dots, X_k) \in B]$, $B \subseteq \mathfrak{R}^k = \mathfrak{R} \times \dots \times \mathfrak{R}$ (k factors), and their *joint d.f.* is: $F_{X_1, \dots, X_k}(x_1, \dots, x_k) = P(X_1 \leq x_1, \dots, X_k \leq x_k)$, $x_1, \dots, x_k \in \mathfrak{R}$. The obvious versions of properties #1 and #3 stated in Section 4.1 hold here too; also, a suitable version of property #2 holds, but we shall not insist on it. The *joint p.d.f.* of X_1, \dots, X_k is denoted by f_{X_1, \dots, X_k} and is defined in an obvious manner. Thus, for the case the r.v.'s X_1, \dots, X_k are discrete taking on respective values x_{1i}, \dots, x_{ki} , we have $f_{X_1, \dots, X_k}(x_{1i}, \dots, x_{ki}) = P(X_1 = x_{1i}, \dots, X_k = x_{ki})$ and 0 otherwise. Then, for $B \subseteq \mathfrak{R}^k$, $P[(X_1, \dots, X_k) \in B] = \sum f_{X_1, \dots, X_k}(x_1, \dots, x_k)$, where the summation extends over all $(x_1, \dots, x_k) \in B$. For the continuous case, the joint p.d.f. is a nonnegative function such that

$F_{X_1, \dots, X_k}(x_1, \dots, x_k) = \int_{-\infty}^{x_k} \cdots \int_{-\infty}^{x_1} f_{X_1, \dots, X_k}(t_1, \dots, t_k) dt_1 \dots dt_k$. It follows that for $B \subseteq \mathfrak{R}^k$ (where you may interpret B as a familiar geometric figure in \mathfrak{R}^k): $P[(X_1, \dots, X_k) \in B] = \underbrace{\int \cdots \int}_B f_{X_1, \dots, X_k}(x_1, \dots, x_k) dx_1 \dots dx_k$. As in the

2-dimensional case, $\frac{\partial^k}{\partial x_1 \dots \partial x_k} F_{X_1, \dots, X_k}(x_1, \dots, x_k) = f_{X_1, \dots, X_k}(x_1, \dots, x_k)$ (for continuity points (x_1, \dots, x_k) of f_{X_1, \dots, X_k}). In the next subsection, three concrete examples will be presented, one for the discrete case, and two for the continuous case. In the present k -dimensional case, there are many marginal d.f.'s and p.d.f.'s. Thus, if in $F_{X_1, \dots, X_k}(x_1, \dots, x_k)$, t of the x 's, x_{j_1}, \dots, x_{j_t} , are replaced by $+\infty$ (in the sense they are let to tend to $+\infty$), then what is left is the *marginal joint d.f.* of the r.v.'s X_{i_1}, \dots, X_{i_s} , $F_{X_{i_1}, \dots, X_{i_s}}$, where $s + t = k$. Likewise, if in $f_{X_1, \dots, X_k}(x_1, \dots, x_k)$, x_{j_1}, \dots, x_{j_t} are eliminated through summation (for the discrete case) or integration (for the continuous case), what is left is the *marginal joint p.d.f.* of the r.v.'s X_{i_1}, \dots, X_{i_s} , $f_{X_{i_1}, \dots, X_{i_s}}$. Combining joint and marginal joint p.d.f.'s, as in the 2-dimensional case, we obtain a variety of *conditional p.d.f.'s*. Thus, for example,

$$f_{X_{j_1}, \dots, X_{j_t} | X_{i_1}, \dots, X_{i_s}}(x_{j_1}, \dots, x_{j_t} | x_{i_1}, \dots, x_{i_s}) = \frac{f_{X_1, \dots, X_k}(x_1, \dots, x_k)}{f_{X_{i_1}, \dots, X_{i_s}}(x_{i_1}, \dots, x_{i_s})}.$$

Utilizing conditional p.d.f.'s, we can define *conditional expectations* and *conditional variances*, as in the 2-dimensional case (see relations (2), (3) and (5), (6)). For a (real-valued) function g defined on \mathfrak{R}^k , the expectation of the r.v. $g(X_1, \dots, X_k)$ is defined in a way analogous to that in (8) for the 2-dimensional case, and the validity of properties (9) and (10) is immediate. In particular, provided the expectations involved exist:

$$E(c_1 X_1 + \cdots + c_k X_k + d) = c_1 E X_1 + \cdots + c_k E X_k + d, \\ c_1, \dots, c_k, d \text{ constants.} \quad (41)$$

By choosing $g(x_1, \dots, x_k) = \exp(t_1 x_1 + \cdots + t_k x_k)$, $t_1, \dots, t_k \in \mathfrak{R}$, the resulting expectation (assuming it is finite) is the *joint m.g.f.* of X_1, \dots, X_k ; i.e.,

$$M_{X_1, \dots, X_k}(t_1, \dots, t_k) = E e^{t_1 X_1 + \cdots + t_k X_k}, \quad (t_1, \dots, t_k) \in C \subseteq \mathfrak{R}^k. \quad (42)$$

The appropriate versions of properties (21) and (23) become here:

$$M_{c_1 X_1 + d_1, \dots, c_k X_k + d_k}(t_1, \dots, t_k) = e^{d_1 t_1 + \cdots + d_k t_k} M_{X_1, \dots, X_k}(c_1 t_1, \dots, c_k t_k), \quad (43)$$

where c_1, \dots, c_k and d_1, \dots, d_k are constants, and:

$$\frac{\partial^{n_1 + \cdots + n_k}}{\partial^{n_1} t_1 \dots \partial^{n_k} t_k} M_{X_1, \dots, X_k}(t_1, \dots, t_k) |_{t_1 = \cdots = t_k = 0} = E(X_1^{n_1} \dots X_k^{n_k}), \quad (44)$$

for ≥ 0 integers n_1, \dots, n_k .

REMARK 3 Relation (44) demonstrates the joint moment generating property of the joint m.g.f. The joint m.g.f. can also be used for recovering the joint distribution of the r.v.'s X_1, \dots, X_k as indicated in Remark 2.

Finally, the appropriate versions of relations (39) and (40) become here, by setting $\sigma_{X_i}^2 = \text{Var}(X_i)$, $i = 1, \dots, k$:

$$\begin{aligned} \text{Var}(X_1 + \dots + X_k) &= \sum_{i=1}^k \sigma_{X_i}^2 + 2 \sum_{1 \leq i < j \leq k} \text{Cov}(X_i, X_j) \\ &= \sum_{i=1}^k \sigma_{X_i}^2 + 2 \sum_{1 \leq i < j \leq k} \sigma_{X_i} \sigma_{X_j} \rho(X_i, X_j), \end{aligned} \quad (45)$$

and

$$\text{Var}(X_1 + \dots + X_k) = \sum_{i=1}^k \sigma_{X_i}^2 \quad \text{if the } X_i\text{'s are pairwise uncorrelated;} \quad (46)$$

i.e., $\rho(X_i, X_j) = 0$ for $i \neq j$.

Exercises

4.1 If the r.v.'s X_1, X_2, X_3 have the joint p.d.f. $f_{X_1, X_2, X_3}(x_1, x_2, x_3) = c^3 e^{-c(x_1 + x_2 + x_3)}$, $x_1 > 0, x_2 > 0, x_3 > 0$ ($c > 0$), determine:

- (i) The constant c .
- (ii) The marginal p.d.f.'s f_{X_1} , f_{X_2} , and f_{X_3} .
- (iii) The joint conditional p.d.f. of X_1 and X_2 , given X_3 .
- (iv) The conditional p.d.f. of X_1 , given X_2 and X_3 .

4.2 Determine the joint m.g.f. of the r.v.'s X_1, X_2, X_3 with p.d.f. $f_{X_1, X_2, X_3}(x_1, x_2, x_3) = c^3 e^{-c(x_1 + x_2 + x_3)}$, $x_1 > 0, x_2 > 0, x_3 > 0$ (c any positive constant, see also Exercise 4.1).

4.3 (*Cramér-Wold devise*) Show that if we know the joint distribution of the r.v.'s X_1, \dots, X_n , then we can determine the distribution of any linear combination $c_1 X_1 + \dots + c_n X_n$ of X_1, \dots, X_n , where c_1, \dots, c_n are constants. Conversely, if we know the distribution of all linear combinations just described, then we can determine the joint distribution of X_1, \dots, X_n .

4.4 If the r.v.'s X_1, \dots, X_m and Y_1, \dots, Y_n have finite second moments, then show that:

$$\text{Cov} \left(\sum_{i=1}^m X_i, \sum_{j=1}^n Y_j \right) = \sum_{i=1}^m \sum_{j=1}^n \text{Cov}(X_i, Y_j).$$

4.5 The Multinomial, the Bivariate Normal, and the Multivariate Normal Distributions

In this section, we introduce and study to some extent three multidimensional distributions; they are the *Multinomial* distribution, the *2-dimensional Normal* or *Bivariate Normal* distribution, and the *k-dimensional Normal* distribution.

4.5.1 Multinomial Distribution

A *multinomial* experiment is a straightforward generalization of a binomial experiment, where, instead of 2, there are k (mutually exclusive) possible outcomes, O_1, \dots, O_k , say, occurring with respective probabilities p_1, \dots, p_k . Simple examples of multinomial experiments are those of rolling a die (with 6 possible outcomes); selecting (with replacement) r balls from a collection of $n_1 + \dots + n_k$ balls, so that n_i balls have the number i written on them, $i = 1, \dots, k$; selecting (with replacement) r objects out of a collection of objects of which n_1 are in good condition, n_2 have minor defects, and n_3 have serious defects, etc. Suppose a multinomial experiment is carried out independently n times and the probabilities p_1, \dots, p_k remain the same throughout. Denote by X_i the r.v. of the number of times outcome O_i occurs, $i = 1, \dots, k$. Then the joint p.d.f. of X_1, \dots, X_k is given by:

$$f_{X_1, \dots, X_k}(x_1, \dots, x_k) = \frac{n!}{x_1! \dots x_k!} p_1^{x_1} \dots p_k^{x_k}, \quad (47)$$

where x_1, \dots, x_k are ≥ 0 integers with $x_1 + \dots + x_k = n$, and, of course, $0 < p_i < 1$, $i = 1, \dots, k$, $p_1 + \dots + p_k = 1$. The distribution given by (47) is the *Multinomial* distribution with *parameters* n and p_1, \dots, p_k , and the r.v.'s X_1, \dots, X_k are said to have the *Multinomial* distribution with these parameters. That the right-hand side of (47) is the right formula for the joint probabilities $P(X_1 = x_1, \dots, X_k = x_k)$ ensues as follows: By independence, the probability that O_i occurs n_i times, $i = 1, \dots, k$, in specified positions, is given by: $p_1^{x_1} \dots p_k^{x_k}$ regardless of the positions of occurrence of O_i 's. The different ways of choosing the n_i positions for the occurrence of O_i , $i = 1, \dots, k$, is equal to: $\binom{n}{n_1} \binom{n-n_1}{n_2} \dots \binom{n-n_1-\dots-n_{k-1}}{n_k}$. Writing out each term in factorial form and making the obvious cancellations, we arrive at: $n!/(x_1! \dots x_k!)$ (see also Exercise 5.1). For illustrative purposes, let us consider the following example.

EXAMPLE 24

A fair die is rolled independently 10 times. Find the probability that faces #1 through #6 occur the following respective number of times: 2, 1, 3, 1, 2, and 1.

DISCUSSION By letting X_i be the r.v. denoting the number of occurrences of face i , $i = 1, \dots, 6$, we have:

$$f_{X_1, \dots, X_6}(2, 1, 3, 1, 2, 1) = \frac{10!}{2!1!3!1!2!1!} (1/6)^{10} = \frac{4,725}{1,889,568} \simeq 0.003.$$

In a Multinomial distribution, all marginal p.d.f.'s and all conditional p.d.f.'s are also Multinomial. More precisely, we have the following result.

THEOREM 4

Let X_1, \dots, X_k be Multinomially distributed with parameters n and p_1, \dots, p_k , and for $1 \leq s < k$, let $1 \leq i_1 < i_2 < \dots < i_s \leq k$, $Y = n - (X_{i_1} + \dots + X_{i_s})$ and $q = 1 - (p_{i_1} + \dots + p_{i_s})$. Then:

- (i) The r.v.'s $X_{i_1}, \dots, X_{i_s}, Y$ are distributed Multinomially with parameters n and $p_{i_1}, \dots, p_{i_s}, q$.
- (ii) The conditional joint distribution of X_{j_1}, \dots, X_{j_t} , given $X_{i_1} = x_{i_1}, \dots, X_{i_s} = x_{i_s}$, is Multinomial with parameters $n - r$ and $p_{j_1}/q, \dots, p_{j_t}/q$, where $r = x_{i_1} + \dots + x_{i_s}$ and $t = k - s$.

PROOF

- (i) For ≥ 0 integers x_{i_1}, \dots, x_{i_s} with $x_{i_1} + \dots + x_{i_s} = r \leq n$, we have:

$$\begin{aligned} f_{X_{i_1}, \dots, X_{i_s}}(x_{i_1}, \dots, x_{i_s}) &= P(X_{i_1} = x_{i_1}, \dots, X_{i_s} = x_{i_s}) \\ &= P(X_{i_1} = x_{i_1}, \dots, X_{i_s} = x_{i_s}, Y = n - r) \\ &= \frac{n!}{x_{i_1}! \dots x_{i_s}! (n - r)!} p_{i_1}^{x_{i_1}} \dots p_{i_s}^{x_{i_s}} q^{n-r}. \end{aligned}$$

- (ii) For ≥ 0 integers x_{j_1}, \dots, x_{j_t} with $x_{j_1} + \dots + x_{j_t} = n - r$, we have:

$$\begin{aligned} f_{X_{j_1}, \dots, X_{j_t} | X_{i_1}, \dots, X_{i_s}}(x_{j_1}, \dots, x_{j_t} | x_{i_1}, \dots, x_{i_s}) &= P(X_{j_1} = x_{j_1}, \dots, X_{j_t} = x_{j_t} | X_{i_1} = x_{i_1}, \dots, X_{i_s} = x_{i_s}) \\ &= P(X_{j_1} = x_{j_1}, \dots, X_{j_t} = x_{j_t}, X_{i_1} = x_{i_1}, \dots, X_{i_s} = x_{i_s}) / \\ &\quad P(X_{i_1} = x_{i_1}, \dots, X_{i_s} = x_{i_s}) \\ &= \frac{n!}{x_{j_1}! \dots x_{j_t}! x_{i_1}! \dots x_{i_s}!} p_{j_1}^{x_{j_1}} \dots p_{j_t}^{x_{j_t}} \times p_{i_1}^{x_{i_1}} \dots p_{i_s}^{x_{i_s}} / \\ &\quad \left(\frac{n!}{x_{i_1}! \dots x_{i_s}! (n - r)!} p_{i_1}^{x_{i_1}} \dots p_{i_s}^{x_{i_s}} q^{n-r} \right) \\ &= \frac{(n - r)!}{x_{j_1}! \dots x_{j_t}!} (p_{j_1}/q)^{x_{j_1}} \dots (p_{j_t}/q)^{x_{j_t}}. \quad \blacktriangle \end{aligned}$$

EXAMPLE 25

In reference to Example 24, calculate: $P(X_2 = X_4 = X_6 = 2)$ and $P(X_1 = X_3 = 1, X_5 = 2 | X_2 = X_4 = X_6 = 2)$.

DISCUSSION Here $n = 10$, $r = 6$, $p_2 = p_4 = p_6 = \frac{1}{6}$ and $q = 1 - \frac{3}{6} = \frac{1}{2}$. Thus:

$$P(X_2 = X_4 = X_6 = 2) = \frac{10!}{2!2!2!4!} \left(\frac{1}{6}\right)^6 \left(\frac{1}{2}\right)^4 = \frac{4,725}{186,624} \simeq 0.025,$$

and:

$$P(X_1 = X_3 = 1, X_5 = 2 | X_2 = X_4 = X_6 = 2) = \frac{4!}{1!1!2!} \left(\frac{1/6}{1/2}\right)^4 = \frac{4}{27} \simeq 0.148.$$

EXAMPLE 26

In a genetic experiment, two different varieties of a certain species are crossed and a specific characteristic of the offspring can occur only at three levels, A, B, and C, say. According to a proposed model, the probabilities for A, B, and C are $\frac{1}{12}$, $\frac{3}{12}$, and $\frac{8}{12}$, respectively. Out of 60 offspring, calculate:

- (i) The probability that 6, 18, and 36 fall into levels A, B, and C, respectively.
- (ii) The (conditional) probability that 6 and 18 fall into levels A and B, respectively, given that 36 falls into level C.

DISCUSSION

- (i) Formula (47) applies with $n = 60$, $k = 3$, $p_1 = \frac{1}{12}$, $p_2 = \frac{3}{12}$, $p_3 = \frac{8}{12}$, $x_1 = 6$, $x_2 = 18$, $x_3 = 36$ and yields:

$$P(X_1 = 6, X_2 = 18, X_3 = 36) = \frac{60!}{6!18!36!} \left(\frac{1}{12}\right)^6 \left(\frac{3}{12}\right)^{18} \left(\frac{8}{12}\right)^{36} \simeq 0.011.$$

- (ii) Here Theorem 4(ii) applies with $s = 1$, $t = 2$, $x_{i_1} = x_3 = 36$, $x_{j_1} = x_1 = 6$, $x_{j_2} = x_2 = 18$, $r = 36$, so that $n - r = 60 - 36 = 24$, $q = 1 - p_3 = 1 - \frac{8}{12} = \frac{4}{12}$, and yields:

$$\begin{aligned} P(X_1 = 6, X_2 = 18 / X_3 = 36) &= \frac{(n-r)!}{x_1!x_2!} \left(\frac{p_1}{q}\right)^{x_1} \left(\frac{p_2}{q}\right)^{x_2} \\ &= \frac{(24)!}{6!18!} \left(\frac{1/12}{4/12}\right)^6 \left(\frac{3/12}{4/12}\right)^{18} \\ &= \binom{24}{6} \left(\frac{1}{4}\right)^6 \left(\frac{3}{4}\right)^{18} \\ &= 0.1852 \quad (\text{from the Binomial tables}). \end{aligned}$$

An application of formula (42) gives the joint m.g.f. of X_1, \dots, X_k as follows, where the summation is over all ≥ 0 integers x_1, \dots, x_k with $x_1 + \dots + x_k = n$:

$$\begin{aligned} M_{X_1, \dots, X_k}(t_1, \dots, t_k) &= \sum e^{t_1 x_1 + \dots + t_k x_k} \frac{n!}{x_1! \dots x_k!} p_1^{x_1} \dots p_k^{x_k} \\ &= \sum \frac{n!}{x_1! \dots x_k!} (p_1 e^{t_1})^{x_1} \dots (p_k e^{t_k})^{x_k} \\ &= (p_1 e^{t_1} + \dots + p_k e^{t_k})^n; \text{ i.e.,} \\ M_{X_1, \dots, X_k}(t_1, \dots, t_k) &= (p_1 e^{t_1} + \dots + p_k e^{t_k})^n, \quad t_1, \dots, t_k \in \mathfrak{R}. \quad (48) \end{aligned}$$

By means of (44) and (48), we can find the $Cov(X_i, X_j)$ and the $\rho(X_i, X_j)$ for any $1 \leq i < j \leq k$. Indeed, $EX_i = np_i$, $EX_j = np_j$, $Var(X_i) = np_i(1 - p_i)$,

$\text{Var}(X_j) = np_j(1 - p_j)$ and $E(X_i X_j) = n(n - 1)p_i p_j$. Therefore:

$$\text{Cov}(X_i, X_j) = -np_i p_j \quad \text{and} \quad \rho(X_i, X_j) = -[p_i p_j / ((1 - p_i)(1 - p_j))]^{1/2} \quad (49)$$

(see Exercise 5.4 for details).

4.5.2 Bivariate Normal Distribution

The joint distribution of the r.v.'s X and Y is said to be the *Bivariate Normal* distribution with *parameters* μ_1, μ_2 in \mathfrak{R} , σ_1, σ_2 positive and $\rho \in [-1, 1]$, if the joint p.d.f. is given by the formula:

$$f_{X,Y}(x, y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} e^{-q/2}, \quad x, y \in \mathfrak{R}, \quad (50)$$


where

$$q = \frac{1}{1-\rho^2} \left[\left(\frac{x-\mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x-\mu_1}{\sigma_1} \right) \left(\frac{y-\mu_2}{\sigma_2} \right) + \left(\frac{y-\mu_2}{\sigma_2} \right)^2 \right]. \quad (51)$$

This distribution is also referred to as *2-dimensional Normal*. The shape of $f_{X,Y}$ looks like a bell sitting on the xy -plane and whose highest point is located at the point $(\mu_1, \mu_2, 1/(2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}))$ (see Figure 4.6).

Figure 4.6

Graphs of the p.d.f. of the Bivariate Normal Distribution: (a) Centered at the Origin; (b) Centered Elsewhere in the (x, y) -plane



That $f_{X,Y}$ integrates to 1 and therefore is a p.d.f. is seen by rewriting it in a convenient way. Specifically,

$$\begin{aligned} & \left(\frac{x-\mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x-\mu_1}{\sigma_1} \right) \left(\frac{y-\mu_2}{\sigma_2} \right) + \left(\frac{y-\mu_2}{\sigma_2} \right)^2 \\ &= \left(\frac{y-\mu_2}{\sigma_2} \right)^2 - 2\left(\rho \frac{x-\mu_1}{\sigma_1} \right) \left(\frac{y-\mu_2}{\sigma_2} \right) + \left(\rho \frac{x-\mu_1}{\sigma_1} \right)^2 + (1-\rho^2) \left(\frac{x-\mu_1}{\sigma_1} \right)^2 \\ &= \left[\left(\frac{y-\mu_2}{\sigma_2} \right) - \left(\rho \frac{x-\mu_1}{\sigma_1} \right) \right]^2 + (1-\rho^2) \left(\frac{x-\mu_1}{\sigma_1} \right)^2. \end{aligned} \quad (52)$$

Furthermore,

$$\begin{aligned} \frac{y - \mu_2}{\sigma_2} - \rho \frac{x - \mu_1}{\sigma_1} &= \frac{y - \mu_2}{\sigma_2} - \frac{1}{\sigma_2} \cdot \rho \sigma_2 \frac{x - \mu_1}{\sigma_1} \\ &= \frac{1}{\sigma_2} \left\{ y - \left[\mu_2 + \frac{\rho \sigma_2}{\sigma_1} (x - \mu_1) \right] \right\} \\ &= \frac{y - b_x}{\sigma_2}, \quad \text{where } b_x = \mu_2 + \frac{\rho \sigma_2}{\sigma_1} (x - \mu_1) \end{aligned}$$

(see also Exercise 5.6).

Therefore, the right-hand side of (52) is equal to:

$$\left(\frac{y - b_x}{\sigma_2} \right)^2 + (1 - \rho^2) \left(\frac{x - \mu_1}{\sigma_1} \right)^2,$$

and hence the exponent becomes:

$$-\frac{(x - \mu_1)^2}{2\sigma_1^2} - \frac{(y - b_x)^2}{2(\sigma_2\sqrt{1 - \rho^2})^2}.$$

Then the joint p.d.f. may be rewritten as follows:

$$f_{X,Y}(x, y) = \frac{1}{\sqrt{2\pi}\sigma_1} e^{-\frac{(x-\mu_1)^2}{2\sigma_1^2}} \cdot \frac{1}{\sqrt{2\pi}(\sigma_2\sqrt{1-\rho^2})} e^{-\frac{(y-b_x)^2}{2(\sigma_2\sqrt{1-\rho^2})^2}}. \quad (53)$$

The first factor on the right-hand side of (53) is the p.d.f. of $N(\mu_1, \sigma_1^2)$ and the second factor is the p.d.f. of $N(b_x, (\sigma_2\sqrt{1 - \rho^2})^2)$. Therefore, integration with respect to y produces the marginal $N(\mu_1, \sigma_1^2)$ distribution, which, of course, integrates to 1. So, we have established the following two facts: $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy = 1$, and

$$X \sim N(\mu_1, \sigma_1^2), \quad \text{and, by symmetry, } Y \sim N(\mu_2, \sigma_2^2). \quad (54)$$

The results recorded in (54) also reveal the special significance of the parameters μ_1, σ_1^2 and μ_2, σ_2^2 . Namely, they are the means and the variances of the (normally distributed) r.v.'s X and Y , respectively. Relations (53) and (54) also provided immediately the conditional p.d.f. $f_{Y|X}$; namely,

$$f_{Y|X}(y/x) = \frac{1}{\sqrt{2\pi}(\sigma_2\sqrt{1-\rho^2})} \exp \left[-\frac{(y-b_x)^2}{2(\sigma_2\sqrt{1-\rho^2})^2} \right].$$

Thus, in obvious notation:

$$Y | X = x \sim N(b_x, (\sigma_2\sqrt{1-\rho^2})^2), \quad b_x = \mu_2 + \frac{\rho\sigma_2}{\sigma_1}(x - \mu_1), \quad (55)$$


and by symmetry:

$$X | Y = y \sim N(b_y, (\sigma_1\sqrt{1-\rho^2})^2), \quad b_y = \mu_1 + \frac{\rho\sigma_1}{\sigma_2}(y - \mu_2). \quad (56)$$

In Figure 4.7, the conditional p.d.f. $f_{Y|X}(\cdot | x)$ is depicted for three values of x : $x = 5, 10$, and 15 .

Figure 4.7

**Conditional
Probability Density
Functions of the
Bivariate Normal
Distribution**



Formulas (53), (54), and (56) also allow us to calculate easily the covariance and the correlation coefficient of X and Y . Indeed, by (53):

$$\begin{aligned} E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x, y) dx dy = \int_{-\infty}^{\infty} x f_X(x) \left[\int_{-\infty}^{\infty} y f_{Y|X}(y|x) dy \right] dx \\ &= \int_{-\infty}^{\infty} x f_X(x) b_x dx = \int_{-\infty}^{\infty} x f_X(x) \left[\mu_2 + \frac{\rho \sigma_2}{\sigma_1} (x - \mu_1) \right] dx \\ &= \mu_1 \mu_2 + \rho \sigma_1 \sigma_2 \end{aligned}$$

(see also Exercise 5.7). Since we already know that $EX = \mu_1$, $EY = \mu_2$, and $\text{Var}(X) = \sigma_1^2$, $\text{Var}(Y) = \sigma_2^2$, we obtain:

$$\text{Cov}(X, Y) = E(XY) - (EX)(EY) = \mu_1 \mu_2 + \rho \sigma_1 \sigma_2 - \mu_1 \mu_2 = \rho \sigma_1 \sigma_2,$$

and therefore $\rho(X, Y) = \frac{\rho \sigma_1 \sigma_2}{\sigma_1 \sigma_2} = \rho$. Thus, we have:

$$\text{Cov}(X, Y) = \rho \sigma_1 \sigma_2 \quad \text{and} \quad \rho(X, Y) = \rho. \quad (57)$$

Relation (57) reveals that the parameter ρ in (50) is, actually, the correlation coefficient of the r.v.'s X and Y .

EXAMPLE 27

If the r.v.'s X_1 and X_2 have the Bivariate Normal distribution with parameters $\mu_1, \mu_2, \sigma_1^2, \sigma_2^2$, and ρ :

- (i) Calculate the quantities: $E(c_1 X_1 + c_2 X_2)$, $\text{Var}(c_1 X_1 + c_2 X_2)$, where c_1, c_2 are constants.
- (ii) How the expression in part (i) becomes for: $\mu_1 = -1$, $\mu_2 = 3$, $\sigma_1^2 = 4$, $\sigma_2^2 = 9$, and $\rho = \frac{1}{2}$?

DISCUSSION

(i) $E(c_1 X_1 + c_2 X_2) = c_1 EX_1 + c_2 EX_2 = c_1 \mu_1 + c_2 \mu_2$, since $X_i \sim N(\mu_i, \sigma_i^2)$, so that $EX_i = \mu_i$, $i = 1, 2$. Also,

$$\begin{aligned} \text{Var}(c_1 X_1 + c_2 X_2) &= c_1^2 \sigma_1^2 + c_2^2 \sigma_2^2 + 2c_1 c_2 \sigma_{X_1} \sigma_{X_2} \rho(X_1, X_2) \quad (\text{by (32)}) \\ &= c_1^2 \sigma_1^2 + c_2^2 \sigma_2^2 + 2c_1 c_2 \sigma_1 \sigma_2 \rho, \end{aligned}$$

since $X_i \sim N(\mu_i, \sigma_i^2)$, so that $\text{Var}(X_i) = \sigma_i^2$, $i = 1, 2$, and $\rho(X_1, X_2) = \rho$, by (57).

(ii) Here $E(c_1X_1 + c_2X_2) = -c_1 + 3c_2$, and $\text{Var}(c_1X_1 + c_2X_2) = 4c_1 + 9c_2 + 2c_1c_2 \times 2 \times 3 \times \frac{1}{2} = 4c_1 + 9c_2 + 6c_1c_2$.

Finally, it can be seen by integration that the joint m.g.f. of X and Y is given by the formula:

$$M_{X,Y}(t_1, t_2) = \exp \left[\mu_1 t_1 + \mu_2 t_2 + \frac{1}{2} (\sigma_1^2 t_1^2 + 2\rho\sigma_1\sigma_2 t_1 t_2 + \sigma_2^2 t_2^2) \right], \quad t_1, t_2 \in \mathfrak{R}; \quad (58)$$

we choose not to pursue its justification (which can be found, e.g., in pages 158–159, in the book “A Course in Mathematical Statistics,” 2nd edition (1997), Academic Press, by G.G. Roussas). We see, however, easily that

$$\frac{\partial}{\partial t_1} M_{X,Y}(t_1, t_2) = (\mu_1 + \sigma_1^2 t_1 + \rho\sigma_1\sigma_2 t_2) M_{X,Y}(t_1, t_2),$$

and hence:

$$\begin{aligned} \frac{\partial^2}{\partial t_1 \partial t_2} M_{X,Y}(t_1, t_2) &= \rho\sigma_1\sigma_2 M_{X,Y}(t_1, t_2) + (\mu_1 + \sigma_1^2 t_1 + \rho\sigma_1\sigma_2 t_2) \\ &\quad \times (\mu_2 + \sigma_2^2 t_2 + \rho\sigma_1\sigma_2 t_1) M_{X,Y}(t_1, t_2), \end{aligned}$$

which, evaluated at $t_1 = t_2 = 0$, yields: $\rho\sigma_1\sigma_2 + \mu_1\mu_2 = E(XY)$, as we have already seen.

4.5.3 Multivariate Normal Distribution

The Multivariate Normal distribution is a generalization of the Bivariate Normal distribution and can be defined in a number of ways; we choose the one given here. To this end, for $k \geq 2$, let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_k)$ be a vector of constants, and let $\boldsymbol{\Sigma}$ be a $k \times k$ nonsingular matrix, so that the inverse $\boldsymbol{\Sigma}^{-1}$ exists and the determinant $|\boldsymbol{\Sigma}| \neq 0$. Finally, set \mathbf{X} for the vector of r.v.'s X_1, \dots, X_k ; i.e., $\mathbf{X} = (X_1, \dots, X_k)$ and $\mathbf{x} = (x_1, \dots, x_k)$ for any point in \mathfrak{R}^k . Then, the joint p.d.f. of the X_i 's, or the p.d.f. of the random vector \mathbf{X} , is said to be *Multivariate Normal*, or *k-Variate Normal*, if it is given by the formula:

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{(2\pi)^{k/2} |\boldsymbol{\Sigma}|^{1/2}} \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu}) \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})' \right], \quad \mathbf{x} \in \mathfrak{R}^k,$$

where, it is to be recalled that “ $'$ ” stands for transpose.

It can be seen that: $EX_i = \mu_i$, $\text{Var}(X_i) = \sigma_i^2$ is the (i, i) th element of $\boldsymbol{\Sigma}$, and $\text{Cov}(X_i, X_j)$ is the (i, j) th element of $\boldsymbol{\Sigma}$, so that $\boldsymbol{\mu} = (EX_1, \dots, EX_k)$ and $\boldsymbol{\Sigma} = (\text{Cov}(X_i, X_j))$, $i, j = 1, \dots, k$. The quantities $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ are called the *parameters* of the distribution. It can also be seen that the joint m.g.f. of the X_i 's, or the m.g.f. of the random vector \mathbf{X} , is given by:

$$M_{\mathbf{X}}(t) = \exp \left(\boldsymbol{\mu} t' + \frac{1}{2} t \boldsymbol{\Sigma} t' \right), \quad t \in \mathfrak{R}^k.$$

The k -Variate Normal distribution has properties similar to those of the 2-dimensional Normal distribution, and the latter is obtained from the former by taking $\boldsymbol{\mu} = (\mu_1, \mu_2)$ and $\boldsymbol{\Sigma} = \begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{pmatrix}$, where $\rho = \rho(X_1, X_2)$.

More relevant information can be found, e.g., in Chapter 18 of the reference cited in the discussion of Example 27.

Exercises

5.1 Show that

$$\binom{n}{n_1} \binom{n-n_1}{n_2} \cdots \binom{n-n_1-\cdots-n_{k-1}}{n_k} = \frac{n!}{n_1!n_2!\cdots n_k!}.$$

5.2 In a store selling TV sets, it is known that 25% of the customers will purchase a TV set of brand A, 40% will purchase a TV set of brand B, and 35% will just be browsing around. For a lot of 10 customers:

- (i) What is the probability that 2 will purchase a TV set of brand A, 3 will purchase a TV set of brand B, and 5 will purchase neither?
- (ii) If it is known that 6 customers did not purchase a TV set, what is the (conditional) probability that 1 of the rest will purchase a TV set of brand A and 3 will purchase a TV set of brand B?

5.3 Human blood occurs in 4 types termed A , B , AB , and O with respective frequencies $p_A = 0.40$, $p_B = 0.10$, $p_{AB} = 0.05$, and $p_O = 0.45$. If n donors participate in a blood drive, denote by X_A , X_B , X_{AB} , and X_O the numbers of donors with respective blood types A , B , AB , and O . Then X_A , X_B , X_{AB} , and X_O are r.v.'s having the Multinomial distribution with parameters n and p_A , p_B , p_{AB} , p_O . Write out the appropriate formulas for the following probabilities:

- (i) $P(X_A = x_A, X_B = x_B, X_{AB} = x_{AB}, X_O = x_O)$ for x_A, x_B, x_{AB} , and x_O nonnegative integers with $x_A + x_B + x_{AB} + x_O = n$.
- (ii) $P(X_A = x_A, X_B = x_B, X_{AB} = x_{AB})$.
- (iii) $P(X_A = x_A, X_B = x_B)$.
- (iv) $P(X_A = x_A)$.
- (v) $P(X_A = x_A, X_B = x_B, X_{AB} = x_{AB} | X_O = x_O)$.
- (vi) $P(X_A = x_A, X_B = x_B | X_{AB} = x_{AB}, X_O = x_O)$.
- (vii) $P(X_A = x_A | X_B = x_B, X_{AB} = x_{AB}, X_O = x_O)$.
- (viii) Give numerical answers to parts (i)–(vii), if $n = 20$, and $x_A = 8$, $x_B = 2$, $x_{AB} = 1$, $x_O = 9$.

5.4 In conjunction with the Multinomial distribution, show that:

$$EX_i = np_i, EX_j = np_j, \text{Var}(X_i) = np_i(1 - p_i), \text{Var}(X_j) = np_j(1 - p_j),$$

$$\text{Cov}(X_i, X_j) = -np_i p_j \quad \text{and} \quad \rho(X_i, X_j) = -\frac{p_i p_j}{[p_i(1 - p_i)p_j(1 - p_j)]^{1/2}}.$$

5.5 Refer to Exercises 5.3 and 5.4, and for $n = 20$, calculate the quantities:

$$EX_A, EX_B, EX_{AB}, EX_O; \text{Var}(X_A), \text{Var}(X_B), \text{Var}(X_{AB}), \\ \text{Var}(X_O); \text{Cov}(X_A, X_B), \text{Cov}(X_A, X_{AB}), \text{Cov}(X_A, X_O); \\ \rho(X_A, X_B), \rho(X_A, X_{AB}), \rho(X_A, X_O).$$

5.6 Elaborate on the expressions in (51), as well as the expressions following (51).

5.7 If the r.v.'s X and Y have the Bivariate Normal distribution with parameters $\mu_1, \mu_2, \sigma_1^2, \sigma_2^2$, and ρ , show that $E(XY) = \mu_1\mu_2 + \rho\sigma_1\sigma_2$.

Hint: Write the joint p.d.f. $f_{X,Y}$ as $f_{Y|X}(y|x)f_X(x)$ and use the fact (see relation (54)) that $E(Y|X=x) = b_x = \mu_2 + \frac{\rho\sigma_2}{\sigma_1}(x - \mu_1)$.

5.8 If the r.v.'s X and Y have the Bivariate Normal distribution, then, by using Exercise 5.7, show that the parameter ρ is, indeed, the correlation coefficient of the r.v.'s X and Y , $\rho = \rho(X, Y)$.

5.9 If the r.v.'s X and Y have the Bivariate Normal distribution, and c_1, c_2 are constants, express the expectation $E(c_1X + c_2Y)$ and the variance $\text{Var}(c_1X + c_2Y)$ in terms of $c_1, c_2, \mu_1 = EX, \mu_2 = EY, \sigma_1^2 = \text{Var}(X), \sigma_2^2 = \text{Var}(Y)$, and $\rho = \rho(X, Y)$.

5.10 If the r.v.'s X and Y have the Bivariate Normal distribution, then it is known (see, e.g., relation (11) on page 158 in the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G.G. Roussas) that the joint m.g.f. of X and Y is given by:

$$M_{X,Y}(t_1, t_2) = \exp\left[\mu_1 t_1 + \mu_2 t_2 + \frac{1}{2}(\sigma_1^2 t_1^2 + 2\rho\sigma_1\sigma_2 t_1 t_2 + \sigma_2^2 t_2^2)\right], \quad t_1, t_2 \in \mathfrak{R}.$$

Use this m.g.f. in order to show that:

$$EX = \mu_1, EY = \mu_2, \text{Var}(X) = \sigma_1^2, \text{Var}(Y) = \sigma_2^2,$$

$$\text{Cov}(X, Y) = \rho\sigma_1\sigma_2, \quad \text{and} \quad \rho(X, Y) = \rho.$$

5.11 Use the joint m.g.f. of the r.v.'s X and Y having a Bivariate Normal distribution (see Exercise 5.10) in order to show that:

- (i) If X and Y have the Bivariate Normal distribution with parameters $\mu_1, \mu_2, \sigma_1^2, \sigma_2^2$, and ρ , then, for any constants c_1 and c_2 , the r.v. $c_1X + c_2Y$ has the Normal distribution with parameters $c_1\mu_1 + c_2\mu_2$, and $c_1^2\sigma_1^2 + 2c_1c_2\rho\sigma_1\sigma_2 + c_2^2\sigma_2^2$.
- (ii) If the r.v. $c_1X + c_2Y$ is Normally distributed, then the r.v.'s X and Y have the Bivariate Normal distribution with parameters $\mu_1 = EX, \mu_2 = EY, \sigma_1^2 = \text{Var}(X), \sigma_2^2 = \text{Var}(Y)$, and $\rho = \rho(X, Y)$.

5.12 Consider the function f defined by:

$$f(x, y) = \begin{cases} \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}}, & \text{for } (x, y) \text{ outside the square } [-1, 1] \times [-1, 1] \\ \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} + \frac{1}{2\pi e} x^3 y^3, & \text{for } (x, y) \text{ in the square } [-1, 1] \times [-1, 1]. \end{cases}$$

- (i) Show that f is a non-Bivariate Normal p.d.f.
- (ii) Also, show that both marginals, call them f_1 and f_2 , are $N(0, 1)$ p.d.f.'s.

Remark: We know that if X, Y have the Bivariate Normal distribution, then the distributions of the r.v.'s X and Y themselves are Normal. This exercise shows that the inverse need not be true.

5.13 Let the r.v.'s X and Y have the Bivariate Normal distribution with parameters $\mu_1, \mu_2, \sigma_1^2, \sigma_2^2$, and ρ , and set $U = X + Y, V = X - Y$. Then show that:

- (i) The r.v.'s U and V also have the Bivariate Normal distribution with parameters $\mu_1 + \mu_2, \mu_1 - \mu_2, \tau_1^2 = \sigma_1^2 + 2\rho\sigma_1\sigma_2 + \sigma_2^2, \tau_2^2 = \sigma_1^2 - 2\rho\sigma_1\sigma_2 + \sigma_2^2$, and $\rho_0 = (\sigma_1^2 - \sigma_2^2)/\tau_1\tau_2$.
- (ii) $U \sim N(\mu_1 + \mu_2, \tau_1^2), V \sim N(\mu_1 - \mu_2, \tau_2^2)$.
- (iii) The r.v.'s U and V are uncorrelated if and only if $\sigma_1^2 = \sigma_2^2$.

Independence of Random Variables and Some Applications

This chapter consists of two sections. In the first section, we introduce the concept of independence of r.v.'s and establish criteria for proving or disproving independence. Also, its relationship to uncorrelatedness is discussed. In the second section, the sample mean and the sample variance are defined, and some of their moments are also produced. The main thrust of this section, however, is the discussion of the reproductive property of certain distributions. As a by-product, we also obtain the distribution of the sample mean and of a certain multiple of the sample variance for independent and Normally distributed r.v.'s.

5.1 Independence of Random Variables and Criteria of Independence

In Section 4 of Chapter 2, the concept of independence of two events was introduced and it was suitably motivated and illustrated by means of examples. This concept was then generalized to more than two events. What is done in this section is, essentially, to carry over the concept of independence from events to r.v.'s. To this end, consider first two r.v.'s X_1 and X_2 and the events induced in the sample space \mathcal{S} by each one of them separately as well as by both of them jointly. That is, for subsets B_1, B_2 of \mathfrak{R} , let:

$$A_1 = (X_1 \in B_1) = X_1^{-1}(B_1) = \{s \in \mathcal{S}; X_1(s) \in B_1\}, \quad (1)$$

$$A_2 = (X_2 \in B_2) = X_2^{-1}(B_2) = \{s \in \mathcal{S}; X_2(s) \in B_2\}, \quad (2)$$

$$\begin{aligned} A_{12} &= ((X_1, X_2) \in B_1 \times B_2) = (X_1 \in B_1 \& X_2 \in B_2) = (X_1, X_2)^{-1}(B_1 \times B_2) \\ &= \{s \in \mathcal{S}; X_1(s) \in B_1 \text{ as } X_2(s) \in B_2\} = A_1 \cap A_2. \end{aligned} \quad (3)$$

Then the r.v.'s X_1, X_2 are said to be independent if, for any B_1 and B_2 as before, the corresponding events A_1 and A_2 are independent; that is, $P(A_1 \cap A_2) = P(A_1)P(A_2)$. By (1)–(3), clearly, this relation is equivalent to:

$$P(X_1 \in B_1, X_2 \in B_2) = P(X_1 \in B_1)P(X_2 \in B_2). \quad (4)$$

This relation states, in effect, that information regarding one r.v. has no effect on the probability distribution of the other r.v. For example,

$$\begin{aligned} P(X_1 \in B_1 | X_2 \in B_2) &= \frac{P(X_1 \in B_1, X_2 \in B_2)}{P(X_2 \in B_2)} \\ &= \frac{P(X_1 \in B_1)P(X_2 \in B_2)}{P(X_2 \in B_2)} = P(X_1 \in B_1). \end{aligned}$$

Relation (4) is taken as the definition of independence of these two r.v.'s, which is then generalized in a straightforward way to k r.v.'s.

DEFINITION 1

Two r.v.'s X_1 and X_2 are said to be *independent (statistically or stochastically or in the probability sense)* if, for any subsets B_1 and B_2 of \mathfrak{R} ,

$$P(X_1 \in B_1, X_2 \in B_2) = P(X_1 \in B_1)P(X_2 \in B_2).$$

The r.v.'s X_1, \dots, X_k are said to be *independent (in the same sense as above)* if, for any subsets B_1, \dots, B_k of \mathfrak{R} ,

$$P(X_i \in B_i, i = 1, \dots, k) = \prod_{i=1}^k P(X_i \in B_i). \quad (5)$$

Nonindependent r.v.'s are said to be *dependent*.

The practical question which now arises is how one checks independence of k given r.v.'s, or lack thereof. This is done by means of the following criterion referred to as the Factorization Theorem because of the form of the expressions involved.

THEOREM 1

(Criterion of independence, Factorization Theorem) For $k \geq 2$, the r.v.'s X_1, \dots, X_k are independent if and only if any one of the following three relations holds:

$$(i) F_{X_1, \dots, X_k}(x_1, \dots, x_k) = F_{X_1}(x_1) \cdots F_{X_k}(x_k) \quad (6)$$

for all x_1, \dots, x_k in \mathfrak{R} .

$$(ii) f_{X_1, \dots, X_k}(x_1, \dots, x_k) = f_{X_1}(x_1) \cdots f_{X_k}(x_k) \quad (7)$$

for all x_1, \dots, x_k in \mathfrak{R} .

$$(iii) M_{X_1, \dots, X_k}(t_1, \dots, t_k) = M_{X_1}(t_1) \cdots M_{X_k}(t_k) \quad (8)$$

for all t_1, \dots, t_k in a non-degenerate interval containing 0.

Before we proceed with the justification of this theorem, let us refer to Example 1 in Chapter 4 and notice that: $f_X(3) = 0.04$, $f_Y(2) = 0.15$, and $f_{X,Y}(3, 2) = 0.02$, so that $f_{X,Y}(3, 2) = 0.02 \neq 0.04 \times 0.15 = 0.006 = f_X(3)f_Y(2)$.

Accordingly, the r.v.'s X and Y are *not* independent. On the other hand, in reference to Example 2 (see also Example 6), we have, for all $x, y > 0$:

$$f_{X,Y}(x, y) = \lambda_1 \lambda_2 e^{-\lambda_1 x - \lambda_2 y} = (\lambda_1 e^{-\lambda_1 x})(\lambda_2 e^{-\lambda_2 y}) = f_X(x)f_Y(y),$$

so that $f_{X,Y}(x, y) = f_X(x)f_Y(y)$ for all x and y , and consequently, the r.v.'s X and Y are independent. Finally, refer to the Bivariate Normal distribution whose p.d.f. is given by (49) of Chapter 4 and set $\rho = 0$. Then, from (49), (50), and (53), we have $f_{X,Y}(x, y) = f_X(x)f_Y(y)$ for all x and y . Therefore, $\rho = 0$ implies that the r.v.'s X and Y are independent.

EXAMPLE 1

Examine the r.v.'s X and Y from an independence viewpoint, if their joint p.d.f. is given by: $f_{X,Y}(x, y) = 4xy$, $0 < x < 1$, $0 < y < 1$ (and 0 otherwise).

DISCUSSION We will use part (ii) of Theorem 1 for which the marginal p.d.f.'s are needed. To this end, we have:

$$f_X(x) = 4x \int_0^1 y dy = 2x, \quad 0 < x < 1;$$

$$f_Y(y) = 4y \int_0^1 x dx = 2y, \quad 0 < y < 1.$$

Hence, for all $0 < x < 1$ and $0 < y < 1$, it holds that: $2x \times 2y = 4xy$, or $f_X(x)f_Y(y) = f_{X,Y}(x, y)$. This relation is also, trivially, true (both sides are equal to 0) for x and y not satisfying the inequalities $0 < x < 1$ and $0 < y < 1$. It follows that X and Y are independent.

Here are two examples where the r.v.'s involved are not independent.

EXAMPLE 2

If the r.v.'s X and Y have joint p.d.f. given by: $f_{X,Y}(x, y) = 2$, $0 < x < y < 1$ (and 0 otherwise), check whether these r.v.'s are independent or not.

DISCUSSION Reasoning as in the previous example, we find:

$$f_X(x) = 2 \int_x^1 dy = 2(1 - x), \quad 0 < x < 1;$$

$$f_Y(y) = 2 \int_0^y dx = 2y, \quad 0 < y < 1.$$

Then independence of X and Y would require that: $4(1 - x)y = 2$ for all $0 < x < y < 1$, which, clearly, need not hold. For example, for $x = \frac{1}{4}$, $y = \frac{1}{2}$, $4(1 - x)y = 4 \times \frac{3}{4} \times \frac{1}{2} = \frac{3}{2} \neq 2$. Thus, the X and Y are not independent.

EXAMPLE 3

In reference to Example 8 in Chapter 4, the r.v.'s X and Y have joint p.d.f. $f_{X,Y}(x, y) = 8xy$, $0 < x \leq y < 1$ (and 0 otherwise), and:

$$f_X(x) = 4x(1 - x^2), \quad 0 < x < 1; \quad f_Y(y) = 4y^3, \quad 0 < y < 1.$$

Independence of X and Y would require that: $4x(1-x^2) \times 4y^3 = 8xy$ or $(1-x^2)y^2 = \frac{1}{2}$, $0 < x \leq y \leq 1$. However, this relation need not be true because, for example, for $x = \frac{1}{4}$ and $y = \frac{1}{2}$, we have: left-hand side = $\frac{15}{64} \neq \frac{1}{2}$ = right-hand side. So, the r.v.'s X and Y are dependent.

REMARK 1 On the basis of Examples 2 and 3, one may surmise the following rule of thumb: If the arguments x and y (for the case of two r.v.'s) do not vary independently of each other, the r.v.'s involved are likely to be dependent.

A special case of the following result will be needed for the proof of Theorem 1.

PROPOSITION 1 Consider the r.v.'s X_1, \dots, X_k , the functions $g_i: \mathfrak{R} \rightarrow \mathfrak{R}$, $i = 1, \dots, k$, and suppose all expectations appearing below are finite. Then independence of the r.v.'s X_1, \dots, X_k implies:

$$E \left[\prod_{i=1}^k g_i(X_i) \right] = \prod_{i=1}^k E g_i(X_i). \quad (9)$$

PROOF Suppose the r.v.'s are of the continuous type (so that we use integrals; replace them by summations, if the r.v.'s are discrete). Then:

$$\begin{aligned} E \left[\prod_{i=1}^k g_i(X_i) \right] &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g_1(x_1) \cdots g_k(x_k) f_{X_1, \dots, X_k}(x_1, \dots, x_k) dx_1 \cdots dx_k \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g_1(x_1) \cdots g_k(x_k) f_{X_1}(x_1) \cdots f_{X_k}(x_k) dx_1 \cdots dx_k \\ &\quad \text{(by independence)} \\ &= \left[\int_{-\infty}^{\infty} g_1(x_1) f_{X_1}(x_1) dx_1 \right] \cdots \left[\int_{-\infty}^{\infty} g_k(x_k) f_{X_k}(x_k) dx_k \right] \\ &= E g_1(X_1) \cdots E g_k(X_k) = \prod_{i=1}^k E g_i(X_i). \quad \blacktriangle \end{aligned}$$

COROLLARY 1 By taking $g_i(X_i) = e^{t_i X_i}$, $t_i \in \mathfrak{R}$, $i = 1, \dots, k$, relation (9) becomes:

$$\begin{aligned} E \exp(t_1 X_1 + \cdots + t_k X_k) &= \prod_{i=1}^k E \exp(t_i X_i), \quad \text{or} \\ M_{X_1, \dots, X_k}(t_1, \dots, t_k) &= \prod_{i=1}^k M_{X_i}(t_i). \quad (10) \end{aligned}$$

COROLLARY 2 If the r.v.'s X and Y are independent, then they are uncorrelated. The converse is also true, if the r.v.'s have the Bivariate Normal distribution.

PROOF In (9), take $k = 2$, identify X_1 and X_2 with X and Y , respectively, and let $g_1(x) = g_2(x) = x$, $x \in \mathfrak{R}$. Then $E(XY) = (EX)(EY)$, which implies $Cov(X, Y) = 0$ and $\rho(X, Y) = 0$. The converse for the Bivariate Normal distribution follows by means of (50) and (53) in Chapter 4. ▲

REMARK 2 That uncorrelated r.v.'s are not, in general, independent may be illustrated by means of examples (see, e.g., Exercise 1.20).

REMARK 3 If X_1, \dots, X_k are independent r.v.'s, then it is intuitively clear that independence should be preserved for suitable functions of the X_i 's. For example, if $Y_i = g(X_i)$, $i = 1, \dots, k$, then the r.v.'s Y_1, \dots, Y_k are also independent. Independence is also preserved if we take different functions of the X_i 's, provided these functions do not include the same X_i 's. For instance, if $Y = g(X_{i_1}, \dots, X_{i_m})$ and $Z = h(X_{j_1}, \dots, X_{j_n})$, where $1 \leq i_1 < \dots < i_m \leq k$, $1 \leq j_1 < \dots < j_n \leq k$ and all i_1, \dots, i_m are distinct from all j_1, \dots, j_n , then the r.v.'s Y and Z are independent. This will be a rule of thumb to be followed in this book.

PROOF OF THEOREM 1 The proof can be only partial but sufficient for the purposes of this book.

- (i) Independence of the r.v.'s X_1, \dots, X_k means that relation (5) is satisfied. In particular, this is true if $B_i = (-\infty, x_i]$, $i = 1, \dots, k$ which is (6). That (6) implies (5) is a deep probabilistic result dealt with at a much higher level.
- (ii) Suppose the r.v.'s are independent and first assume they are discrete. Then, by taking $B_i = \{x_i\}$, $i = 1, \dots, k$ in (5), we obtain (7). If the r.v.'s are continuous, then consider (6) and differentiate both sides with respect to x_1, \dots, x_k , which, once again, leads to (7) (for continuity points x_1, \dots, x_k). For the converse, suppose that (7) is true; that is, for all t_1, \dots, t_k in \mathfrak{R} ,

$$f_{X_1, \dots, X_k}(t_1, \dots, t_k) = f_{X_1}(t_1) \cdots f_{X_k}(t_k).$$

Then, if the r.v.'s are discrete, sum over the t_i 's from $-\infty$ to x_i , $i = 1, \dots, k$ to obtain (6); if the r.v.'s are continuous, replace the summation operations by integrations in order to obtain (6) again. In either case, independence follows.

- (iii) Independence of X_1, \dots, X_k implies (8) by means of Corollary 1 to Proposition 1 above.

The converse is also true but its proof will not be pursued here (it requires the use of the so-called inversion formula as indicated in Section 1 of Chapter 3 and Remarks 1 and 2 of Chapter 4). ▲

Part (ii) of Theorem 1 has the following corollary, which provides still another useful criterion for independence of k r.v.'s.

COROLLARY 3 The r.v.'s X_1, \dots, X_k are independent if and only if $f_{X_1, \dots, X_k}(x_1, \dots, x_k) = h_1(x_1) \cdots h_k(x_k)$ for all x_1, \dots, x_k in \mathfrak{R} , where h_i is a non-negative function of x_i alone, $i = 1, \dots, k$.

PROOF Suppose the r.v.'s X_1, \dots, X_k are independent. Then, by (7), $f_{X_1, \dots, X_k}(x_1, \dots, x_k) = f_{X_1}(x_1) \cdots f_{X_k}(x_k)$ for all x_1, \dots, x_k in \mathfrak{R} , so that the above factorization holds with $h_i = f_{X_i}$, $i = 1, \dots, k$. Next, assume that the factorization holds, and suppose that the r.v.'s are continuous. For each fixed $i = 1, \dots, k$, set

$$\begin{aligned} c_i &= \int_{-\infty}^{\infty} h_i(x_i) dx_i, \\ \text{so that } c_1 \cdots c_k &= \int_{-\infty}^{\infty} h_1(x_1) dx_1 \cdots \int_{-\infty}^{\infty} h_k(x_k) dx_k \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} h_1(x_1) \cdots h_k(x_k) dx_1 \cdots dx_k \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f_{X_1, \dots, X_k}(x_1, \dots, x_k) dx_1 \cdots dx_k \\ &= 1. \end{aligned}$$

Then, integrating $f_{X_1, \dots, X_k}(x_1, \dots, x_k)$ with respect to all x_j 's with $j \neq i$, we get

$$\begin{aligned} f_{X_i}(x_i) &= c_1 \cdots c_{i-1} c_{i+1} \cdots c_k h_i(x_i) \\ &= \frac{1}{c_i} h_i(x_i). \end{aligned}$$

Hence

$$\begin{aligned} f_{X_1}(x_1) \cdots f_{X_k}(x_k) &= \frac{1}{c_1 \cdots c_k} h_1(x_1) \cdots h_k(x_k) \\ &= h_1(x_1) \cdots h_k(x_k) = f_{X_1, \dots, X_k}(x_1, \dots, x_k), \end{aligned}$$

or $f_{X_1, \dots, X_k}(x_1, \dots, x_k) = f_{X_1}(x_1) \cdots f_{X_k}(x_k)$, for all x_1, \dots, x_k in \mathfrak{R} , so that the r.v.'s X_1, \dots, X_k are independent. The same conclusion holds in case the r.v.'s are discrete by using summations rather than integrations. \blacktriangle

The significance of Corollary 3 is that, in order to check for independence of the r.v.'s X_1, \dots, X_k all one has to do is to establish a factorization of f_{X_1, \dots, X_k} as stated in the corollary. One does not have to verify that the factors are the marginal p.d.f.'s (that will follow as indicated previously).

This section is concluded with the definition of what is known as a random sample. Namely, n independent and identically distributed (i.i.d.) r.v.'s are referred to as forming a *random sample* of size n . Some of their properties are discussed in the next section.

Exercises

- 1.1** In reference to Exercise 2.5 in Chapter 4, determine whether or not the r.v.'s X and Y are independent. Justify your answer.
- 1.2** In reference to Exercises 1.1 and 2.1 in Chapter 4, determine whether or not the r.v.'s X and Y are independent.
- 1.3** The r.v.'s X , Y , and Z have the joint p.d.f. given by: $f_{X,Y,Z}(x, y, z) = \frac{1}{4}$ if $x = 1, y = z = 0; x = 0, y = 1, z = 0; x = y = 0, z = 1; x = y = z = 1$.
- Derive the marginal joint p.d.f.'s $f_{X,Y}, f_{X,Z}, f_{Y,Z}$.
 - Derive the marginal p.d.f.'s f_X, f_Y , and f_Z .
 - Show that any two of the r.v.'s X, Y , and Z are independent.
 - Show that the r.v.'s X, Y , and Z are dependent.
- 1.4** In reference to Exercise 2.8 in Chapter 4, decide whether or not the r.v.'s X and Y are independent. Justify your answer.
- 1.5** In reference to Examples 4 and 7 in Chapter 4, investigate whether or not the r.v.'s X and Y are independent and justify your answer.
- 1.6** Let X and Y be r.v.'s with joint p.d.f. given by:

$$f_{X,Y}(x, y) = \frac{6}{5}(x^2 + y), \quad 0 \leq x \leq 1, \quad 0 \leq y \leq 1.$$

- Determine the marginal p.d.f.'s f_X and f_Y .
 - Investigate whether or not the r.v.'s X and Y are independent. Justify your answer.
- 1.7** The r.v.'s X , and Y have joint p.d.f. given by:

$$f_{X,Y}(x, y) = 1, \quad 0 < x < 1, \quad 0 < y < 1.$$

Then:

- Derive the marginal p.d.f.'s f_X , and f_Y .
 - Show that X and Y are independent.
 - Calculate the probability $P(X + Y < c)$.
 - Give the numerical value of the probability in part (iii) for $c = 1/4$.
- 1.8** The r.v.'s X, Y , and Z have joint p.d.f. given by:

$$f_{X,Y,Z}(x, y, z) = 8xyz, \quad 0 < x < 1, \quad 0 < y < 1, \quad 0 < z < 1.$$

- Derive the marginal p.d.f.'s f_X, f_Y , and f_Z .
 - Show that the r.v.'s X, Y , and Z are independent.
 - Calculate the probability $P(X < Y < Z)$.
- 1.9** The r.v.'s X and Y have joint p.d.f. given by:

$$f_{X,Y}(x, y) = c, \quad \text{for } x^2 + y^2 \leq 9.$$

- Determine the constant c .
- Derive the marginal p.d.f.'s f_X and f_Y .
- Show that the r.v.'s X and Y are dependent.

1.10 The r.v.'s X , Y , and Z have joint p.d.f. given by:

$$f_{X,Y,Z}(x, y, z) = c^3 e^{-c(x+y+z)}, \quad x > 0, \quad y > 0, \quad z > 0.$$

- (i) Determine the constant c .
- (ii) Derive the marginal joint p.d.f.'s $f_{X,Y}$, $f_{X,Z}$, and $f_{Y,Z}$.
- (iii) Derive the marginal p.d.f.'s f_X , f_Y , and f_Z .
- (iv) Show that any two of the r.v.'s X , Y , and Z , as well as all three r.v.'s are independent.

1.11 The r.v.'s X and Y have joint p.d.f. given by the following product: $f_{X,Y}(x, y) = g(x)h(y)$, where g and h are nonnegative functions.

- (i) Derive the marginal p.d.f.'s f_X and f_Y as functions of g and h , respectively.
- (ii) Show that the r.v.'s X and Y are independent.
- (iii) If $h = g$, then the r.v.'s are identically distributed.
- (iv) From part (iii), conclude that $P(X > Y) = 1/2$, provided the distribution is of the continuous type.

1.12 The life of a certain part in a new automobile is a r.v. X whose p.d.f. is Negative Exponential with parameter $\lambda = 0.005$ days.

- (i) What is the expected life of the part in question?
- (ii) If the automobile comes with a spare part whose life is a r.v. Y distributed as X and independent of it, find the p.d.f. of the combined life of the part and its spare.
- (iii) What is the probability that $X + Y \geq 500$ days?

1.13 Let the r.v. X be distributed as $U(0, 1)$ and set $Y = -\log X$.

- (i) Determine the d.f. of Y and then its p.d.f.
- (ii) If the r.v.'s Y_1, \dots, Y_n are independently distributed as Y , and $Z = Y_1 + \dots + Y_n$, determine the distribution of the r.v. Z .

1.14 Let the independent r.v.'s X and Y be distributed as $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$, respectively, and define the r.v.'s U and V by: $U = aX + b$, $V = cY + d$, where a, b, c , and d are constants.

- (i) Use the m.g.f. approach in order to show that:

$$U \sim N(a\mu_1 + b, (a\sigma_1)^2), \quad V \sim N(c\mu_2 + d, (c\sigma_2)^2).$$

- (ii) Determine the joint m.g.f. of U and V .
- (iii) From parts (i) and (ii), conclude that U and V are independent.

1.15 Let X and Y be independent r.v.'s denoting the lifetimes of two batteries and having the Negative Exponential distribution with parameter λ . Set $T = X + Y$ and:

- (i) Determine the d.f. of T by integration, and then the corresponding p.d.f.
- (ii) Determine the p.d.f. of T by using the m.g.f.'s approach.
- (iii) For $\lambda = 1/3$, calculate the probability $P(T \leq 6)$.

1.16 Let X_1, \dots, X_n be i.i.d. r.v.'s with m.g.f. M , and let $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$. Express the m.g.f. $M_{\bar{X}}$ in terms of M .

1.17 In reference to Exercise 3.1 in Chapter 4:

- (i) Calculate the $\text{Var}(X + Y)$ and the s.d. of $X + Y$.
- (ii) Use the Tchebichev inequality to determine a lower bound for the probability: $P(X + Y \leq 10)$.

1.18 Let p be the proportion of defective computer chips in a very large lot of chips produced over a period of time by a certain manufacturing process.

For $i = 1, \dots, n$, associated with the i th chip the r.v. X_i , where $X_i = 1$ if the i th chip is defective, and $X_i = 0$ otherwise. Then X_1, \dots, X_n are independent r.v.'s distributed as $B(1, p)$, and let $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$.

- (i) Calculate the $E\bar{X}$ and the $\text{Var}(\bar{X})$ in terms of p and $q = 1 - p$.
- (ii) Use the Tchebichev inequality to determine the smallest value of n for which $P(|\bar{X} - p| < 0.1\sqrt{pq}) \geq 0.99$.

1.19 Let the independent r.v.'s X_1, \dots, X_n be distributed as $P(\lambda)$, and set $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$.

- (i) Calculate the $E\bar{X}$ and the $\text{Var}(\bar{X})$ in terms of λ and n .
- (ii) Use the Tchebichev inequality to determine the smallest n , in terms of λ and c , for which $P(|\bar{X} - \lambda| < c) \geq 0.95$, for some $c > 0$.
- (iii) Give the numerical value of n for $c = \sqrt{\lambda}$ and $c = 0.1\sqrt{\lambda}$.

1.20 The joint distribution of the r.v.'s X and Y is given by:

$y \backslash x$	-1	0	1
-1	α	β	α
0	β	0	β
1	α	β	α

where $\alpha, \beta > 0$ with $\alpha + \beta = 1/4$.

- (i) Derive the marginal p.d.f.'s f_X and f_Y .
- (ii) Calculate the EX , EY , and $E(XY)$.
- (iii) Show that $\text{Cov}(X, Y) = 0$.
- (iv) Show that the r.v.'s X and Y are dependent.

Remark: Whereas independent r.v.'s are always uncorrelated, this exercise shows that the converse need not be true.

1.21 Refer to Exercise 1.10 and calculate the following quantities without any integration: $E(XY)$, $E(XYZ)$, $\text{Var}(X + Y)$, $\text{Var}(X + Y + Z)$.

1.22 The i.i.d. r.v.'s X_1, \dots, X_n have expectation $\mu \in \Re$ and variance $\sigma^2 < \infty$, and set $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$.

- (i) Determine the $E\bar{X}$ and the $\text{Var}(\bar{X})$ in terms of μ and σ .
- (ii) Use the Tchebichev inequality to determine the smallest value of n for which $P(|\bar{X} - \mu| < k\sigma)$ is at least 0.99; take $k = 1, 2, 3$.

1.23 A piece of equipment works on a battery whose lifetime is a r.v. X with expectation μ and s.d. σ . If n such batteries are used successively and independently of each other, denote by X_1, \dots, X_n their respective lifetimes, so that $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$ is the average lifetime of the batteries.

Use the Tchebichev inequality to determine the smallest value of n for which $P(|\bar{X} - \mu| < 0.5\sigma) \geq 0.99$.

1.24 Let X_1, \dots, X_n be i.i.d. r.v.'s with $EX_1 = \mu \in \mathfrak{R}$ and $Var(X_1) = \sigma^2 < \infty$, and set $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$.

(i) Calculate the $E\bar{X}$ and the $Var(\bar{X})$ in terms of μ and σ .

(ii) Use the Tchebichev inequality in order to determine the smallest value of n , in terms of the positive constant c and α , so that

$$P(|\bar{X} - \mu| < c\sigma) \geq \alpha \quad (0 < \alpha < 1).$$

(iii) What is the numerical value of n in part (ii) if $c = 0.1$ and $\alpha = 0.90$, $\alpha = 0.95$, $\alpha = 0.99$?

1.25 In reference to Exercise 5.13(iii) in Chapter 4, show that the r.v.'s U and V are independent if and only if $\sigma_1^2 = \sigma_2^2$.

5.2 The Reproductive Property of Certain Distributions

Independence plays a decisive role in the reproductive property of certain r.v.'s. Specifically, if X_1, \dots, X_k are r.v.'s having certain distributions, then, if they are also *independent*, it follows that the r.v. $X_1 + \dots + X_k$ is of the same kind. This is, basically, the content of this section. The tool used in order to establish this assertion is the m.g.f., and the basic result employed is relation (8), characterizing independence of r.v.'s. The conditions of applicability of (8) hold in all cases considered here.

First, we derive some general results regarding the sample mean and the sample variance of k r.v.'s, which will be used, in particular, in the Normal distribution case discussed below. To this end, for any k r.v.'s X_1, \dots, X_k , their *sample mean*, denoted by \bar{X}_k or just \bar{X} , is defined by:

$$\bar{X} = \frac{1}{k} \sum_{i=1}^k X_i. \quad (11)$$

The *sample variance* of the X_i 's, denoted by S_k^2 or just S^2 , is defined by:

$$S^2 = \frac{1}{k} \sum_{i=1}^k (X_i - EX_i)^2,$$

provided the EX_i 's are finite. In particular, if $EX_1 = \dots = EX_k = \mu$, say, then S^2 becomes:

$$S^2 = \frac{1}{k} \sum_{i=1}^k (X_i - \mu)^2. \quad (12)$$

The r.v.'s defined by (11) and (12) are most useful when the underlying r.v.'s form a random sample; that is, they are i.i.d.

PROPOSITION 2 Let X_1, \dots, X_k be i.i.d. r.v.'s with (finite) mean μ . Then $E\bar{X} = \mu$. Furthermore, if the X_i 's also have (finite) variance σ^2 , then $Var(\bar{X}) = \frac{\sigma^2}{k}$ and $ES^2 = \sigma^2$.

PROOF The first result follows from (40) in Chapter 4 by taking $c_1 = \dots = c_k = 1/k$. The second result follows from (44) in the same chapter, by way of Corollary 2 to Proposition 1 here, because independence of X_i and X_j , for $i \neq j$, implies $\rho(X_i, X_j) = 0$. In order to check the third result, observe first that:

$$\sum_{i=1}^k (X_i - \mu)^2 = \sum_{i=1}^k X_i^2 + k\mu^2 - 2\mu \sum_{i=1}^k X_i,$$

so that

$$\begin{aligned} E \sum_{i=1}^k (X_i - \mu)^2 &= \sum_{i=1}^k EX_i^2 + k\mu^2 - 2\mu \cdot k\mu \\ &= \sum_{i=1}^k (\sigma^2 + \mu^2) + k\mu^2 - 2k\mu^2 = k\sigma^2. \end{aligned}$$

Then $ES^2 = \frac{1}{k} E \sum_{i=1}^k (X_i - \mu)^2 = \frac{1}{k} k\sigma^2 = \sigma^2$. \blacktriangle

The general thrust of the following four results is to the effect that, if X_1, \dots, X_k are independent and have certain distributions, then their sum $X_1 + \dots + X_k$ has a distribution of the same respective kind. The proof of this statement relies on relation (8), which is validated on account of (10).

THEOREM 2

Let the r.v.'s X_1, \dots, X_k be independent and let $X_i \sim B(n_i, p)$ (the same p), $i = 1, \dots, k$. Then $\sum_{i=1}^k X_i \sim B(\sum_{i=1}^k n_i, p)$.

PROOF By independence, relation (20) in Chapter 3, and $t \in \mathfrak{R}$:

$$M_{\sum_{i=1}^k X_i}(t) = \prod_{i=1}^k M_{X_i}(t) = \prod_{i=1}^k (pe^t + q)^{n_i} = (pe^t + q)^{\sum_{i=1}^k n_i},$$

which is the m.g.f. of $B(\sum_{i=1}^k n_i, p)$. Then $\sum_{i=1}^k X_i \sim B(\sum_{i=1}^k n_i, p)$. \blacktriangle

THEOREM 3

Let the r.v.'s X_1, \dots, X_k be independent and let $X_i \sim P(\lambda_i)$, $i = 1, \dots, k$. Then $\sum_{i=1}^k X_i \sim P(\sum_{i=1}^k \lambda_i)$.

PROOF As above, employ independence and relation (24) in Chapter 3 in order to obtain:

$$M_{\sum_{i=1}^k X_i}(t) = \prod_{i=1}^k M_{X_i}(t) = \prod_{i=1}^k \exp(\lambda_i e^t - \lambda_i) = \exp \left[\left(\sum_{i=1}^k \lambda_i \right) e^t - \left(\sum_{i=1}^k \lambda_i \right) \right],$$

which is the m.g.f. of $P(\sum_{i=1}^k \lambda_i)$, so that $\sum_{i=1}^k X_i \sim P(\sum_{i=1}^k \lambda_i)$. \blacktriangle

THEOREM 4

Let the r.v.'s X_1, \dots, X_k be independent and let $X_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, \dots, k$. Then $\sum_{i=1}^k X_i \sim N(\sum_{i=1}^k \mu_i, \sum_{i=1}^k \sigma_i^2)$. In particular, if $\mu_1 = \dots = \mu_k = \mu$ and $\sigma_1 = \dots = \sigma_k = \sigma$, then $\sum_{i=1}^k X_i \sim N(k\mu, k\sigma^2)$.

PROOF Use independence and formula (44) in Chapter 3, for $t \in \mathfrak{R}$, in order to obtain:

$$\begin{aligned} M_{\sum_{i=1}^k X_i}(t) &= \prod_{i=1}^k M_{X_i}(t) = \prod_{i=1}^k \exp\left(\mu_i t + \frac{\sigma_i^2}{2} t^2\right) \\ &= \exp\left[\left(\sum_{i=1}^k \mu_i\right)t + \frac{\sum_{i=1}^k \sigma_i^2}{2} t^2\right], \end{aligned}$$

which is the m.g.f. of $N(\sum_{i=1}^k \mu_i, \sum_{i=1}^k \sigma_i^2)$, so that $\sum_{i=1}^k X_i \sim N(\sum_{i=1}^k \mu_i, \sum_{i=1}^k \sigma_i^2)$. The special case is immediate. \blacktriangle

To this theorem, there are the following two corollaries.

COROLLARY 1 If the r.v.'s X_1, \dots, X_k are independent and distributed as $N(\mu, \sigma^2)$, then their sample mean $\bar{X} \sim N(\mu, \frac{\sigma^2}{k})$, and $\frac{\sqrt{k}(\bar{X}-\mu)}{\sigma} \sim N(0, 1)$.

PROOF Here $\bar{X} = Y_1 + \dots + Y_k$, where $Y_i = \frac{X_i}{k}$, $i = 1, \dots, k$ are independent and $Y_i \sim N(\frac{\mu}{k}, \frac{\sigma^2}{k^2})$ by Theorem 2 in Chapter 3, applied with $c = 1/k$ and $d = 0$. Then the conclusion follows by Theorem 4. The second conclusion is immediate by the part just established and Proposition 1 in Chapter 3, since $\frac{\sqrt{k}(\bar{X}-\mu)}{\sigma} = \frac{\bar{X}-\mu}{\sqrt{\sigma^2/k}}$. \blacktriangle

COROLLARY 2 Let the r.v.'s X_1, \dots, X_k be independent, let $X_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, \dots, k$, and let $c_i, i = 1, \dots, k$ be constants. Then $\sum_{i=1}^k c_i X_i \sim N(\sum_{i=1}^k c_i \mu_i, \sum_{i=1}^k c_i^2 \sigma_i^2)$.

PROOF As in Corollary 1, $X_i \sim N(\mu_i, \sigma_i^2)$ implies $c_i X_i \sim N(c_i \mu_i, c_i^2 \sigma_i^2)$, and the r.v.'s $c_i X_i, i = 1, \dots, k$ are independent. Then the conclusion follows from the theorem. \blacktriangle

THEOREM 5

Let the r.v.'s X_1, \dots, X_k be independent and let $X_i \sim \chi_{r_i}^2$, $i = 1, \dots, k$. Then $\sum_{i=1}^k X_i \sim \chi_{r_1+\dots+r_k}^2$.

PROOF Use independence and formula (37) in Chapter 3, for $t < \frac{1}{2}$, to obtain:

$$M_{\sum_{i=1}^k X_i}(t) = \prod_{i=1}^k M_{X_i}(t) = \prod_{i=1}^k \frac{1}{(1-2t)^{r_i/2}} = \frac{1}{(1-2t)^{(r_1+\dots+r_k)/2}},$$

which is the m.g.f. of $\chi_{r_1+\dots+r_k}^2$. \blacktriangle

COROLLARY Let the r.v.'s X_1, \dots, X_k be independent and let $X_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, \dots, k$. Then $\sum_{i=1}^k (\frac{X_i - \mu_i}{\sigma_i})^2 \sim \chi_k^2$, and, in particular, if $\mu_1 = \dots = \mu_k = \mu$ and $\sigma_1^2 = \dots = \sigma_k^2 = \sigma^2$, then $\frac{kS^2}{\sigma^2} \sim \chi_k^2$, where S^2 is given in (12).

PROOF The assumption $X_i \sim N(\mu_i, \sigma_i^2)$ implies that $\frac{X_i - \mu_i}{\sigma_i} \sim N(0, 1)$ by Proposition 1 in Chapter 3, and $(\frac{X_i - \mu_i}{\sigma_i})^2 \sim \chi_1^2$ by Proposition 2 in the same chapter. Since independence of X_i , $i = 1, \dots, k$ implies that of $(\frac{X_i - \mu_i}{\sigma_i})^2$, $i = 1, \dots, k$, the theorem applies and yields the first assertion. The second assertion follows from the first by taking $\mu_1 = \dots = \mu_k = \mu$ and $\sigma_1 = \dots = \sigma_k = \sigma$, and using (12) to obtain $\frac{kS^2}{\sigma^2} = \sum_{i=1}^k (\frac{X_i - \mu}{\sigma})^2$. ▲

REMARK 4 From the fact that $\frac{kS^2}{\sigma^2} \sim \chi_k^2$ and formula (37) in Chapter 3, we have $E(\frac{kS^2}{\sigma^2}) = k$, $Var(\frac{kS^2}{\sigma^2}) = 2k$, or $ES^2 = \sigma^2$ and $Var(S^2) = 2\sigma^4/k$.

REMARK 5 Knowing the distribution of $\sum_{i=1}^k X_i$ is of considerable practical importance. For instance, if X_i is the number of defective items among n_i in the i th lot of certain items, $i = 1, \dots, k$, then $\sum_{i=1}^k X_i$ is the total number of defective items in the k lots (and Theorem 2 applies). Likewise, if X_i is the number of particles emitted by the i th radioactive source, $i = 1, \dots, k$, then $\sum_{i=1}^k X_i$ is the total number of particles emitted by all k radioactive sources (and Theorem 3 applies). Also, if X_i is the rain (in inches, for example) which fell in the i th location over a specified period of time, $i = 1, \dots, k$, then $\sum_{i=1}^k X_i$ is the total rainfall in all of k locations under consideration over the specified period of time (and Theorem 4 applies). Finally, if Y_i denotes the lifetime of the i th battery in a lot of k identical batteries, whose lifetime is assumed to be normally distributed, then $X_i = [(Y_i - \mu)/\sigma]^2$ measures a deviation from the mean lifetime μ , and $\sum_{i=1}^k X_i$ is the totality of such deviations for the k batteries (and Theorem 5 applies).

Here are some numerical applications.

EXAMPLE 4

The defective items in two lots of sizes $n_1 = 10$ and $n_2 = 15$ occur independently at the rate of 6.25%. Calculate the probabilities that the total number of defective items: (i) Does not exceed 2; (ii) Is more than 5.

DISCUSSION If X_1 and X_2 are the r.v.'s denoting the numbers of defective items in the two lots, then $X_1 \sim B(10, 0.0625)$, $X_2 \sim B(15, 0.0625)$ and they are independent. Then $X = X_1 + X_2 \sim B(25, 0.0625)$ and therefore: (i) $P(X \leq 2) = 0.7968$ and (ii) $P(X > 5) = 1 - P(X \leq 5) = 0.0038$ (from the Binomial tables).

EXAMPLE 5

Five radioactive sources independently emit particles at the rate of 0.08 per certain time unit. What is the probability that the total number of particles does not exceed 3 in the time unit considered?

DISCUSSION In obvious notation, we have here the independent r.v.'s X_i distributed as $P(0.08)$, $i = 1, \dots, 5$. Then $X = \sum_{i=1}^5 X_i \sim P(0.4)$, and the required probability is: $P(X \leq 3) = 0.999224$ (from the Poisson tables).

EXAMPLE 6

The rainfall in two locations is measured (in inches over a certain time unit) by two independent and Normally distributed r.v.'s X_1 and X_2 as follows: $X_1 \sim N(10, 9)$ and $X_2 \sim N(15, 25)$. What is the probability that the total rainfall: (i) Will exceed 30 inches (which may result in flooding)? (ii) Will be less than 8 inches (which will mean a drought)?

DISCUSSION If $X = X_1 + X_2$, then $X \sim N(25, 34)$, so that: (i) $P(X > 30) = 1 - P(X \leq 30) = 1 - P(Z \leq \frac{30-25}{\sqrt{34}}) \simeq 1 - \Phi(0.86) = 1 - 0.805105 = 0.194895$, and (ii) $P(X < 8) = P(Z < \frac{8-25}{\sqrt{34}}) \simeq \Phi(-2.92) = 1 - \Phi(2.92) = 1 - 0.99825 = 0.00175$.

In the definition of S^2 by (12), we often replace μ by the sample mean \bar{X} ; this is done habitually in statistics as μ is not really known. Let us denote by \bar{S}^2 the resulting quantity; that is,

$$\bar{S}^2 = \frac{1}{k} \sum_{i=1}^k (X_i - \bar{X})^2. \quad (13)$$

Then it is easy to establish the following identity:

$$\sum_{i=1}^k (X_i - \mu)^2 = \sum_{i=1}^k (X_i - \bar{X})^2 + k(\bar{X} - \mu)^2, \quad (14)$$

or

$$kS^2 = k\bar{S}^2 + [\sqrt{k}(\bar{X} - \mu)]^2. \quad (15)$$

Indeed,

$$\sum_{i=1}^k (X_i - \mu)^2 = \sum_{i=1}^k [(X_i - \bar{X}) + (\bar{X} - \mu)]^2 = \sum_{i=1}^k (X_i - \bar{X})^2 + k(\bar{X} - \mu)^2,$$

since $\sum_{i=1}^k (X_i - \bar{X})(\bar{X} - \mu) = (\bar{X} - \mu)(k\bar{X} - k\bar{X}) = 0$.

From (15), we have, dividing through by σ^2 :

$$\frac{kS^2}{\sigma^2} = \frac{k\bar{S}^2}{\sigma^2} + \left[\frac{\sqrt{k}(\bar{X} - \mu)}{\sigma} \right]^2. \quad (16)$$

Now $\frac{kS^2}{\sigma^2} \sim \chi_k^2$ and $[\frac{\sqrt{k}(\bar{X}-\mu)}{\sigma}]^2 \sim \chi_1^2$ (by Propositions 1 and 2 in Chapter 3) when the r.v.'s X_1, \dots, X_k are independently distributed as $N(\mu, \sigma^2)$. Therefore, from (16), it appears quite feasible that $\frac{k\bar{S}^2}{\sigma^2} \sim \chi_{k-1}^2$. This is, indeed, the case and is the content of the following theorem. This theorem is presently established under an assumption to be justified later on (see Theorem 9 in Chapter 6). The *assumption* is this: If the r.v.'s X_1, \dots, X_k are independent and distributed as $N(\mu, \sigma^2)$, then the r.v.'s \bar{X} and \bar{S}^2 are independent. (The independence of \bar{X} and \bar{S}^2 implies then that of $[\frac{\sqrt{k}(\bar{X}-\mu)}{\sigma}]^2$ and $\frac{k\bar{S}^2}{\sigma^2}$.)

THEOREM 6

Let the r.v.'s X_1, \dots, X_k be independent and distributed as $N(\mu, \sigma^2)$, and let \bar{S}^2 be defined by (13). Then $\frac{k\bar{S}^2}{\sigma^2} \sim \chi_{k-1}^2$. Consequently, $E\bar{S}^2 = \frac{k-1}{k}\sigma^2$ and $Var(\bar{S}^2) = \frac{2(k-1)\sigma^4}{k^2}$.

PROOF Consider relation (16), take the m.g.f.'s of both sides, and use the corollary to Theorem 5 and the assumption of independence made previously in order to obtain:

$$M_{kS^2/\sigma^2}(t) = M_{k\bar{S}^2/\sigma^2}(t)M_{[\sqrt{k}(\bar{X}-\mu)/\sigma]^2}(t),$$

so that

$$M_{k\bar{S}^2/\sigma^2}(t) = M_{kS^2/\sigma^2}(t)/M_{[\sqrt{k}(\bar{X}-\mu)/\sigma]^2}(t),$$

or

$$M_{k\bar{S}^2/\sigma^2}(t) = \frac{1/(1-2t)^{k/2}}{1/(1-2t)^{1/2}} = \frac{1}{(1-2t)^{(k-1)/2}},$$

which is the m.g.f. of the χ_{k-1}^2 distribution. The second assertion follows immediately from the first and formula (37) in Chapter 3. ▲

This chapter is concluded with the following comment. Theorems 2–5 may be misleading in the sense that the sum of independent r.v.'s always has a distribution of the same kind as the summands. That this is definitely not so is illustrated by examples. For instance, if the independent r.v.'s X and Y are $U(0, 1)$, then their sum $X + Y$ is *not* uniform; rather, it is triangular (see Example 4 (continued) in Chapter 6).

Exercises

2.1 For any r.v.'s X_1, \dots, X_n , set

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \quad \text{and} \quad S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2,$$

and show that:

(i)

$$nS^2 = \sum_{i=1}^n (X_i - \bar{X})^2 = \sum_{i=1}^n X_i^2 - n\bar{X}^2.$$

(ii) If the r.v.'s have common (finite) expectation μ , then

$$\sum_{i=1}^n (X_i - \mu)^2 = \sum_{i=1}^n (X_i - \bar{X})^2 + n(\bar{X} - \mu)^2 = nS^2 + n(\bar{X} - \mu)^2.$$

- 2.2** In reference to Exercise 3.1 in Chapter 4, specify the distribution of the sum $X + Y$, and write out the expression for the exact probability $P(X + Y \leq 10)$.
- 2.3** If the independent r.v.'s X and Y are distributed as $B(m, p)$ and $B(n, p)$, respectively:
- What is the distribution of the r.v. $X + Y$?
 - If $m = 8$, $n = 12$, and $p = 0.25$, what is the numerical value of the probability: $P(5 \leq X + Y \leq 15)$?
- 2.4** The independent r.v.'s X_1, \dots, X_n are distributed as $B(1, p)$, and let $S_n = X_1 + \dots + X_n$.
- Determine the distribution of the r.v. S_n .
 - What is the EX_i and the $Var(X_i)$, $i = 1, \dots, n$?
 - From part (ii) and the definition of S_n , compute the ES_n and $Var(S_n)$.
- 2.5** Let X_1, \dots, X_n be i.i.d. r.v.'s with p.d.f. f , and let I be an interval in \Re . Let $p = P(X_1 \in I)$.
- Express p in terms of the p.d.f. f .
 - For k with $1 \leq k \leq n$, express the probability that at least k of X_1, \dots, X_n take values in the interval I in terms of p .
 - Simplify the expression in part (ii), if f is the Negative Exponential p.d.f. with parameter λ and $I = (\frac{1}{\lambda}, \infty)$.
 - Find the numerical value of the probability in part (iii) for $n = 4$ and $k = 2$.
- 2.6** The breakdown voltage of a randomly chosen diode of a certain type is known to be Normally distributed with mean value 40V and s.d. 1.5V.
- What is the probability that the voltage of a single diode is between 39 and 42?
 - If 5 diodes are independently chosen, what is the probability that at least one has a voltage exceeding 42?
- 2.7** Refer to Exercise 1.18 and set $X = X_1 + \dots + X_n$.
- Justify the statement that $X \sim B(n, p)$.
 - Suppose that n is large and p is small (both assumptions quite appropriate in the framework of Exercise 1.18), so that:

$$f(x) = \binom{n}{x} p^x q^{n-x} \simeq e^{-np} \frac{(np)^x}{x!}, \quad x = 0, 1, \dots$$

If $np = 2$, calculate the approximate values of the probabilities $f(x)$ for $x = 0, 1, 2, 3$, and 4.

- 2.8** The r.v.'s X_1, \dots, X_n are independent and $X_i \sim P(\lambda_i)$:
- What is the distribution of the r.v. $X = X_1 + \dots + X_n$?
 - If $\bar{X} = \frac{1}{n}(X_1 + \dots + X_n)$, calculate the $E\bar{X}$ and the $Var(\bar{X})$ in terms of $\lambda_1, \dots, \lambda_n$, and n .

(iii) What do the $E\bar{X}$ and the $Var(\bar{X})$ become when the X_i 's in part (i) are distributed as $P(\lambda)$?

2.9 Suppose that the number of no-shows for a scheduled airplane flight is a r.v. X distributed as $P(\lambda)$, and it is known from past experience that, on the average, there are 2 no-shows. If there are 5 flights scheduled, compute the following probabilities for the total number of no-shows $X = X_1 + \cdots + X_5$:

- (i) 0. (v) At most 10. (ix) 15.
 (ii) At most 5. (vi) 10. (x) At least 15.
 (iii) 5. (vii) At least 10.
 (iv) At least 5. (viii) At most 15.

2.10 The r.v.'s X_1, \dots, X_n are independent and $X_i \sim P(\lambda_i)$, $i = 1, \dots, n$. Set $T = \sum_{i=1}^n X_i$ and $\lambda = \sum_{i=1}^n \lambda_i$, and show that:

- (i) The conditional p.d.f. of X_i , given $T = t$, is $B(t, \lambda_i/\lambda)$, $i = 1, \dots, n$.
 (ii) What does the distribution in part (i) become for $\lambda_1 = \cdots = \lambda_n = c$, say?

2.11 If the independent r.v.'s X and Y are distributed as $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$, respectively:

- (i) Specify the distribution of $X - Y$.
 (ii) Calculate the probability $P(X > Y)$ in terms of μ_1, μ_2, σ_1 , and σ_2 .
 (iii) If $\mu_1 = \mu_2$, conclude that $P(X > Y) = 0.5$.

2.12 The $m + n$ r.v.'s X_1, \dots, X_m and Y_1, \dots, Y_n are independent and $X_i \sim N(\mu_1, \sigma_1^2)$, $i = 1, \dots, m$, $Y_j \sim N(\mu_2, \sigma_2^2)$, $j = 1, \dots, n$. Set $\bar{X} = \frac{1}{m} \sum_{i=1}^m X_i$, $\bar{Y} = \frac{1}{n} \sum_{j=1}^n Y_j$ and:

- (i) Calculate the probability $P(\bar{X} > \bar{Y})$ in terms of $m, n, \mu_1, \mu_2, \sigma_1$, and σ_2 .
 (ii) Give the numerical value of the probability in part (i) when $\mu_1 = \mu_2$ unspecified.

2.13 Let the independent r.v.'s X_1, \dots, X_n be distributed as $N(\mu, \sigma^2)$ and set $X = \sum_{i=1}^n \alpha_i X_i$, $Y = \sum_{j=1}^n \beta_j X_j$, where the α_i 's and the β_j 's are constants. Then:

- (i) Determine the p.d.f.'s of the r.v.'s X and Y .
 (ii) Show that the joint m.g.f. of X and Y is given by:

$$M_{X,Y}(t_1, t_2) = \exp\left[\mu_1 t_1 + \mu_2 t_2 + \frac{1}{2}(\sigma_1^2 t_1^2 + 2\rho\sigma_1\sigma_2 t_1 t_2 + \sigma_2^2 t_2^2)\right],$$

where $\mu_1 = \mu \sum_{i=1}^n \alpha_i$, $\mu_2 = \mu \sum_{j=1}^n \beta_j$, $\sigma_1^2 = \sigma^2 \sum_{i=1}^n \alpha_i^2$, $\sigma_2^2 = \sigma^2 \sum_{j=1}^n \beta_j^2$, $\rho = (\sum_{i=1}^n \alpha_i \beta_i) / \sigma_1 \sigma_2$.

- (iii) From part (ii), conclude that X and Y have the Bivariate Normal distribution with correlation coefficient

$$\rho(X, Y) = \rho = \left(\sum_{i=1}^n \alpha_i \beta_i\right) / \sigma_1 \sigma_2.$$

(iv) From part (iii), conclude that X and Y are independent if and only if $\sum_{i=1}^n \alpha_i \beta_i = 0$.

2.14 Let X and Y be independent r.v.'s distributed as $N(0, \sigma^2)$.

- (i) Set $R = \sqrt{X^2 + Y^2}$ and determine the probability: $P(R \leq r)$, for $r > 0$.
- (ii) What is the numerical value of $P(R \leq r)$ for $\sigma = 1$ and $r = 1.665$, $r = 2.146$, $r = 2.448$, $r = 2.716$, $r = 3.035$, and $r = 3.255$?



Chapter 6

Transformation of Random Variables

This chapter is devoted to transforming a given set of r.v.'s to another set of r.v.'s. The practical need for such transformations will become apparent by means of concrete examples to be cited and/or discussed. The chapter consists of five sections. In the first section, a single r.v. is transformed into another single r.v. In the following section, the number of available r.v.'s is at least two, and they are to be transformed into another set of r.v.'s of the same or smaller number. Two specific applications produce two new distributions, the t -distribution and the F -distribution, which are of great applicability in statistics. A brief account of specific kinds of transformations is given in the subsequent two sections, and the chapter is concluded with a section on order statistics.

6.1 Transforming a Single Random Variable

EXAMPLE 1

Suppose that the r.v.'s X and Y represent the temperature in a certain locality measured in degrees Celsius and Fahrenheit, respectively. Then it is known that X and Y are related as follows: $Y = \frac{9}{5}X + 32$.

This simple example illustrates the need for transforming a r.v. X into another r.v. Y , if Celsius degrees are to be transformed into Fahrenheit degrees.

EXAMPLE 2

As another example, let the r.v. X denote the velocity of a molecule of mass m . Then it is known that the kinetic energy of the molecule is a r.v. Y related to X in the following manner: $Y = \frac{1}{2}mX^2$.

Thus, determining the distribution of the kinetic energy of the molecule involves transforming the r.v. X as indicated above.

The formulation of the general problem is as follows: Let X be a r.v. of the continuous type with p.d.f. f_X , and let h be a real-valued function defined on \mathfrak{R} . Define the r.v. Y by $Y = h(X)$ and determine its p.d.f. f_Y . Under suitable regularity conditions, this problem can be resolved in two ways. One is to determine first the d.f. F_Y and then obtain f_Y by differentiation, and the other is to obtain f_Y directly.

THEOREM 1

Let $S \subseteq \mathfrak{R}$ be the set over which f_X is strictly positive, let $h : S \rightarrow T$ (the image of S under h) $\subseteq \mathfrak{R}$ be one-to-one (that is, to distinct x 's in S there correspond distinct y 's in T) and strictly monotone, and let $Y = h(X)$. For $x \in S$, set $y = h(x) \in T$. Then $F_Y(y) = F_X[h^{-1}(y)]$, if h is increasing, and $F_Y(y) = 1 - F_X[h^{-1}(y)]$, if h is decreasing.

PROOF Inverting the function $y = h(x)$, we get $x = h^{-1}(y)$. Then for increasing h (which implies increasing h^{-1}), we have:

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P[h(X) \leq y] = P\{h^{-1}[h(X)] \leq h^{-1}(y)\} \\ &= P[X \leq h^{-1}(y)] = F_X[h^{-1}(y)]. \end{aligned} \quad (1)$$

If h is decreasing, then so is h^{-1} and therefore:

$$\begin{aligned} F_Y(y) &= P[h(X) \leq y] = P\{h^{-1}[h(X)] \geq h^{-1}(y)\} \\ &= P[X \geq h^{-1}(y)] = 1 - P[X < h^{-1}(y)] \\ &= 1 - P[X \leq h^{-1}(y)] = 1 - F_X[h^{-1}(y)]. \quad \blacktriangle \end{aligned} \quad (2)$$

As an illustration, consider the case $Y = \frac{9}{5}X + 32$ in Example 1 above. Here $y = h(x) = \frac{9}{5}x + 32$ is one-to-one and strictly increasing. Therefore

$$F_Y(y) = F_X\left[\frac{5}{9}(y - 32)\right] \quad \text{and} \quad f_Y(y) = \frac{5}{9}f_X\left[\frac{5}{9}(y - 32)\right]. \quad (3)$$

The function $y = h(x)$ may not be one-to-one and strictly increasing on the entire S but it is so on subsets of it. Then F_Y can still be determined. Example 2 above illustrates the point. Let $Y = \frac{1}{2}mX^2$ as mentioned above. Then proceed as follows: For $y > 0$:

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P\left(\frac{1}{2}mX^2 \leq y\right) = P\left(X^2 \leq \frac{2y}{m}\right) \\ &= P\left(-\sqrt{\frac{2y}{m}} \leq X \leq \sqrt{\frac{2y}{m}}\right) = P\left(X \leq \sqrt{\frac{2y}{m}}\right) - P\left(X < -\sqrt{\frac{2y}{m}}\right) \\ &= P\left(X \leq \sqrt{\frac{2y}{m}}\right) - P\left(X \leq -\sqrt{\frac{2y}{m}}\right) = F_X\left(\sqrt{\frac{2y}{m}}\right) - F_X\left(-\sqrt{\frac{2y}{m}}\right). \end{aligned} \quad (4)$$

Differentiating in (1) or (2) (depending on whether h is increasing or decreasing), we obtain the p.d.f. of Y , namely,

$$f_Y(y) = \frac{d}{dy} F_Y(y) = \frac{d}{dx} F_X(x) \Big|_{x=h^{-1}(y)} \times \left| \frac{dx}{dy} \right| = f_X[h^{-1}(y)] \left| \frac{d}{dy} h^{-1}(y) \right|, \quad y \in T. \quad (5)$$

In the case of formula (3), relation (5) gives: $f_Y(y) = \frac{5}{9} f_X[\frac{5}{9}(y - 32)]$, as has already been seen. In the case of formula (4),

$$f_Y(y) = \left[f_X\left(\sqrt{\frac{2y}{m}}\right) + f_X\left(-\sqrt{\frac{2y}{m}}\right) \right] (1/\sqrt{2my}). \quad (6)$$

Instead of going through the d.f. (which process requires monotonicity of the transformation $y = h(x)$), under certain conditions, f_Y may be obtained directly from f_X . Such conditions are described in the following theorem.

THEOREM 2

Let X be a r.v. with positive and continuous p.d.f. on the set $S \subseteq \mathfrak{R}$, and let $h : S \rightarrow T$ (the image of S under h) be a one-to-one transformation, so that the inverse $x = h^{-1}(y)$, $y \in T$, exists. Suppose that, for $y \in T$, the derivative $\frac{d}{dy} h^{-1}(y)$ exists, is continuous, and $\neq 0$. Then the p.d.f. of the r.v. $Y = h(X)$ is given by:

$$f_Y(y) = f_X[h^{-1}(y)] \left| \frac{d}{dy} h^{-1}(y) \right|, \quad y \in T \text{ (and } = 0 \text{ for } y \notin T). \quad (7)$$

PROOF (rough outline) Let $B = [c, d]$ be an interval in T and suppose B is transformed into the interval $A = [a, b]$ by the inverse transformation $x = h^{-1}(y)$. Then:

$$P(Y \in B) = P[h(X) \in B] = P(X \in A) = \int_A f_X(x) dx.$$

When transforming x into y through the transformation $x = h^{-1}(y)$, $\int_A f_X(x) dx = \int_B f_X[h^{-1}(y)] \left| \frac{d}{dy} h^{-1}(y) \right| dy$, according to the theory of changing variables in integrals. Thus,

$$P(Y \in B) = \int_B f_X[h^{-1}(y)] \left| \frac{d}{dy} h^{-1}(y) \right| dy,$$

which implies that the integrand is the p.d.f. of Y . \blacktriangle

Relation (7) has already been illustrated by Example 1. A slightly more general case is the following one.

EXAMPLE 3

Determine the p.d.f. of the r.v. Y defined by: $Y = aX + b$ ($a \neq 0$). In particular, determine f_Y , if $X \sim N(\mu, \sigma^2)$.

DISCUSSION The transformation $y = ax + b$ gives $x = h^{-1}(y) = \frac{y-b}{a}$, so that $\frac{dx}{dy} = \frac{d}{dy}h^{-1}(y) = \frac{1}{a}$. Therefore: $f_Y(y) = f_X(\frac{y-b}{a})\frac{1}{|a|}$. For the special case:

$$f_Y(y) = \frac{1}{\sqrt{2\pi}|a|\sigma} \exp\left[-\frac{(\frac{y-b}{a} - \mu)^2}{2\sigma^2}\right] = \frac{1}{\sqrt{2\pi}|a|\sigma} \exp\left\{-\frac{[y - (a\mu + b)]^2}{2(a\sigma)^2}\right\}.$$

Thus, if $X \sim N(\mu, \sigma^2)$, then $Y = aX + b \sim N(a\mu + b, (a\sigma)^2)$.

A modification of Theorem 2 when the assumption that $h : S \rightarrow T$ is one-to-one is not satisfied, but a version of it is, is stated in the following result. This result has already been illustrated by (6) in connection with Example 2.

THEOREM 3

Let X be a r.v. with positive and continuous p.d.f. on the set $S \subseteq \mathfrak{R}$, and suppose that the transformation $h : S \rightarrow T$ is *not* one-to-one. Suppose further that when S is partitioned into the pairwise disjoint subsets S_1, \dots, S_r and h is restricted to S_j and takes values in T_j (the image of S_j under h), then h is one-to-one. Denoting by h_j this restriction, we have then: $h_j : S_j \rightarrow T_j$ is one-to-one, so that the inverse $x = h_j^{-1}(y)$, $y \in T_j$, exists, $j = 1, \dots, r$. Finally, we suppose that, for any $y \in T_j$, $j = 1, \dots, r$, the derivatives $\frac{d}{dy}h_j^{-1}(y)$ exist, are continuous, and $\neq 0$. Then the p.d.f. of the r.v. $Y = h(X)$ is determined as follows: Set

$$f_{Y_j} = f_X[h_j^{-1}(y)] \left| \frac{d}{dy}h_j^{-1}(y) \right|, \quad y \in T_j, \quad j = 1, \dots, r,$$

and for $y \in T$, suppose that y belongs to k of the r T_j 's, $1 \leq k \leq r$. Then $f_Y(y)$ is the sum of the corresponding k $f_{Y_j}(y)$'s. Alternatively,

$$f_Y(y) = \sum_{j=1}^r \delta_j(y) f_{Y_j}(y), \quad y \in T \quad (\text{and } = 0 \text{ for } y \notin T), \quad (8)$$

where $\delta_j(y) = 1$, if $y \in T_j$ and $\delta_j(y) = 0$, if $y \notin T_j$, $j = 1, \dots, r$.

REMARK 1 It is to be noticed that, whereas the subsets S_1, \dots, S_r are pairwise disjoint, their images T_1, \dots, T_r need not be so. For instance, in Example 2, $S_1 = (0, \infty)$, $S_2 = (-\infty, 0)$ but $T_1 = T_2 = (0, \infty)$.

Exercises

- 1.1 The r.v. X has p.d.f. $f_X(x) = (1 - \alpha)\alpha^x$, $x = 0, 1, \dots$ ($0 < \alpha < 1$), and set $Y = X^3$. Determine the p.d.f. f_Y .
- 1.2 Let the r.v.'s X and Y represent the temperature of a certain object in degrees Celsius and Fahrenheit, respectively. Then, it is known that $Y = \frac{9}{5}X + 32$ and $X = \frac{5}{9}Y - \frac{160}{9}$.

- (i) If $Y \sim N(\mu, \sigma^2)$, determine the distribution of X .
- (ii) If $P(90 \leq Y \leq 95) = 0.95$, then also $P(a \leq X \leq b) = 0.95$, for some $a < b$. Determine the numbers a and b .
- (iii) We know that: $P(\mu - \sigma \leq Y \leq \mu + \sigma) \simeq 0.6827 = p_1$, $P(\mu - 2\sigma \leq Y \leq \mu + 2\sigma) \simeq 0.9545 = p_2$, and $P(\mu - 3\sigma \leq Y \leq \mu + 3\sigma) \simeq 0.9973 = p_3$. Calculate the intervals $[a_k, b_k]$, $k = 1, 2, 3$ for which $P(a_k \leq X \leq b_k)$ is, respectively, equal to p_k , $k = 1, 2, 3$.
- 1.3** Let the r.v. X have p.d.f. f_X positive on the set $S \subseteq \mathfrak{R}$, and set $U = aX + b$, where a and b are constants and $a > 0$.
- (i) Use Theorem 2 in order to derive the p.d.f. f_U .
- (ii) If X has the Negative Exponential distribution with parameter λ , show that U has the same kind of distribution with parameter λ/a .
- (iii) If $X \sim U(c, d)$, then show that $U \sim U(ac + b, ad + b)$.
- 1.4** If the r.v. X has the Negative Exponential distribution with parameter λ , set $Y = e^X$ and $Z = \log X$ and determine the p.d.f.'s f_Y and f_Z .
- 1.5** Let $X \sim U(\alpha, \beta)$ and set $Y = e^X$. Then determine the p.d.f. f_Y . If $\alpha > 0$, set $Z = \log X$ and determine the p.d.f. f_Z .
- 1.6** (i) If the r.v. X is distributed as $U(0, 1)$ and $Y = -2 \log X$, show that Y is distributed as χ_2^2 .
- (ii) If X_1, \dots, X_n is a random sample from the $U(0, 1)$ distribution and $Y_i = -2 \log X_i$, use part (i) and the m.g.f. approach in order to show that $\sum_{i=1}^n Y_i$ is distributed as χ_{2n}^2 .
- 1.7** If the r.v. X has the p.d.f. $f_X(x) = \frac{1}{\sqrt{2\pi}} x^{-2} e^{-1/2x^2}$, $x \in \mathfrak{R}$, show that the r.v. $Y = \frac{1}{X} \sim N(0, 1)$.
- 1.8** Suppose that the velocity of a molecule of mass m is a r.v. X with p.d.f. $f_X(x) = \sqrt{\frac{2}{\pi}} x^2 e^{-x^2/2}$, $x > 0$ (the so-called Maxwell distribution). Derive the p.d.f. of the r.v. $Y = \frac{1}{2} m X^2$, which is the kinetic energy of the molecule.
- 1.9** If the r.v. $X \sim N(0, 1)$, use Theorem 3 in order to show that the r.v. $Y = X^2 \sim \chi_1^2$.
- 1.10** Let X_r be a r.v. distributed as t with r degrees of freedom: $X_r \sim t_r$ ($r = 1, 2, \dots$) whose p.d.f. is given in relation (10) below. Then show that:
- (i) EX_r does not exist for $r = 1$.
- (ii) $EX_r = 0$ for $r \geq 2$.
- (iii) $Var(X_r) = \frac{r}{r-2}$ for $r \geq 3$.

Hint: That EX_r does not exist for $r = 1$ is, actually, reduced to Exercise 1.16 in Chapter 3. That $EX_r = 0$ for $r \geq 2$ follows by a simple integration. So, all that remains to calculate is EX_r^2 . For this purpose, first reduce the original integral to an integral over the interval $(0, \infty)$, by symmetry of the region of integration and the fact that the integrand is an even function. Then, use the transformation $\frac{t^2}{r} = x$, and

next the transformation $\frac{1}{1+x} = y$. Except for constants, the integral is then reduced to the form

$$\int_0^1 y^{\alpha-1}(1-y)^{\beta-1}dy \quad (\alpha > 0, \beta > 0).$$

At this point, use the following fact:

$$\int_0^1 y^{\alpha-1}(1-y)^{\beta-1}dy = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}.$$

(A proof of this fact may be found, e.g., in pages 70–71, of the book *A Course in Mathematical Statistics*, 2nd edition (1967), Academic Press, by G. G. Roussas.) The proof is concluded by using the recursive relation of the Gamma function ($\Gamma(\gamma) = (\gamma - 1)\Gamma(\gamma - 1)$) and the fact that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$.

6.2 Transforming Two or More Random Variables

Often the need arises to transform two or more given r.v.'s to another set of r.v.'s. The following examples illustrate the point.

EXAMPLE 4

The times of arrival of a bus at two successive bus stops are r.v.'s X_1 and X_2 distributed as $U(\alpha, \beta)$, for two time points $\alpha < \beta$. Calculate the probabilities $P(X_1 + X_2 > x)$ for $2\alpha < x < 2\beta$.

Clearly, this question calls for the determination of the distribution of the r.v. $X_1 + X_2$.

Or more generally (and more realistically), suppose that a bus makes k stops between its depot and its terminal, and that the arrival time at the i th stop is a r.v. $X_i \sim U(\alpha_i, \beta_i)$, $\alpha_i < \beta_i$, $i = 1, \dots, k+1$ (where X_{k+1} is the time of arrival at the terminal). Determine the distribution of the duration of the trip $X_1 + \dots + X_{k+1}$.

EXAMPLE 5

Consider certain events occurring in every time interval $[t_1, t_2]$ ($0 < t_1 < t_2$) according to the Poisson distribution $P(\lambda(t_2 - t_1))$. Then the waiting times between successive occurrences are independent r.v.'s distributed according to the Negative Exponential distribution with parameter λ . Let X_1 and X_2 be two such times. What is the probability that one would have to wait at least twice as long for the second occurrence than the first? That is, what is the probability $P(X_2 > 2X_1)$?

Here one would have to compute the distribution of the r.v. $X_2 - 2X_1$.

Below, a brief outline of the theory underpinning the questions posed in the examples is presented. First, consider the case of two r.v.'s X_1 and X_2 having the joint p.d.f. f_{X_1, X_2} . Often the question posed is that of determining the distribution of a function of X_1 and X_2 , $h_1(X_1, X_2)$. The general approach is to set $Y_1 = h_1(X_1, X_2)$ and also consider another (convenient) transformation $Y_2 = h_2(X_1, X_2)$. Next, determine the joint p.d.f. of Y_1 and Y_2 , f_{Y_1, Y_2} , and, finally, compute the (marginal) p.d.f. f_{Y_1} . Conditions under which f_{Y_1, Y_2} is determined by way of f_{X_1, X_2} are given below.

THEOREM 4

Consider the r.v.'s X_1 and X_2 with joint p.d.f. f_{X_1, X_2} positive and continuous on the set $S \subseteq \mathfrak{R}^2$, and let h_1, h_2 be two real-valued transformations defined on S ; that is, $h_1, h_2: S \rightarrow \mathfrak{R}$, and let T be the image of S under the transformation (h_1, h_2) . Suppose that (h_1, h_2) is one-to-one from S onto T . Thus, if we set $y_1 = h_1(x_1, x_2)$ and $y_2 = h_2(x_1, x_2)$, we can solve uniquely for $x_1, x_2: x_1 = g_1(y_1, y_2)$, $x_2 = g_2(y_1, y_2)$. Suppose further that the partial derivatives $g_{1i}(y_1, y_2) = \frac{\partial}{\partial y_i} g_1(y_1, y_2)$ and $g_{2i}(y_1, y_2) = \frac{\partial}{\partial y_i} g_2(y_1, y_2)$, $i = 1, 2$ exist and are continuous for $(y_1, y_2) \in T$. Finally, suppose that the Jacobian $J = \begin{vmatrix} g_{11}(y_1, y_2) & g_{12}(y_1, y_2) \\ g_{21}(y_1, y_2) & g_{22}(y_1, y_2) \end{vmatrix}$ is $\neq 0$ on T . Then the joint p.d.f. of the r.v.'s $Y_1 = h_1(X_1, X_2)$ and $Y_2 = h_2(X_1, X_2)$, f_{Y_1, Y_2} , is given by:

$$f_{Y_1, Y_2}(y_1, y_2) = f_{X_1, X_2}[g_1(y_1, y_2), g_2(y_1, y_2)]|J|, \quad (y_1, y_2) \in T \quad (9)$$

(and = 0 for $(y_1, y_2) \notin T$).

The justification of this theorem is entirely analogous to that of Theorem 2 and will be omitted.

In applying Theorem 4, one must be careful in checking that the underlying assumptions hold and in determining correctly the set T . As an illustration, let us discuss the first part of Example 4.

EXAMPLE 4

(continued) **Discussion** We have $y_1 = x_1 + x_2$ and let $y_2 = x_2$. Then $x_1 = y_1 - y_2$ and $x_2 = y_2$, so that $\frac{\partial x_1}{\partial y_1} = 1$, $\frac{\partial x_1}{\partial y_2} = -1$, $\frac{\partial x_2}{\partial y_1} = 0$, $\frac{\partial x_2}{\partial y_2} = 1$, and $J = \begin{vmatrix} 1 & -1 \\ 0 & 1 \end{vmatrix} = 1$. For the determination of S and T , see Figures 6.1 and 6.2.

Since $f_{X_1, X_2}(x_1, x_2) = \frac{1}{(\beta - \alpha)^2}$ for $(x_1, x_2) \in S$, we have $f_{Y_1, Y_2}(y_1, y_2) = \frac{1}{(\beta - \alpha)^2}$ for $(y_1, y_2) \in T$; that is, for $2\alpha < y_1 < 2\beta$, $\alpha < y_2 < \beta$, $\alpha < y_1 - y_2 < \beta$ (and = 0 for $(y_1, y_2) \notin T$).

Thus, we get:

$$f_{Y_1, Y_2}(y_1, y_2) = \begin{cases} \frac{1}{(\beta - \alpha)^2}, & 2\alpha < y_1 < 2\beta, \alpha < y_2 < \beta, \alpha < y_1 - y_2 < \beta \\ 0, & \text{otherwise.} \end{cases}$$

Figure 6.1

$$S = \{(x_1, x_2) \in \mathfrak{R}^2; f_{x_1, x_2}(x_1, x_2) > 0\}$$





Figure 6.2

$T =$ Image of S
Under the
Transformation
Used



Therefore:

$$f_{Y_1}(y_1) = \begin{cases} \frac{1}{(\beta-\alpha)^2} \int_{\alpha}^{y_1-\alpha} dy_2 = \frac{y_1-2\alpha}{(\beta-\alpha)^2}, & \text{for } 2\alpha < y_1 \leq \alpha + \beta \\ \frac{1}{(\beta-\alpha)^2} \int_{y_1-\beta}^{\beta} dy_2 = \frac{2\beta-y_1}{(\beta-\alpha)^2}, & \text{for } \alpha + \beta < y_1 \leq 2\beta \\ 0, & \text{otherwise.} \end{cases}$$


The graph of f_{Y_1} is given in Figure 6.3.

EXAMPLE 5

(continued) **Discussion** Here $y_1 = x_2 - 2x_1 = -2x_1 + x_2$ and let $y_2 = x_2$. Then $x_1 = -\frac{1}{2}y_1 + \frac{1}{2}y_2$ and $x_2 = y_2$, so that $J = \begin{vmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 \end{vmatrix} = -\frac{1}{2}$ and $|J| = \frac{1}{2}$. Clearly, S is the first quadrant. As for T , we have $y_2 = x_2$, so that $y_2 > 0$.


Figure 6.3

This Density Is Known as the *Triangular p.d.f.*



Also, $-\frac{1}{2}y_1 + \frac{1}{2}y_2 = x_1$, so that $-\frac{1}{2}y_1 + \frac{1}{2}y_2 > 0$ or $-y_1 + y_2 > 0$ or $y_2 > y_1$. The conditions $y_2 > 0$ and $y_2 > y_1$ determine T (see Figure 6.4).

Figure 6.4



T is the part of the plane above the y_1 -axis and also above the main diagonal $y_1 = y_2$.

Since $f_{X_1, X_2}(x_1, x_2) = \lambda^2 e^{-\lambda(x_1+x_2)}$ ($x_1, x_2 > 0$), we have $f_{Y_1, Y_2}(y_1, y_2) = \frac{\lambda^2}{2} e^{\frac{\lambda}{2}y_1 - \frac{3\lambda}{2}y_2}$, $(y_1, y_2) \in T$ (and = 0 otherwise). Therefore $f_{Y_1}(y_1)$ is taken by integrating out y_2 . More precisely, for $y_1 < 0$:

$$\begin{aligned} f_{Y_1}(y_1) &= \frac{\lambda^2}{2} e^{\frac{\lambda}{2}y_1} \int_0^{\infty} e^{-\frac{3\lambda}{2}y_2} dy_2 = -\frac{\lambda^2}{2} \times \frac{2}{3\lambda} e^{\frac{\lambda}{2}y_1} \times e^{-\frac{3\lambda}{2}y_2} \Big|_0^{\infty} \\ &= -\frac{\lambda}{3} e^{\frac{\lambda}{2}y_1} (0 - 1) = \frac{\lambda}{3} e^{\frac{\lambda}{2}y_1}, \end{aligned}$$

whereas for $y_1 > 0$:

$$\begin{aligned} f_{Y_1}(y_1) &= \frac{\lambda^2}{2} e^{\frac{\lambda}{2} y_1} \int_{y_1}^{\infty} e^{-\frac{3\lambda}{2} y_2} dy_2 = -\frac{\lambda}{3} e^{\frac{\lambda}{2} y_1} \times e^{-\frac{3\lambda}{2} y_2} \Big|_{y_1}^{\infty} \\ &= -\frac{\lambda}{3} e^{\frac{\lambda}{2} y_1} (0 - e^{-\frac{3\lambda}{2} y_1}) = \frac{\lambda}{3} e^{-\lambda y_1}. \end{aligned}$$

To summarize:

$$f_{Y_1}(y_1) = \begin{cases} \frac{\lambda}{3} e^{\frac{\lambda}{2} y_1}, & y_1 < 0 \\ \frac{\lambda}{3} e^{-\lambda y_1}, & y_1 \geq 0. \end{cases}$$

Therefore $P(X_2 > 2X_1) = P(X_2 - 2X_1 > 0) = P(Y_1 > 0) = \frac{\lambda}{3} \int_0^{\infty} e^{-\lambda y_1} dy_1 = \frac{1}{3}$.

REMARK 2 To be sure, the preceding probability is also calculated as follows:

$$\begin{aligned} P(X_2 > 2X_1) &= \int \int_{(x_2 > 2x_1)} \lambda^2 e^{-\lambda x_1 - \lambda x_2} dx_1 dx_2 \\ &= \int_0^{\infty} \lambda e^{-\lambda x_2} \left(\int_0^{x_2/2} \lambda e^{-\lambda x_1} dx_1 \right) dx_2 \\ &= \int_0^{\infty} \lambda e^{-\lambda x_2} (1 - e^{-\frac{\lambda}{2} x_2}) dx_2 \\ &= \int_0^{\infty} \lambda e^{-\lambda x_2} dx_2 - \frac{2}{3} \int_0^{\infty} \frac{3\lambda}{2} e^{-\frac{3\lambda}{2} x_2} dx_2 = 1 - \frac{2}{3} = \frac{1}{3}. \end{aligned}$$

Applications of Theorem 4 lead to two new distributions, which are of great importance in statistics. They are the t -distribution and the F -distribution.

DEFINITION 1

Let X and Y be two independent r.v.'s distributed as follows: $X \sim N(0, 1)$ and $Y \sim \chi_r^2$, and define the r.v. T by: $T = X/\sqrt{Y/r}$. The r.v. T is said to have the (Student's) t -distribution with r degrees of freedom (d.f.). The notation used is: $T \sim t_r$.


The p.d.f. of T , f_T , is given by the formula:

$$f_T(t) = \frac{\Gamma[\frac{1}{2}(r+1)]}{\sqrt{\pi r} \Gamma(r/2)} \times \frac{1}{[1 + (t^2/r)]^{(1/2)(r+1)}}, \quad t \in \Re, \quad (10)$$

and its graph (for $r = 5$) is presented in Figure 6.5.

From formula (10), it is immediate that f_T is symmetric about 0 and tends to 0 as $t \rightarrow \pm\infty$. It can also be seen (see Exercise 2.10) that $f_T(t)$ tends to the p.d.f. of the $N(0, 1)$ distribution as the number r of d.f. tends to ∞ . This is

Figure 6.5
Two Curves of the t
Probability Density
Function



depicted in Figure 6.5 by means of the curve denoted by t_∞ . Also, it is seen (see Exercise 2.9) that $ET = 0$ for $r \geq 2$, and $\text{Var}(T) = \frac{r}{r-2}$ for $r \geq 3$. Finally, the probabilities $P(T \leq t)$, for selected values of t and r , are given by tables (the t -tables). For $r \geq 91$, one may use the tables for the standard Normal distribution.

Regarding the derivation of f_T , we have:

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-(1/2)x^2}, \quad x \in \mathfrak{R},$$

$$f_Y(y) = \begin{cases} \frac{1}{\Gamma(\frac{1}{2}r)2^{(1/2)r}} y^{(r/2)-1} e^{-y/2}, & y > 0 \\ 0, & y \leq 0. \end{cases}$$

Set $U = Y$ and consider the transformation

$$(h_1, h_2): \begin{cases} t = \frac{x}{\sqrt{y/r}}; \\ u = y \end{cases}; \quad \text{then} \quad \begin{cases} x = \frac{1}{\sqrt{r}} t \sqrt{u} \\ y = u, \end{cases}$$

and

$$J = \begin{vmatrix} \frac{\sqrt{u}}{\sqrt{r}} & \frac{t}{2\sqrt{u}\sqrt{r}} \\ 0 & 1 \end{vmatrix} = \frac{\sqrt{u}}{\sqrt{r}}.$$

Therefore, for $t \in \mathfrak{R}$, $u > 0$, we get

$$\begin{aligned} f_{T,U}(t, u) &= \frac{1}{\sqrt{2\pi}} e^{-t^2 u/2r} \times \frac{1}{\Gamma(r/2)2^{r/2}} u^{(r/2)-1} e^{-u/2} \times \frac{\sqrt{u}}{\sqrt{r}} \\ &= \frac{1}{\sqrt{2\pi r} \Gamma(r/2)2^{r/2}} u^{(1/2)(r+1)-1} \exp \left[-\frac{u}{2} \left(1 + \frac{t^2}{r} \right) \right]. \end{aligned}$$

Hence

$$f_T(t) = \int_0^\infty \frac{1}{\sqrt{2\pi r} \Gamma(r/2) 2^{r/2}} u^{(1/2)(r+1)-1} \exp\left[-\frac{u}{2}\left(1 + \frac{t^2}{r}\right)\right] du.$$

Set

$$\frac{u}{2}\left(1 + \frac{t^2}{r}\right) = z, \quad \text{so that} \quad u = 2z\left(1 + \frac{t^2}{r}\right)^{-1}, \quad du = 2\left(1 + \frac{t^2}{r}\right)^{-1} dz,$$

and $z \in [0, \infty)$. Therefore we continue as follows:

$$\begin{aligned} f_T(t) &= \int_0^\infty \frac{1}{\sqrt{2\pi r} \Gamma(r/2) 2^{r/2}} \left[\frac{2z}{1 + (t^2/r)}\right]^{(1/2)(r+1)-1} e^{-z} \frac{2}{1 + (t^2/r)} dz \\ &= \frac{1}{\sqrt{2\pi r} \Gamma(r/2) 2^{r/2}} \frac{2^{(1/2)(r+1)}}{[1 + (t^2/r)]^{(1/2)(r+1)}} \int_0^\infty z^{(1/2)(r+1)-1} e^{-z} dz \\ &= \frac{1}{\sqrt{\pi r} \Gamma(r/2)} \frac{1}{[1 + (t^2/r)]^{(1/2)(r+1)}} \Gamma\left[\frac{1}{2}(r+1)\right], \end{aligned}$$

since $\frac{1}{\Gamma[\frac{1}{2}(r+1)]} z^{(1/2)(r+1)-1} e^{-z}$ ($z > 0$) is the p.d.f. of the Gamma distribution with parameters $\alpha = \frac{r+1}{2}$ and $\beta = 1$; that is,

$$f_T(t) = \frac{\Gamma[\frac{1}{2}(r+1)]}{\sqrt{\pi r} \Gamma(r/2)} \times \frac{1}{[1 + (t^2/r)]^{(1/2)(r+1)}}, \quad t \in \mathfrak{R}.$$

Now, we proceed with the definition of the F -distribution.

DEFINITION 2


Let X and Y be two independent r.v.'s distributed as follows: $X \sim \chi_{r_1}^2$ and $Y \sim \chi_{r_2}^2$, and define the r.v. F by: $F = \frac{X/r_1}{Y/r_2}$. The r.v. F is said to have the F -distribution with r_1 and r_2 degrees of freedom (d.f.). The notation often used is: $F \sim F_{r_1, r_2}$.

The p.d.f. of F , f_F , is given by the formula:

$$f_F(f) = \begin{cases} \frac{\Gamma[\frac{1}{2}(r_1+r_2)](r_1/r_2)^{r_1/2}}{\Gamma(\frac{1}{2}r_1)\Gamma(\frac{1}{2}r_2)} \times \frac{f^{r_1/2-1}}{[1+(r_1/r_2)f]^{(1/2)(r_1+r_2)}}, & \text{for } f > 0 \\ 0, & \text{for } f \leq 0, \end{cases} \quad (11)$$

and its graphs (for $r_1 = 10, r_2 = 4$ and $r_1 = r_2 = 10$) are given in Figure 6.6. The probabilities $P(F \leq f)$, for selected values of f and r_1, r_2 , are given by tables (the F -tables).

Figure 6.6
Two Curves of the F
Probability Density
Function



The derivation of f_F is based on Theorem 4 and is as follows. For x and $y > 0$, we have:

$$f_X(x) = \frac{1}{\Gamma(\frac{1}{2}r_1)2^{r_1/2}} x^{(r_1/2)-1} e^{-x/2}, \quad x > 0,$$

$$f_Y(y) = \frac{1}{\Gamma(\frac{1}{2}r_2)2^{r_2/2}} y^{(r_2/2)-1} e^{-y/2}, \quad y > 0.$$

We set $Z = Y$, and consider the transformation

$$(h_1, h_2) : \begin{cases} f = \frac{x/r_1}{y/r_2}; \\ z = y \end{cases} \quad \text{then} \quad \begin{cases} x = \frac{r_1}{r_2} f z \\ y = z, \end{cases}$$

and

$$J = \begin{vmatrix} \frac{r_1}{r_2} z & \frac{r_1}{r_2} f \\ 0 & 1 \end{vmatrix} = \frac{r_1}{r_2} z, \quad \text{so that } |J| = \frac{r_1}{r_2} z.$$

For $f, z > 0$, we get:

$$\begin{aligned} f_{F,Z}(f, z) &= \frac{1}{\Gamma(\frac{1}{2}r_1)\Gamma(\frac{1}{2}r_2)2^{(1/2)(r_1+r_2)}} \left(\frac{r_1}{r_2}\right)^{(r_1/2)-1} f^{(r_1/2)-1} z^{(r_1/2)-1} z^{(r_2/2)-1} \\ &\quad \times \exp\left(-\frac{r_1}{2r_2}\right) f z e^{-z/2} \frac{r_1}{r_2} z \\ &= \frac{(r_1/r_2)^{r_1/2} f^{(r_1/2)-1}}{\Gamma(\frac{1}{2}r_1)\Gamma(\frac{1}{2}r_2)2^{(1/2)(r_1+r_2)}} z^{(1/2)(r_1+r_2)-1} \exp\left[-\frac{z}{2}\left(\frac{r_1}{r_2}f + 1\right)\right]. \end{aligned}$$

Therefore

$$\begin{aligned} f_F(f) &= \int_0^\infty f_{F,Z}(f, z) dz \\ &= \frac{(r_1/r_2)^{r_1/2} f^{(r_1/2)-1}}{\Gamma(\frac{1}{2}r_1)\Gamma(\frac{1}{2}r_2)2^{(1/2)(r_1+r_2)}} \int_0^\infty z^{(1/2)(r_1+r_2)-1} \exp\left[-\frac{z}{2}\left(\frac{r_1}{r_2}f + 1\right)\right] dz. \end{aligned}$$

Set

$$\frac{z}{2}\left(\frac{r_1}{r_2}f + 1\right) = t, \quad \text{so that } z = 2t\left(\frac{r_1}{r_2}f + 1\right)^{-1},$$

$$dz = 2\left(\frac{r_1}{r_2}f + 1\right)^{-1} dt, \quad t \in [0, \infty).$$

Thus continuing, we have

$$\begin{aligned} f_F(f) &= \frac{(r_1/r_2)^{r_1/2} f^{(r_1/2)-1}}{\Gamma(\frac{1}{2}r_1)\Gamma(\frac{1}{2}r_2)2^{(1/2)(r_1+r_2)}} 2^{(1/2)(r_1+r_2)-1} \left(\frac{r_1}{r_2}f + 1\right)^{-(1/2)(r_1+r_2)+1} \\ &\quad \times 2\left(\frac{r_1}{r_2}f + 1\right)^{-1} \int_0^\infty t^{(1/2)(r_1+r_2)-1} e^{-t} dt \\ &= \frac{\Gamma[\frac{1}{2}(r_1 + r_2)](r_1/r_2)^{r_1/2}}{\Gamma(\frac{1}{2}r_1)\Gamma(\frac{1}{2}r_2)} \times \frac{f^{(r_1/2)-1}}{[1 + (r_1/r_2)f]^{(1/2)(r_1+r_2)}}, \end{aligned}$$

since $\frac{1}{\Gamma[\frac{1}{2}(r_1+r_2)]} t^{(1/2)(r_1+r_2)-1} e^{-t}$ ($t > 0$) is the p.d.f. of the Gamma distribution with parameters $\alpha = \frac{r_1+r_2}{2}$ and $\beta = 1$. Therefore

$$f_F(f) = \begin{cases} \frac{\Gamma[\frac{1}{2}(r_1+r_2)](r_1/r_2)^{r_1/2}}{\Gamma(\frac{1}{2}r_1)\Gamma(\frac{1}{2}r_2)} \times \frac{f^{(r_1/2)-1}}{[1+(r_1/r_2)f]^{(1/2)(r_1+r_2)}}, & \text{for } f > 0 \\ 0, & \text{for } f \leq 0. \end{cases}$$

REMARK 3

- (i) From the definition of the F -distribution, it follows that, if $F \sim F_{r_1, r_2}$, then $\frac{1}{F} \sim F_{r_2, r_1}$.
- (ii) If $T \sim t_r$, then $T^2 \sim F_{1, r}$. Indeed, $T = X/\sqrt{Y/r}$, where X and Y are independent, and $X \sim N(0, 1)$, $Y \sim \chi_r^2$. But then $T^2 = \frac{X^2}{Y/r} = \frac{X^2/1}{Y/r} \sim F_{1, r}$, since $X^2 \sim \chi_1^2$ and X^2 and Y are independent.
- (iii) If $F \sim F_{r_1, r_2}$, then it can be shown (see Exercise 2.10) that

$$EF = \frac{r_2}{r_2 - 2}, \quad \text{for } r_2 \geq 3, \quad \text{and } \text{Var}(F) = \frac{2r_2^2(r_1 + r_2 - 2)}{r_1(r_2 - 2)^2(r_2 - 4)}, \quad \text{for } r_2 \geq 5.$$

One can formulate a version of Theorem 4 for $k(>2)$ r.v.'s X_1, \dots, X_k , as well as a version of Theorem 3. In the following, such versions are formulated for reference purposes.

THEOREM 5

Consider the r.v.'s X_1, \dots, X_k with joint p.d.f. f_{X_1, \dots, X_k} positive and continuous on the set $S \subseteq \mathfrak{R}^k$, and let h_1, \dots, h_k be real-valued transformations defined on S ; that is, $h_1, \dots, h_k : S \rightarrow \mathfrak{R}$, and let T be the image of S under the transformation (h_1, \dots, h_k) . Suppose that (h_1, \dots, h_k) is one-to-one from S onto T . Thus, if we set $y_i = h_i(x_1, \dots, x_k)$, $i = 1, \dots, k$, then we can solve uniquely for x_i , $i = 1, \dots, k$: $x_i = g_i(y_1, \dots, y_k)$, $i = 1, \dots, k$. Suppose further that the partial derivatives $g_{ij}(y_1, \dots, y_k) = \frac{\partial}{\partial y_j} g_i(y_1, \dots, y_k)$, $i, j = 1, \dots, k$ exist and are continuous for $(y_1, \dots, y_k) \in T$. Finally, suppose that the Jacobian

$$J = \begin{vmatrix} g_{11}(y_1, \dots, y_k) & \cdots & g_{1k}(y_1, \dots, y_k) \\ \dots & \cdots & \dots \\ g_{k1}(y_1, \dots, y_k) & \cdots & g_{kk}(y_1, \dots, y_k) \end{vmatrix}$$

is $\neq 0$ on T . Then the joint p.d.f. of the r.v.'s $Y_i = h_i(X_1, \dots, X_k)$, $i = 1, \dots, k$, f_{Y_1, \dots, Y_k} , is given by:

$$f_{Y_1, \dots, Y_k}(y_1, \dots, y_k) = f_{X_1, \dots, X_k}[g_1(y_1, \dots, y_k), \dots, g_k(y_1, \dots, y_k)] \cdot |J|, \\ (y_1, \dots, y_k) \in T \quad (\text{and } = 0 \text{ for } (y_1, \dots, y_k) \notin T). \quad (12)$$

A suitable version of the previous result when the transformations h_1, \dots, h_k are not one-to-one is stated below; it will be employed in Theorem 12 in Section 5.

THEOREM 6

Let X_1, \dots, X_k be r.v.'s with joint p.d.f. f_{X_1, \dots, X_k} positive and continuous on the set $S \subseteq \mathfrak{R}^k$, and let h_1, \dots, h_k be real-valued transformations defined on S ; that is, $h_1, \dots, h_k : S \rightarrow \mathfrak{R}$, and let T be the image of S under the transformation (h_1, \dots, h_k) . Suppose that (h_1, \dots, h_k) is *not* one-to-one from S onto T but there is a partition of S into (pairwise disjoint) subsets S_1, \dots, S_r such that when (h_1, \dots, h_k) is restricted to S_j and takes values in T_j (the image of S_j under (h_1, \dots, h_k)), $j = 1, \dots, r$, then (h_1, \dots, h_k) is one-to-one. Denoting by (h_{1j}, \dots, h_{kj}) this restriction, we have then: $(h_{1j}, \dots, h_{kj}) : S_j \rightarrow T_j$ is one-to-one, so that we can solve uniquely for x_i , $i = 1, \dots, k$: $x_i = g_{ji}(y_1, \dots, y_k)$, $i = 1, \dots, k$, for each $j = 1, \dots, r$. Suppose further that the partial derivatives $g_{jil}(y_1, \dots, y_k) = \frac{\partial}{\partial y_l} g_{ji}(y_1, \dots, y_k)$, $i, l = 1, \dots, k$, $j = 1, \dots, r$ exist and are continuous for $(y_1, \dots, y_k) \in T_j$, $j = 1, \dots, r$, and the Jacobian

$$J_j = \begin{vmatrix} g_{j11}(y_1, \dots, y_k) & \cdots & g_{j1k}(y_1, \dots, y_k) \\ \dots & \cdots & \dots \\ g_{jk1}(y_1, \dots, y_k) & \cdots & g_{jkk}(y_1, \dots, y_k) \end{vmatrix}$$

is $\neq 0$ on T_j for $j = 1, \dots, r$.

Set

$$f_{Y_j}(y_1, \dots, y_k) = f_{X_1, \dots, X_k}[g_{j1}(y_1, \dots, y_k), \dots, g_{jk}(y_1, \dots, y_k)]|J_j|, \\ (y_1, \dots, y_k) \in T_j, \quad j = 1, \dots, r.$$

Then the joint p.d.f. of the r.v.'s $Y_i = h_i(X_1, \dots, X_k)$, $i = 1, \dots, k$, f_{Y_1, \dots, Y_k} , is given by:

$$f_{Y_1, \dots, Y_k}(y_1, \dots, y_k) = \sum_{j=1}^r \delta_j(y_1, \dots, y_k) f_{Y_j}(y_1, \dots, y_k), \quad (y_1, \dots, y_k) \in T \\ (\text{and } = 0 \text{ for } (y_1, \dots, y_k) \notin T), \quad (13)$$

where $\delta_j(y_1, \dots, y_k) = 1$, if $(y_1, \dots, y_k) \in T_j$ and $\delta_j(y_1, \dots, y_k) = 0$, if $(y_1, \dots, y_k) \notin T_j$, $j = 1, \dots, r$.

Exercises

- 2.1** The r.v.'s X and Y denote the outcomes of one independent throw of two fair dice, and let $Z = X + Y$. Determine the distribution of Z .
- 2.2** Let the independent r.v.'s X and Y have the Negative Exponential distribution with $\lambda = 1$, and set $U = X + Y$, $V = X/Y$.
- Derive the joint p.d.f. $f_{U,V}$.
 - Then derive the marginal p.d.f.'s f_U and f_V .
 - Show that the r.v.'s U and V are independent.
- 2.3** Let the independent r.v.'s X and Y have the Negative Exponential distribution with $\lambda = 1$, and set $U = \frac{1}{2}(X + Y)$, $V = \frac{1}{2}(X - Y)$.
- Show that the joint p.d.f. of the r.v.'s U and V is given by:

$$f_{U,V}(u, v) = 2e^{-2u}, \quad -u < v < u, \quad u > 0.$$

- Also, show that the marginal p.d.f.'s f_U and f_V are given by:

$$f_U(u) = 4ue^{-2u}, \quad u > 0; \quad f_V(v) = e^{-2v}, \quad \text{for } v > 0,$$

$$f_V(v) = e^{2v}, \quad \text{for } v < 0.$$

- 2.4** Let the independent r.v.'s X and Y have the joint p.d.f. $f_{X,Y}$ positive on a set S , subset of \mathfrak{R}^2 , and set $U = aX + b$, $V = cY + d$, where a , b , c , and d are constants with $ac \neq 0$.

- (i) Use Theorem 4 in order to show that the joint p.d.f. of U and V is given by:

$$\begin{aligned} f_{U,V}(u, v) &= \frac{1}{|ac|} f_{X,Y}\left(\frac{u-b}{a}, \frac{v-c}{d}\right) \\ &= \frac{1}{|ac|} f_X\left(\frac{u-b}{c}\right) f_Y\left(\frac{v-c}{d}\right), \quad (u, v) \in T, \end{aligned}$$

the image of S under the transformations $u = ax + b$, $v = cy + d$.

- (ii) If $X \sim N(\mu_1, \sigma_1^2)$ and $Y \sim N(\mu_2, \sigma_2^2)$, show that U and V are independently distributed as $N(a\mu_1 + b, (a\sigma_1^2))$ and $N(c\mu_2 + d, (c\sigma_2^2))$, respectively.

2.5 If the independent r.v.'s X and Y are distributed as $N(0, 1)$, set $U = X + Y$, $V = X - Y$, and:

- (i) Determine the p.d.f.'s of U and V .
(ii) Show that U and V are independent.
(iii) Compute the probability $P(U < 0, V > 0)$.

2.6 Let X and Y be independent r.v.'s distributed as $N(0, 1)$, and set

$$U = \frac{1}{\sqrt{2}}(X + Y), \quad V = \frac{1}{\sqrt{2}}(X - Y).$$

- (i) Determine the joint p.d.f. of U and V .
(ii) From the joint p.d.f. $f_{U,V}$, infer f_U and f_V without integration.
(iii) Conclude that U and V are also independent.
(iv) How else could you arrive at the p.d.f.'s f_U and f_V ?

2.7 Let X and Y be independent r.v.'s distributed as $N(0, \sigma^2)$. Then show that the r.v. $U = X^2 + Y^2$ has the Negative Exponential distribution with parameter $\lambda = 1/2\sigma^2$.

2.8 The independent r.v.'s X and Y have p.d.f. given by: $f_{X,Y}(x, y) = \frac{1}{\pi}$, for $x, y \in \mathfrak{R}$ with $x^2 + y^2 \leq 1$, and let $Z^2 = X^2 + Y^2$. Use polar coordinates to determine the p.d.f. f_{Z^2} .

Hint: Let $Z = +\sqrt{Z^2}$ and set $X = Z \cos \Theta$, $Y = Z \sin \Theta$, where $Z \geq 0$ and $0 < \Theta \leq 2\pi$. First, determine the joint p.d.f. $f_{Z,\Theta}$ and then the marginal p.d.f. f_Z . Finally, by means of f_Z and the transformation $U = Z^2$, determine the p.d.f. $f_U = f_{Z^2}$.

2.9 If the r.v. $X_r \sim t_r$, then the t -tables (at least the ones in this book) do not give probabilities for $r > 90$. For such values, we can use instead the Normal tables. The reason for this is that the p.d.f. of X_r converges to

the p.d.f. of the $N(0, 1)$ distribution as $r \rightarrow \infty$. More precisely,

$$f_{X_r}(t) = \frac{\Gamma\left(\frac{r+1}{2}\right)}{\sqrt{\pi r} \Gamma\left(\frac{r}{2}\right)} \times \frac{1}{\left(1 + \frac{t^2}{r}\right)^{(r+1)/2}} \xrightarrow{r \rightarrow \infty} \frac{1}{\sqrt{2\pi}} e^{-t^2/2} \quad (t > 0).$$

Hint: In proving this convergence, first observe that

$$\left(1 + \frac{t^2}{r}\right)^{(r+1)/2} = \left[\left(1 + \frac{t^2}{r}\right)^r\right]^{1/2} \times \left(1 + \frac{t^2}{r}\right)^{1/2} \xrightarrow{r \rightarrow \infty} e^{t^2/2},$$

and then show that

$$\frac{\Gamma\left(\frac{r+1}{2}\right)}{\Gamma\left(\frac{r}{2}\right)} \xrightarrow{r \rightarrow \infty} \frac{1}{\sqrt{2}},$$

by utilizing the *Stirling formula*. This formula states that:

$$\frac{\Gamma(n)}{\sqrt{2\pi n} n^{(2n-1)/2} e^{-n}} \rightarrow 1 \quad \text{as } n \rightarrow \infty.$$

2.10 Let X_{r_1, r_2} be a r.v. having the F -distribution with parameters r_1 and r_2 ; i.e., $X_{r_1, r_2} \sim F_{r_1, r_2}$. Then show that:

$$EX_{r_1, r_2} = \frac{r_2}{r_2 - 2}, \quad r_2 \geq 3; \quad \text{Var}(X_{r_1, r_2}) = \frac{2r_2^2(r_1 + r_2 - 2)}{r_1(r_2 - 2)^2(r_2 - 4)}, \quad r_2 \geq 5.$$

Hint: Start out with the k th moment EX_{r_1, r_2}^k , use first the transformation $\frac{r_1}{r_2} f = x$, and second the transformation $\frac{1}{1+x} = y$. Then observe that the integrand is the p.d.f. of a Gamma distribution (except for suitable constants). Thus, the EX_{r_1, r_2}^k is expressed in terms of the Gamma function without carrying out any integrations. Specifically, we find:

$$EX_{r_1, r_2}^k = \left(\frac{r_2}{r_1}\right)^k \frac{\Gamma\left(\frac{r_1+2k}{2}\right) \Gamma\left(\frac{r_2-2k}{2}\right)}{\Gamma\left(\frac{r_1}{2}\right) \Gamma\left(\frac{r_2}{2}\right)}, \quad r_2 > 2k.$$

Applying this formula for $k = 1$ (which requires that $r_2 \geq 3$), and $k = 2$ (which requires that $r_2 \geq 5$), and using the recursive property of the Gamma function, we determine the required expressions.

6.3 Linear Transformations

In this section, a brief discussion is presented for a specific kind of transformation, linear transformations. The basic concepts and results used here can be found in any textbook on linear algebra.

DEFINITION 3

Suppose the variables x_1, \dots, x_k are transformed into the variables y_1, \dots, y_k in the following manner:

$$y_i = \sum_{j=1}^k c_{ij}x_j, \quad i = 1, \dots, k, \quad (14)$$

where the c_{ij} 's are real constants. Such a transformation is called a *linear transformation* (all the x_i 's enter into the transformation in a linear way, in the first power).

Some terminology and elementary facts from matrix algebra will be used here. Denote by \mathbf{C} the $k \times k$ matrix of the c_{ij} , $i, j = 1, \dots, k$ constants; that is, $\mathbf{C} = (c_{ij})$, and by $|\mathbf{C}|$ or Δ its determinant. Then it is well known that if $\Delta \neq 0$, one can uniquely solve for x_i in (14):

$$x_i = \sum_{j=1}^k d_{ij}y_j, \quad i = 1, \dots, k, \quad (15)$$

for suitable constants d_{ij} . Denote by \mathbf{D} the $k \times k$ matrix of the d_{ij} 's and by Δ^* its determinant: $\mathbf{D} = (d_{ij})$, $\Delta^* = |\mathbf{D}|$. Then it is known that $\Delta^* = 1/\Delta$. Among the linear transformations, a specific class is of special importance; it is the class of orthogonal transformations.

A linear transformation is said to be *orthogonal*, if

$$\sum_{j=1}^k c_{ij}^2 = 1 \quad \text{and} \quad \sum_{j=1}^k c_{ij}c_{i'j} = 0, \quad i, i' = 1, \dots, k, \quad i \neq i',$$

or, equivalently,

$$\sum_{i=1}^k c_{ij}^2 = 1 \quad \text{and} \quad \sum_{i=1}^k c_{ij}c_{ij'} = 0, \quad j, j' = 1, \dots, k, \quad j \neq j'. \quad (16)$$

Relations (16) simply state that the row (column) vectors of the matrix \mathbf{C} have norm (length) 1, and any two of them are perpendicular. The matrix \mathbf{C} itself is also called *orthogonal*. For an orthogonal matrix \mathbf{C} , it is known that $|\mathbf{C}| = \pm 1$. Also, in the case of an orthogonal matrix \mathbf{C} , it happens that $d_{ij} = c_{ji}$, $i, j = 1, \dots, k$; or in matrix notation: $\mathbf{D} = \mathbf{C}'$, where \mathbf{C}' is the *transpose* of \mathbf{C} (the rows of \mathbf{C}' are the same as the columns of \mathbf{C}). Thus, in this case:

$$x_i = \sum_{j=1}^k c_{ji}y_j, \quad i = 1, \dots, k. \quad (17)$$

Also, under orthogonality, the vectors of the x_i 's and of the y_j 's have the same norm. To put it differently:

$$\sum_{i=1}^k x_i^2 = \sum_{j=1}^k y_j^2. \quad (18)$$

Some of these concepts and results are now to be used in connection with r.v.'s.

THEOREM 7

Suppose the r.v.'s X_1, \dots, X_k are transformed into the r.v.'s Y_1, \dots, Y_k through a linear transformation with the matrix $\mathbf{C} = (c_{ij})$ and $|\mathbf{C}| = \Delta \neq 0$. Let $S \subseteq \mathfrak{R}^k$ be the set over which the joint p.d.f. of X_1, \dots, X_k , f_{X_1, \dots, X_k} , is positive, and let T be the image of S under the linear transformation. Then:

(i) The joint p.d.f. of Y_1, \dots, Y_k , f_{Y_1, \dots, Y_k} , is given by:

$$f_{Y_1, \dots, Y_k}(y_1, \dots, y_k) = f_{X_1, \dots, X_k} \left(\sum_{j=1}^k d_{1j} y_j, \dots, \sum_{j=1}^k d_{kj} y_j \right) \frac{1}{|\Delta|}, \quad (19)$$

for $(y_1, \dots, y_k) \in T$ (and = 0 otherwise), where the d_{ij} 's are as in (15).

(ii) In particular, if \mathbf{C} is orthogonal, then:

$$f_{Y_1, \dots, Y_k}(y_1, \dots, y_k) = f_{X_1, \dots, X_k} \left(\sum_{j=1}^k c_{j1} y_j, \dots, \sum_{j=1}^k c_{jk} y_j \right), \quad (20)$$

for $(y_1, \dots, y_k) \in T$ (and = 0 otherwise); also,

$$\sum_{j=1}^k Y_j^2 = \sum_{i=1}^k X_i^2. \quad (21)$$

PROOF

(i) Relation (19) follows from Theorem 5.

(ii) Relation (20) follows from (19) and (17), and (21) is a restatement of (18). ▲

Next, we specialize this result to the case that the r.v.'s X_1, \dots, X_k are Normally distributed and independent.

THEOREM 8

Let the independent r.v.'s X_1, \dots, X_k be distributed as follows: $X_i \sim N(\mu_i, \sigma^2)$, $i = 1, \dots, k$, and suppose they are transformed into the r.v.'s Y_1, \dots, Y_k by means of an orthogonal transformation \mathbf{C} . Then the r.v.'s Y_1, \dots, Y_k are also independent and Normally distributed as follows:

$$Y_i \sim N \left(\sum_{j=1}^k c_{ij} \mu_j, \sigma^2 \right), \quad i = 1, \dots, k. \quad (22)$$

PROOF From the transformations $Y_i = \sum_{j=1}^k c_{ij} X_j$, it is immediate that each Y_i is Normally distributed with mean $EY_i = \sum_{j=1}^k c_{ij} \mu_j$ and variance $\text{Var}(Y_i) = \sum_{j=1}^k c_{ij}^2 \sigma^2 = \sigma^2 \sum_{j=1}^k c_{ij}^2 = \sigma^2$. So the only thing to be justified is the assertion

of independence. From the Normality assumption on the X_i 's, we have:

$$f_{X_1, \dots, X_k}(x_1, \dots, x_k) = \left(\frac{1}{\sqrt{2\pi}\sigma} \right)^k \exp \left[-\frac{1}{2\sigma^2} \sum_{i=1}^k (x_i - \mu_i)^2 \right]. \quad (23)$$

Then, since \mathbf{C} is orthogonal, (20) applies and gives, by means of (23):

$$f_{Y_1, \dots, Y_k}(y_1, \dots, y_k) = \left(\frac{1}{\sqrt{2\pi}\sigma} \right)^k \exp \left[-\frac{1}{2\sigma^2} \sum_{i=1}^k \left(\sum_{j=1}^k c_{ji} y_j - \mu_i \right)^2 \right]. \quad (24)$$

Thus, the proof is completed by establishing the following algebraic relation:

$$\sum_{i=1}^k \left(\sum_{j=1}^k c_{ji} y_j - \mu_i \right)^2 = \sum_{i=1}^k \left(y_i - \sum_{j=1}^k c_{ij} \mu_j \right)^2 \quad (25)$$

(see Exercise 3.1). ▲

Finally, suppose the orthogonal matrix \mathbf{C} in Theorem 8 is chosen to be as follows:

$$\mathbf{C} = \begin{pmatrix} 1/\sqrt{k} & 1/\sqrt{k} & \dots & \dots & \dots & 1/\sqrt{k} \\ 1/\sqrt{2 \times 1} & -1/\sqrt{2 \times 1} & 0 & \dots & \dots & 0 \\ 1/\sqrt{3 \times 2} & 1/\sqrt{3 \times 2} & -2/\sqrt{3 \times 2} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 1/\sqrt{k(k-1)} & 1/\sqrt{k(k-1)} & \dots & \dots & 1/\sqrt{k(k-1)} & -(k-1)/\sqrt{k(k-1)} \end{pmatrix}.$$

That is, the elements of \mathbf{C} are given by the expressions:

$$\begin{aligned} c_{1j} &= 1/\sqrt{k}, & j &= 1, \dots, k, \\ c_{ij} &= 1/\sqrt{i(i-1)}, & \text{for } i &= 2, \dots, k \text{ and } j = 1, \dots, i-1, \\ & & & \text{and } 0 \text{ for } j = i+1, \dots, k, \\ c_{ii} &= -(i-1)/\sqrt{i(i-1)}, & i &= 2, \dots, k. \end{aligned}$$

From these expressions, it readily follows that $\sum_{j=1}^k c_{ij}^2 = 1$ for all $i = 1, \dots, k$, and $\sum_{j=1}^k c_{ij} c_{i'j} = 0$ for all $i, i' = 1, \dots, k$, with $i \neq i'$, so that \mathbf{C} is, indeed, orthogonal (see also Exercise 3.2). Next, let Z_1, \dots, Z_k be independent r.v.'s distributed as $N(0, 1)$, and transform them into the r.v.'s Y_1, \dots, Y_k by means of \mathbf{C} ; that is,

$$\begin{aligned} Y_1 &= \frac{1}{\sqrt{k}} Z_1 + \frac{1}{\sqrt{k}} Z_2 + \dots + \frac{1}{\sqrt{k}} Z_k \\ Y_2 &= \frac{1}{\sqrt{2 \times 1}} Z_1 - \frac{1}{\sqrt{2 \times 1}} Z_2 \end{aligned}$$

$$\begin{aligned}
Y_3 &= \frac{1}{\sqrt{3 \times 2}} Z_1 + \frac{1}{\sqrt{3 \times 2}} Z_2 - \frac{2}{\sqrt{3 \times 2}} Z_3 \\
&\vdots \\
Y_k &= \frac{1}{\sqrt{k(k-1)}} Z_1 + \frac{1}{\sqrt{k(k-1)}} Z_2 + \cdots + \frac{1}{\sqrt{k(k-1)}} Z_{k-1} - \frac{k-1}{\sqrt{k(k-1)}} Z_k.
\end{aligned}$$

Then, by Theorem 8, the r.v.'s Y_1, \dots, Y_k are independently distributed as $N(0, 1)$, whereas by (21)

$$\sum_{j=1}^k Y_j^2 = \sum_{i=1}^k Z_i^2.$$

However, $Y_1 = \sqrt{k}\bar{Z}$, so that

$$\sum_{j=2}^k Y_j^2 = \sum_{j=1}^k Y_j^2 - Y_1^2 = \sum_{i=1}^k Z_i^2 - (\sqrt{k}\bar{Z})^2 = \sum_{i=1}^k Z_i^2 - k\bar{Z}^2 = \sum_{i=1}^k (Z_i - \bar{Z})^2.$$

On the other hand, $\sum_{j=2}^k Y_j^2$ and Y_1 are independent; equivalently, $\sum_{i=1}^k (Z_i - \bar{Z})^2$ and $k\bar{Z}$ are independent, or

$$\bar{Z} \quad \text{and} \quad \sum_{i=1}^k (Z_i - \bar{Z})^2 \quad \text{are independent.} \quad (26)$$

This last conclusion is now applied as follows.

THEOREM 9

Let X_1, \dots, X_k be independent r.v.'s distributed as $N(\mu, \sigma^2)$. Then the sample mean $\bar{X} = \frac{1}{k} \sum_{i=1}^k X_i$ and the sample variance $S^2 = \frac{1}{k} \sum_{i=1}^k (X_i - \bar{X})^2$ are independent.

PROOF The assumption that $X_i \sim N(\mu, \sigma^2)$ implies that $\frac{X_i - \mu}{\sigma} \sim N(0, 1)$. By setting $Z_i = (X_i - \mu)/\sigma$, $i = 1, \dots, k$, the Z_i 's are as in the preceding derivations and therefore (26) applies. Since

$$\bar{Z} = \frac{1}{k} \sum_{i=1}^k \left(\frac{X_i - \mu}{\sigma} \right) = \frac{1}{\sigma} (\bar{X} - \mu), \text{ and}$$

$$\sum_{i=1}^k (Z_i - \bar{Z})^2 = \sum_{i=1}^k \left(\frac{X_i - \mu}{\sigma} - \frac{\bar{X} - \mu}{\sigma} \right)^2 = \frac{1}{\sigma^2} \sum_{i=1}^k (X_i - \bar{X})^2,$$

it follows that $\frac{1}{\sigma} (\bar{X} - \mu)$ and $\frac{1}{\sigma^2} \sum_{i=1}^k (X_i - \bar{X})^2$ are independent or that \bar{X} and $\frac{1}{k} \sum_{i=1}^k (X_i - \bar{X})^2$ are independent. \blacktriangle

Exercises

3.1 Establish relation (25) in the proof of Theorem 8.

Hint: Expand the left-hand side and the right-hand side in (25), use orthogonality, and show that the common value of both sides is:

$$\sum_{j=1}^k y_j^2 + \sum_{j=1}^k \mu_j^2 - 2 \sum_{j=1}^k \sum_{i=1}^k c_{ji} \mu_i \mu_j.$$

3.2 Show that the matrix with row elements given by:

$$\begin{aligned} c_{1j} &= 1/\sqrt{k}, \quad j = 1, \dots, k, \\ c_{ij} &= 1/\sqrt{i(i-1)}, \quad i = 2, \dots, k \text{ and } j = 1, \dots, i-1, \\ &\text{and } 0 \text{ for } j = i+1, \dots, k, \\ c_{ii} &= -(i-1)/\sqrt{i(i+1)}, \quad i = 2, \dots, k \text{ is orthogonal.} \end{aligned}$$

3.3 Let X_1, X_2, X_3 be independent r.v.'s such that $X_i \sim N(\mu_i, \sigma^2)$, $i = 1, 2, 3$, and set

$$\begin{aligned} Y_1 &= -\frac{1}{\sqrt{2}}X_1 + \frac{1}{\sqrt{2}}X_2, \\ Y_2 &= -\frac{1}{\sqrt{3}}X_1 - \frac{1}{\sqrt{3}}X_2 + \frac{1}{\sqrt{3}}X_3, \\ Y_3 &= \frac{1}{\sqrt{6}}X_1 + \frac{1}{\sqrt{6}}X_2 + \frac{2}{\sqrt{6}}X_3. \end{aligned}$$

Then:

(i) Show that the r.v.'s Y_1, Y_2, Y_3 are independent Normally distributed with variance σ^2 and respective means:

$$\begin{aligned} EY_1 &= \frac{1}{\sqrt{2}}(-\mu_1 + \mu_2), & EY_2 &= \frac{1}{\sqrt{3}}(-\mu_1 - \mu_2 + \mu_3), \\ EY_3 &= \frac{1}{\sqrt{6}}(\mu_1 + \mu_2 + 2\mu_3). \end{aligned}$$

(ii) If $\mu_1 = \mu_2 = \mu_3 = 0$, then show that $\frac{1}{\sigma^2}(Y_1^2 + Y_2^2 + Y_3^2) \sim \chi_3^2$,

Hint: For part (i), prove that the transformation employed is orthogonal and then use Theorem 8 to conclude independence of Y_1, Y_2, Y_3 . That the means and the variance are as described follows either from Theorem 8 or directly. Part (ii) follows from part (i) and the assumption that $\mu_1 = \mu_2 = \mu_3 = 0$.

3.4 If the r.v.'s X and Y have the Bivariate Normal distribution with parameters $\mu_1, \mu_2, \sigma_1^2, \sigma_2^2$, and ρ , then the r.v.'s $U = \frac{X-\mu_1}{\sigma_1}$, $V = \frac{Y-\mu_2}{\sigma_2}$ have the Bivariate Normal distribution with parameters 0, 0, 1, 1, and ρ ; and vice versa.

- 3.5** If the r.v.'s X and Y have the Bivariate Normal distribution with parameters $0, 0, 1, 1,$ and ρ , then the r.v.'s cX and dY have the Bivariate Normal distribution with parameters $0, 0, c^2, d^2,$ and ρ_0 , where $\rho_0 = 1$ if $cd > 0$, and $\rho_0 = -1$ if $cd < 0$; c and d are constants with $cd \neq 0$.
- 3.6** Let the r.v.'s X and Y have the Bivariate Normal distribution with parameters $0, 0, 1, 1,$ and ρ , and set: $U = X + Y, V = X - Y$. Then show that:
- The r.v.'s U and V also have the Bivariate Normal distribution with parameters $0, 0, 2(1 + \rho), 2(1 - \rho),$ and 0 .
 - From part (i), conclude that the r.v.'s U and V are independent.
 - From part (i), also conclude that: $U \sim N(0, 2(1 + \rho)), V \sim N(0, 2(1 - \rho))$.
- 3.7** Let the r.v.'s X and Y have the Bivariate Normal distribution with parameters $\mu_1, \mu_2, \sigma_1^2, \sigma_2^2,$ and ρ , and set:

$$U = \frac{X - \mu_1}{\sigma_1}, \quad V = \frac{Y - \mu_2}{\sigma_2}.$$

Then:

- Determine the joint distribution of the r.v.'s U and V .
- Show that $U + V$ and $U - V$ have the Bivariate Normal distribution with parameters $0, 0, 2(1 + \rho), 2(1 - \rho),$ and 0 and are independent. Also, $U + V \sim N(0, 2(1 + \rho)), U - V \sim N(0, 2(1 - \rho))$.
- For $\sigma_1^2 = \sigma_2^2 = \sigma^2$, say, conclude that the r.v.'s $X + Y$ and $X - Y$ are independent.

REMARK 4 Actually, the converse of part (iii) is also true; namely, if X and Y have the Bivariate Normal distribution $N(\mu_1, \mu_2, \sigma_1^2, \sigma_2^2, \rho)$, then independence of $X + Y$ and $X - Y$ implies $\sigma_1^2 = \sigma_2^2$. The justification of this statement is easier by means of m.g.f.'s, and it was, actually, discussed in Exercise 5.13 of Chapter 4.

- 3.8** Let the independent r.v.'s X_1, \dots, X_n be distributed as $N(\mu, \sigma^2)$ and suppose that $\mu = k\sigma$ ($k > 0$). Set

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i, \quad S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2.$$

Then:

- Determine an expression for the probability:

$$P(a\mu < \bar{X} < b\mu, \quad 0 < S^2 < c\sigma^2),$$

where $a, b,$ and c are constants, $a < b$ and $c > 0$.

- Give the numerical value of the probability in part (i) if $a = \frac{1}{2}, b = \frac{3}{2}, c = 1.487, k = 1.5,$ and $n = 16$.

Hint: Use independence of \bar{X} and S^2 provided by Theorem 9. Also, use the fact that $\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$ by Theorem 6 in Chapter 5 (where S^2 is denoted by \bar{S}^2).

6.4 The Probability Integral Transform

In this short section, a very special type of transformation is considered, the so-called *probability integral transform*. By means of this transformation, two results are derived. Roughly, these results state that, if $X \sim F$ and $Y = F(X)$, then, somewhat surprisingly, Y is always distributed as $U(0, 1)$. Furthermore, for a given d.f. F , there is always a r.v. $X \sim F$; this r.v. is given by $X = F^{-1}(Y)$, where $Y \sim U(0, 1)$ and F^{-1} is the inverse function of F . To facilitate the derivations, F will be assumed to be (strictly) increasing.

THEOREM 10

For an increasing d.f. F , let $X \sim F$ and set $Y = F(X)$. Then $Y \sim U(0, 1)$.

PROOF Since $0 \leq F(X) \leq 1$, it suffices to consider $y \in [0, 1]$. Then

$$\begin{aligned} P(Y \leq y) &= P[F(X) \leq y] = P\{F^{-1}[F(X)] \leq F^{-1}(y)\} \\ &= P[X \leq F^{-1}(y)] = F[F^{-1}(y)] = y, \end{aligned}$$

so that $Y \sim U(0, 1)$. ▲

THEOREM 11

Let F be a given increasing d.f., and let the r.v. $Y \sim U(0, 1)$. Define the r.v. X by: $X = F^{-1}(Y)$. Then $X \sim F$.

PROOF For $x \in \Re$,

$$\begin{aligned} P(X \leq x) &= P[F^{-1}(Y) \leq x] = P\{F[F^{-1}(Y)] \leq F(x)\} \\ &= P[Y \leq F(x)] = F(x), \end{aligned}$$

as was to be seen. ▲

In the form of a verification of Theorems 10 and 11, consider the following simple examples.

EXAMPLE 6

Let the r.v. X have the Negative Exponential distribution with parameter λ . Then, for $x > 0$, $F(x) = 1 - e^{-\lambda x}$. Let Y be defined by: $Y = 1 - e^{-\lambda X}$. Then Y should be $\sim U(0, 1)$.

DISCUSSION Indeed, for $0 < y < 1$,

$$\begin{aligned} P(Y \leq y) &= P(1 - e^{-\lambda X} \leq y) = P(e^{-\lambda X} \geq 1 - y) = P[-\lambda X \geq \log(1 - y)] \\ &= P\left[X \leq -\frac{1}{\lambda} \log(1 - y)\right] \end{aligned}$$

$$\begin{aligned}
&= 1 - \exp \left\{ (-\lambda) \left[-\frac{1}{\lambda} \log(1 - y) \right] \right\} \\
&= 1 - \exp[\log(1 - y)] = 1 - (1 - y) = y,
\end{aligned}$$

as was to be seen.

EXAMPLE 7

Let F be the d.f. of the Negative Exponential distribution with parameter λ , so that $F(x) = 1 - e^{-\lambda x}$, $x > 0$. Let $y = 1 - e^{-\lambda x}$ and solve for x to obtain $x = -\frac{1}{\lambda} \log(1 - y)$, $0 < y < 1$. Let $Y \sim U(0, 1)$ and define the r.v. X by: $X = -\frac{1}{\lambda} \log(1 - Y)$. Then X should be $\sim F$.

DISCUSSION Indeed,

$$\begin{aligned}
P(X \leq x) &= P \left[-\frac{1}{\lambda} \log(1 - Y) \leq x \right] = P[\log(1 - Y) \geq -\lambda x] \\
&= P(1 - Y \geq e^{-\lambda x}) = P(Y \leq 1 - e^{-\lambda x}) = 1 - e^{-\lambda x},
\end{aligned}$$

as was to be seen.

Exercise

- 4.1 (i) Let X be a r.v. with continuous and (strictly) increasing d.f. F , and define the r.v. Y by $Y = F(X)$. Then use Theorem 2 in order to show that $Z = -2 \log(1 - Y) \sim \chi_2^2$.
- (ii) If X_1, \dots, X_n is a random sample with d.f. F as described in part (i) and if $Y_i = F(X_i)$, $i = 1, \dots, n$, then show that the r.v. $U = \sum_{i=1}^n Z_i \sim \chi_{2n}^2$, where $Z_i = -2 \log(1 - Y_i)$, $i = 1, \dots, n$.

Hint: For part (i), use Theorem 10, according to which $Y \sim U(0, 1)$.

6.5 Order Statistics

In this section, an unconventional kind of transformation is considered, which, when applied to r.v.'s, leads to the so-called *order statistics*. For the definition of the transformation, consider n distinct numbers x_1, \dots, x_n and order them in ascending order. Denote by $x_{(1)}$ the smallest number: $x_{(1)} =$ smallest of x_1, \dots, x_n ; by $x_{(2)}$ the second smallest, and so on until $x_{(n)}$ is the n th smallest or, equivalently, the largest of the x_i 's. In a summary form, we write: $x_{(j)} =$ the j th smallest of the numbers x_1, \dots, x_n , where $j = 1, \dots, n$. Then, clearly, $x_{(1)} < x_{(2)} < \dots < x_{(n)}$. For simplicity, set $y_j = x_{(j)}$, $j = 1, \dots, n$, so that again $y_1 < y_2 < \dots < y_n$. The transformation under consideration is the one which transforms the x_i 's into the y_j 's in the way just described.

This transformation now applies to n r.v.'s as follows.

Let X_1, X_2, \dots, X_n be i.i.d. r.v.'s with d.f. F . The j th order statistic of X_1, X_2, \dots, X_n is denoted by $X_{(j)}$, or Y_j for easier writing, and is defined as follows:

$$Y_j = j\text{th smallest of the } X_1, X_2, \dots, X_n, \quad j = 1, \dots, n;$$

(that is, for each $s \in \mathcal{S}$, look at $X_1(s), X_2(s), \dots, X_n(s)$, and then $Y_j(s)$ is defined to be the j th smallest among the numbers $X_1(s), X_2(s), \dots, X_n(s)$, $j = 1, 2, \dots, n$). It follows that $Y_1 \leq Y_2 \leq \dots \leq Y_n$, and, in general, the Y_j 's are not independent.

We assume now that the X_i 's are of the continuous type with p.d.f. f such that $f(x) > 0$, $(-\infty \leq) a < x < b (\leq \infty)$ and zero otherwise. One of the problems we are concerned with is that of finding the joint p.d.f. of the Y_j 's. By means of Theorem 6, it will be established that:

THEOREM 12

If X_1, \dots, X_n are i.i.d. r.v.'s with p.d.f. f which is positive for $a < x < b$ and 0 otherwise, then the joint p.d.f. of the order statistics Y_1, \dots, Y_n is given by:

$$g(y_1, \dots, y_n) = \begin{cases} n! f(y_1) \cdots f(y_n), & a < y_1 < y_2 < \cdots < y_n < b \\ 0, & \text{otherwise.} \end{cases} \quad (27)$$

PROOF The proof is carried out explicitly for $n = 3$, but it is easily seen, with the proper change in notation, to be valid in the general case as well. In the first place, since for $i \neq j$,

$$P(X_i = X_j) = \iint_{(x_i=x_j)} f(x_i)f(x_j) dx_i dx_j = \int_a^b \int_a^{x_j} f(x_i)f(x_j) dx_i dx_j = 0,$$

and therefore $P(X_i = X_j = X_k) = 0$ for $i \neq j \neq k$, we may assume that the joint p.d.f., $f(\cdot, \cdot, \cdot)$, of X_1, X_2, X_3 is zero, if at least two of the arguments x_1, x_2, x_3 are equal. Thus, we have:

$$f(x_1, x_2, x_3) = \begin{cases} f(x_1)f(x_2)f(x_3), & a < x_1 \neq x_2 \neq x_3 < b \\ 0, & \text{otherwise.} \end{cases}$$

Therefore $f(x_1, x_2, x_3)$ is positive on the set S , where

$$S = \{(x_1, x_2, x_3) \in \mathfrak{R}^3; \quad a < x_i < b, \quad i = 1, 2, 3, \quad x_1, x_2, x_3 \text{ all different}\}.$$

Let $S_{ijk} \subset S$ be defined by:

$$S_{ijk} = \{(x_1, x_2, x_3); \quad a < x_i < x_j < x_k < b\}, \quad i, j, k = 1, 2, 3, \quad i \neq j \neq k.$$

Then we have that these six events are pairwise disjoint and (essentially)

$$S = S_{123} \cup S_{132} \cup S_{213} \cup S_{231} \cup S_{312} \cup S_{321}.$$

Now on each one of the S_{ijk} 's there exists a one-to-one transformation from the x_i 's to the y_i 's defined as follows:

$$S_{123} : y_1 = x_1, y_2 = x_2, y_3 = x_3$$

$$S_{132} : y_1 = x_1, y_2 = x_3, y_3 = x_2$$

$$S_{213} : y_1 = x_2, y_2 = x_1, y_3 = x_3$$

$$S_{231} : y_1 = x_2, y_2 = x_3, y_3 = x_1$$

$$S_{312} : y_1 = x_3, y_2 = x_1, y_3 = x_2$$

$$S_{321} : y_1 = x_3, y_2 = x_2, y_3 = x_1.$$

Solving for the x_i 's, we have then:

$$S_{123} : x_1 = y_1, x_2 = y_2, x_3 = y_3$$

$$S_{132} : x_1 = y_1, x_2 = y_3, x_3 = y_2$$

$$S_{213} : x_1 = y_2, x_2 = y_1, x_3 = y_3$$

$$S_{231} : x_1 = y_3, x_2 = y_1, x_3 = y_2$$

$$S_{312} : x_1 = y_2, x_2 = y_3, x_3 = y_1$$

$$S_{321} : x_1 = y_3, x_2 = y_2, x_3 = y_1.$$

The Jacobians are thus given by:

$$\begin{aligned} S_{123} : J_{123} &= \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 1, & S_{231} : J_{231} &= \begin{vmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{vmatrix} = 1, \\ S_{132} : J_{132} &= \begin{vmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{vmatrix} = -1, & S_{312} : J_{312} &= \begin{vmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{vmatrix} = 1, \\ S_{213} : J_{213} &= \begin{vmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{vmatrix} = -1, & S_{321} : J_{321} &= \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{vmatrix} = -1. \end{aligned}$$

Hence $|J_{123}| = \cdots = |J_{321}| = 1$, and Theorem 6 gives

$$g(y_1, y_2, y_3) = \begin{cases} f(y_1)f(y_2)f(y_3) + f(y_1)f(y_3)f(y_2) + f(y_2)f(y_1)f(y_3) \\ \quad + f(y_3)f(y_1)f(y_2) + f(y_2)f(y_3)f(y_1) + f(y_3)f(y_2)f(y_1), & a < y_1 < y_2 < y_3 < b \\ 0, & \text{otherwise.} \end{cases}$$

That is,

$$g(y_1, y_2, y_3) = \begin{cases} 3!f(y_1)f(y_2)f(y_3), & a < y_1 < y_2 < y_3 < b \\ 0, & \text{otherwise.} \end{cases} \quad \blacktriangle$$

Notice that the proof in the general case is exactly the same. One has $n!$ regions forming S , one for each permutation of the integers 1 through n . From the definition of a determinant and the fact that each row and column contains exactly one 1 and the rest all 0, it follows that the $n!$ Jacobians are either 1 or -1 and the remaining part of the proof is identical to the one just given except that one adds up $n!$ like terms instead of $3!$.

The theorem is illustrated by the following two examples.

EXAMPLE 8

Let X_1, \dots, X_n be i.i.d. r.v.'s distributed as $N(\mu, \sigma^2)$. Then the joint p.d.f. of the order statistics Y_1, \dots, Y_n is given by

$$g(y_1, \dots, y_n) = n! \left(\frac{1}{\sqrt{2\pi}\sigma} \right)^n \exp \left[-\frac{1}{2\sigma^2} \sum_{j=1}^n (y_j - \mu)^2 \right],$$

if $-\infty < y_1 < \dots < y_n < \infty$, and zero otherwise.

EXAMPLE 9

Let X_1, \dots, X_n be i.i.d. r.v.'s distributed as $U(\alpha, \beta)$. Then the joint p.d.f. of the order statistics Y_1, \dots, Y_n is given by

$$g(y_1, \dots, y_n) = \frac{n!}{(\beta - \alpha)^n},$$

if $\alpha < y_1 < \dots < y_n < \beta$, and zero otherwise.

From the joint p.d.f. in (27), it is relatively easy to derive the p.d.f. of Y_j for any j , as well as the joint p.d.f. of Y_i and Y_j for any $1 \leq i < j \leq n$. We restrict ourselves to the derivation of the distributions of Y_1 and Y_n alone.

THEOREM 13

Let X_1, \dots, X_n be i.i.d. r.v.'s with d.f. F and p.d.f. f which is positive and continuous for $(-\infty \leq) a < x < b (\leq \infty)$ and zero otherwise, and let Y_1, \dots, Y_n be the order statistics. Then the p.d.f.'s g_1 and g_n of Y_1 and Y_n , respectively, are given by:

$$g_1(y_1) = \begin{cases} n[1 - F(y_1)]^{n-1} f(y_1), & a < y_1 < b \\ 0, & \text{otherwise,} \end{cases} \quad (28)$$

and

$$g_n(y_n) = \begin{cases} n[F(y_n)]^{n-1} f(y_n), & a < y_n < b \\ 0, & \text{otherwise.} \end{cases} \quad (29)$$

PROOF First, derive the d.f.'s involved and then differentiate them to obtain the respective p.d.f.'s. To this end,

$$\begin{aligned} G_n(y_n) &= P(Y_n \leq y_n) = P[\max(X_1, \dots, X_n) \leq y_n] \\ &= P(\text{all } X_1, \dots, X_n \leq y_n) = P(X_1 \leq y_n, \dots, X_n \leq y_n) \\ &= P(X_1 \leq y_n) \cdots P(X_n \leq y_n) \quad (\text{by the independence of the } X_i\text{'s}) \\ &= [F(y_n)]^n. \end{aligned}$$

That is, $G_n(y_n) = [F(y_n)]^n$, so that

$$g_n(y_n) = \frac{d}{dy_n} G_n(y_n) = n[F(y_n)]^{n-1} \frac{d}{dy_n} F(y_n) = n[F(y_n)]^{n-1} f(y_n).$$

Likewise,

$$\begin{aligned} 1 - G_1(y_1) &= P(Y_1 > y_1) = P[\min(X_1, \dots, X_n) > y_1] \\ &= P(\text{all } X_1, \dots, X_n > y_1) = P(X_1 > y_1, \dots, X_n > y_1) \\ &= P(X_1 > y_1) \cdots P(X_n > y_1) \quad (\text{by the independence of the } X_i\text{'s}) \\ &= [1 - P(X_1 \leq y_1)] \cdots [1 - P(X_1 \leq y_1)] = [1 - F(y_1)]^n. \end{aligned}$$

That is, $1 - G_1(y_1) = [1 - F(y_1)]^n$, so that

$$\begin{aligned} -g_1(y_1) &= \frac{d}{dy_1} [1 - G_1(y_1)] = n[1 - F(y_1)]^{n-1} \frac{d}{dy_1} [1 - F(y_1)] \\ &= n[1 - F(y_1)]^{n-1} [-f(y_1)] = -n[1 - F(y_1)]^{n-1} f(y_1), \end{aligned}$$

and hence

$$g_1(y_1) = n[1 - F(y_1)]^{n-1} f(y_1). \quad \blacktriangle$$

As an illustration of the theorem, consider the following example.

EXAMPLE 10

Let the independent r.v.'s X_1, \dots, X_n be distributed as $U(0, 1)$. Then, for $0 < y_1, y_n < 1$:

$$g_1(y_1) = n(1 - y_1)^{n-1} \quad \text{and} \quad g_n(y_n) = ny_n^{n-1}.$$

DISCUSSION Here, for $0 < x < 1$, $f(x) = 1$ and $F(x) = x$. Therefore relations (28) and (29) give, for $0 < y_1, y_n < 1$:

$$g_1(y_1) = n(1 - y_1)^{n-1} \cdot 1 = n(1 - y_1)^{n-1} \quad \text{and} \quad g_n(y_n) = ny_n^{n-1},$$

as asserted.

As a further illustration of the theorem, consider the following example, which is of interest in its own right.

EXAMPLE 11

If X_1, \dots, X_n are independent r.v.'s having the Negative Exponential distribution with parameter λ , then Y_1 has also the Negative Exponential distribution with parameter $n\lambda$.

DISCUSSION Here $f(x) = \lambda e^{-\lambda x}$ and $F(x) = 1 - e^{-\lambda x}$ for $x > 0$. Then, for $y_1 > 0$, formula (28) yields:

$$g_1(y_1) = n(e^{-\lambda y_1})^{n-1} \times \lambda e^{-\lambda y_1} = (n\lambda)e^{-(n-1)y_1} e^{-\lambda y_1} = (n\lambda)e^{-(n\lambda)y_1},$$

as was to be seen.

EXAMPLE 12

- (i) In a complex system, n identical components are connected serially, so that the system works, if and only if all n components function. If the lifetime of said components is described by a r.v. X with d.f. F and p.d.f. f , write out the expression for the probability that the system functions for at least t time units.
- (ii) Do the same as in part (i), if the components are connected in parallel, so that the system functions, if and only if at least one of the components works.
- (iii) Simplify the expressions in parts (i) and (ii), if f is the Negative Exponential with parameter λ .

DISCUSSION

- (i) Clearly, $P(\text{system works for at least } t \text{ time units})$

$$\begin{aligned} &= P(X_1 \geq t, \dots, X_n \geq t) && \text{(where } X_i \text{ is the lifetime of the } \\ & && \textit{ith} \text{ component)} \\ &= P(Y_1 \geq t) && \text{(where } Y_1 \text{ is the smallest order statistic)} \\ &= \int_t^\infty g_1(y) dy && \text{(where } g_1 \text{ is the p.d.f. of } Y_1) \\ &= \int_t^\infty n[1 - F(y)]^{n-1} f(y) dy && \text{(by (28)).} \end{aligned} \tag{30}$$

- (ii) Here

$P(\text{system works for at least } t \text{ time units})$

$$\begin{aligned} &= P(\text{at least one of } X_1, \dots, X_n \geq t) \\ &= P(Y_n \geq t) && \text{(where } Y_n \text{ is the largest order statistic)} \\ &= \int_t^\infty g_n(y) dy && \text{(where } g_n \text{ is the p.d.f. of } Y_n) \\ &= \int_t^\infty n[F(y)]^{n-1} f(y) dy && \text{(by (29)).} \end{aligned} \tag{31}$$

- (iii) Here $F(y) = 1 - e^{-\lambda y}$ and $f(y) = \lambda e^{-\lambda y}$ ($y > 0$) from Example 11. Also, from the same example, the p.d.f. of Y_1 is $g_1(y) = (n\lambda)e^{-(n\lambda)y}$, so that (30) gives:

$$\begin{aligned} P(Y_1 \geq t) &= \int_t^\infty (n\lambda)e^{-(n\lambda)y} dy \\ &= - \int_t^\infty de^{-(n\lambda)y} dy \\ &= -e^{-(n\lambda)y} \Big|_t^\infty = e^{-n\lambda t}, \end{aligned}$$

and, by (31),

$$P(Y_n \geq t) = \int_t^\infty n(1 - e^{-\lambda y})^{n-1} \lambda e^{-\lambda y} dy.$$

For example, for $n = 2$, this last probability is equal to:

$$\begin{aligned} \int_t^\infty 2(1 - e^{-\lambda y})\lambda e^{-\lambda y} dy &= 2 \int_t^\infty \lambda e^{-\lambda y} dy - \int_t^\infty 2\lambda e^{-2\lambda y} dy \\ &= -2 \int_t^\infty de^{-\lambda y} + \int_t^\infty de^{-2\lambda y} \\ &= -2e^{-\lambda y} \Big|_t^\infty + e^{-2\lambda y} \Big|_t^\infty \\ &= 2e^{-\lambda t} - e^{-2\lambda t}. \end{aligned}$$

Exercises

- 5.1** Let X_1, \dots, X_n be independent r.v.'s with p.d.f. $f(x) = cx^{-(c+1)}$, $x > 1$ ($c > 0$), and set $U = Y_1 = \min(X_1, \dots, X_n)$, $V = Y_n = \max(X_1, \dots, X_n)$.
- Determine the d.f. F corresponding to the p.d.f. f .
 - Use Theorem 13 to determine the p.d.f.'s f_U and f_V .
- 5.2** Refer to Example 10 and calculate the expectations EY_1 and EY_n , and also determine the $\lim EY_n$ as $n \rightarrow \infty$.
- 5.3** Let Y_1 and Y_n be the smallest and the largest order statistics based on a random sample X_1, \dots, X_n from the $U(\alpha, \beta)$ ($\alpha < \beta$) distribution.
- For $n = 3$ and $n = 4$, show that the joint p.d.f. of Y_1 and Y_n is given, respectively, by:

$$g_{13}(y_1, y_3) = \frac{3 \times 2}{(\beta - \alpha)^2} (y_3 - y_1), \quad \alpha < y_1 < y_3 < \beta,$$

$$g_{14}(y_1, y_4) = \frac{4 \times 3}{(\beta - \alpha)^3} (y_4 - y_1)^2, \quad \alpha < y_1 < y_4 < \beta.$$

- Generalize the preceding results and show that:

$$g_{1n}(y_1, y_n) = \frac{n(n-1)}{(\beta - \alpha)^n} (y_n - y_1)^{n-2}, \quad \alpha < y_1 < y_n < \beta.$$

Hint: For part (ii), all one has to do is to calculate the integrals:

$$\int_{y_1}^{y_n} \int_{y_1}^{y_{n-1}} \cdots \int_{y_1}^{y_2} \int_{y_1}^{y_3} dy_2 dy_3 \cdots dy_{n-2} dy_{n-1},$$

which is done one at a time; also, observe the pattern emerging.

5.4 Let Y_1 and Y_n be the smallest and the largest order statistics based on a random sample X_1, \dots, X_n from the $U(0, 1)$ distribution. Then show that:

$$\text{Cov}(Y_1, Y_n) = \frac{1}{(n+1)^2(n+2)}.$$

Hint: Use the joint p.d.f. taken from Exercise 5.3(ii) for $\alpha = 0$ and $\beta = 1$.

5.5 If Y_1 and Y_n are the smallest and the largest order statistics based on a random sample X_1, \dots, X_n from the $U(0, 1)$ distribution:

(i) Show that the p.d.f. of the *sample range* $R = Y_n - Y_1$ is given by:

$$f_R(r) = n(n-1)r^{n-2}(1-r), \quad 0 < r < 1.$$

(ii) Also, calculate the expectation ER .

5.6 Refer to Example 11 and set $Z = nY_1$. Then show that Z is distributed as the X_i 's.

5.7 The lifetimes of two batteries are independent r.v.'s X and Y with the Negative Exponential distribution with parameter λ . Suppose that the two batteries are connected serially, so that the system works if and only if both work.

(i) Use Example 11 (with $n = 2$) to calculate the probability that the system works beyond time $t > 0$.

(ii) What is the expected lifetime of the system?

(iii) What do parts (i) and (ii) become for $\lambda = 1/3$?

5.8 Let Y_1 and Y_n be the smallest and the largest order statistics based on a random sample X_1, \dots, X_n from the Negative Exponential distribution with parameter λ . Then, by Example 11, $g_1(y_1) = (n\lambda)e^{-(n\lambda)y_1}$, $y_1 > 0$.

(i) Use relation (29) (with $a = 0$ and $b = \infty$) to determine the p.d.f. g_n of the r.v. Y_n .

(ii) Calculate the EY_n for $n = 2$ and $n = 3$.

5.9 (i) Refer to Exercise 5.8(i) and show that:

$$EY_n = \frac{n}{\lambda} \sum_{r=0}^{n-1} (-1)^{n-r-1} \frac{\binom{n-1}{r}}{(n-r)^2}.$$

(ii) Apply part (i) for $n = 2$ and $n = 3$ to recover the values found in Exercise 5.8 (ii).

Hint: Consider the binomial expansion: $(a+b)^k = \sum_{r=0}^k \binom{k}{r} a^r b^{k-r}$ and apply it to: $(1 - e^{-\lambda y})^{n-1}$ for $a = 1$, $b = -e^{-\lambda y}$, and $k = n - 1$. Then carry out the multiplications indicated and integrate term by term.

5.10 Let X_1, \dots, X_n be a random sample of size n of the continuous type with d.f. F and p.d.f. f , positive in $-\infty \leq a < x < b \leq \infty$, and let Y_1 and Y_n be the smallest and the largest order statistics of the X_i 's. Use relation (27) in order to show that the joint p.d.f. g_{1n} of the r.v.'s Y_1 and Y_n is given by the expression:

$$g_{1n}(y_1, y_n) = n(n-1)[F(y_n) - F(y_1)]^{n-2} f(y_1) f(y_n), \quad a < y_1 < y_n < b.$$

Hint: The p.d.f. g_{1n} is obtained by integrating $g(y_1, \dots, y_n)$ in (27) with respect to $y_{n-1}, y_{n-2}, \dots, y_2$ as indicated below:

$$g_{1n}(y_1, y_n) = n! f(y_1) f(y_n) \int_{y_1}^{y_n} \cdots \int_{y_{n-3}}^{y_n} \int_{y_{n-2}}^{y_n} f(y_{n-1}) f(y_{n-2}) \times \\ \cdots f(y_2) dy_{n-1} dy_{n-2} \cdots dy_2.$$

However,

$$\int_{y_{n-2}}^{y_n} f(y_{n-1}) dy_{n-1} = F(y_n) - F(y_{n-2}) = \frac{[F(y_n) - F(y_{n-2})]^1}{1!}, \\ \int_{y_{n-3}}^{y_n} \frac{[F(y_n) - F(y_{n-2})]^1}{1!} f(y_{n-2}) dy_{n-2} \\ = - \int_{y_{n-3}}^{y_n} \frac{[F(y_n) - F(y_{n-2})]^1}{1!} d[F(y_n) \\ - F(y_{n-2})] = - \frac{[F(y_n) - F(y_{n-2})]^2}{2!} \Big|_{y_{n-3}}^{y_n} = \frac{[F(y_n) - F(y_{n-3})]^2}{2!},$$

and continuing on like this, we finally get:

$$\int_{y_1}^{y_n} \frac{[F(y_n) - F(y_2)]^{n-3}}{(n-3)!} f(y_2) dy_2 \\ = - \int_{y_1}^{y_n} \frac{[F(y_n) - F(y_2)]^{n-3}}{(n-3)!} d[F(y_n) - F(y_2)] \\ = - \frac{[F(y_n) - F(y_2)]^{n-2}}{(n-2)!} \Big|_{y_1}^{y_n} = \frac{[F(y_n) - F(y_1)]^{n-2}}{(n-2)!}.$$

Since $\frac{n!}{(n-2)!} = n(n-1)$, the result follows.



Chapter 7

Some Modes of Convergence of Random Variables, Applications

The first thing which is done in this chapter is to introduce two modes of convergence for sequences of r.v.'s, convergence in distribution and convergence in probability, and then to investigate their relationship.

A suitable application of these convergences leads to the most important results in this chapter, which are the Weak Law of Large Numbers and the Central Limit Theorem. These results are illustrated by concrete examples, including numerical examples in the case of the Central Limit Theorem.

In the final section of the chapter, it is shown that convergence in probability is preserved under continuity. This is also the case, in a limited sense, for convergence in distribution. These statements are illustrated by two general results and a specific application.

The proofs of some of the theorems stated are given in considerable detail; in some cases, only a rough outline is presented, whereas in other cases, we restrict ourselves to the statements of the theorems alone.

7.1 Convergence in Distribution or in Probability and their Relationship

In all that follows, X_1, \dots, X_n are i.i.d. r.v.'s, which may be either discrete or continuous. In applications, these r.v.'s represent n independent observations on a r.v. X , associated with an underlying phenomenon which is of importance to us. In a probabilistic/statistical environment, our interest lies in knowing the distribution of X , whether it is represented by the probabilities


$P(X \in B)$, $B \subseteq \mathfrak{R}$, or the d.f. F of the X_i 's, or their p.d.f. f . In practice, this distribution is unknown to us. Something then that would be desirable would be to approximate the unknown distribution, in some sense, by a known distribution. In this section, the foundation is set for such an approximation.

DEFINITION 1

Let Y_1, \dots, Y_n be r.v.'s with respective d.f.'s F_1, \dots, F_n . The r.v.'s may be either discrete or continuous and need be neither independent nor identically distributed. Also, let Y be a r.v. with d.f. G . We say that the sequence of r.v.'s $\{Y_n\}$, $n \geq 1$, *converges in distribution* to the r.v. Y as $n \rightarrow \infty$ and write $Y_n \xrightarrow[n \rightarrow \infty]{d} Y$, if $F_n(x) \xrightarrow[n \rightarrow \infty]{} G(x)$ for all continuity points x of G .

Figure 7.1

The d.f. Represented by the Solid Curve Is Approximated by the d.f.'s Represented by the $\cdots\cdots\cdots$, $-\cdot-\cdot-\cdot-$, and $- - - -$ Curves



The following example illustrates the definition.

EXAMPLE 1


For $n \geq 1$, let the d.f.'s F_n and the d.f. G be given by:

$$F_n(x) = \begin{cases} 0, & \text{if } x < 1 - \frac{1}{n} \\ \frac{1}{2}, & \text{if } 1 - \frac{1}{n} \leq x < 1 + \frac{1}{n} \\ 1, & \text{if } x \geq 1 + \frac{1}{n} \end{cases}, \quad G(x) = \begin{cases} 0, & \text{if } x < 1 \\ 1, & \text{if } x \geq 1, \end{cases}$$

and discuss whether or not $F_n(x)$ converges to $G(x)$ as $n \rightarrow \infty$.

Figure 7.2

The d.f. G Is
Approximated by the
d.f.'s F_n at all Points
 $x \neq 1$



DISCUSSION The d.f. G is continuous everywhere except for the point $x = 1$. For $x < 1$, let $n_0 > 1/(1 - x)$. Then $x < 1 - \frac{1}{n_0}$ and also $x < 1 - \frac{1}{n}$ for all $n \geq n_0$. Thus, $F_n(x) = 0$, $n \geq n_0$. For $x > 1$, let $n_0 \geq 1/(x - 1)$. Then $x \geq 1 + \frac{1}{n_0}$ and also $x \geq 1 + \frac{1}{n}$ for all $n \geq n_0$, so that $F_n(x) = 1$, $n \geq n_0$. Thus, for $x \neq 1$, $F_n(x) \rightarrow G(x)$, so, if Y_n and Y are r.v.'s such that $Y_n \sim F_n$ and $Y \sim G$, then $Y_n \xrightarrow[n \rightarrow \infty]{d} Y$.

REMARK 1 The example also illustrates the point that, if x is a discontinuity point of G , then $F_n(x)$ need not converge to $G(x)$. In Example 1, $F_n(1) = \frac{1}{2}$ for all n , and $G(1) = 1$.

The idea, of course, behind Definition 1 is the approximation of the (presumably unknown) probability $P(Y \leq x) = G(x)$ by the (presumably known) probabilities $P(Y_n \leq x) = F_n(x)$, for large enough n . Convergence in distribution also allows the approximation of probabilities of the form $P(x < Y \leq y)$ by the probabilities $P(x < Y_n \leq y)$, for x and y continuity points of G . This is so because

$$\begin{aligned} P(x < Y_n \leq y) &= P(Y_n \leq y) - P(Y_n \leq x) = F_n(y) - F_n(x) \\ &\xrightarrow[n \rightarrow \infty]{} G(y) - G(x) = P(x < Y \leq y). \end{aligned}$$


Whereas convergence in distribution allows the comparison of certain probabilities, calculated in terms of the *individual* r.v.'s Y_n and Y , it does not provide evaluation of probabilities calculated on the *joint* behavior of Y_n and Y . This is taken care of to a satisfactory extent by the following mode of convergence.

DEFINITION 2

The sequence of r.v.'s $\{Y_n\}$, $n \geq 1$, converges in probability to the r.v. Y as $n \rightarrow \infty$, if, for every $\varepsilon > 0$, $P(|Y_n - Y| > \varepsilon) \xrightarrow[n \rightarrow \infty]{} 0$; equivalently, $P(|Y_n - Y| \leq \varepsilon) \xrightarrow[n \rightarrow \infty]{} 1$. The notation used is: $Y_n \xrightarrow[n \rightarrow \infty]{P} Y$.

Thus, if the event $A_n(\varepsilon)$ is defined by: $A_n(\varepsilon) = \{s \in \mathcal{S}; Y(s) - \varepsilon \leq Y_n(s) \leq Y(s) + \varepsilon\}$, (that is, the event for which the r.v. Y_n is within ε from

the r.v. Y), then $P(A_n(\varepsilon)) \xrightarrow{n \rightarrow \infty} 1$ for every $\varepsilon > 0$. Equivalently, $P(A_n^c(\varepsilon)) = P(\{s \in \mathcal{S}; Y_n(s) < Y(s) - \varepsilon \text{ or } Y_n(s) > Y(s) + \varepsilon\}) \xrightarrow{n \rightarrow \infty} 0$.



The probability that Y_n lies within a small neighborhood around Y , such as $(Y_n - \varepsilon, Y_n + \varepsilon)$, is as close to 1 as one pleases, provided n is sufficiently large.

It is rather clear that convergence in probability is stronger than convergence in distribution. That this is, indeed, the case is illustrated by the following example, where we have convergence in distribution but not in probability.

EXAMPLE 2

Let $\mathcal{S} = \{1, 2, 3, 4\}$, and on the subsets of \mathcal{S} , let P be the discrete uniform probability function. Define the following r.v.'s:

$$X_n(1) = X_n(2) = 1, \quad X_n(3) = X_n(4) = 0, \quad n = 1, 2, \dots,$$

and

$$X(1) = X(2) = 0, \quad X(3) = X(4) = 1.$$

DISCUSSION Then

$$|X_n(s) - X(s)| = 1 \quad \text{for all } s \in \mathcal{S}.$$

Hence X_n does *not* converge in probability to X , as $n \rightarrow \infty$. Now,

$$F_n(x) = \begin{cases} 0, & x < 0 \\ \frac{1}{2}, & 0 \leq x < 1, \\ 1, & x \geq 1 \end{cases} \quad G(x) = \begin{cases} 0, & x < 0 \\ \frac{1}{2}, & 0 \leq x < 1 \\ 1, & x \geq 1, \end{cases}$$

so that $F_n(x) = G(x)$ for all $x \in \mathfrak{R}$. Thus, trivially, $F_n(x) \xrightarrow{n \rightarrow \infty} G(x)$ for all continuity points of G ; that is, $X_n \xrightarrow[n \rightarrow \infty]{d} X$, but X_n does not converge in probability to X .

The precise relationship between convergence in distribution and convergence in probability is stated in the following theorem.

THEOREM 1

Let $\{Y_n\}$, $n \geq 1$, be a sequence of r.v.'s and let Y be a r.v. Then $Y_n \xrightarrow[n \rightarrow \infty]{P} Y$ always implies $Y_n \xrightarrow[n \rightarrow \infty]{d} Y$. The converse is not true in general (as illustrated by Example 2). However, it is true if $P(Y=c)=1$, where c is a constant. That is, $Y_n \xrightarrow[n \rightarrow \infty]{d} c$ implies $Y_n \xrightarrow[n \rightarrow \infty]{P} c$, so that $Y_n \xrightarrow[n \rightarrow \infty]{P} c$ if and only if $Y_n \xrightarrow[n \rightarrow \infty]{d} c$.

PROOF (outline) That $Y_n \xrightarrow[n \rightarrow \infty]{P} Y$ implies $Y_n \xrightarrow[n \rightarrow \infty]{d} Y$ is established by employing the concepts of \liminf (limit inferior) and \limsup (limit superior) of a sequence of numbers, and we choose not to pursue it. For the proof of the fact that $Y_n \xrightarrow[n \rightarrow \infty]{d} c$ implies $Y_n \xrightarrow[n \rightarrow \infty]{P} c$, observe that $F(x)=0$, for $x < c$ and $F(x)=1$ for $x \geq c$, where F is the d.f. of c so that $c - \varepsilon$ and $c + \varepsilon$ are continuity points of F for all $\varepsilon > 0$. But $P(|Y_n - c| \leq \varepsilon) = P(c - \varepsilon \leq Y_n \leq c + \varepsilon) = P(Y_n \leq c + \varepsilon) - P(Y_n < c - \varepsilon) = F_n(c + \varepsilon) - P(Y_n < c - \varepsilon)$. However, $F_n(c + \varepsilon) \xrightarrow[n \rightarrow \infty]{} 1$ and $P(Y_n < c - \varepsilon) \leq P(Y_n \leq c - \varepsilon) = F_n(c - \varepsilon) \xrightarrow[n \rightarrow \infty]{} 0$, so that $P(Y_n < c - \varepsilon) \xrightarrow[n \rightarrow \infty]{} 0$. Thus, $P(|Y_n - c| \leq \varepsilon) \xrightarrow[n \rightarrow \infty]{} 1$ or $Y_n \xrightarrow[n \rightarrow \infty]{P} c$. \blacktriangle

According to Definition 1, in order to establish that $Y_n \xrightarrow[n \rightarrow \infty]{d} Y$, all one has to do is to prove the (pointwise) convergence $F_n(x) \xrightarrow[n \rightarrow \infty]{} F(x)$ for every continuity point x of F . As is often the case, however, definitions do not lend themselves to checking the concepts defined. This also holds here. Accordingly, convergence in distribution is delegated to convergence of m.g.f.'s, which, in general, is a much easier task to perform. That this can be done is based on the following deep probabilistic result. Its justification is omitted entirely.

THEOREM 2

(Continuity Theorem) For $n = 1, 2, \dots$, let Y_n and Y be r.v.'s with respective d.f.'s F_n and F , and respective m.g.f.'s M_n and M (which are assumed to be finite at least in an interval $(-c, c)$, some $c > 0$). Then:

(i) If $F_n(x) \xrightarrow[n \rightarrow \infty]{} F(x)$ for all continuity points x of F , it follows that $M_n(t) \xrightarrow[n \rightarrow \infty]{} M(t)$ for all $t \in (-c, c)$.

(ii) Let $M_n(t) \xrightarrow[n \rightarrow \infty]{} g(t)$, $t \in (-c, c)$, some function g , which is continuous at $t = 0$. Then g is, actually, a m.g.f. and let F be the corresponding d.f. It follows that $F_n(x) \xrightarrow[n \rightarrow \infty]{} F(x)$ for all continuity points x of F .

Thus, according to this result, $Y_n \xrightarrow[n \rightarrow \infty]{d} Y$ or, equivalently, $F_n(x) \xrightarrow[n \rightarrow \infty]{} F(x)$ for all continuity points x of F , if and only if $M_n(t) \xrightarrow[n \rightarrow \infty]{} M(t)$, $t \in (-c, c)$, some $c > 0$. The fact that convergence of m.g.f.'s implies convergence of the respective d.f.'s is the most useful part from a practical viewpoint.

Exercises

- 1.1** For $n = 1, 2, \dots$, let X_n be a r.v. with d.f. F_n defined by: $F_n(x) = 0$ for $x < n$, and $F_n(x) = 1$ for $x \geq n$. Then show that $F_n(x) \xrightarrow[n \rightarrow \infty]{} F(x)$, which is identically 0 in \Re and hence it is *not* a d.f. of a r.v.
- 1.2** Let $\{X_n\}$, $n \geq 1$, be r.v.'s with X_n taking the values 1 and 0 with respective probabilities p_n and $1 - p_n$; i.e., $P(X_n = 1) = p_n$ and $P(X_n = 0) = 1 - p_n$. Then show that $X_n \xrightarrow[n \rightarrow \infty]{P} 0$, if and only if $p_n \xrightarrow[n \rightarrow \infty]{} 0$.
- 1.3** For $n = 1, 2, \dots$, let X_n be a r.v. distributed as $B(n, p_n)$ and suppose that $np_n \xrightarrow[n \rightarrow \infty]{} \lambda \in (0, \infty)$. Then show that $X_n \xrightarrow[n \rightarrow \infty]{d} X$, where X is a r.v. distributed as $P(\lambda)$, by showing that $M_{X_n}(t) \xrightarrow[n \rightarrow \infty]{} M_X(t)$, $t \in \Re$.
- 1.4** Let $Y_{1,n}$ and $Y_{n,n}$ be the smallest and the largest order statistics based on the random sample X_1, \dots, X_n from the $U(0, 1)$ distribution. Then show that:
- (i) $Y_{1,n} \xrightarrow[n \rightarrow \infty]{P} 0$; (ii) $Y_{n,n} \xrightarrow[n \rightarrow \infty]{P} 1$.
- Hint:** For $\varepsilon > 0$, calculate the probabilities: $P(|Y_{1,n}| > \varepsilon)$ and $P(|Y_{n,n} - 1| > \varepsilon)$ and show that they tend to 0 as $n \rightarrow \infty$. Use the p.d.f.'s of $Y_{1,n}$ and $Y_{n,n}$ determined in Example 10 of Chapter 6.
- 1.5** Refer to Exercise 1.4 and set: $U_n = nY_{1,n}$, $V_n = n(1 - Y_{n,n})$, and let U and V be r.v.'s having the Negative Exponential distribution with parameter $\lambda = 1$. Then:
- (i) Derive the p.d.f.'s of the r.v.'s U_n and V_n .
- (ii) Derive the d.f.'s of the r.v.'s U_n and V_n , and show that $U_n \xrightarrow[n \rightarrow \infty]{d} U$ by showing that

$$F_{U_n}(u) \xrightarrow[n \rightarrow \infty]{} F_U(u), \quad u \in \Re.$$

Likewise for V_n .

- 1.6** We say that a sequence $\{X_n\}$, $n \geq 1$, of r.v.'s converges to a r.v. X in *quadratic mean* and write

$$X_n \xrightarrow[n \rightarrow \infty]{q.m.} X \quad \text{or} \quad X_n \xrightarrow[n \rightarrow \infty]{(2)} X, \quad \text{if } E(X_n - X)^2 \xrightarrow[n \rightarrow \infty]{} 0.$$

Now, if X_1, \dots, X_n are i.i.d. r.v.'s with (finite) expectation μ and (finite) variance σ^2 , show that the sample mean $\bar{X}_n \xrightarrow[n \rightarrow \infty]{q.m.} \mu$.

- 1.7** In Theorem 1 of Chapter 4, the following, version of the *Cauchy-Schwarz* inequality was established: For any two r.v.'s X and Y with $EX = EY = 0$ and $\text{Var}(X) = \text{Var}(Y) = 1$, it holds: $|E(XY)| \leq 1$. (This is, actually, part only of said inequality.) Another more general version of this inequality is the following: For any two r.v.'s X and Y with finite expectations and variances, it holds: $|E(XY)| \leq E|XY| \leq E^{1/2}|X|^2 \times E^{1/2}|Y|^2$.

- (i) Prove the inequality in this setting.
(ii) For any r.v. X , show that $|EX| \leq E|X| \leq E^{1/2}|X|^2$.

Hint: For part (i), use the obvious result $(x \pm y)^2 = x^2 + y^2 \pm 2xy \geq 0$ in order to conclude that $\pm xy \leq \frac{1}{2}(x^2 + y^2)$ and hence $|xy| \leq \frac{1}{2}(x^2 + y^2)$. Next, replace x by $X/E^{1/2}|X|^2$, and y by $Y/E^{1/2}|Y|^2$ (assuming, of course, that $E|X|^2 > 0$, $E|Y|^2 > 0$, because otherwise the inequality is, trivially, true), and take the expectations of both sides to arrive at the desirable result.

- 1.8 Let $\{X_n\}$ and $\{Y_n\}$, $n \geq 1$, be two sequences of r.v.'s such that: $X_n \xrightarrow[n \rightarrow \infty]{q.m.} X$, some r.v., and $X_n - Y_n \xrightarrow[n \rightarrow \infty]{q.m.} 0$. Then show that $Y_n \xrightarrow[n \rightarrow \infty]{q.m.} X$.

Hint: Use appropriately the Cauchy–Schwarz inequality discussed in Exercise 1.7.

7.2 Some Applications of Convergence in Distribution: The Weak Law of Large Numbers and the Central Limit Theorem

As a first application of the concept of convergence in distribution, we have the so-called *Weak Law of Large Numbers* (WLLN). This result is stated and proved, an interpretation is provided, and then a number of specific applications are presented.

THEOREM 3

(Weak Law of Large Numbers, WLLN) Let X_1, X_2, \dots be i.i.d. r.v.'s with (common) finite expectation μ , and let \bar{X}_n be the sample mean of X_1, \dots, X_n . Then $\bar{X}_n \xrightarrow[n \rightarrow \infty]{d} \mu$, or (on account of Theorem 1) $\bar{X}_n \xrightarrow[n \rightarrow \infty]{P} \mu$.

PROOF The proof is a one-line proof, if it happens that the X_i 's also have a (common) finite variance σ^2 (which they are not required to have for the validity of the theorem). Since $E\bar{X}_n = \mu$ and $\text{Var}(\bar{X}_n) = \frac{\sigma^2}{n}$, the Tchebichev inequality gives, for every $\varepsilon > 0$, $P(|\bar{X}_n - \mu| > \varepsilon) \leq \frac{1}{\varepsilon^2} \times \frac{\sigma^2}{n} \xrightarrow[n \rightarrow \infty]{} 0$, so that $\bar{X}_n \xrightarrow[n \rightarrow \infty]{P} \mu$.

Without reference to the variance, one would have to show that $M_{\bar{X}_n}(t) \xrightarrow[n \rightarrow \infty]{} M_\mu(t)$ (for $t \in (-c, c)$, some $c > 0$). Let M stand for the (common) m.g.f. of the X_i 's. Then use familiar properties of the m.g.f. and independence of the X_i 's in order to obtain:

$$M_{\bar{X}_n}(t) = M_{\sum_{i=1}^n X_i} \left(\frac{t}{n} \right) = \prod_{i=1}^n M_{X_i} \left(\frac{t}{n} \right) = \left[M \left(\frac{t}{n} \right) \right]^n.$$

Consider the function $M(z)$, and expand it around $z=0$ according to Taylor's formula up to terms of first order to get:

$$\begin{aligned} M(z) &= M(0) + \frac{z}{1!} \frac{d}{dz} M(z)|_{z=0} + R(z) \quad \left(\frac{1}{z} R(z) \rightarrow 0 \text{ as } z \rightarrow 0 \right) \\ &= 1 + z\mu + R(z), \end{aligned}$$

since $M(0) = 1$ and $\frac{d}{dz}M(z)|_{z=0} = EX_1 = \mu$. Replacing z by t/n , for fixed t , the last formula becomes:

$$M\left(\frac{t}{n}\right) = 1 + \frac{t}{n}\mu + R\left(\frac{t}{n}\right), \quad \text{where } nR\left(\frac{t}{n}\right) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore

$$M_{\bar{X}_n}(t) = \left[1 + \frac{\mu t + nR\left(\frac{t}{n}\right)}{n}\right]^n,$$

and this converges to $e^{\mu t}$, as $n \rightarrow \infty$, by Remark 2 below. Since $e^{\mu t}$ is the m.g.f. of (the degenerate r.v.) μ , we have shown that $M_{\bar{X}_n}(t) \xrightarrow[n \rightarrow \infty]{} M_\mu(t)$, as was to be seen. \blacktriangle

REMARK 2 For every $z \in \mathfrak{R}$, one way of defining the exponential function e^z is: $e^z = \lim_{n \rightarrow \infty} (1 + \frac{z}{n})^n$. It is a consequence of this result that, as $n \rightarrow \infty$, also $(1 + \frac{z_n}{n})^n \rightarrow e^z$ whenever $z_n \rightarrow z$.

The interpretation and most common use of the WLLN is that, if μ is an unknown entity, which is typically the case in statistics, then μ may be approximated (in the sense of distribution or probability) by the known entity \bar{X}_n , for sufficiently large n .

7.2.1 Applications of the WLLN

1. If the independent X_i 's are distributed as $B(1, p)$, then $EX_i = p$ and therefore $\bar{X}_n \xrightarrow[n \rightarrow \infty]{P} p$.
2. If the independent X_i 's are distributed as $P(\lambda)$, then $EX_i = \lambda$ and therefore $\bar{X}_n \xrightarrow[n \rightarrow \infty]{P} \lambda$.
3. If the independent X_i 's are distributed as $N(\mu, \sigma^2)$, then $EX_i = \mu$ and therefore $\bar{X}_n \xrightarrow[n \rightarrow \infty]{P} \mu$.
4. If the independent X_i 's are distributed as Negative Exponential with parameter λ , $f(x) = \lambda e^{-\lambda x}$, $x > 0$, then $EX_i = 1/\lambda$ and therefore $\bar{X}_n \xrightarrow[n \rightarrow \infty]{P} 1/\lambda$.

A somewhat more involved application is that of the approximation of an entire d.f. by the so-called empirical d.f. To this effect:

5. Let X_1, X_2, \dots, X_n be i.i.d. r.v.'s with d.f. F , and define the *empirical* d.f. F_n as follows. For each $x \in \mathfrak{R}$ and each $s \in \mathcal{S}$,

$$F_n(x, s) = \frac{1}{n}[\text{number of } X_1(s), \dots, X_n(s) \leq x].$$

From this definition, it is immediate that, for each fixed $x \in \mathfrak{R}$, $F_n(x, s)$ is a r.v. as a function of s , and for each fixed $s \in \mathcal{S}$, $F_n(x, s)$ is a d.f. as a function of x . Actually, if we set $Y_i(x, s) = 1$ when $X_i(s) \leq x$, and $Y_i(x, s) = 0$ when $X_i(s) > x$, then $Y_i(x, \cdot), \dots, Y_i(x, \cdot)$ are r.v.'s which are independent and distributed as $B(1, F(x))$, since $P[Y_i(x, \cdot) = 1] = P(X_i \leq x) = F(x)$. Also, $EY_i(x, \cdot) = F(x)$.

Then $F_n(x, s)$ may be rewritten as:

$$F_n(x, s) = \frac{1}{n} \sum_{i=1}^n Y_i(x, s), \quad \text{the sample mean of } Y_1(x, s), \dots, Y_n(x, s).$$

By omitting the sample point s , as is usually the case, we write $F_n(x)$ and $Y_i(x)$, $i = 1, \dots, n$ rather than $F_n(x, s)$ and $Y_i(x, s)$, $i = 1, \dots, n$, respectively. Then $F_n(x) \xrightarrow{P} F(x)$ for each $x \in \mathfrak{R}$. Thus, for every $x \in \mathfrak{R}$, the value of $F(x)$ of the (potentially unknown) d.f. F is approximated by the (known) values $F_n(x)$ of the r.v.'s $F_n(x)$.

REMARK 3 Actually, it can be shown that the convergence $F_n(x) \xrightarrow{P} F(x)$ is *uniform* in $x \in \mathfrak{R}$. This implies that, for every $\varepsilon > 0$, there is a positive integer $N(\varepsilon)$ such that $F_n(x) - \varepsilon < F(x) < F_n(x) + \varepsilon$ with probability as close to 1 as one pleases *simultaneously* for all $x \in \mathfrak{R}$, provided $n > N(\varepsilon)$.

As another application of the concept of convergence in distribution, we obtain, perhaps, the most celebrated theorem of Probability Theory; it is the so-called *Central Limit Theorem* (CLT), which is stated and proved below. Comments on the significance of the CLT follow, and the section is concluded by applications and numerical examples.

THEOREM 4

(Central Limit Theorem, CLT) Let X_1, X_2, \dots be i.i.d. r.v.'s with finite expectation μ and finite and positive variance σ^2 , and let \bar{X}_n be the sample mean of X_1, \dots, X_n . Then:

$$\frac{\bar{X}_n - E\bar{X}_n}{\sqrt{\text{Var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu}{\frac{\sigma}{\sqrt{n}}} = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1),$$

or

$$P\left[\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \leq z\right] \xrightarrow[n \rightarrow \infty]{} \Phi(z) = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx, \quad z \in \mathfrak{R}. \quad (1)$$

(Also, see Remark 4(iii).)

REMARK 4

- (i) Denote by S_n the partial sum $\sum_{i=1}^n X_i$, $S_n = \sum_{i=1}^n X_i$, so that $ES_n = n\mu$ and $\text{Var}(S_n) = n\sigma^2$. Then:

$$\frac{S_n - ES_n}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - n\mu}{\sigma\sqrt{n}} = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma}.$$

Therefore, by (1):

$$P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq z\right) \xrightarrow[n \rightarrow \infty]{} \Phi(z), \quad z \in \mathfrak{R}. \quad (2)$$

(Although the notation S_n has been used before (relation (12) in Chapter 5) to denote the sample standard deviation of X_1, \dots, X_n , there should be no confusion; from the context, it should be clear what S_n stands for.)

(ii) An interpretation of (1) and (2) is that, for sufficiently large n :

$$P\left[\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \leq z\right] = P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq z\right) \simeq \Phi(z), \quad z \in \mathfrak{R}. \quad (3)$$

Often this approximation is also denoted (rather loosely) as follows:

$$\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \simeq N(0, 1) \quad \text{or} \quad \bar{X}_n \simeq N\left(\mu, \frac{\sigma^2}{n}\right) \quad \text{or} \quad S_n \simeq N(n\mu, n\sigma^2). \quad (4)$$

(iii) Actually, it can be shown that the convergence in (1) or (2) is *uniform* in $z \in \mathfrak{R}$. That is to say, if we set

$$F_n(z) = P\left[\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \leq z\right] = P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq z\right), \quad (5)$$

then

$$F_n(z) \xrightarrow[n \rightarrow \infty]{} \Phi(z) \quad \text{uniformly in } z \in \mathfrak{R}. \quad (6)$$

To be more precise, for every $\varepsilon > 0$, there exists a positive integer $N(\varepsilon)$ *independent* of $z \in \mathfrak{R}$, such that

$$|F_n(z) - \Phi(z)| < \varepsilon \quad \text{for } n \geq N(\varepsilon) \text{ and all } z \in \mathfrak{R}. \quad (7)$$

(iv) The approximation of the probability $F_n(z)$ by $\Phi(z)$, provided by the CLT, is also referred to as *Normal approximation* for obvious reasons.

(v) On account of (3), the CLT also allows for the approximation of probabilities of the form $P(a < S_n \leq b)$ for any $a < b$. Indeed,

$$\begin{aligned} P(a < S_n \leq b) &= P(S_n \leq b) - P(S_n \leq a) \\ &= P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq \frac{b - n\mu}{\sigma\sqrt{n}}\right) - P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq \frac{a - n\mu}{\sigma\sqrt{n}}\right) \\ &= P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq b_n^*\right) - P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq a_n^*\right), \end{aligned}$$

where

$$a_n^* = \frac{a - n\mu}{\sigma\sqrt{n}} \quad \text{and} \quad b_n^* = \frac{b - n\mu}{\sigma\sqrt{n}}. \quad (8)$$

By (3),

$$P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq b_n^*\right) \simeq \Phi(b_n^*) \quad \text{and} \quad P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq a_n^*\right) \simeq \Phi(a_n^*),$$

so that

$$P(a < S_n \leq b) \simeq \Phi(b_n^*) - \Phi(a_n^*). \quad (9)$$

The uniformity referred to in Remark 3(iii) is what, actually, validates many of the applications of the CLT. This is the case, for instance, in Remark 3(v).

(vi) So, the convergence in (i) is a special case of the convergence depicted in Figure 7.1, where the limiting d.f. is Φ and F_n is the d.f. of $\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma}$. This convergence holds for all $x \in \mathfrak{R}$ since Φ is a continuous function in \mathfrak{R} .

EXAMPLE 3

From a large collection of bolts which is known to contain 3% defective bolts, 1,000 are chosen at random. If X is the number of the defective bolts among those chosen, what is the (approximate) probability that X does not exceed 5% of 1,000?

DISCUSSION With the selection of the i th bolt, associate the r.v. X_i to take the value 1, if the bolt is defective, and 0 otherwise. Then it may be assumed that the r.v.'s $X_i, i = 1, \dots, 1,000$ are independently distributed as $B(1, 0.03)$. Furthermore, it is clear that $X = \sum_{i=1}^{1,000} X_i$. Since 5% of 1,000 is 50, the required probability is: $P(X \leq 50)$. Since $EX_i = 0.03$, $Var(X_i) = 0.03 \times 0.97 = 0.0291$, the CLT gives:

$$\begin{aligned} P(X \leq 50) &= P(0 \leq X \leq 50) = P(-0.5 < X \leq 50) \\ &= P(X \leq 50) - P(X \leq -0.5) \simeq \Phi(b_n^*) - \Phi(a_n^*), \end{aligned}$$

$$\text{where } a_n^* = \frac{-0.5 - 1,000 \times 0.03}{\sqrt{1,000 \times 0.03 \times 0.97}} = -\frac{30.5}{\sqrt{29.1}} \simeq -\frac{30.5}{5.394} \simeq -5.65,$$

$$b_n^* = \frac{50 - 1,000 \times 0.03}{\sqrt{1,000 \times 0.03 \times 0.97}} = \frac{20}{\sqrt{29.1}} \simeq \frac{20}{5.394} \simeq 3.71,$$

so that

$$P(X \leq 50) \simeq \Phi(3.71) - \Phi(-5.65) = \Phi(3.71) = 0.999896.$$

EXAMPLE 4

A certain manufacturing process produces vacuum tubes whose lifetimes in hours are independent r.v.'s with Negative Exponential distribution with mean 1,500 hours. What is the probability that the total life of 50 tubes will exceed 80,000 hours?

DISCUSSION If X_i is the r.v. denoting the lifetime of the i th vacuum tube, then $X_i, i = 1, \dots, 50$ are independent Negative Exponentially distributed with $EX_i = \frac{1}{\lambda} = 1,500$ and $Var(X_i) = \frac{1}{\lambda^2} = 1,500^2$. Since $nEX_i = 50 \times 1,500 = 75,000$, $\sigma\sqrt{n} = 1,500\sqrt{50}$, if we set $S_{50} = \sum_{i=1}^{50} X_i$, then the required probability is:

$$\begin{aligned} P(S_{50} > 80,000) &= 1 - P(S_{50} \leq 80,000) \simeq 1 - \Phi\left(\frac{80,000 - 75,000}{1,500\sqrt{50}}\right) \\ &= 1 - \Phi\left(\frac{\sqrt{50}}{15}\right) \simeq 1 - \Phi(0.47) \\ &= 1 - 0.680822 = 0.319178 \simeq 0.319. \end{aligned}$$

The proof of the theorem is based on the same ideas as those used in the proof of the WLLN and goes as follows.

PROOF OF THEOREM 4 Set $Z_i = \frac{X_i - \mu}{\sigma}$, so that Z_1, \dots, Z_n are i.i.d. r.v.'s with $EZ_i = 0$ and $\text{Var}(Z_i) = 1$. Also,

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n Z_i = \frac{1}{\sigma\sqrt{n}}(S_n - n\mu) = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma}. \quad (10)$$

With F_n defined by (5), we wish to show that (6) holds (except for the uniformity assertion, with which we will not concern ourselves). Its justification is provided by Lemma 1, pages 206–207, in the book “A Course in Mathematical Statistics,” 2nd edition (1997), Academic Press, by G. G. Roussas. By Theorem 2, it suffices to show that, for all t ,

$$M_{\sqrt{n}(\bar{X}_n - \mu)/\sigma}(t) \xrightarrow{n \rightarrow \infty} M_Z(t) = e^{t^2/2}. \quad (11)$$

By means of (10), and with M standing for the (common) m.g.f. of the Z_i 's, we have:

$$\begin{aligned} M_{\sqrt{n}(\bar{X}_n - \mu)/\sigma}(t) &= M_{\frac{1}{\sqrt{n}} \sum_{i=1}^n z_i}(t) = M_{\sum_{i=1}^n z_i}\left(\frac{t}{\sqrt{n}}\right) \\ &= \prod_{i=1}^n M_{Z_i}\left(\frac{t}{\sqrt{n}}\right) = \left[M\left(\frac{t}{\sqrt{n}}\right)\right]^n. \end{aligned} \quad (12)$$

Expand the function $M(z)$ around $z = 0$ according to Taylor's formula up to terms of second order to get:

$$\begin{aligned} M(z) &= M(0) + \frac{z}{1!} \frac{d}{dz} M(z)|_{z=0} + \frac{z^2}{2!} \frac{d^2}{dz^2} M(z)|_{z=0} + R(z) \\ &= 1 + zEZ_1 + \frac{z^2}{2} EZ_1^2 + R(z) \\ &= 1 + \frac{z^2}{2} + R(z), \quad \text{where } \frac{1}{z^2} R(z) \rightarrow 0 \text{ as } z \rightarrow 0. \end{aligned}$$

In this last formula, replace z by t/\sqrt{n} , for fixed t , in order to obtain:

$$M\left(\frac{t}{\sqrt{n}}\right) = 1 + \frac{t^2}{2n} + R\left(\frac{t}{\sqrt{n}}\right), \quad nR\left(\frac{t}{\sqrt{n}}\right) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore (12) becomes:

$$M_{\sqrt{n}(\bar{X}_n - \mu)/\sigma}(t) = \left[1 + \frac{t^2}{2n} + R\left(\frac{t}{\sqrt{n}}\right)\right]^n = \left\{1 + \frac{t^2 \left[1 + \frac{2n}{t^2} R\left(\frac{t}{\sqrt{n}}\right)\right]}{n}\right\}^n,$$

and this converges to $e^{t^2/2}$, as $n \rightarrow \infty$, by Remark 2. This completes the proof of the theorem. \blacktriangle

7.2.2 Applications of the CLT

In all of the following applications, it will be assumed that n is sufficiently large, so that the CLT will apply.

1. Let the independent X_i 's be distributed as $B(1, p)$, set $S_n = \sum_{i=1}^n X_i$, and let a, b be integers such that $0 \leq a < b \leq n$. By an application of the CLT, we wish to find an approximate value to the probability $P(a < S_n \leq b)$.

If p denotes the proportion of defective items in a large lot of certain items, then S_n is the number of actually defective items among the n sampled. Then approximation of the probability $P(a < S \leq b)$ is meaningful when the Binomial tables are not usable (either because of p or because of n or, perhaps, because of both).

Here $EX_i = p$, $Var(X_i) = pq$ ($q = 1 - p$), and therefore by (9):

$$P(a < S_n \leq b) \simeq \Phi(b_n^*) - \Phi(a_n^*), \quad \text{where } a_n^* = \frac{a - np}{\sqrt{npq}}, \quad b_n^* = \frac{b - np}{\sqrt{npq}}. \quad (13)$$

REMARK 5 If the required probability is of any one of the forms: $P(a \leq S_n \leq b)$ or $P(a \leq S_n < b)$ or $P(a < S_n \leq b)$, then formula (9) applies again, provided the necessary adjustments are first made; namely, $P(a \leq S_n \leq b) = P(a - 1 < S_n \leq b)$, $P(a \leq S_n < b) = P(a - 1 < S_n \leq b - 1)$, $P(a < S_n \leq b) = P(a < S_n \leq b - 1)$. However, if the underlying distribution is continuous, then $P(a < S_n \leq b) = P(a \leq S_n \leq b) = P(a \leq S_n < b) = P(a < S_n < b)$, and no adjustments are required for the approximation in (9) to hold.

EXAMPLE 5

(Numerical) For $n = 100$ and $p = \frac{1}{2}$ or $p = \frac{5}{16}$, find the probability $P(45 \leq S_n \leq 55)$.

DISCUSSION

- (i) For $p = \frac{1}{2}$, it is seen (from tables) that the exact value is equal to: 0.7288. For the Normal approximation, we have: $P(45 \leq S_n \leq 55) = P(44 < S_n \leq 55)$ and, by (13):

$$a^* = \frac{44 - 100 \times \frac{1}{2}}{\sqrt{100 \times \frac{1}{2} \times \frac{1}{2}}} = -\frac{6}{5} = -1.2, \quad b^* = \frac{55 - 100 \times \frac{1}{2}}{\sqrt{100 \times \frac{1}{2} \times \frac{1}{2}}} = \frac{5}{5} = 1.$$

Therefore $\Phi(b^*) - \Phi(a^*) = \Phi(1) - \Phi(-1.2) = \Phi(1) + \Phi(1.2) - 1 = 0.841345 + 0.884930 - 1 = 0.7263$. So:

$$\text{Exact value: } 0.7288, \quad \text{Approximate value: } 0.7263,$$

and the exact probability is underestimated by 0.0025, or the approximate probability is about 99.66% of the exact probability.

- (ii) For $p = \frac{5}{16}$, the exact probability is almost 0; 0.0000. For the approximate probability, we find $a^* = 2.75$ and $b^* = 4.15$, so that $\Phi(b^*) - \Phi(a^*) = 0.0030$. Thus:

$$\text{Exact value: } 0.0000, \quad \text{Approximate value: } 0.0030,$$

and the exact probability is overestimated by 0.0030.

2. If the underlying distribution is $P(\lambda)$, then $ES_n = Var(S_n) = n\lambda$ and formulas (8) and (9) become:

$$P(a < S_n \leq b) \simeq \Phi(b_n^*) - \Phi(a_n^*), \quad a_n^* = \frac{a - n\lambda}{\sqrt{n\lambda}}, \quad b_n^* = \frac{b - n\lambda}{\sqrt{n\lambda}}.$$

The comments made in Remark 4 apply here also.

EXAMPLE 6

(Numerical) In the Poisson distribution $P(\lambda)$, let n and λ be so that $n\lambda = 16$ and find the probability $P(12 \leq S_n \leq 21)$ ($= P(11 < S_n \leq 21)$).

DISCUSSION The exact value (found from tables) is: 0.7838. For the Normal approximation, we have:

$$a^* = \frac{11 - 16}{\sqrt{16}} = -\frac{5}{4} = -1.25, \quad b^* = \frac{21 - 16}{\sqrt{16}} = \frac{5}{4} = 1.25,$$

so that $\Phi(b^*) - \Phi(a^*) = \Phi(1.25) - \Phi(-1.25) = 2\Phi(1.25) - 1 = 2 \times 0.894350 - 1 = 0.7887$. So:

$$\text{Exact value: } 0.7838, \quad \text{Approximate value: } 0.7887,$$


and the exact probability is overestimated by 0.0049, or the approximating probability is about 100.63% of the exact probability.

7.2.3 The Continuity Correction

When a discrete distribution is approximated by the Normal distribution, the error committed is easy to see in a geometric picture. This is done, for instance in Figure 7.3, where the p.d.f. of the $B(10, 0.2)$ distribution is approximated by the p.d.f. of the $N(10 \times 0.2, 10 \times 0.2 \times 0.8) = N(2, 1.6)$ distribution (see relation (4)). From the same figure, it is also clear how the approximation may be improved.

Figure 7.3

Exact and Approximate Values for the Probability
 $P(a \leq S_n \leq b_n) =$
 $P(a - 1 < S_n \leq$
 $b_n) = P(1 < S_n \leq 3)$



Now

$$\begin{aligned} P(1 < S_n \leq 3) &= P(2 \leq S_n \leq 3) = f_n(2) + f_n(3) \\ &= \text{shaded area,} \end{aligned}$$

while the approximation without correction is the area bounded by the Normal curve, the horizontal axis, and the abscissas 1 and 3. Clearly, the correction, given by the area bounded by the Normal curve, the horizontal axis, and the abscissas 1.5 and 3.5, is closer to the exact area.

To summarize, under the conditions of the CLT, and for discrete r.v.'s, $P(a < S_n \leq b) \simeq \Phi(b^*) - \Phi(a^*)$, where $a^* = \frac{a - n\mu}{\sigma\sqrt{n}}$ and $b^* = \frac{b - n\mu}{\sigma\sqrt{n}}$ without continuity correction, and $P(a < S_n \leq b) \simeq \Phi(b') - \Phi(a')$, where $a' = \frac{a + 0.5 - n\mu}{\sigma\sqrt{n}}$ and $b' = \frac{b + 0.5 - n\mu}{\sigma\sqrt{n}}$ with continuity correction.

For integer-valued r.v.'s and probabilities of the form $P(a \leq S_n \leq b)$, we first rewrite the expression as follows:

$$P(a \leq S_n \leq b) = P(a - 1 < S_n \leq b),$$

and then apply the preceding approximations in order to obtain:

$$P(a \leq S_n \leq b) \simeq \Phi(b^*) - \Phi(a^*), \text{ where}$$

$a^* = \frac{a-1-n\mu}{\sigma\sqrt{n}}$ and $b^* = \frac{b-n\mu}{\sigma\sqrt{n}}$ without continuity correction, and $P(a \leq S_n \leq b) \simeq \Phi(b') - \Phi(a')$, where $a' = \frac{a-0.5-n\mu}{\sigma\sqrt{n}}$ and $b' = \frac{b+0.5-n\mu}{\sigma\sqrt{n}}$ with continuity correction. Similarly for the intervals $[a, b)$ and (a, b) .

The improvement brought about by the continuity correction is demonstrated by the following numerical examples.

EXAMPLE 5

(continued)

DISCUSSION

(i) For $p = \frac{1}{2}$, we get:

$$a' = \frac{44 + 0.5 - 100 \times \frac{1}{2}}{\sqrt{100 \times \frac{1}{2} \times \frac{1}{2}}} = -\frac{5.5}{5} = -1.1,$$

$$b' = \frac{55 + 0.5 - 100 \times \frac{1}{2}}{\sqrt{100 \times \frac{1}{2} \times \frac{1}{2}}} = \frac{5.5}{5} = 1.1,$$

so that:

$$\begin{aligned} \Phi(b') - \Phi(a') &= \Phi(1.1) - \Phi(-1.1) = 2\Phi(1.1) - 1 \\ &= 2 \times 0.864334 - 1 = 0.7286. \end{aligned}$$

Thus, we have:

Exact value: 0.7288,

Approximate value with continuity correction: 0.7286,

and the approximation underestimates the probability by only 0.0002, or the approximating probability (with continuity correction) is about 99.97% of the exact probability.

(ii) For $p = \frac{5}{16}$, we have $a' = 2.86$, $b' = 5.23$ and $\Phi(b') - \Phi(a') = 0.0021$. Then:

Exact value: 0.0000,

Approximate value with continuity correction: 0.0021,

and the probability is overestimated by only 0.0021.

EXAMPLE 6

(continued)

DISCUSSION Here:

$$a' = \frac{11 + 0.5 - 16}{\sqrt{16}} = -\frac{4.5}{4} = -1.125, \quad b' = \frac{21 + 0.5 - 16}{\sqrt{16}} = \frac{5.5}{4} = 1.375,$$

so that:

$$\Phi(b') - \Phi(a') = \Phi(1.375) - \Phi(-1.125) = \Phi(1.375) + \Phi(1.125) - 1 = 0.7851.$$

Thus:

Exact value: 0.7838,*Approximate value with continuity correction:* 0.7851,

and the approximation overestimates the probability by only 0.0013, or the approximating probability (with continuity correction) is about 100.17% of the exact probability.

Exercises

2.1 Let X_1, \dots, X_n be i.i.d. r.v.'s, and for a positive integer k , suppose that EX_1^k is finite. Form the k th sample mean $\bar{X}_n^{(k)}$ defined by

$$\bar{X}_n^{(k)} = \frac{1}{n} \sum_{i=1}^n X_i^k.$$

Then show that:

$$\bar{X}_n^{(k)} \xrightarrow[n \rightarrow \infty]{P} EX_1^k.$$

2.2 Let X be a r.v. with p.d.f. $f_X(x) = c\alpha^x$, $x = 0, 1, \dots$ ($0 < \alpha < 1$). Then $c = 1 - \alpha$ by Exercise 2.8 in Chapter 2.

- (i) Show that the m.g.f. of X is: $M_X(t) = \frac{1-\alpha}{1-\alpha e^t}$, $t < -\log \alpha$.
- (ii) Use the m.g.f. to show that $EX = \frac{\alpha}{1-\alpha}$.
- (iii) If X_1, \dots, X_n is a random sample from f_X , show that the WLLN holds by showing that

$$M_{\bar{X}_n}(t) \xrightarrow[n \rightarrow \infty]{} e^{\alpha t/(1-\alpha)} = M_{EX}(t), \quad t < -\log \alpha.$$

Hint: Expand e^t around 0 up to second term, according to Taylor's formula, $e^t = 1 + t + R(t)$, where $\frac{1}{t}R(t) \xrightarrow[t \rightarrow 0]{} 0$, replace t by $\frac{t}{n}$, and use the fact that $(1 + \frac{x_n}{n})^n \rightarrow e^x$, if $x_n \rightarrow x$ as $n \rightarrow \infty$.

2.3 Let the r.v. X be distributed as $B(150, 0.6)$. Then:

- (i) Write down the formula for the exact probability $P(X \leq 80)$.
- (ii) Use the CLT in order to find an approximate value for the above probability. (Do not employ the continuity correction.)

2.4 A binomial experiment with probability p of a success is repeated independently 1,000 times, and let X be the r.v. denoting the number of successes. For $p = \frac{1}{2}$ and $p = \frac{1}{4}$, find:

- (i) The exact probability $P(1,000p - 50 \leq X \leq 1,000p + 50)$
- (ii) Use the CLT to find an approximate value for this probability.

2.5 Let X_1, \dots, X_{100} be independent r.v.'s distributed as $B(1, p)$. Then:

- (i) Write out the expression for the exact probability $P(\sum_{i=1}^{100} X_i = 50)$.
- (ii) Use of CLT in order to find an approximate value for this probability.
- (iii) What is the numerical value of the probability in part (ii) for $p = 0.5$?

Hint: For part (ii), first observe that $P(X = 50) = P(49.5 < X \leq 50)$, and then apply the CLT.

2.6 Fifty balanced dice are tossed once, and let X be the r.v. denoting the sum of the upturned spots. Use the CLT to find an approximate value of the probability $P(150 \leq X \leq 200)$.

Hint: With the i th die, associate the r.v. X_i which takes on the values 1 through 6, each with probability $1/6$. These r.v.'s may be assumed to be independent and $X = \sum_{i=1}^{50} X_i$.

2.7 One thousand cards are drawn (with replacement) from a standard deck of 52 playing cards, and let X be the r.v. denoting the total number of aces drawn. Use the CLT to find an approximate value of the probability $P(65 \leq X \leq 90)$.

2.8 From a large collection of bolts which is known to contain 3% defective bolts, 1,000 are chosen at random, and let X be the r.v. denoting the number of defective bolts among those chosen. Use the CLT to find an approximate value of the probability that X does not exceed 5% of 1,000.

Hint: With the i th bolt drawn, associate the r.v. X_i which takes on the value 1, if the bolt drawn is defective, and 0 otherwise. Since the collection of bolts is large, we may assume that after each drawing, the proportion of the remaining defective bolts remains (approximately) the same. This implies that the independent r.v.'s $X_1, \dots, X_{1,000}$ are distributed as $B(1, 0.03)$ and that $X = \sum_{i=1}^{1,000} X_i \sim B(1,000, 0.3)$.

2.9 A manufacturing process produces defective items at the constant (but unknown to us) proportion p . Suppose that n items are sampled independently, and let X be the r.v. denoting the number of defective items among the n , so that $X \sim B(n, p)$. Determine the smallest value of the sample size n , so that

$$P\left(\left|\frac{X}{n} - p\right| < 0.05\sqrt{pq}\right) \geq 0.95 \quad (q = 1 - p):$$

- (i) By utilizing the CLT.
- (ii) By using the Tchebichev inequality.
- (iii) Compare the answers in parts (i) and (ii).

2.10 Suppose that 53% of the voters favor a certain legislative proposal. How many voters must be sampled so that the observed relative frequency of those favoring the proposal will not differ from the assumed frequency by more than 2% with probability 0.99?

Hint: With the i th voter sampled, associate the r.v. X_i which takes on the value 1, if the voter favors the proposal, and 0 otherwise. Then it may be assumed that the r.v.'s X_1, \dots, X_n are independent and their common distribution is $B(1, 0.53)$. Furthermore, the number of the voters favoring the proposal is $X = \sum_{i=1}^n X_i$. Use the CLT in order to find the required probability.

2.11 In playing a game, you win or lose \$1 with probability 0.5, and you play the game independently 1,000 times. Use the CLT to find an approximate value of the probability that your fortune (i.e., the total amount you won or lost) is at least \$10.

Hint: With the i th game, associate the r.v. X_i which takes on the value 1 if \$1 is won, and -1 if \$1 is lost. Then the r.v.'s $X_1, \dots, X_{1,000}$ are independent, and the fortune X is given by $\sum_{i=1}^{1,000} X_i$.

2.12 It is known that the number of misprints in a page of a certain publication is a r.v. X having the Poisson distribution with parameter λ . If X_1, \dots, X_n are the misprints counted in n pages, use the CLT to determine the (approximate) probability that the total number of misprints is:

- (i) Not more than λn .
- (ii) At least λn .
- (iii) Between $\lambda n/2$ and $3\lambda n/2$.
- (iv) Give the numerical values in parts (i)–(iii) for $\lambda n = 100$ (which may be interpreted, e.g., as one misprint per 4 pages ($\lambda = 0.25$) in a book of 400 pages).

2.13 Let the r.v. X be distributed as $P(100)$. Then:

- (i) Write down the formula for the exact probability $P(X \leq 116)$.
- (ii) Use the CLT appropriately in order to find an approximate value for the above probability. (Do not use the continuity correction.)

Hint: Select n large and λ small, so that $n\lambda = 100$ and look at X as the sum $\sum_{i=1}^n X_i$ of n independent r.v.'s X_1, \dots, X_n distributed as $P(\lambda)$.

2.14 A certain manufacturing process produces vacuum tubes whose lifetimes in hours are independently distributed r.v.'s with Negative Exponential distribution with mean 1,500 hours. Use the CLT in order to find an

approximate value for the probability that the total life of 50 tubes will exceed 80,000 hours.

- 2.15** The lifespan of an electronic component in a (complicated) system is a r.v. X having the Negative Exponential distribution with parameter λ .
- What is the probability that said lifespan will be at least t time units?
 - If the independent r.v.'s X_1, \dots, X_n represent the lifespans of n spare items such as the one described above, then $Y = \sum_{i=1}^n X_i$ is the combined lifespan of these n items. Use the CLT in order to find an approximate value of the probability $P(t_1 \leq Y \leq t_2)$, where $0 < t_1 < t_2$ are given time units.
 - Compute the numerical answer in part (i), if $t = -\log(0.9)/\lambda$.
 - Do the same for part (ii), if $\lambda = 1/10$, $n = 36$, $t_1 = 300$, and $t_2 = 420$.
- 2.16** Let the independent r.v.'s X_1, \dots, X_n be distributed as $U(0, 1)$.
- Use the CLT to find an approximate value for the probability $P(a \leq \bar{X} \leq b)$ ($a < b$).
 - What is the numerical value of this probability for $n = 12$, $a = 7/16$, and $b = 9/16$?
- 2.17** If the independent r.v.'s X_1, \dots, X_{12} are distributed as $U(0, \theta)$ ($\theta > 0$), use the CLT to show that the probability $P(\frac{\theta}{4} < \bar{X} < \frac{3\theta}{4})$ is approximately equal to 0.9973.
- 2.18** Refer to Exercise 3.42 in Chapter 3 and let $X_i, i = 1, \dots, n$ be the diameters of n ball bearings. If $EX_i = \mu = 0.5$ inch and s.d. $(X_i) = \sigma = 0.0005$ inch, use the CLT to determine the smallest value of n for which $P(|\bar{X} - \mu| \leq 0.0001) = 0.99$, where \bar{X} is the sample mean of the X_i 's.
- 2.19** The i.i.d. r.v.'s X_1, \dots, X_{100} have (finite) mean μ and variance $\sigma^2 = 4$. Use the CLT to determine the value of the constant c for which $P(|\bar{X} - \mu| \leq c) = 0.90$, where \bar{X} is the sample means of the X_i 's.
- 2.20** Let X_1, \dots, X_n be i.i.d. r.v.'s with (finite) expectation μ and (finite and positive) variance σ^2 , and let \bar{X}_n be the sample mean of the X_i 's. Determine the smallest value of the sample size n , in terms of k and p , for which $P(|\bar{X}_n - \mu| < k\sigma) \geq p$, where $p \in (0, 1)$, $k > 0$. Do so by using:
- The CLT.
 - The Tchebichev inequality.
 - Find the numerical values of n in parts (i) and (ii) for $p = 0.90, 0.95, 0.99$ and $k = 0.50, 0.25, 0.10$ for each value of p .
- 2.21** Refer to Exercise 3.41 in Chapter 3, and suppose that the r.v. X considered there has $EX = 2,000$ and s.d. $(X) = 200$, but is not necessarily Normally distributed. Also, consider another manufacturing process producing light bulbs whose mean lifespan is claimed to be 10% higher than the mean lifespan of the bulbs produced by the existing process; it is assumed

that the s.d. remains the same for the new process. How many bulbs manufactured by the new process must be examined to establish the claim of their superiority (should that be the case) with probability 0.95?

Hint: Let Y be the r.v. denoting the lifespan of a light bulb manufactured by the new process. We do not necessarily assume that Y is Normally distributed. If the claim made is correct, then $EY = 2,000 + 10\% \times 2,000 = 2,200$, whereas $\text{s.d.}(Y) = 200$. A random sample from Y produces the sample mean \bar{Y}_n for which $E\bar{Y}_n = 2,200$ (under the claim) and $\text{Var}(\bar{Y}_n) = 200^2/n$, and we must determine n , so that $P(\bar{Y}_n > 2,000) = 0.95$. If the new process were the same as the old one, then, for all sufficiently large n , $P(\bar{Y}_n > 2,000) \simeq 0.50$. So, if $P(\bar{Y}_n > 2,000) = 0.95$, the claim made would draw support.

- 2.22** (i) Consider the i.i.d. r.v.'s X_1, \dots, X_n and Y_1, \dots, Y_n with expectation μ and variance σ^2 , both finite, and let \bar{X}_n and \bar{Y}_n be the respective sample means. Use the CLT in order to determine the sample size n , so that $P(|\bar{X}_n - \bar{Y}_n| \leq 0.25\sigma) = 0.95$.
- (ii) Let the random samples X_1, \dots, X_n and Y_1, \dots, Y_n be as in part (i), but we do not assume that they are coming from the same distribution. We do assume, however, that they have the same mean and the same variance σ^2 , both finite. Then determine n as required above by using the Tchebichev inequality.

Hint: Set $Z_i = X_i - Y_i$ and then work as in Exercise 2.20(ii) with the i.i.d. r.v.'s Z_1, \dots, Z_n . Finally, revert to the X_i 's and the Y_i 's.

- 2.23** Let $X_i, i = 1, \dots, n, Y_i, i = 1, \dots, n$ be independent r.v.'s such that the X_i 's are identically distributed with $EX_i = \mu_1, \text{Var}(X_i) = \sigma^2$, both finite, and the Y_i 's are identically distributed with $EY_i = \mu_2$ and $\text{Var}(Y_i) = \sigma^2$, both finite. If \bar{X}_n and \bar{Y}_n are the respective sample means of the X_i 's and the Y_i 's, then:

- (i) Show that $E(\bar{X}_n - \bar{Y}_n) = \mu_1 - \mu_2, \text{Var}(\bar{X}_n - \bar{Y}_n) = \frac{2\sigma^2}{n}$.
- (ii) Use the CLT in order to show that $\frac{\sqrt{n}[(\bar{X}_n - \bar{Y}_n) - (\mu_1 - \mu_2)]}{\sigma\sqrt{2}}$ is asymptotically distributed as $N(0, 1)$.

Hint: Set $Z_i = X_i - Y_i$ and work with the i.i.d. r.v.'s Z_1, \dots, Z_n ; then revert to the X_i 's and the Y_i 's.

- 2.24** An academic department in a university wishes to admit 20 first-year graduate students. From past experience, it follows that, on the average, 40% of the students admitted will, actually, accept the admission offer. It may be assumed that acceptance and rejection of admission offers by the various students are independent events, and let Y_n be the r.v. denoting the number of those students, actually, accepting admission.

- (i) Use the CLT in order to determine n , so that the probability $P(|Y_n - 20| \leq 2)$ is maximum. Use a rough geometric argument.
- (ii) Compute the (approximate) probability $P(|Y_n - 20| \leq 2)$ once n is determined.
- (iii) Derive in a more rigorous way a relation through which n is to be determined subject to the maximization of the probability $P(|Y_n - 20| \leq 2)$.

Hint: With each one of the n students offered admission, associate a r.v. X_i , which takes on the value 1, if the i th student accepts the offer, and 0 otherwise. Then the r.v.'s X_1, \dots, X_n are independent, and also assume that they have the same distribution; i.e., $P(X_i = 1) = p (= 0.40 \text{ here})$ for all i 's. Then the X_i 's are distributed as $B(1, p)$ and $Y_n = \sum_{i=1}^n X_i$. Then, for part (i), draw the $N(0, 1)$ p.d.f. curve, and by symmetry and geometric considerations conclude that seemingly (but not precisely) the required probability is maximized for the value of n for which $18 - np = -(22 - np)$. (Do not use any continuity correction.)

For part (iii), pretend that n is a continuous variable, differentiate with respect to it, and equate to 0 in order to arrive at the following relationship, after some cancellations and modifications:

$$\frac{80 - 1.6n}{0.24n} = \log_8 \frac{0.4n + 22}{0.4n + 18}.$$

Remark: The problem may also be posed by replacing 20, 2, and 40% by c , d , and $100p\%$, say.

7.3 Further Limit Theorems

Convergence in probability enjoys some of the familiar properties of the usual pointwise convergence. One such property is stated below in the form of a theorem whose proof is omitted.

THEOREM 5

- (i) For $n \geq 1$, let X_n and X be r.v.'s such that $X_n \xrightarrow[n \rightarrow \infty]{P} X$, and let g be a continuous real-valued function; that is, $g : \mathfrak{R} \rightarrow \mathfrak{R}$ continuous. Then the r.v.'s $g(X_n)$, $n \geq 1$, also converge in probability to $g(X)$; that is, $g(X_n) \xrightarrow[n \rightarrow \infty]{P} g(X)$. More generally:
- (ii) For $n \geq 1$, let X_n, Y_n, X , and Y be r.v.'s such that $X_n \xrightarrow[n \rightarrow \infty]{P} X$, $Y_n \xrightarrow[n \rightarrow \infty]{P} Y$, and let g be a continuous real-valued function; that is, $g : \mathfrak{R}^2 \rightarrow \mathfrak{R}$ continuous. Then the r.v.'s $g(X_n, Y_n)$, $n \geq 1$, also converge in probability to $g(X, Y)$; that is, $g(X_n, Y_n) \xrightarrow[n \rightarrow \infty]{P} g(X, Y)$. (This part

also generalizes in an obvious manner to k sequences $\{X_n^{(i)}\}$,
 $n \geq 1$, $i = 1, \dots, k$.)

To this theorem, there is the following important corollary.

COROLLARY If $X_n \xrightarrow[n \rightarrow \infty]{P} X$ and $Y_n \xrightarrow[n \rightarrow \infty]{P} Y$, then:

- (i) $aX_n + bY_n \xrightarrow[n \rightarrow \infty]{P} aX + bY$, where a and b are constants; and, in particular,
 $X_n + Y_n \xrightarrow[n \rightarrow \infty]{P} X + Y$.
- (ii) $X_n Y_n \xrightarrow[n \rightarrow \infty]{P} XY$.
- (iii) $\frac{X_n}{Y_n} \xrightarrow[n \rightarrow \infty]{P} \frac{X}{Y}$, provided $P(Y_n \neq 0) = P(Y \neq 0) = 1$.

PROOF Although the proof of the theorem was omitted, the corollary can be proved. Indeed, all one has to do is to take: $g : \Re^2 \rightarrow \Re$ as follows, respectively, for parts (i)–(iii) and observe that it is continuous: $g(x, y) = ax + by$ (and, in particular, $g(x, y) = x + y$); $g(x, y) = xy$; $g(x, y) = x/y$, $y \neq 0$. ▲

Actually, a special case of the preceding corollary also holds for convergence in distribution. Specifically, we have

THEOREM 6

(Slutsky) Let $X_n \xrightarrow[n \rightarrow \infty]{d} X$ and let $Y_n \xrightarrow[n \rightarrow \infty]{d} c$, a constant c rather than a (proper) r.v. Y . Then:

- (i) $X_n + Y_n \xrightarrow[n \rightarrow \infty]{d} X + c$; (ii) $X_n Y_n \xrightarrow[n \rightarrow \infty]{d} cX$; (iii) $\frac{X_n}{Y_n} \xrightarrow[n \rightarrow \infty]{d} \frac{X}{c}$, provided $P(Y_n \neq 0) = 1$ and $c \neq 0$.

In terms of d.f.'s, these convergences are written as follows, always as $n \rightarrow \infty$ and for all $z \in \Re$ for which: $z - c$ is a continuity point of F_X for part (i); z/c is a continuity point of F_X for part (ii); cz is a continuity point of F_X for part (iii):

$$P(X_n + Y_n \leq z) \rightarrow P(X + c \leq z) = P(X \leq z - c), \quad \text{or}$$

$$F_{X_n + Y_n}(z) \rightarrow F_X(z - c);$$

$$P(X_n Y_n \leq z) \rightarrow P(cX \leq z) = \begin{cases} P(X \leq \frac{z}{c}), & c > 0 \\ P(X \geq \frac{z}{c}), & c < 0 \end{cases}, \quad \text{or}$$

$$F_{X_n Y_n}(z) \rightarrow \begin{cases} F_X(\frac{z}{c}), & c > 0 \\ 1 - P(X < \frac{z}{c}) (= 1 - F_X(\frac{z}{c}), \text{ if } F_X \text{ is continuous}), & c < 0 \end{cases};$$

$$P\left(\frac{X_n}{Y_n} \leq z\right) \rightarrow P\left(\frac{X}{c} \leq z\right) = \begin{cases} P(X \leq cz), & c > 0 \\ P(X \geq cz), & c < 0 \end{cases}, \quad \text{or}$$

$$F_{\frac{X_n}{Y_n}}(z) \rightarrow \begin{cases} F_X(cz), & c > 0 \\ 1 - P(X < cz) (= 1 - F_X(cz), \text{ if } F_X \text{ is continuous}), & c < 0. \end{cases}$$

The proof of this theorem, although conceptually not complicated, is, nevertheless, long and is omitted. Recall, however, that $Y_n \xrightarrow[n \rightarrow \infty]{d} c$ if and only if $Y_n \xrightarrow[n \rightarrow \infty]{P} c$, and this is another way the convergence of Y_n is stated.

As a simple concrete application of Theorem 5, consider the following example.

EXAMPLE 7

Suppose $X_n \xrightarrow{d} X \sim N(\mu, \sigma^2)$, and let c_n, c, d_n , and d be constants such that $c_n \rightarrow c$ and $d_n \rightarrow d$. Then $c_n X_n + d_n \xrightarrow{d} Y \sim N(c\mu + d, c^2\sigma^2)$.

DISCUSSION Trivially, $c_n \xrightarrow{d} c$ and $d_n \xrightarrow{d} d$, so that, by Theorem 6(ii), $c_n X_n \xrightarrow{d} cX$, and by Theorem 6(i), $c_n X_n + d_n \xrightarrow{d} cX + d$. However, $X \sim N(\mu, \sigma^2)$ implies that $cX + d \sim N(c\mu + d, c^2\sigma^2)$. Thus, $c_n X_n + d_n \xrightarrow{d} cX + d = Y \sim N(c\mu + d, c^2\sigma^2)$.

The following result is an application of Theorems 5 and 6 and is of much use in statistical inference. For its formulation, let X_1, \dots, X_n be i.i.d. r.v.'s with finite mean μ and finite and positive variance σ^2 , and let \bar{X}_n and S_n^2 be the sample mean and the "adjusted" (in the sense that μ is replaced by \bar{X}_n) sample variance (which we have denoted by \bar{S}_n^2 in relation (13) of Chapter 5); that is, $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$, $S_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2$.

THEOREM 7

Under the assumptions just made and the notation introduced, it holds:

$$(i) S_n^2 \xrightarrow[n \rightarrow \infty]{P} \sigma^2; (ii) \frac{\sqrt{n}(\bar{X}_n - \mu)}{S_n} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1).$$

PROOF (i) Recall that $\sum_{i=1}^n (X_i - \bar{X}_n)^2 = \sum_{i=1}^n X_i^2 - n\bar{X}_n^2$, so that $S_n^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}_n^2$. Since $EX_i^2 = \text{Var}(X_i) + (EX_i)^2 = \sigma^2 + \mu^2$, the WLLN applies to the i.i.d. r.v.'s X_1^2, \dots, X_n^2 and gives: $\frac{1}{n} \sum_{i=1}^n X_i^2 \xrightarrow[n \rightarrow \infty]{P} \sigma^2 + \mu^2$. Also, $\bar{X}_n \xrightarrow[n \rightarrow \infty]{P} \mu$, by the WLLN again, and then $\bar{X}_n^2 \xrightarrow[n \rightarrow \infty]{P} \mu^2$ by Theorem 5(i). Then, by Theorem 5(ii),

$$\frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}_n^2 \xrightarrow[n \rightarrow \infty]{P} (\sigma^2 + \mu^2) - \mu^2 = \sigma^2,$$

which is what part (i) asserts.

(ii) Part (i) and Theorem 5(i) imply that $S_n \xrightarrow[n \rightarrow \infty]{P} \sigma$, or $\frac{S_n}{\sigma} \xrightarrow[n \rightarrow \infty]{P} 1$. By Theorem 4, $\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1)$. Then Theorem 6(iii) applies and

gives:

$$\frac{\sqrt{n}(\bar{X}_n - \mu)/\sigma}{S_n/\sigma} = \frac{\sqrt{n}(\bar{X}_n - \mu)}{S_n} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1). \quad \blacktriangle$$

REMARK 6 Part (ii) of the theorem states, in effect, that for sufficiently large n , σ may be replaced in the CLT by the adjusted sample standard deviation S_n and the resulting expression still has a distribution which is close to the $N(0, 1)$ distribution.

The WLLN states that $\bar{X}_n \xrightarrow[n \rightarrow \infty]{d} \mu$, which, for a real-valued continuous function g , implies that

$$g(\bar{X}_n) \xrightarrow[n \rightarrow \infty]{P} g(\mu). \quad (14)$$

On the other hand, the CLT states that:

$$\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow[n \rightarrow \infty]{d} N(0, 1) \quad \text{or} \quad \sqrt{n}(\bar{X}_n - \mu) \xrightarrow[n \rightarrow \infty]{d} N(0, \sigma^2).$$

The question then arises what happens to the distribution of $g(\bar{X}_n)$. In other words, is there a result analogous to (14) when the distribution of $g(\bar{X}_n)$ is involved? The question is answered by the following result.

THEOREM 8

Let X_1, \dots, X_n be i.i.d. r.v.'s with finite mean μ and variance $\sigma^2 \in (0, \infty)$, and let $g : \Re \rightarrow \Re$ be differentiable with derivative g' continuous at μ . Then:

$$\sqrt{n}[g(\bar{X}_n) - g(\mu)] \xrightarrow[n \rightarrow \infty]{d} N(0, [\sigma g'(\mu)]^2). \quad (15)$$

The proof of this result involves the employment of some of the theorems established in this chapter, including the CLT, along with a Taylor expansion. The proof itself will not be presented, and this section will be concluded with an application to Theorem 8. The method of establishing asymptotic normality for $g(\bar{X}_n)$ is often referred to as the *delta method*, and it also applies in cases more general than the one described here.

APPLICATION Let the independent r.v.'s X_1, \dots, X_n be distributed as $B(1, p)$. Then:

$$\sqrt{n}[\bar{X}_n(1 - \bar{X}_n) - pq] \xrightarrow[n \rightarrow \infty]{d} N(0, pq(1 - 2p)^2) \quad (q = 1 - p). \quad (16)$$

PROOF Here $\mu = p$, $\sigma^2 = pq$, and $g(x) = x(1 - x)$, so that $g'(x) = 1 - 2x$ continuous for all x . Since $g(\bar{X}_n) = \bar{X}_n(1 - \bar{X}_n)$, $g(\mu) = p(1 - p) = pq$, and $g'(\mu) = 1 - 2p$, the convergence in (15) becomes as stated in (16). \blacktriangle

Exercise

- 3.1** Let X_1, \dots, X_n be i.i.d. r.v.'s with finite $EX_i = \mu$, and $\text{Var}(X_i) = \sigma^2 \in (0, \infty)$ so that the CLT holds; that is,

$$\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1), \quad \text{where } \bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i.$$

Then use Theorem 6 in order to show that the WLLN also holds.



Chapter 8

An Overview of Statistical Inference

A review of the previous chapters reveals that the main objectives throughout have been those of calculating probabilities or certain summary characteristics of a distribution, such as mean, variance, median, and mode. However, for these calculations to result in numerical answers, it is a prerequisite that the underlying distribution be completely known. Typically, this is rarely, if ever, the case. The reason for this is that the parameters which appear, for example, in the functional form of the p.d.f. of a distribution are simply unknown to us. The only thing known about them is that they lie in specified sets of possible values for these parameters, the *parameter space*.

It is at this point where statistical inference enters the picture. Roughly speaking, the aim of statistical inference is to make certain determinations with regard to the unknown constants (*parameters*) figuring in the underlying distribution. This is to be done on the basis of data, represented by the observed values of a random sample drawn from said distribution. Actually, this is the so-called *parametric statistical inference* as opposed to the *nonparametric statistical inference*. The former is applicable to distributions, which are completely determined by the knowledge of a finite number of parameters. The latter applies to distributions not determined by any finite number of parameters.

The remaining part of this book is, essentially, concerned with statistical inference and mostly with parametric statistical inference. Within the framework of parametric statistical inference, there are three main objectives, depending on what kind of determinations we wish to make with regard to the parameters. If the objective is to arrive at a number, by means of the available data, as the value of an unknown parameter, then we are talking about *point estimation*. If, on the other hand, we are satisfied with the statement that an unknown parameter lies within a known random interval (that is, an interval with r.v.'s as its end-points) with high prescribed probability, then we

are dealing with *interval estimation* or *confidence intervals*. Finally, if the objective is to decide that an unknown parameter lies in a specified subset of the parameter space, then we are in the area of *testing hypotheses*.

These three subjects — point estimation, interval estimation, and testing hypotheses — are briefly discussed in the following three sections. In the subsequent three sections, it is pointed out what the statistical inference issues are in specific models — a *regression model* and two *analysis of variance models*. The final section touches upon some aspects of *nonparametric statistical inference*.

8.1 The Basics of Point Estimation

The problem here, briefly stated, is as follows. Let X be a r.v. with a p.d.f. f which, however, involves a parameter. This is the case, for instance, in the Binomial distribution $B(1, p)$, the Poisson distribution $P(\lambda)$, the Negative Exponential $f(x) = \lambda e^{-\lambda x}$, $x > 0$ distribution, the Uniform distribution $U(0, \alpha)$, and the Normal distribution $N(\mu, \sigma^2)$ with one of the quantities μ and σ^2 known. The *parameter* is usually denoted by θ , and the set of its possible values is denoted by Ω and is called the *parameter space*. In order to emphasize the fact that the p.d.f. depends on θ , we write $f(\cdot; \theta)$. Thus, in the distributions mentioned above, we have for the respective p.d.f.'s and the parameter spaces:

$$f(x; \theta) = \theta^x(1 - \theta)^{1-x}, \quad x = 0, 1, \quad \theta \in \Omega = (0, 1).$$

The situations described in Examples 5, 6, 8, 9, and 10 of Chapter 1 may be described by a Binomial distribution.

$$f(x; \theta) = \frac{e^{-\theta} \theta^x}{x!}, \quad x = 0, 1, \dots, \quad \theta \in \Omega = (0, \infty).$$

The Poisson distribution can be used appropriately in the case described in Example 12 of Chapter 1.

$$f(x; \theta) = \theta e^{-\theta x}, \quad x > 0, \quad \theta \in \Omega = (0, \infty).$$

$$f(x; \theta) = \begin{cases} \frac{1}{\theta}, & 0 < x < \theta \\ 0, & \text{otherwise,} \end{cases} \quad \theta \in \Omega = (0, \infty).$$

$$f(x; \theta) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\theta)^2}{2\sigma^2}}, \quad x \in \mathfrak{R}, \quad \theta \in \Omega = \mathfrak{R}, \quad \sigma^2 \text{ known,}$$

and

$$f(x; \theta) = \frac{1}{\sqrt{2\pi}\theta} e^{-\frac{(x-\mu)^2}{2\theta}}, \quad x \in \mathfrak{R}, \quad \theta \in \Omega = (0, \infty), \quad \mu \text{ known.}$$

Normal distributions are suitable for modeling the situations described in Examples 16 and 17 of Chapter 1.

Our objective is to draw a random sample of size n , X_1, \dots, X_n , from the underlying distribution, and on the basis of it to construct a *point estimate* (or *estimator*) for θ , that is, a statistic $\hat{\theta} = \hat{\theta}(X_1, \dots, X_n)$, which is used for

estimating θ , where a *statistic* is a known function of the random sample X_1, \dots, X_n . If x_1, \dots, x_n are the actually observed values of the r.v.'s X_1, \dots, X_n , respectively, then the observed value of our estimate has the numerical value $\hat{\theta}(x_1, \dots, x_n)$. The observed values x_1, \dots, x_n are also referred to as *data*. Then, on the basis of the available data, it is declared that the value of θ is $\hat{\theta}(x_1, \dots, x_n)$ from among all possible points in Ω . A point estimate is often referred to just as an estimate, and the notation $\hat{\theta}$ is used indiscriminately, both for the estimate $\hat{\theta}(X_1, \dots, X_n)$ (which is a r.v.) and for its observed value $\hat{\theta}(x_1, \dots, x_n)$ (which is just a number).

The only obvious restriction on $\hat{\theta}(x_1, \dots, x_n)$ is that it lies in Ω for all possible values of X_1, \dots, X_n . Apart from it, there is any number of estimates one may construct — thus, the need to assume certain principles and/or invent methods for constructing $\hat{\theta}$. Perhaps, the most widely accepted principle is the so-called principle of *Maximum Likelihood* (ML). This principle dictates that we form the joint p.d.f. of the x_i 's, for the observed values of the X_i 's, look at this joint p.d.f. as a function of θ (and call it the *likelihood function*), and maximize the likelihood function with respect to θ . The maximizing point (assuming it exists and is unique) is a function of x_1, \dots, x_n , and is what we call the *Maximum Likelihood Estimate* (MLE) of θ . The notation used for the likelihood function is $L(\theta | x_1, \dots, x_n)$. Then, we have that:

$$L(\theta | x_1, \dots, x_n) = f(x_1; \theta) \cdots f(x_n; \theta), \quad \theta \in \Omega.$$

The MLE will be studied fairly extensively in Chapter 9.

Another principle often used in constructing an estimate for θ is the principle of *unbiasedness*. In this context, an estimate is usually denoted by $U = U(X_1, \dots, X_n)$. Then the principle of unbiasedness dictates that U should be constructed so as to be *unbiased*; that is, its expectation (mean value) should always be θ , no matter what the value of θ in Ω . More formally, $E_\theta U = \theta$ for all $\theta \in \Omega$. (In the expectation sign E , the parameter θ was inserted to indicate that this expectation does depend on θ , since it is calculated by using the p.d.f. $f(\cdot; \theta)$.) Now, it is intuitively clear that, in comparing two unbiased estimates, one would pick the one with the smaller variance, since it would be more closely concentrated around its mean θ . Envision the case that, within the class of all unbiased estimates, there exists one which has the smallest variance (and that is true for all $\theta \in \Omega$). Such an estimate is called a *Uniformly Minimum Variance Unbiased* (UMVU) estimate and is, clearly, a desirable estimate. In the next chapter, we will see how we go about constructing such estimates.

The principle (or rather the method) based on sample moments is another way of constructing estimates. The *method of moments*, in the simplest case, dictates to form the sample mean \bar{X} and equate it with the (theoretical) mean $E_\theta X$. Then solve for θ (assuming it can be done, and, indeed, uniquely) in order to arrive at a *moment estimate* of θ .

A much more sophisticated method of constructing estimates of θ is the so-called *decision-theoretic* method. This method calls for the introduction of a host of concepts, terminology, and notation, and it will be taken up in the next chapter.

Finally, another relatively popular method (in particular, in the context of certain models) is the method of *Least Squares* (LS). The method of LS leads to the construction of an estimate for θ , the *Least Squares Estimate* (LSE) of θ , through a minimization (with respect to θ) of the sum of certain squares. This sum of squares represents squared deviations between what we actually observe after experimentation is completed and what we would expect to have on the basis of an assumed model. Once again, details will be presented later on, more specifically, in Chapter 13.

In all of the preceding discussion, it was assumed that the underlying p.d.f. depended on a single parameter, which was denoted by θ . It may very well be the case that there are two or more parameters involved. This may happen, for instance, in the Uniform distribution $U(\alpha, \beta)$, $-\infty < \alpha < \beta < \infty$, where both α and β are unknown; the Normal distribution, $N(\mu, \sigma^2)$, where both μ and σ^2 are unknown; and it does happen in the Multinomial distribution, where the number of parameters is k , p_1, \dots, p_k (or more precisely, $k - 1$, since the k th parameter, for example, $p_k = 1 - p_1 - \dots - p_{k-1}$). For instance, Examples 20 and 21 of Chapter 1 refer to situations where a Multinomial distribution is appropriate. In such multiparameter cases, one simply applies to each parameter separately what was said above for a single parameter. The alternative option to use the vector notation for the parameters involved does simplify things in a certain way but also introduces some complications in other ways.

8.2 The Basics of Interval Estimation

Suppose we are interested in constructing a point estimate of the mean μ in the Normal distribution $N(\mu, \sigma^2)$ with known variance; this is to be done on the basis of a random sample of size n , X_1, \dots, X_n , drawn from the underlying distribution. This amounts to constructing a suitable statistic of the X_i 's, call it $V = V(X_1, \dots, X_n)$, which for the observed values x_i of X_i , $i = 1, \dots, n$ is a numerical entity, and declare it to be the (unknown) value of μ . This looks somewhat presumptuous, since from the set of possible values for μ , $-\infty < \mu < \infty$, just one is selected as its value. Thinking along these lines, it might be more reasonable to aim instead at a random interval which will contain the (unknown) value of μ with high (prescribed) probability. This is exactly what a confidence interval does.

To be more precise and in casting the problem in a general setting, let X_1, \dots, X_n be a random sample from the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \Re$, and let $L = L(X_1, \dots, X_n)$ and $U = U(X_1, \dots, X_n)$ be two statistics of the X_i 's such that $L < U$. Then the interval with end-points L and U , $[L, U]$, is called a *random interval*. Let α be a small number in $(0, 1)$, such as 0.005, 0.01, 0.05, and suppose that the random interval $[L, U]$ contains θ with probability equal to $1 - \alpha$ (such as 0.995, 0.99, 0.95) no matter what the true value of θ in Ω is. In other words, suppose that:

$$P_\theta(L \leq \theta \leq U) = 1 - \alpha \quad \text{for all } \theta \in \Omega. \quad (1)$$

If relation (1) holds, then we say that the random interval $[L, U]$ is a *confidence interval* for θ with *confidence coefficient* $1 - \alpha$.

The interpretation of the significance of a confidence interval is based on the relative frequency interpretation of the concept of probability, and it goes like this: Suppose n independent r.v.'s are drawn from the p.d.f. $f(\cdot; \theta)$, and let x_1, \dots, x_n be their observed values. Also, let $[L_1, U_1]$ be the interval resulting from the observed values of $L = L(X_1, \dots, X_n)$ and $U = U(X_1, \dots, X_n)$; that is, $L_1 = L(x_1, \dots, x_n)$ and $U_1 = U(x_1, \dots, x_n)$. Proceed to draw independently a second set of n r.v.'s as above, and let $[L_2, U_2]$ be the resulting interval. Repeat this process independently a large number of times, N , say, with the corresponding interval being $[L_N, U_N]$. Then the interpretation of (1) is that, on the average, about $100(1 - \alpha)\%$ of the above N intervals will, actually, contain the true value of θ . For example, for $\alpha = 0.05$ and $N = 1,000$, the proportion of such intervals will be 95%; that is, one would expect 950 out of the 1,000 intervals constructed as above to contain the true value of θ . Empirical evidence shows that such an expectation is valid.

We may also define an *upper confidence limit* for θ , $U = U(X_1, \dots, X_n)$, and a *lower confidence limit* for θ , $L = L(X_1, \dots, X_n)$, both with *confidence coefficient* $1 - \alpha$, if, respectively, the intervals $(-\infty, U]$ and $[L, \infty)$ are confidence intervals for θ with confidence coefficient $1 - \alpha$. That is to say:

$$P_\theta(-\infty < \theta \leq U) = 1 - \alpha, \quad P_\theta(L \leq \theta < \infty) = 1 - \alpha \quad \text{for all } \theta \in \Omega. \quad (2)$$

Confidence intervals and upper and/or lower confidence limits can be sought, for instance, in Examples 5, 6, 8, 9, and 10 (Binomial distribution), 12 (Poisson distribution), and 16 and 17 (Normal distribution) in Chapter 1.

There are some variations of (1) and (2). For example, when the underlying p.d.f. is discrete, then equalities in (1) and (2) rarely obtain for given α and have to be replaced by inequalities \geq . Also, except for special cases, equalities in (1) and (2) are valid only approximately for large values of the sample size n (even in cases where the underlying r.v.'s are continuous). In such cases, we say that the respective confidence intervals (confidence limits) have *confidence coefficient approximately* $1 - \alpha$.

Finally, the parameters of interest may be two (or more) rather than one, as we assumed so far. In such cases, the concept of a confidence interval is replaced by that of a *confidence region* (in the multidimensional parameter space Ω). This concept will be illustrated by an example in Chapter 10. In the same chapter, we will also expand considerably on what was briefly discussed here.

8.3 The Basics of Testing Hypotheses

Often, we are not interested in a point estimate of a parameter θ or even a confidence interval for it, but rather whether said parameter lies or does not lie in a specified subset ω of the parameter space Ω . To clarify this point, we refer to some of the examples described in Chapter 1. Thus, in Example 5, all

we might be interested in is whether Jones has ESP at all or not and not to what degree he does. In statistical terms, this amounts to taking n independent observations from a $B(1, \theta)$ distribution and, on the basis of these observations, deciding whether $\theta \in \omega = (0, 0.5]$ (as opposed to $\theta \in \omega^c = (0.5, 1)$); here θ is the probability that Jones correctly identifies the picture. The situation in Example 6 is similar, and the objective might be to decide whether or not $\theta \in \omega = (\theta_0, 1)$; here θ is the true proportion of unemployed workers and θ_0 is a certain desirable or guessed value of θ . Examples 8, 9, and 10 in Chapter 1 fall into the same category.

In Example 12, the stipulated model is a Poisson distribution $P(\theta)$ and, on the basis of n independent observations, we might wish to decide whether or not $\theta \in (\theta_0, \infty)$, where θ_0 is a known value of θ .

In Example 16, the stipulated underlying models may be Normal distributions $N(\mu_1, \sigma^2)$ and $N(\mu_2, \sigma^2)$ for the survival times X and Y , respectively, and then the question of interest may be to decide whether or not $\mu_2 \leq \mu_1$; σ^2 may be assumed to be either known or unknown. Of course, we are going to arrive at the desirable decision on the basis of two independent random samples drawn from the underlying distributions. Example 17 is of the same type.

In Example 20, the statistical problem is that of comparing two Multinomial populations, by making appropriate statements about the probabilities p_{AE}, p_{AA}, p_{AP} and p_{BE}, p_{BA}, p_{BP} ; here p_{AE} is the probability that any one of the 80 infants, subjected to diet A , is of "excellent" health, and similarly for the remaining probabilities. Example 21 is of a similar type.

On the basis of the preceding discussion and examples, we may now proceed with the formulation of the general problem. To this effect, let X_1, \dots, X_n be i.i.d. r.v.'s with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}^r$, $r \geq 1$, and by means of this random sample, suppose we are interested in checking whether $\theta \in \omega$, a proper subset of Ω , or $\theta \in \omega^c$, the complement of ω with respect to Ω . The statements that $\theta \in \omega$ and $\theta \in \omega^c$ are called (*statistical*) *hypotheses* (about θ), and are denoted thus: $H_0 : \theta \in \omega$, $H_A : \theta \in \omega^c$. The hypothesis H_0 is called a *null hypothesis* and the hypothesis H_A is called *alternative* (to H_0) hypothesis. The hypotheses H_0 and H_A are called *simple*, if they contain a single point, and *composite* otherwise. The procedure of checking whether H_0 is true or not, on the basis of the observed values x_1, \dots, x_n of X_1, \dots, X_n , is called *testing* the hypothesis H_0 against the alternative H_A .

In the special case that $\Omega \subseteq \mathfrak{R}$, some null hypotheses and the respective alternatives are as follows:

$$\begin{aligned} H_0 : \theta = \theta_0 \text{ against } H_A : \theta > \theta_0; & \quad H_0 : \theta = \theta_0 \text{ against } H_A : \theta < \theta_0; \\ H_0 : \theta \leq \theta_0 \text{ against } H_A : \theta > \theta_0; & \quad H_0 : \theta \geq \theta_0 \text{ against } H_A : \theta < \theta_0; \\ H_0 : \theta = \theta_0 \text{ against } H_A : \theta \neq \theta_0. & \end{aligned}$$

The testing is carried out by means of a function $\varphi : \mathfrak{R}^n \rightarrow [0, 1]$ which is called a *test function* or just a *test*. The number $\varphi(x_1, \dots, x_n)$ represents the probability of rejecting H_0 , given that $X_i = x_i$, $i = 1, \dots, n$. In its simplest form,

φ is the indicator of a set B in \mathfrak{N}^n , which is called the *critical* or *rejection region*; its complement B^c is called the *acceptance region*. Thus, $\varphi(x_1, \dots, x_n) = 1$ if x_1, \dots, x_n are in B , and $\varphi(x_1, \dots, x_n) = 0$, otherwise. Actually, such a test is called a *nonrandomized* test as opposed to tests which also take values strictly between 0 and 1 and are called *randomized* tests. In the case of continuous distributions, nonrandomized tests suffice, but in discrete distributions, a test will typically be required to take on one or two values strictly between 0 and 1.

By using a test φ , suppose that our data x_1, \dots, x_n lead us to the rejection of H_0 . This will happen, for instance, if the test φ is nonrandomized with rejection region B , and the x_i 's lie in B . By rejecting the hypothesis H_0 , we may be doing the correct thing, because H_0 is false (that is, $\theta \notin \omega$). On the other hand, we may be taking the wrong action because it may happen that H_0 is, indeed, true (that is, $\theta \in \omega$), only the test and the data do not reveal it. Clearly, in so doing, we commit an error, which is referred to as *type I error*. Of course, we would like to find ways of minimizing the frequency of committing this error. To put it more mathematically, this means searching for a rejection region B , which will minimize the above frequency. In our framework, frequencies are measured by probabilities, and this leads to a determination of B so that

$$\begin{aligned} P(\text{of type I error}) &= P(\text{of rejecting } H_0 \text{ whereas } H_0 \text{ is true}) \\ &= P_\theta(X_1, \dots, X_n \text{ lie in } B \text{ whereas } \theta \in \omega) \\ &= P_\theta(X_1, \dots, X_n \text{ lie in } B \mid \theta \in \omega) \stackrel{\text{def}}{=} \alpha(\theta) \text{ is minimum.} \quad (3) \end{aligned}$$

Clearly, the probabilities $\alpha(\theta)$ in (3) must be minimized for each $\theta \in \omega$, since we don't know which value in ω is the true θ . This will happen if we minimize the $\max_{\theta \in \omega} \alpha(\theta) \stackrel{\text{def}}{=} \alpha$. This maximum probability of type I error is called the *level of significance* of the test employed. Thus, we are led to selecting the rejection region B so that its level of significance α will be minimum. Since $\alpha \geq 0$, its minimum value would be 0, and this would happen if (essentially) $B = \emptyset$. But then (essentially) the x_i 's would always be in $B^c = \mathfrak{N}^n$, and this would happen with probability

$$P_\theta(X_1, \dots, X_n \text{ in } \mathfrak{N}^n) = 1 \quad \text{for all } \theta. \quad (4)$$

This, however, creates a problem for the following reason. If the rejection region B is \emptyset , then the acceptance region is \mathfrak{N}^n ; that is, we always accept H_0 . As long as H_0 is true (that is, $\theta \in \omega$), this is exactly what we wish to do, but what about the case that H_0 is false (that is, $\theta \in \omega^c$)? When we accept a false hypothesis H_0 , we commit an error, which is called the *type II error*. As in (3), this error is also measured in terms of probabilities; namely,

$$\begin{aligned} P(\text{of type II error}) &= P(\text{of accepting } H_0 \text{ whereas } H_0 \text{ is false}) \\ &= P_\theta(X_1, \dots, X_n \text{ lie in } B^c \text{ whereas } \theta \in \omega^c) \\ &= P_\theta(X_1, \dots, X_n \text{ lie in } B^c \mid \theta \in \omega^c) \\ &\stackrel{\text{def}}{=} \beta(\theta). \quad (5) \end{aligned}$$

According to (5), these probabilities would be 1 for all $\theta \in \omega^c$ (actually, for all $\theta \in \Omega$), if $B = \emptyset$. Clearly, this is undesirable. The preceding discussion then leads to the conclusion that the rejection region B must be different from \emptyset and then α will be > 0 . The objective then becomes that of choosing B so that α will have a preassigned acceptable value (such as 0.005, 0.01, 0.05) and, subject to this restriction, the probabilities of type II error are minimized. That is,

$$\beta(\theta) = P_{\theta}(X_1, \dots, X_n \text{ lie in } B^c) \text{ is minimum for each } \theta \in \omega^c. \quad (6)$$

Since $P_{\theta}(X_1, \dots, X_n \text{ lie in } B^c) = 1 - P_{\theta}(X_1, \dots, X_n \text{ lie in } B)$, the minimization in (6) is equivalent to the maximization of

$$P_{\theta}(X_1, \dots, X_n \text{ lie in } B) = 1 - P_{\theta}(X_1, \dots, X_n \text{ lie in } B^c) \quad \text{for all } \theta \in \omega^c.$$

The function $\pi(\theta)$, $\theta \in \omega^c$, defined by:

$$\pi(\theta) = P_{\theta}(X_1, \dots, X_n \text{ lie in } B), \quad \theta \in \omega^c, \quad (7)$$

is called the *power* of the test employed. So, power of a test = 1 – probability of a type II error, and we may summarize our objective as follows: Choose a test with a preassigned level of significance α , which has maximum power among all tests with level of significance $\leq \alpha$. In other words, if φ is the desirable test, then it should satisfy the requirements:

The level of significance of φ is α , and its power, to be denoted by $\pi_{\varphi}(\theta)$, $\theta \in \omega^c$, satisfies the inequality $\pi_{\varphi}(\theta) \geq \pi_{\varphi^*}(\theta)$ for all $\theta \in \omega^c$ and any test φ^* with level of significance $\leq \alpha$.

Such a test φ , should it exist, is called *Uniformly Most Powerful* (UMP) for obvious reasons. (The term “most powerful” is explained by the inequality $\pi_{\varphi}(\theta) \geq \pi_{\varphi^*}(\theta)$, and the term “uniformly” is due to the fact that this inequality must hold for *all* $\theta \in \omega^c$.) If ω^c consists of a single point, then the concept of uniformity is void, and we talk simply of a *Most Powerful* (MP) test.

The concepts introduced so far hold for a parameter of any dimensionality. However, UMP tests can be constructed only when θ is a real-valued parameter, and then only for certain forms of H_0 and H_A and specific p.d.f.'s $f(\cdot; \theta)$. If the parameter is multidimensional, desirable tests can still be constructed; they are not going to be, in general, UMP tests, but they are derived, nevertheless, on the basis of principles which are intuitively satisfactory. Preeminent among such tests are the so-called *Likelihood Ratio* (LR) tests. Another class of tests are the so-called *goodness-of-fit* tests, and still others are constructed on the basis of *decision-theoretic* concepts. Some of the tests mentioned above will be discussed more extensively in Chapters 11 and 12. Here, we conclude this section with the introduction of a LR test.

On the basis of the random sample X_1, \dots, X_n with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}^r$, $r \geq 1$, suppose we wish to test the hypothesis $H_0 : \theta \in \omega$ (a proper) subset of Ω . It is understood that the alternative is $H_A : \theta \in \omega^c$, but in the present framework it is not explicitly stated. Let x_1, \dots, x_n be the observed values of X_1, \dots, X_n and form the likelihood function $L(\theta) = L(\theta | x_1, \dots, x_n) =$

$\prod_{i=1}^n f(x_i; \theta)$. Maximize $L(\theta)$ and denote the resulting maximum by $L(\hat{\Omega})$. This maximization happens when θ is equal to the MLE $\hat{\theta} = \hat{\theta}(x_1, \dots, x_n)$, so that $L(\hat{\Omega}) = L(\hat{\theta})$. Next, maximize the likelihood $L(\theta)$ under the restriction that $\theta \in \omega$, and denote the resulting maximum by $L(\hat{\omega})$. Denote by $\hat{\theta}_\omega$ the MLE of θ subject to the restriction that $\theta \in \omega$. Then $L(\hat{\omega}) = L(\hat{\theta}_\omega)$. Assume now that $L(\theta)$ is continuous (in θ), and suppose that the true value of θ , call it θ_0 , is in ω . It is a property of a MLE that it gets closer and closer to the true parameter as the sample size n increases. Under the assumption that $\theta_0 \in \omega$, it follows that both $\hat{\theta}$ and $\hat{\theta}_\omega$ will be close to θ_0 and therefore close to each other. Then, by the assumed continuity of $L(\theta)$, the quantities $L(\hat{\theta})$ and $L(\hat{\theta}_\omega)$ are close together, so that the ratio

$$\lambda(x_1, \dots, x_n) = \lambda = L(\hat{\theta}_\omega)/L(\hat{\theta}) \quad (8)$$

(which is always ≤ 1) is close to 1. On the other hand, if $\theta_0 \in \omega^c$, then $\hat{\theta}$ and $\hat{\theta}_\omega$ are not close together, and therefore $L(\hat{\theta})$ and $L(\hat{\theta}_\omega)$ need not be close either. Thus, the ratio $L(\hat{\theta}_\omega)/L(\hat{\theta})$ need not be close to 1. These considerations lend to the following test:

$$\text{Reject } H_0 \text{ when } \lambda < \lambda_0, \quad \text{where } \lambda_0 \text{ is a constant to be determined.} \quad (9)$$

By the monotonicity of the function $y = \log x$, the inequality $\lambda < \lambda_0$ is equivalent to $-2 \log \lambda(X_1, \dots, X_n) > C (= -2 \log \lambda_0)$. It is seen in Chapter 11 that an approximate determination of C is made by the fact that, under certain conditions, the distribution of $-2 \log \lambda(X_1, \dots, X_n)$ is χ_f^2 , where $f = \text{dimension of } \Omega - \text{dimension of } \omega$. Namely:

$$\text{Reject } H_0 \text{ when } -2 \log \lambda > C, \quad \text{where } C \simeq \chi_{f; \alpha}^2. \quad (10)$$

In closing this section, it is to be mentioned that the concept of P -value is another way of looking at a test in an effort to assess how strong (or weak) the rejection of a hypothesis is. The P -value (probability value) of a test is defined to be the smallest probability at which the hypothesis tested would be rejected for the data at hand. Roughly put, the P -value of a test is the probability, calculated under the null hypothesis, when the observed value of the test statistic is used as if it were the cut-off point of the test. The P -value of a test often accompanies a null hypothesis which is rejected, as an indication of the strength or weakness of rejection. The smaller the P -value, the stronger the rejection of the null hypothesis, and vice versa. More about it in Chapter 11.

8.4 The Basics of Regression Analysis

In the last three sections, we discussed the general principles of point estimation, interval estimation, and testing hypotheses in a general setup. These principles apply, in particular, in specific models. Two such models are *Regression models* and *Analysis of Variance models*.

A regression model arises in situations such as those described in Examples 22 and 23 in Chapter 1. Its simplest form is as follows: At fixed points x_1, \dots, x_n ,

respective measurements y_1, \dots, y_n are taken, which may be subject to an assortment of random errors e_1, \dots, e_n . Thus, the y_i 's are values of r.v.'s Y_i 's, which may often be assumed to have the structure: $Y_i = \beta_1 + \beta_2 x_i + e_i$, $i = 1, \dots, n$; here β_1 and β_2 are parameters (unknown constants) of the model. For the random errors e_i , it is not unreasonable to assume that $Ee_i = 0$; we also assume that they have the same variance, $\text{Var}(e_i) = \sigma^2 \in (0, \infty)$. Furthermore, it is reasonable to assume that the e_i 's are i.i.d. r.v.'s, which implies independence of the r.v.'s Y_1, \dots, Y_n . It should be noted, however, that the Y_i 's are *not* identically distributed, since, for instance, they have different expectations: $EY_i = \beta_1 + \beta_2 x_i$, $i = 1, \dots, n$. Putting these assumptions together, we arrive at the following simple *linear regression model*.

$$Y_i = \beta_1 + \beta_2 x_i + e_i, \quad \text{the } e_i\text{'s are i.i.d. with } Ee_i = 0 \text{ and} \\ \text{Var}(e_i) = \sigma^2, \quad i = 1, \dots, n. \quad (11)$$

The quantities β_1 , β_2 , and σ^2 are the *parameters* of the model; the Y_i 's are independent but not identically distributed; also, $EY_i = \beta_1 + \beta_2 x_i$ and $\text{Var}(Y_i) = \sigma^2$, $i = 1, \dots, n$.

The term “regression” derives from the way the Y_i 's are produced from the x_i 's, and the term “linear” indicates that the parameters β_1 and β_2 enter into the model raised to the first power.

The main problems in connection with model (11) are to estimate the parameters β_1 , β_2 , and σ^2 ; construct confidence intervals for β_1 and β_2 ; test hypotheses about β_1 and β_2 ; and predict the expected value EY_{i_0} (or the value itself Y_{i_0}) corresponding to an x_{i_0} , distinct, in general, from x_1, \dots, x_n . Estimates of β_1 and β_2 , the *Least Squares Estimates* (LSE's), can be constructed without any further assumptions; the same for an estimate of σ^2 . For the remaining parts, however, there is a need to stipulate a distribution for the e_i 's. Since the e_i 's are random errors, it is reasonable to assume that they are Normally distributed; this then implies Normal distribution for the Y_i 's. Thus, model (11) now becomes:

$$Y_i = \beta_1 + \beta_2 x_i + e_i, \quad \text{the } e_i\text{'s are independently} \\ \text{distributed as } N(0, \sigma^2), \quad i = 1, \dots, n. \quad (12)$$

Under model (12), the MLE's of β_1 , β_2 , and σ^2 are derived, and their distributions are determined. This allows us to pursue the resolution of the parts of constructing confidence intervals, testing hypotheses, and of prediction. The relevant discussion is presented in Chapter 13.

8.5 The Basics of Analysis of Variance

Analysis of Variance (ANOVA) is a powerful technique, which provides the means of assessing and/or comparing several entities. ANOVA can be used effectively in many situations; in particular, it can be used in assessing and/or comparing crop yields corresponding to different soil treatments; crop yields

corresponding to different soils and fertilizers; for the comparison of a certain brand of gasoline with or without an additive by using it in several cars; the comparison of different brands of gasoline by using them in several cars; the comparison of the wearing of different materials; the comparison of the effect of different types of oil on the wear of several piston rings, etc.; the comparison of the yields of a chemical substance by using different catalytic methods; the comparison of the strengths of certain objects made of different batches of some material; the comparison of test scores from different schools and different teachers, etc.; and identification of the melting point of a metal by using different thermometers. Example 24 in Chapter 1 provides another case where ANOVA techniques are appropriate.

Assessment and comparisons are done by way of point estimation, interval estimation, and testing hypotheses, as these techniques apply to the specific ANOVA models to be considered. The more factors involved in producing an outcome, the more complicated the model becomes. However, the basic ideas remain the same throughout.

For the sake of illustrating the issues involved, consider the so-called *one-way layout* or *one-way classification* model. Consider one kind of gasoline, for example, unleaded regular gasoline, and suppose we supply ourselves with amounts of this gasoline, purchased from I different companies. The objective is to compare these I brands of gasoline from yield viewpoint. To this end, a car (or several but pretty similar cars) operates under each one of the I brands of gasoline for J runs in each case. Let Y_{ij} be the number of miles per hour for the j th run when the i th brand of gasoline is used. Then the Y_{ij} 's are r.v.'s for which the following structure is assumed: For a given i , the actual number of miles per hour for the j th run varies around a *mean value* μ_i , and these variations are due to an assortment of random errors e_{ij} . In other words, it makes sense to assume that $Y_{ij} = \mu_i + e_{ij}$. It is also reasonable to assume that the random errors e_{ij} are independent r.v.'s distributed as $N(0, \sigma^2)$, some unknown variance σ^2 . Thus, we have stipulated the following model:

$$Y_{ij} = \mu_i + e_{ij}, \quad \text{where the } e_{ij}\text{'s are independently} \\ \sim N(0, \sigma^2), \quad i = 1, \dots, I(\geq 2), \quad j = 1, \dots, J(\geq 2). \quad (13)$$

The quantities μ_i , $i = 1, \dots, I$, and σ^2 are the *parameters* of the model.

It follows that the r.v.'s Y_{ij} are independent and $Y_{ij} \sim N(\mu_i, \sigma^2)$, $j = 1, \dots, J$, $i = 1, \dots, I$.

The issues of interest here are those of estimating the μ_i 's (mean number of miles per hour for the i th brand of gasoline) and σ^2 . Also, we wish to test the hypothesis that there is really no difference between these I different brands of gasoline; in other words, test $H_0 : \mu_1 = \dots = \mu_I (= \mu, \text{ say, unknown})$. Should this hypothesis be rejected, we would wish to identify the brands of gasoline which cause the rejection. This can be done by constructing a confidence interval for certain linear combinations of the μ_i 's called *contrasts*. That is, $\sum_{i=1}^I c_i \mu_i$, where c_1, \dots, c_I are constants with $\sum_{i=1}^I c_i = 0$.

Instead of having one factor (gasoline brand) affecting the outcome (number of miles per hour), there may be two (or more) such factors. For example,

there might be some chemical additives meant to enhance the mileage. In this framework, suppose there are J such chemical additives, and let us combine each one of the I brands of gasoline with each one of the J chemical additives. For simplicity, suppose we take just one observation, Y_{ij} , on each one of the IJ pairs. Then it makes sense to assume that the r.v. Y_{ij} is the result of the following additive components: A basic quantity (*grand mean*) μ , the same for all i and j ; an *effect* α_i due to the i th brand of gasoline (the *ith row effect*); an *effect* β_j due to the j th chemical additive (the *ith column effect*); and, of course, the random error e_{ij} due to a host of causes. So, the assumed model is then: $Y_{ij} = \mu + \alpha_i + \beta_j + e_{ij}$. As usually, we assume that the e_{ij} 's are independent $\sim N(0, \sigma^2)$ with some (unknown) variance σ^2 , which implies that the Y_{ij} 's are independent r.v.'s and $Y_{ij} \sim N(\mu + \alpha_i + \beta_j, \sigma^2)$. We further assume that some of α_i effects are ≥ 0 , some are < 0 , and on the whole $\sum_{i=1}^I \alpha_i = 0$; and likewise for the β_j effects: $\sum_{j=1}^J \beta_j = 0$. Summarizing these assumptions, we have then:

$$\begin{aligned} Y_{ij} &= \mu + \alpha_i + \beta_j + e_{ij}, & \text{where the } e_{ij}\text{'s are independently} \\ &\sim N(0, \sigma^2), & i = 1, \dots, I(\geq 2), \quad j = 1, \dots, J(\geq 2), \\ \sum_{i=1}^I \alpha_i &= 0, & \sum_{j=1}^J \beta_j = 0. \end{aligned} \quad (14)$$

The quantities $\mu, \alpha_i, i = 1, \dots, I, \beta_j, j = 1, \dots, J$ and σ^2 are the *parameters* of the model.

As already mentioned, the implication is that the r.v.'s Y_{ij} are independent and $Y_{ij} \sim N(\mu + \alpha_i + \beta_j, \sigma^2), i = 1, \dots, I, j = 1, \dots, J$.

The model described by (14) is called *two-way layout* or *two-way classification*, as the observations are affected by two factors.

The main statistical issues are those of estimating the parameters involved and testing irrelevance of either one of the factors involved—that is, testing $H_{0A} : \alpha_1 = \dots = \alpha_I = 0, H_{0B} : \beta_1 = \dots = \beta_J = 0$. Details will be presented in Chapter 14. There, an explanation of the term “ANOVA” will also be given.

8.6 The Basics of Nonparametric Inference

All of the problems discussed in the previous sections may be summarized as follows: On the basis of a random sample of size n, X_1, \dots, X_n , drawn from the p.d.f. $f(\cdot; \theta), \theta \in \Omega \subseteq \mathfrak{R}$, construct a point estimate for θ , a confidence interval for θ , and test hypotheses about θ . In other words, the problems discussed were those of making (*statistical*) *inference* about θ . These problems are suitably modified for a multidimensional parameter. The fundamental assumption in this framework is that the functional form of the p.d.f. $f(\cdot; \theta)$ is known; the only thing which does not render $f(\cdot; \theta)$ completely known is the presence of the (unknown constant) parameter θ .

In many situations, stipulating a functional form for $f(\cdot; \theta)$ either is dictated by circumstances or is the product of accumulated experience. In the absence

of these, we must still proceed with the problems of estimating important quantities, either by points or by intervals, and testing hypotheses about them. However, the framework now is *nonparametric*, and the relevant inference is referred to as *nonparametric inference*.

Actually, there have been at least three cases so far where *nonparametric estimation* was made without referring to it as such. Indeed, if X_1, \dots, X_n are i.i.d. r.v.'s with unknown mean μ , then the sample mean \bar{X}_n may be taken as an estimate of μ , regardless of what the underlying distribution of the X_i 's is. This estimate is recommended on the basis of at least three considerations. First, it is unbiased, $E\bar{X}_n = \mu$ no matter what the underlying distribution is; second, \bar{X}_n is the moment estimate of μ ; and third, by the WLLN, $\bar{X}_n \xrightarrow{P} \mu$, so that \bar{X}_n is close to μ , in the sense of probability, for all sufficiently large n . Now suppose that the X_i 's also have (an unknown) variance $\sigma^2 \in (0, \infty)$. Then the sample variance S_n^2 can be used as an estimate of σ^2 , because it is unbiased (Section 8.1) and also $S_n^2 \xrightarrow{P} \sigma^2$. Furthermore, by combining \bar{X}_n and S_n^2 and using Theorem 7(ii) in Chapter 7, we have that $\sqrt{n}(\bar{X}_n - \mu)/S_n \simeq N(0, 1)$ for large n . Then, for such n , $[\bar{X}_n - z_{\alpha/2} \frac{S_n}{\sqrt{n}}, \bar{X}_n + z_{\alpha/2} \frac{S_n}{\sqrt{n}}]$ is a confidence interval for μ with confidence coefficient approximately $1 - \alpha$.

Also, the (unknown) d.f. F of the X_i 's has been estimated at every point $x \in \mathfrak{R}$ by the empirical d.f. F_n (see relation (1) in Chapter 7). The estimate F_n has at least two desirable properties. For all $x \in \mathfrak{R}$ and regardless of the form of the d.f. F : $EF_n(x) = F(x)$ and $F_n(x) \xrightarrow{P} F(x)$.

What has not been done so far is to estimate the p.d.f. $f(x)$ at each $x \in \mathfrak{R}$, under certain regularity conditions, which do not include postulation of a functional form for f . There are several ways of doing this; in Chapter 15, we are going to adopt the so-called *kernel method* of estimating f . Some desirable results of the proposed estimate will be stated without proofs.

Regarding testing hypotheses, the problems to be addressed in Chapter 15 will be to test the hypothesis that the (unknown) d.f. F is, actually, equal to a known one F_0 ; that is $H_0 : F = F_0$, the alternative H_A being that $F(x) \neq F_0(x)$ for at least one $x \in \mathfrak{R}$. Actually, from a practical viewpoint, it is more important to compare two (unknown) d.f.'s F and G , by stipulating $H_0 : F = G$. The alternative can be any one of the following: $H_A : F \neq G$, $H'_A : F > G$, $H''_A : F < G$, in the sense that $F(x) \geq G(x)$ or $F(x) \leq G(x)$, respectively, for all $x \in \mathfrak{R}$, and strict inequality for at least one x . In carrying out the appropriate tests, one has to use some pretty sophisticated asymptotic results regarding empirical d.f.'s. An alternative approach to using empirical d.f.'s is to employ the concept of a *rank* test or the concept of a *sign* test. These things will be discussed to some extent in Chapter 15. That chapter is concluded with the basics of regression estimation but in a nonparametric framework. In such a situation, what is estimated is an entire function rather than a few parameters. Some basic results are stated in Chapter 15.



Chapter 9

Point Estimation

In the previous chapter, the basic terminology and concepts of parametric point estimation were introduced briefly. In the present chapter, we are going to elaborate extensively on this matter. For brevity, we will use the term estimation rather than parametric point estimation. The methods of estimation to be discussed here are those listed in the first section of the previous chapter; namely, maximum likelihood estimation, estimation through the concepts of unbiasedness and minimum variance (which lead to uniformly minimum variance estimates), estimation based on decision-theoretic concepts, and estimation by the method of moments. The method of estimation by way of the principle of least squares is commonly used in the so-called linear models. Accordingly, it is deferred to Chapter 13.

Before we embark on the mathematical derivations, it is imperative to keep in mind the big picture; namely, why do we do what we do? A brief description is as follows. Let X be a r.v. with p.d.f. $f(\cdot; \theta)$, where θ is a parameter lying in a parameter space Ω . It is assumed that the functional form of the p.d.f. is completely known. So, if θ were known, then the p.d.f. would be known, and consequently we could calculate, in principle, all probabilities related to X , the expectation of X , its variance, etc. The problem, however, is that most often in practice (and in the present context) θ is not known. Then the objective is to estimate θ on the basis of a random sample of size n from $f(\cdot; \theta)$, X_1, \dots, X_n . Then, replacing θ in $f(\cdot; \theta)$ by a “good” estimate of it, one would expect to be able to use the resulting p.d.f. for the purposes described above to a satisfactory degree.

9.1 Maximum Likelihood Estimation: Motivation and Examples

The following simple example is meant to shed light to the intuitive, yet quite logical, principle of Maximum Likelihood Estimation.

EXAMPLE 1

Let X_1, \dots, X_{10} be i.i.d. r.v.'s from the $B(1, \theta)$ distribution, $0 < \theta < 1$, and let x_1, \dots, x_{10} be the respective observed values. For convenience, set $t = x_1 + \dots + x_{10}$. Further, suppose that in the 10 trials, 6 resulted in successes, so that $t = 6$. Then the likelihood function involved is: $L(\theta | \mathbf{x}) = \theta^6(1 - \theta)^4$, $0 < \theta < 1$, $\mathbf{x} = (x_1, \dots, x_{10})$. Thus, $L(\theta | \mathbf{x})$ is the probability of observing exactly 6 successes in 10 independent Binomial trials, the successes occurring on those trials for which $x_i = 1$, $i = 1, \dots, 10$; this probability is a function of the (unknown) parameter θ . Let us calculate the values of this probability for θ ranging from 0.1 to 0.9. We find:

Values of θ	Values of $L(\theta \mathbf{x})$
0.1	0.000006656
0.2	0.000026200
0.3	0.000175000
0.4	0.000531000
0.5	0.000976000
0.6	0.003320000
0.7	0.003010000
0.8	0.000419000
0.9	0.000053000

We observe that the values of $L(\theta | \mathbf{x})$ keep increasing, it attains its maximum value at $\theta = 0.6$, and then the values keep decreasing. Thus, if these 9 values were the only possible values for θ (which they are not!), one would reasonably enough choose the value of 0.6 as the value of θ . The value $\theta = 0.6$ has the distinction of maximizing (among the 9 values listed) the probability of attaining the 6 already observed successes.

We observe that $0.6 = \frac{6}{10} = \frac{t}{n}$, where n is the number of trials and t is the number of successes. It will be seen in Example 2 that the value $\frac{t}{n}$, actually, maximizes the likelihood function among *all* values of θ with $0 < \theta < 1$. Then $\frac{t}{n}$ will be the Maximum Likelihood Estimate of θ to be denoted by $\hat{\theta}$; i.e., $\hat{\theta} = \frac{t}{n}$.

In a general setting, let X_1, \dots, X_n be i.i.d. r.v.'s with p.d.f. $f(\cdot; \theta)$ with $\theta \in \Omega$, and let x_1, \dots, x_n be the respective observed values and $\mathbf{x} = (x_1, \dots, x_n)$. The likelihood function, $L(\theta | \mathbf{x})$, is given by $L(\theta | \mathbf{x}) = \prod_{i=1}^n f(x_i; \theta)$, and a value of θ which maximizes $L(\theta | \mathbf{x})$ is called a *Maximum Likelihood Estimate* (MLE) of θ . Clearly, the MLE depends on \mathbf{x} , and we usually write $\hat{\theta} = \hat{\theta}(\mathbf{x})$. Thus,

$$L(\hat{\theta} | \mathbf{x}) = \max\{L(\theta | \mathbf{x}); \theta \in \Omega\}. \quad (1)$$

The justification for choosing an estimate as the value of the parameter which maximizes the likelihood function is the same as that given in Example 1, when the r.v.'s are discrete. The same interpretation holds true for r.v.'s of the continuous type, by considering small intervals around the observed values.

Once we decide to adopt the *Maximum Likelihood Principle* (i.e., the principle of choosing an estimate of the parameter through the process of maximizing the likelihood function), the actual identification of an MLE is a purely mathematical problem; namely, that of maximizing a function. This maximization, if possible at all, often (but not always) is done through differentiation. Examples to be discussed below will illustrate various points.

Before embarking on specific examples, it must be stressed that, whenever a maximum is sought by differentiation, the second-order derivative(s) must also be examined in search of a maximum. Also, it should be mentioned that maximization of the likelihood function, which is the product of n factors, is equivalent to maximization of its logarithm (always with base e), which is the sum of n summands, thus much easier to work with.

REMARK 1 Let us recall that a function $y = g(x)$ attains a maximum at a point $x = x_0$, if $\frac{d}{dx}g(x)|_{x=x_0} = 0$ and $\frac{d^2}{dx^2}g(x)|_{x=x_0} < 0$.

EXAMPLE 2

In terms of a random sample of size n , X_1, \dots, X_n from the $B(1, \theta)$ distribution with observed values x_1, \dots, x_n , determine the MLE $\hat{\theta} = \hat{\theta}(\mathbf{x})$ of $\theta \in (0, 1)$, $\mathbf{x} = (x_1, \dots, x_n)$.

DISCUSSION Since $f(x_i; \theta) = \theta^{x_i}(1 - \theta)^{1-x_i}$, $x_i = 0$ or 1 , $i = 1, \dots, n$, the likelihood function is

$$L(\theta | \mathbf{x}) = \prod_{i=1}^n f(x_i; \theta) = \theta^t(1 - \theta)^{n-t}, \quad t = x_1 + \dots + x_n,$$

so that $t = 0, 1, \dots, n$. Hence $\log L(\theta | \mathbf{x}) = t \log \theta + (n - t) \log(1 - \theta)$. From the *likelihood equation* $\frac{\partial}{\partial \theta} \log L(\theta | \mathbf{x}) = \frac{t}{\theta} - \frac{n-t}{1-\theta} = 0$, we obtain $\theta = \frac{t}{n}$. Next, $\frac{\partial^2}{\partial \theta^2} \log L(\theta | \mathbf{x}) = -\frac{t}{\theta^2} - \frac{n-t}{(1-\theta)^2}$, which is negative for all θ and hence for $\theta = t/n$. Therefore, the MLE of θ is $\hat{\theta} = \frac{t}{n} = \bar{x}$.

EXAMPLE 3

Determine the MLE $\hat{\theta} = \hat{\theta}(\mathbf{x})$ of $\theta \in (0, \infty)$ in the $P(\theta)$ distribution in terms of the random sample X_1, \dots, X_n with observed values x_1, \dots, x_n .

DISCUSSION Here $f(x_i; \theta) = \frac{e^{-\theta} \theta^{x_i}}{x_i!}$, $x_i = 0, 1, \dots$, $i = 1, \dots, n$, so that

$$\begin{aligned} \log L(\theta | \mathbf{x}) &= \log \left(\prod_{i=1}^n \frac{e^{-\theta} \theta^{x_i}}{x_i!} \right) = \log \left(e^{-n\theta} \prod_{i=1}^n \frac{\theta^{x_i}}{x_i!} \right) \\ &= -n\theta + (\log \theta) \sum_{i=1}^n x_i - \log \left(\prod_{i=1}^n x_i! \right) \\ &= -n\theta + (n \log \theta) \bar{x} - \log \left(\prod_{i=1}^n x_i! \right). \end{aligned}$$

Hence $\frac{\partial}{\partial \theta} \log L(\theta | \mathbf{x}) = -n + \frac{n\bar{x}}{\theta} = 0$, which gives $\theta = \bar{x}$, and $\frac{\partial^2}{\partial \theta^2} \log L(\theta | \mathbf{x}) = -\frac{n\bar{x}}{\theta^2} < 0$ for all θ and hence for $\theta = \bar{x}$. Therefore the MLE of θ is $\hat{\theta} = \bar{x}$.

EXAMPLE 4

Determine the MLE $\hat{\theta} = \hat{\theta}(\mathbf{x})$ of $\theta \in (0, \infty)$ in the Negative Exponential distribution $f(x; \theta) = \theta e^{-\theta x}$, $x > 0$, on the basis of the random sample X_1, \dots, X_n with observed values x_1, \dots, x_n .

DISCUSSION Since $f(x_i; \theta) = \theta e^{-\theta x_i}$, $x_i > 0$, $i = 1, \dots, n$, we have

$$\log L(\theta | \mathbf{x}) = \log(\theta^n e^{-n\theta \bar{x}}) = n \log \theta - n\bar{x}\theta, \text{ so that}$$

$\frac{\partial}{\partial \theta} \log L(\theta | \mathbf{x}) = \frac{n}{\theta} - n\bar{x} = 0$, and hence $\theta = 1/\bar{x}$. Furthermore, $\frac{\partial^2}{\partial \theta^2} \log L(\theta | \mathbf{x}) = -\frac{n}{\theta^2} < 0$ for all θ and hence for $\theta = 1/\bar{x}$. It follows that $\hat{\theta} = 1/\bar{x}$.

EXAMPLE 5

Let X_1, \dots, X_n be a random sample from the $N(\mu, \sigma^2)$ distribution, where only one of the parameters is known. Determine the MLE of the other (unknown) parameter.

DISCUSSION With x_1, \dots, x_n being the observed values of X_1, \dots, X_n , we have:

(i) *Let μ be unknown.* Then

$$\begin{aligned} \log L(\mu | \mathbf{x}) &= \log \left\{ \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp \left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right] \right\} \\ &= -n \log(\sqrt{2\pi}\sigma) - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2, \text{ so that} \end{aligned}$$

$\frac{\partial}{\partial \mu} \log L(\mu | \mathbf{x}) = \frac{n(\bar{x} - \mu)}{\sigma^2} = 0$, and hence $\mu = \bar{x}$. Furthermore, $\frac{\partial^2}{\partial \mu^2} \log L(\mu | \mathbf{x}) = -\frac{n}{\sigma^2} < 0$ for all μ and hence for $\mu = \bar{x}$. It follows that the MLE of μ is $\hat{\mu} = \bar{x}$.

(ii) *Let σ^2 be unknown.* Then

$$\begin{aligned} \log L(\sigma^2 | \mathbf{x}) &= \log \left\{ \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right] \right\} \\ &= -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2, \text{ so that} \end{aligned}$$

$\frac{\partial}{\partial \sigma^2} \log L(\sigma^2 | \mathbf{x}) = -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n (x_i - \mu)^2 = 0$, and hence $\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2$; set $\frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2 = s^2$. Then


$$\begin{aligned} \frac{\partial^2}{\partial (\sigma^2)^2} \log L(\sigma^2 | \mathbf{x}) &= \frac{n}{2\sigma^4} - \frac{2n}{2\sigma^6} \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2 \\ &= \frac{n}{2(\sigma^2)^2} - \frac{2n}{2(\sigma^2)^3} s^2, \text{ so that} \end{aligned}$$

$\frac{\partial^2}{\partial (\sigma^2)^2} \log L(\sigma^2 | \mathbf{x})|_{\sigma^2=s^2} = \frac{n}{2(s^2)^2} - \frac{2ns^2}{2(s^2)^3} = -\frac{n}{2s^4} < 0$. It follows that the MLE of σ^2 is $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2$.

In all of the preceding examples, the MLE's were determined through differentiation. Below is a case where this method does not apply because,

simply, the derivative does not exist. As an introduction to the problem, let $X \sim U(0, \theta)$ ($\theta > 0$), so that the likelihood function is $L(\theta | x) = \frac{1}{\theta} I_{[0, \theta]}(x)$ where it is to be recalled that the *indicator* function I_A is defined by $I_A(x) = 1$ if $x \in A$, and $I_A(x) = 0$ if $x \in A^c$. The picture of $L(\cdot | x)$ is shown in Figure 9.1.

Figure 9.1



EXAMPLE 6

Let X_1, \dots, X_n be a random sample from the Uniform $U(\alpha, \beta)$ ($\alpha < \beta$) distribution, where only one of α and β is unknown. Determine the MLE of the (unknown) parameter.

DISCUSSION

(i) *Let α be unknown.* Since

$$f(x_i; \alpha) = \frac{1}{\beta - \alpha} I_{[\alpha, \beta]}(x_i), \quad i = 1, \dots, n, \text{ it follows that}$$

$$L(\alpha | \mathbf{x}) = \frac{1}{(\beta - \alpha)^n} \prod_{i=1}^n I_{[\alpha, \beta]}(x_i) = \frac{1}{(\beta - \alpha)^n} I_{[\alpha, \beta]}(x_{(1)}) I_{[\alpha, \beta]}(x_{(n)}),$$

where $x_{(1)} = \min(x_1, \dots, x_n)$, $x_{(n)} = \max(x_1, \dots, x_n)$; or

$$L(\alpha | \mathbf{x}) = \frac{1}{(\beta - \alpha)^n} I_{[\alpha, \infty)}(x_{(1)}) I_{(-\infty, \beta]}(x_{(n)}). \quad (2)$$

Maximization of $L(\alpha | \mathbf{x})$ with respect to α means two things: maximization of $I_{[\alpha, \infty)}(x_{(1)})$ and maximization of $1/(\beta - \alpha)^n$. The maximum value of the former quantity is 1 and occurs as long as $\alpha \leq x_{(1)}$. The latter quantity gets larger and larger as α gets closer and closer to β . But always $\alpha \leq x_{(1)} \leq \beta$, and α is subject to the restriction $\alpha \leq x_{(1)}$. Thus, α gets closest to β , if $\alpha = x_{(1)}$. In other words, the MLE of α is $\hat{\alpha} = x_{(1)}$.

(ii) *Let β be unknown.* Relation (2) then becomes

$$L(\beta | \mathbf{x}) = \frac{1}{(\beta - \alpha)^n} I_{[\alpha, \infty)}(x_{(1)}) I_{(-\infty, \beta]}(x_{(n)}),$$

whereas always $\alpha \leq x_{(n)} \leq \beta$. Then, arguing as in the first case, we have that the MLE of β is $\hat{\beta} = x_{(n)}$.

In the examples discussed so far, there was a single parameter to be estimated. In the examples presented below, the parameters to be estimated will be two or more. If the maximization is to be done through differentiation, then the following remark reminds us how this method is implemented.

REMARK 2 The function $y = g(x_1, \dots, x_k)$ attains a maximum at a point (x_{01}, \dots, x_{0k}) , if the point (x_{01}, \dots, x_{0k}) satisfies the system of the k equations $\frac{\partial}{\partial x_i} g(x_1, \dots, x_k) = 0$, $i = 1, \dots, k$, and, in addition, the point (x_{01}, \dots, x_{0k}) renders the $k \times k$ matrix of the second-order derivatives $(\frac{\partial^2}{\partial x_i \partial x_j} g(x_1, \dots, x_k))$, $i, j = 1, \dots, k$, negative definite. What is meant by the term “negative definite” is that the real-valued quantity below is < 0 for all nonzero vectors $(\lambda_1, \dots, \lambda_k)$ ($\lambda_1^2 + \dots + \lambda_k^2 \neq 0$); namely,

$$(\lambda_1, \dots, \lambda_k) \left(\frac{\partial^2}{\partial x_i \partial x_j} g(x_1, \dots, x_k) \Big|_{(x_1, \dots, x_k) = (x_{01}, \dots, x_{0k})} \right) \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_k \end{pmatrix} < 0.$$

EXAMPLE 7

Refer to Example 5 and suppose that both μ and σ^2 are unknown. Determine their MLE's.

DISCUSSION Here

$$\log L(\mu, \sigma^2 | \mathbf{x}) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2,$$

and then the two likelihood equations produce the unique solution $\mu = \bar{x}$ and $\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$, which we may denote by s^2 . Next, the 2×2 matrix of the second-order derivatives, evaluated at (\bar{x}, s^2) , becomes $\begin{pmatrix} -\frac{n}{s^2} & 0 \\ 0 & -\frac{n}{2s^4} \end{pmatrix}$, which is negative definite (see Exercise 1.2 below). Thus, $\hat{\mu} = \bar{x}$ and $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$ are the MLE's of μ and σ^2 , respectively.

EXAMPLE 8

A Multinomial experiment is carried out independently n times, so that the likelihood function is

$$L(p_1, \dots, p_r | \mathbf{x}) = \frac{n!}{x_1! \cdots x_r!} p_1^{x_1} \cdots p_r^{x_r},$$

where $x_i \geq 0$, $i = 1, \dots, r$ integers, with $x_1 + \dots + x_r = n$, and $0 < p_i < 1$, $i = 1, \dots, r$ with $p_1 + \dots + p_r = 1$. Determine the MLE's of p_i , $i = 1, \dots, r$.

DISCUSSION The number of independent parameters is $r - 1$, since, for example, $p_r = 1 - p_1 - \dots - p_{r-1}$. Looking at the $\log L(p_1, \dots, p_r | \mathbf{x})$ and taking partial derivatives with respect to p_i , $i = 1, \dots, r - 1$ (and remembering that $p_r = 1 - p_1 - \dots - p_{r-1}$) we obtain

$$x_i \frac{1}{p_i} - x_r \frac{1}{p_r} = 0, \quad i = 1, \dots, r - 1.$$

From these relations, the unique solution $p_i = \frac{x_i}{n}$, $i = 1, \dots, r$ follows.

Next, the $(r-1) \times (r-1)$ matrix of the second-order derivatives, evaluated at $p_i = x_i/n$, $i = 1, \dots, r$, is given by

$$\begin{pmatrix} -\frac{n^2}{x_1} - \frac{n^2}{x_r} & -\frac{n^2}{x_r} & \dots & -\frac{n^2}{x_r} & -\frac{n^2}{x_r} \\ -\frac{n^2}{x_r} & -\frac{n^2}{x_2} - \frac{n^2}{x_r} & \dots & -\frac{n^2}{x_r} & -\frac{n^2}{x_r} \\ \dots & \dots & \dots & \dots & \dots \\ -\frac{n^2}{x_r} & -\frac{n^2}{x_r} & \dots & -\frac{n^2}{x_r} & -\frac{n^2}{x_{r-1}} - \frac{n^2}{x_r} \end{pmatrix},$$

which is seen to be negative definite. Consequently, the MLE's of p_i , $i = 1, \dots, r$ are $\hat{p}_i = \frac{x_i}{n}$, $i = 1, \dots, r$. (Also, see Exercise 1.3 below.)

EXAMPLE 9

Refer to Example 6, assume that both α and β are unknown, and determine their MLE's.

DISCUSSION Expression (1) becomes here as follows:

$$L(\alpha, \beta | \mathbf{x}) = \frac{1}{(\beta - \alpha)^n} I_{[\alpha, \infty)}(x_{(1)}) I_{(-\infty, \beta]}(x_{(n)}). \quad (3)$$

Since always $\alpha \leq x_{(1)} \leq x_{(n)} \leq \beta$, the right-hand side of (3) is maximized if $I_{[\alpha, \infty)}(x_{(1)}) = 1$ and $I_{(-\infty, \beta]}(x_{(n)}) = 1$, which happen if $\alpha \leq x_{(1)}$, $x_{(n)} \leq \beta$ and also if α and β are as close together as possible. Clearly, this happens for $\alpha = x_{(1)}$ and $\beta = x_{(n)}$. In other words, the MLE's of α and β are $\hat{\alpha} = x_{(1)}$ and $\hat{\beta} = x_{(n)}$.

Exercises

- 1.1 Refer to Example 6(ii) and justify the statement made there that $x_{(n)}$ is, indeed, the MLE of β .
- 1.2 Show that the matrix $\begin{pmatrix} -n/s^2 & 0 \\ 0 & -n/2s^4 \end{pmatrix}$ in Example 7 is, indeed, negative definite.
- 1.3 In reference to Example 8, show that:
 - (i) $p_i = \frac{x_i}{n}$, $i = 1, \dots, r$, is, indeed, the unique solution of the system of equations considered there.
 - (ii) The $(r-1) \times (r-1)$ matrix exhibited there is the matrix of the second-order derivatives as stated.
 - (iii) The matrix in part (ii) is negative definite.
- 1.4 In reference to Example 18 below, show that $\text{Var}_\theta(S^2) = \frac{2\sigma^4}{n-1}$ as stated there.
- 1.5 If X_1, \dots, X_n are independent r.v.'s distributed as $B(k, \theta)$, $\theta \in \Omega = (0, 1)$, with respective observed values x_1, \dots, x_n , show that $\hat{\theta} = \frac{\bar{x}}{k}$ is the MLE of θ , where \bar{x} is the sample mean of the x_i 's.

- 1.6** If the independent r.v.'s X_1, \dots, X_n have the Geometric p.d.f. $f(x; \theta) = \theta(1 - \theta)^{x-1}$, $x = 1, 2, \dots$, $\theta \in \Omega = (0, 1)$, and respective observed values x_1, \dots, x_n , then show that $\hat{\theta} = 1/\bar{x}$ is the MLE of θ .
- 1.7** On the basis of a random sample of size n from the p.d.f. $f(x; \theta) = (\theta + 1)x^\theta$, $0 < x < 1$, $\theta \in \Omega = (-1, \infty)$, derive the MLE of θ .
- 1.8** On the basis of a random sample of size n from the p.d.f. $f(x; \theta) = \theta x^{\theta-1}$, $0 < x < 1$, $\theta \in \Omega = (0, \infty)$, derive the MLE of θ .
- 1.9** (i) Show that the function $f(x; \theta) = \frac{1}{2\theta} e^{-|x|/\theta}$, $x \in \mathfrak{R}$, $\theta \in \Omega = (0, \infty)$ is a p.d.f. (the so-called *Double Exponential* p.d.f.), and draw its picture.
(ii) On the basis of a random sample from this p.d.f., derive the MLE of θ .
- 1.10** (i) Verify that the function $f(x; \theta) = \theta^2 x e^{-\theta x}$, $x > 0$, $\theta \in \Omega = (0, \infty)$ is a p.d.f., by observing that it is the Gamma p.d.f. with parameters $\alpha = 2$, $\beta = 1/\theta$.
(ii) On the basis of a random sample of size n from this p.d.f., derive the MLE of θ .
- 1.11** (i) Show that the function $f(x; \alpha, \beta) = \frac{1}{\beta} e^{-(x-\alpha)/\beta}$, $x \geq \alpha$, $\alpha \in \mathfrak{R}$, $\beta > 0$, is a p.d.f., and draw its picture.
On the basis of a random sample of size n from this p.d.f., determine the MLE of:
(ii) α when β is known.
(iii) β when α is known.
(iv) α and β when both are unknown.
- 1.12** Refer to the Bivariate Normal distribution discussed in Chapter 4, Section 5, whose p.d.f. is given by:

$$f_{X,Y}(x, y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} e^{-q/2}, \quad x, y \in \mathfrak{R},$$

where

$$q = \frac{1}{1-\rho^2} \left[\left(\frac{x-\mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x-\mu_1}{\sigma_1} \right) \left(\frac{y-\mu_2}{\sigma_2} \right) + \left(\frac{y-\mu_2}{\sigma_2} \right)^2 \right];$$

$\mu_1, \mu_2 \in \mathfrak{R}$, $\sigma_1^2, \sigma_2^2 > 0$ and $-1 \leq \rho \leq 1$ are the parameters of the distribution. The objective here is to find the MLE's of these parameters. This is done in two stages, in the present exercise and the exercises following.

For convenient writing, set $\theta = (\mu_1, \mu_2, \sigma_1^2, \sigma_2^2, \rho)$, and form the likelihood function for a sample of size n , (X_i, Y_i) , $i = 1, \dots, n$, from the underlying distribution; i.e.,

$$L(\theta | \mathbf{x}, \mathbf{y}) = \left(\frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \right)^n \exp\left(-\frac{1}{2} \sum_{i=1}^n q_i\right),$$

where

$$q_i = \frac{1}{1 - \rho^2} \left[\left(\frac{x_i - \mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x_i - \mu_1}{\sigma_1} \right) \left(\frac{y_i - \mu_2}{\sigma_2} \right) + \left(\frac{y_i - \mu_2}{\sigma_2} \right)^2 \right],$$

and $\mathbf{x} = (x_1, \dots, x_n)$, $\mathbf{y} = (y_1, \dots, y_n)$, the observed values of the X_i 's and the Y_i 's. Also, set

$$\begin{aligned} \lambda(\boldsymbol{\theta}) &= \lambda(\boldsymbol{\theta} \mid \mathbf{x}, \mathbf{y}) = \log L(\boldsymbol{\theta} \mid \mathbf{x}, \mathbf{y}) \\ &= -n \log(2\pi) - \frac{n}{2} \log \sigma_1^2 - \frac{n}{2} \log \sigma_2^2 - \frac{n}{2} \log(1 - \rho^2) - \frac{1}{2} \sum_{i=1}^n q_i. \end{aligned}$$

- (i) Show that the first order partial derivatives of q given above are provided by the following expressions:

$$\begin{aligned} \frac{\partial q}{\partial \mu_1} &= -\frac{2(x - \mu_1)}{\sigma_1^2(1 - \rho^2)} + \frac{2\rho(y - \mu_2)}{\sigma_1\sigma_2(1 - \rho^2)}, \\ \frac{\partial q}{\partial \mu_2} &= -\frac{2(y - \mu_2)}{\sigma_2^2(1 - \rho^2)} + \frac{2\rho(x - \mu_1)}{\sigma_1\sigma_2(1 - \rho^2)}, \\ \frac{\partial q}{\partial \sigma_1^2} &= -\frac{(x - \mu_1)^2}{\sigma_1^4(1 - \rho^2)} + \frac{\rho(x - \mu_1)(y - \mu_2)}{\sigma_1^3\sigma_2(1 - \rho^2)}, \\ \frac{\partial q}{\partial \sigma_2^2} &= -\frac{(y - \mu_2)^2}{\sigma_2^4(1 - \rho^2)} + \frac{\rho(x - \mu_1)(y - \mu_2)}{\sigma_1\sigma_2^3(1 - \rho^2)}, \\ \frac{\partial q}{\partial \rho} &= \frac{2}{(1 - \rho^2)^2} \left\{ \rho \left[\left(\frac{x - \mu_1}{\sigma_1} \right)^2 + \left(\frac{y - \mu_2}{\sigma_2} \right)^2 \right] \right. \\ &\quad \left. - (1 + \rho^2) \left(\frac{x - \mu_1}{\sigma_1} \right) \left(\frac{y - \mu_2}{\sigma_2} \right) \right\}. \end{aligned}$$

- (ii) Use the above obtained expressions and $\lambda(\boldsymbol{\theta})$ in order to show that:

$$\begin{aligned} \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \mu_1} &= \frac{n}{\sigma_1^2(1 - \rho^2)} (\bar{x} - \mu_1) - \frac{n\rho}{\sigma_1\sigma_2(1 - \rho^2)} (\bar{y} - \mu_2), \\ \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \mu_2} &= \frac{n}{\sigma_2^2(1 - \rho^2)} (\bar{y} - \mu_2) - \frac{n\rho}{\sigma_1\sigma_2(1 - \rho^2)} (\bar{x} - \mu_1), \\ \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \sigma_1^2} &= -\frac{n}{2\sigma_1^2} + \frac{\sum_{i=1}^n (x_i - \mu_1)^2}{2\sigma_1^4(1 - \rho^2)} - \frac{\rho \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{2\sigma_1^3\sigma_2(1 - \rho^2)}, \\ \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \sigma_2^2} &= -\frac{n}{2\sigma_2^2} + \frac{\sum_{i=1}^n (y_i - \mu_2)^2}{2\sigma_2^4(1 - \rho^2)} - \frac{\rho \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{2\sigma_1\sigma_2^3(1 - \rho^2)}, \\ \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \rho} &= \frac{n\rho}{1 - \rho^2} - \frac{\rho \sum_{i=1}^n (x_i - \mu_1)^2}{\sigma_1^2(1 - \rho^2)} - \frac{\rho \sum_{i=1}^n (y_i - \mu_2)^2}{\sigma_2^2(1 - \rho^2)} \\ &\quad + \frac{(1 + \rho^2) \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{\sigma_1\sigma_2(1 - \rho^2)^2}. \end{aligned}$$

- (iii) Setting $\frac{\partial \lambda(\boldsymbol{\theta})}{\partial \mu_1} = \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \mu_2} = 0$, and solving for μ_1 and μ_2 , show that there is a unique solution given by: $\tilde{\mu}_1 = \bar{x}$ and $\tilde{\mu}_2 = \bar{y}$.
- (iv) By setting $\frac{\partial \lambda(\boldsymbol{\theta})}{\partial \sigma_1^2} = \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \sigma_2^2} = \frac{\partial \lambda(\boldsymbol{\theta})}{\partial \rho} = 0$ and replacing μ_1 and μ_2 by the respective expressions \bar{x} and \bar{y} , show that we arrive at the equations:

$$\frac{S_x}{\sigma_1^2} - \frac{\rho S_{xy}}{\sigma_1 \sigma_2} = 1 - \rho^2, \quad \frac{S_y}{\sigma_2^2} - \frac{\rho S_{xy}}{\sigma_1 \sigma_2} = 1 - \rho^2,$$

$$\frac{S_x}{\sigma_1^2} + \frac{S_y}{\sigma_2^2} - \frac{(1 + \rho^2) S_{xy}}{\rho \sigma_1 \sigma_2} = 1 - \rho^2,$$

where $S_x = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$, $S_y = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2$ and $S_{xy} = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})$.

- (v) In the equations obtained in part (iv), solve for σ_1^2 , σ_2^2 , and ρ in order to obtain the unique solution:

$$\tilde{\sigma}_1^2 = S_x, \quad \tilde{\sigma}_2^2 = S_y, \quad \tilde{\rho} = S_{xy}/S_x^{1/2}S_y^{1/2}.$$

1.13 The purpose of this exercise is to show that the values $\tilde{\mu}_1$, $\tilde{\mu}_2$, $\tilde{\sigma}_1^2$, $\tilde{\sigma}_2^2$, and $\tilde{\rho}$ are actually the MLE's of the respective parameters. To this end:

- (i) Take the second-order partial derivatives of $\lambda(\boldsymbol{\theta})$, as indicated below, and show that they are given by the following expressions:

$$\frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_1^2} = -\frac{n}{\sigma_1^2(1 - \rho^2)} \stackrel{\text{def}}{=} d_{11}, \quad \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_1 \partial \mu_2} = \frac{n\rho}{\sigma_1 \sigma_2(1 - \rho^2)} \stackrel{\text{def}}{=} d_{12},$$

$$\frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_1 \partial \sigma_1^2} = -\frac{n(\bar{x} - \mu_1)}{\sigma_1^4(1 - \rho^2)} + \frac{n\rho(\bar{y} - \mu_2)}{2\sigma_1^3 \sigma_2(1 - \rho^2)} \stackrel{\text{def}}{=} d_{13},$$

$$\frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_1 \partial \sigma_2^2} = \frac{n\rho(\bar{y} - \mu_2)}{2\sigma_1 \sigma_2^3(1 - \rho^2)} \stackrel{\text{def}}{=} d_{14},$$

$$\frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_1 \partial \rho} = \frac{2\rho n(\bar{x} - \mu_1)}{\sigma_1^2(1 - \rho^2)^2} - \frac{n(1 + \rho^2)(\bar{y} - \mu_2)}{\sigma_1 \sigma_2(1 - \rho^2)^2} \stackrel{\text{def}}{=} d_{15}.$$

- (ii) In d_{1i} , $i = 1, \dots, 5$, replace the parameters involved by their respective estimates, and denote the resulting expressions by \tilde{d}_{1i} , $i = 1, \dots, 5$. Then show that:

$$\tilde{d}_{11} = -\frac{nS_y}{S_x S_y - S_{xy}^2}, \quad \tilde{d}_{12} = \frac{nS_{xy}}{S_x S_y - S_{xy}^2}, \quad \tilde{d}_{13} = \tilde{d}_{14} = \tilde{d}_{15} = 0,$$

where S_x , S_y , and S_{xy} are given in Exercise 1.12(iv).

- (iii) Work as in part (i) in order to show that:

$$d_{21} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_2 \partial \mu_1} = \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_1 \partial \mu_2} = d_{12}, \quad d_{22} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_2^2} = -\frac{n}{\sigma_2^2(1 - \rho^2)},$$

$$d_{23} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_2 \partial \sigma_1^2} = \frac{n\rho(\bar{x} - \mu_1)}{2\sigma_1^3 \sigma_2(1 - \rho^2)},$$

$$d_{24} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_2 \partial \sigma_2^2} = -\frac{n(\bar{y} - \mu_2)}{\sigma_2^4(1 - \rho^2)} + \frac{n\rho(\bar{x} - \mu_1)}{2\sigma_1\sigma_2^3(1 - \rho^2)},$$

$$d_{25} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \mu_2 \partial \rho} = \frac{2\rho n(\bar{y} - \mu_2)}{\sigma_2^2(1 - \rho^2)^2} - \frac{n(1 + \rho^2)(\bar{x} - \mu_1)}{\sigma_1\sigma_2(1 - \rho^2)^2}.$$

(iv) In part (iii), replace the parameters by their respective estimates and denote by $\tilde{d}_{2i}, i = 1, \dots, 5$ the resulting expressions. Then, show that:

$$\tilde{d}_{21} = \tilde{d}_{12}, \quad \tilde{d}_{22} = -\frac{nS_x}{S_x S_y - S_{xy}^2}, \quad \text{and} \quad \tilde{d}_{23} = \tilde{d}_{24} = \tilde{d}_{25} = 0.$$

(v) Work as in parts (i) and (iii), and use analogous notation in order to obtain:

$$d_{31} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_1^2 \partial \mu_1} = d_{13}, \quad d_{32} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_1^2 \partial \mu_2} = d_{23},$$

$$d_{33} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial (\sigma_1^2)^2} = \frac{n}{2\sigma_1^4} - \frac{\sum_{i=1}^n (x_i - \mu_1)^2}{\sigma_1^6(1 - \rho^2)} + \frac{3\rho \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{4\sigma_1^5\sigma_2(1 - \rho^2)},$$

$$d_{34} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_1^2 \partial \sigma_2^2} = \frac{\rho \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{4\sigma_1^3\sigma_2^3(1 - \rho^2)},$$

$$d_{35} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_1^2 \partial \rho} = \frac{\rho \sum_{i=1}^n (x_i - \mu_1)^2}{\sigma_1^4(1 - \rho^2)^2} - \frac{(1 + \rho^2) \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{2\sigma_1^3\sigma_2(1 - \rho^2)^2}.$$

(vi) Work as in parts (ii) and (iv), and use analogous notation in order to obtain: $\tilde{d}_{31} = \tilde{d}_{32} = 0$, and

$$\tilde{d}_{33} = -\frac{n(2S_x S_y - S_{xy}^2)}{4S_x^2(S_x S_y - S_{xy}^2)}, \quad \tilde{d}_{34} = \frac{nS_{xy}^2}{4S_x S_y(S_x S_y - S_{xy}^2)},$$

$$\tilde{d}_{35} = \frac{nS_y^{1/2} S_{xy}}{2S_x^{1/2}(S_x S_y - S_{xy}^2)}.$$

(vii) Work as in part (v) in order to obtain:

$$d_{41} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_2^2 \partial \mu_1} = d_{14}, \quad d_{42} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_2^2 \partial \mu_2} = d_{24}, \quad d_{43} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_2^2 \partial \sigma_1^2} = d_{34},$$

$$d_{44} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial (\sigma_2^2)^2} = \frac{n}{2\sigma_2^4} - \frac{\sum_{i=1}^n (y_i - \mu_2)^2}{\sigma_2^6(1 - \rho^2)} + \frac{3\rho \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{4\sigma_1\sigma_2^5(1 - \rho^2)},$$

$$d_{45} \stackrel{\text{def}}{=} \frac{\partial^2 \lambda(\boldsymbol{\theta})}{\partial \sigma_2^2 \partial \rho} = \frac{\rho \sum_{i=1}^n (y_i - \mu_2)^2}{\sigma_2^4(1 - \rho^2)^2} - \frac{(1 + \rho^2) \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2)}{2\sigma_1\sigma_2^3(1 - \rho^2)^2}.$$

(viii) Work as in part (vi) in order to get:

$$\tilde{d}_{41} = \tilde{d}_{42} = 0, \quad \tilde{d}_{43} = \tilde{d}_{34}, \quad \text{and}$$

$$\tilde{d}_{44} = -\frac{n(2S_x S_y - S_{xy}^2)}{4S_y^2(S_x S_y - S_{xy}^2)}, \quad \tilde{d}_{45} = \frac{nS_x^{1/2} S_{xy}}{2S_y^{1/2}(S_x S_y - S_{xy}^2)}.$$

(ix) Work as in part (v), and use analogous notation in order to get:

$$d_{51} = d_{15}, \quad d_{52} = d_{25}, \quad d_{53} = d_{35}, \quad d_{54} = d_{45}, \quad \text{and}$$

$$d_{55} = \frac{n(1 + \rho^2)}{(1 - \rho^2)^2} - \frac{1 + 3\rho^2}{(1 - \rho^2)^3} \left[\frac{1}{\sigma_1^2} \sum_{i=1}^n (x_i - \mu_1)^2 + \frac{1}{\sigma_2^2} \sum_{i=1}^n (y_i - \mu_2)^2 \right. \\ \left. + \frac{2\rho(3 + \rho^2)}{\sigma_1\sigma_2(1 - \rho^2)^3} \sum_{i=1}^n (x_i - \mu_1)(y_i - \mu_2) \right].$$

(x) Work as in part (vi) in order to obtain:

$$\tilde{d}_{51} = \tilde{d}_{15}, \quad \tilde{d}_{52} = \tilde{d}_{25}, \quad \tilde{d}_{53} = \tilde{d}_{35}, \quad \tilde{d}_{54} = \tilde{d}_{45}, \quad \text{and}$$

$$\tilde{d}_{55} = -\frac{nS_xS_y(S_xS_y + S_{xy}^2)}{(S_xS_y - S_{xy}^2)^2}.$$

1.14 In this exercise, it is shown that the solution values of $\tilde{\mu}_1 = \bar{x}$, $\tilde{\mu}_2 = \bar{y}$, $\tilde{\sigma}_1^2 = S_x$, $\tilde{\sigma}_2^2 = S_y$, and $\tilde{\rho} = S_{xy}/S_x^{1/2}S_y^{1/2}$ are, indeed, the MLE's of the respective parameters. To this effect, set \tilde{D} for the determinant

$$\tilde{D} = \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} & \tilde{d}_{13} & \tilde{d}_{14} & \tilde{d}_{15} \\ \tilde{d}_{21} & \tilde{d}_{22} & \tilde{d}_{23} & \tilde{d}_{24} & \tilde{d}_{25} \\ \tilde{d}_{31} & \tilde{d}_{32} & \tilde{d}_{33} & \tilde{d}_{34} & \tilde{d}_{35} \\ \tilde{d}_{41} & \tilde{d}_{42} & \tilde{d}_{43} & \tilde{d}_{44} & \tilde{d}_{45} \\ \tilde{d}_{51} & \tilde{d}_{52} & \tilde{d}_{53} & \tilde{d}_{54} & \tilde{d}_{55} \end{vmatrix},$$

and let D_i be the determinants taken from \tilde{D} by eliminating the last $5 - i$, $i = 1, \dots, 5$ rows and columns; also, set $\tilde{D}_0 = 1$. Thus,

$$\tilde{D}_1 = \tilde{d}_{11}, \quad \tilde{D}_2 = \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} \\ \tilde{d}_{21} & \tilde{d}_{22} \end{vmatrix},$$

$$\tilde{D}_3 = \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} & \tilde{d}_{13} \\ \tilde{d}_{21} & \tilde{d}_{22} & \tilde{d}_{23} \\ \tilde{d}_{31} & \tilde{d}_{32} & \tilde{d}_{33} \end{vmatrix}, \quad \tilde{D}_4 = \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} & \tilde{d}_{13} & \tilde{d}_{14} \\ \tilde{d}_{21} & \tilde{d}_{22} & \tilde{d}_{23} & \tilde{d}_{24} \\ \tilde{d}_{31} & \tilde{d}_{32} & \tilde{d}_{33} & \tilde{d}_{34} \\ \tilde{d}_{41} & \tilde{d}_{42} & \tilde{d}_{43} & \tilde{d}_{44} \end{vmatrix},$$

and $\tilde{D}_5 = \tilde{D}$.

(i) Use parts (ii), (iv), (vi), (viii), and (x) in Exercise 1.13 in order to conclude that the determinants $\tilde{D}(=\tilde{D}_5)$ and \tilde{D}_i , $i = 1, \dots, 4$ take the following forms:

$$\tilde{D} = \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} & 0 & 0 & 0 \\ \tilde{d}_{21} & \tilde{d}_{22} & 0 & 0 & 0 \\ 0 & 0 & \tilde{d}_{33} & \tilde{d}_{34} & \tilde{d}_{35} \\ 0 & 0 & \tilde{d}_{43} & \tilde{d}_{44} & \tilde{d}_{45} \\ 0 & 0 & \tilde{d}_{53} & \tilde{d}_{54} & \tilde{d}_{55} \end{vmatrix},$$

$$\begin{aligned} \tilde{D}_1 &= \tilde{d}_{11}, & \tilde{D}_2 &= \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} \\ \tilde{d}_{21} & \tilde{d}_{22} \end{vmatrix}, \\ \tilde{D}_3 &= \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} & 0 \\ \tilde{d}_{21} & \tilde{d}_{22} & 0 \\ 0 & 0 & \tilde{d}_{33} \end{vmatrix}, & \tilde{D}_4 &= \begin{vmatrix} \tilde{d}_{11} & \tilde{d}_{12} & 0 & 0 \\ \tilde{d}_{21} & \tilde{d}_{22} & 0 & 0 \\ 0 & 0 & \tilde{d}_{33} & \tilde{d}_{34} \\ 0 & 0 & \tilde{d}_{43} & \tilde{d}_{44} \end{vmatrix}. \end{aligned}$$

- (ii) Expand the determinants \tilde{D}_i , $i = 1, \dots, 4$, and also use parts (iv) and (viii) of Exercise 1.13 in order to obtain:

$$\begin{aligned} \tilde{D}_1 &= \tilde{d}_{11}, & \tilde{D}_2 &= \tilde{d}_{11}\tilde{d}_{22} - (\tilde{d}_{12})^2, & \tilde{D}_3 &= \tilde{d}_{33}\tilde{D}_2, \\ \tilde{D}_4 &= [\tilde{d}_{33}\tilde{d}_{44} - (\tilde{d}_{34})^2]\tilde{D}_2. \end{aligned}$$

- (iii) Expand the determinant $\tilde{D}_5 (= \tilde{D})$, and also use parts (viii) and (x) of Exercise 1.13 in order to get:

$$\tilde{D}_5 = \tilde{D}_2(\tilde{d}_{33}A - \tilde{d}_{34}B + \tilde{d}_{35}C),$$

where

$$A = \tilde{d}_{44}\tilde{d}_{55} - (\tilde{d}_{45})^2, \quad B = \tilde{d}_{34}\tilde{d}_{55} - \tilde{d}_{45}\tilde{d}_{35}, \quad C = \tilde{d}_{34}\tilde{d}_{45} - \tilde{d}_{44}\tilde{d}_{35}.$$

- (iv) For convenience, set: $S_x = \alpha$, $S_y = \beta$, $S_{xy} = \gamma$, and $S_x S_y - S_{xy}^2 = \alpha\beta - \gamma^2 = \delta$, so that $\alpha, \beta > 0$ and also $\delta > 0$ by the Cauchy–Schwarz inequality (see Theorem 1(ii) in Chapter 4). Then use parts (ii), (vi), and (viii) in Exercise 1.13 in order to express the determinants \tilde{D}_i , $i = 1, \dots, 4$ in part (ii) in terms of α, β, γ , and δ and obtain:

$$\begin{aligned} \tilde{D}_1 &= -\frac{n\beta}{\delta}, & \tilde{D}_2 &= \frac{n^2}{\delta}, & \tilde{D}_3 &= -\frac{n^3(2\alpha\beta - \gamma^2)}{4\alpha^2\delta^2} = -\frac{n^3(\alpha\beta + \delta)}{4\alpha^2\delta^2}, \\ \tilde{D}_4 &= \frac{n^4}{4\alpha\beta\delta^2}. \end{aligned}$$

- (v) Use the definition of A, B , and C in part (iii), as well as the expressions of $\tilde{d}_{34}, \tilde{d}_{35}, \tilde{d}_{44}, \tilde{d}_{45}$, and \tilde{d}_{55} given in parts (vi), (viii), and (x) of Exercise 1.13, in conjunction with the notation introduced in part (iv) of the present exercise, in order to show that:

$$A = \frac{\alpha^3\beta n^2}{2\delta^3}, \quad B = -\frac{\alpha\beta\gamma^2 n^2}{2\delta^3}, \quad C = \frac{\alpha^{1/2}\gamma n^2}{4\beta^{1/2}\delta^2}.$$

- (vi) Use parts (iii) and (v) here, and parts (ii) and (iv) in Exercise 1.13 in order to obtain:

$$\tilde{D}_5 = \tilde{D}_2(\tilde{d}_{33}A - \tilde{d}_{34}B + \tilde{d}_{35}C) = \frac{n^2}{\delta} \left(-\frac{\alpha\beta n^3}{4\delta^3} \right) = -\frac{\alpha\beta n^5}{4\delta^4}.$$

(vii) From parts (iv) and (vi) and the fact that $\bar{D}_0 = 1$, conclude that:

$$\bar{D}_0 > 0, \quad \bar{D}_1 < 0, \quad \bar{D}_2 > 0, \quad \bar{D}_3 < 0, \quad \bar{D}_4 > 0, \quad \text{and} \quad \bar{D}_5 < 0.$$

Then use a calculus result about the maximum of a function in more than one variable (see, e.g., Theorem 7.9, pages 151–152, in the book *Mathematical Analysis*, Addison-Wesley (1957), by T. M. Apostol) in order to conclude that $\bar{\mu}_1, \bar{\mu}_2, \bar{\sigma}_1^2, \bar{\sigma}_2^2$, and $\bar{\rho}$ are, indeed, the MLE's of the respective parameters; i.e.,

$$\hat{\mu}_1 = \bar{x}, \quad \hat{\mu}_2 = \bar{y}, \quad \hat{\sigma}_1^2 = S_x, \quad \hat{\sigma}_2^2 = S_y, \quad \hat{\rho} = S_{xy}/S_x^{1/2}S_y^{1/2},$$

where

$$S_x = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2, \quad S_y = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2,$$

$$S_{xy} = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}).$$

9.2 Some Properties of Maximum Likelihood Estimates

Refer to Example 3 and suppose that we are interested in estimating the probability that 0 events occur; that is, $P_\theta(X = 0) = \frac{e^{-\theta} \theta^0}{0!} = e^{-\theta}$, call it $g_1(\theta)$. Thus, the estimated quantity is a function of θ rather than θ itself. Next, refer to Example 4 and recall that, if $X \sim f(x; \theta) = \theta e^{-\theta x}$, $x > 0$, then $E_\theta X = 1/\theta$. Thus, in this case it would be, perhaps, more reasonable to estimate $1/\theta$ and call it $g_2(\theta)$, rather than θ . Finally, refer to Examples 5(ii) and 7, and consider the problem of estimating the s.d. $\sigma = +\sqrt{\sigma^2}$ rather than the variance σ^2 . This is quite meaningful, since, as we know, the s.d. is used as the yardstick for measuring distances from the mean. In this last case, set $g_3(\sigma) = +\sqrt{\sigma^2}$.

The functions g_1, g_2 , and g_3 have the common characteristic that they are one-to-one functions of the parameter involved. The estimation problems described above are then formulated in a unified way as follows.

THEOREM 1

Let $\hat{\theta} = \hat{\theta}(\mathbf{x})$ be the MLE of θ on the basis of the observed values x_1, \dots, x_n of the random sample X_1, \dots, X_n from the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \Re$. Also, let $\theta^* = g(\theta)$ be a one-to-one function defined on Ω onto $\Omega^* \subseteq \Re$. Then the MLE of θ^* , $\hat{\theta}^*(\mathbf{x})$, is given by $\hat{\theta}^*(\mathbf{x}) = g[\hat{\theta}(\mathbf{x})]$.

PROOF The equation $\theta^* = g(\theta)$ can be solved for θ , on the basis of the assumption made, and let $\theta = g^{-1}(\theta^*)$. Then

$$L(\theta | \mathbf{x}) = L[g^{-1}(\theta^*) | \mathbf{x}] = L^*(\theta^* | \mathbf{x}), \text{ say.}$$

Thus,

$$\max\{L(\theta | \mathbf{x}); \theta \in \Omega\} = \max\{L^*(\theta^* | \mathbf{x}); \theta^* \in \Omega^*\}. \quad (4)$$

Since the left-hand side in (4) is maximized for $\hat{\theta} = \hat{\theta}(\mathbf{x})$, clearly, the right-hand side is maximized for $\hat{\theta}^* = g(\hat{\theta})$. ▲

REMARK 3 On the basis of Theorem 1, then, we have: The MLE of $\exp(-\theta)$ is $\exp(-\bar{x})$; the MLE of $1/\theta$ is $1/\bar{x}$; and the MLE of σ is $[\frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2]^{1/2}$ for Example 5(ii), and $[\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2]^{1/2}$ for Example 7.

However, in as simple a case as that of the $B(1, \theta)$ distribution, the function g in Theorem 1 may not be one-to-one, and yet we can construct the MLE of $\theta^* = g(\theta)$. This is the content of the next theorem.

THEOREM 2

Let $\hat{\theta} = \hat{\theta}(\mathbf{x})$ be the MLE of θ on the basis of the observed values x_1, \dots, x_n of the random sample X_1, \dots, X_n from the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$. Also, let $\theta^* = g(\theta)$ be an arbitrary function defined on Ω into $\Omega^* \subseteq \mathfrak{R}$, where, without loss of generality, we may assume that Ω^* is the range of g , so that the function g is defined on Ω onto Ω^* . Then the MLE of θ^* , $\hat{\theta}^*(\mathbf{x})$, is still given by $\hat{\theta}^*(\mathbf{x}) = g[\hat{\theta}(\mathbf{x})]$.

PROOF For each $\theta^* \in \Omega^*$, there may be several θ in Ω mapped to the same θ^* under g . Let Ω_{θ^*} be the set of all such θ 's; i.e.,

$$\Omega_{\theta^*} = \{\theta \in \Omega; g(\theta) = \theta^*\}.$$

On Ω^* , define the real-valued function L^* by:

$$L^*(\theta^*) = \sup\{L(\theta); \theta \in \Omega_{\theta^*}\}.$$

The function L^* may be called the *likelihood function induced by g* . Now, since g is a function, it follows that $g(\hat{\theta}) = \hat{\theta}^*$ for a unique $\hat{\theta}^*$ in Ω^* , and $L^*(\hat{\theta}^*) = L(\hat{\theta})$ from the definition of L^* . Finally, for every $\theta^* \in \Omega^*$,

$$L^*(\theta^*) = \sup\{L(\theta); \theta \in \Omega_{\theta^*}\} \leq \max\{L(\theta); \theta \in \Omega\} = L(\hat{\theta}) = L^*(\hat{\theta}^*).$$

This last inequality justifies calling $\hat{\theta}^* = g(\hat{\theta})$ the MLE of θ^* . ▲

(Theorem 2 was adapted from a result established by Peter W. Zehna in the *Annals of Mathematical Statistics*, Vol. 37 (1966), page 744.)

EXAMPLE 10

Refer to Example 2 and determine the MLE $\hat{g}(\theta)$ of the function $g(\theta) = \theta(1 - \theta)$.

DISCUSSION Here the function $g: (0, 1) \rightarrow (0, \frac{1}{4})$ is not one-to-one. However, by Theorem 2, the MLE $\hat{g}(\theta) = \bar{x}(1 - \bar{x})$, since $\hat{\theta} = \bar{x}$.

REMARK 4 Theorem 1 is, of course, a special case of Theorem 2. A suitable version of Theorem 2 holds for multidimensional parameters. This property of the MLE is referred to as the *invariance property* of the MLE for obvious reasons.

Reviewing the examples in the previous section, we see that the data x_1, \dots, x_n are entering into the MLE's in a compactified form, more, precisely,

as a real-valued quantity. From this point on, this is all we have at our disposal; or, perhaps, that is all that was revealed to us by those who collected the data. In contemplating this situation, one cannot help but wonder what we are missing by knowing, for example, only \bar{x} rather than the complete array of data x_1, \dots, x_n . The almost shocking fact of the matter is that, in general, we are missing absolutely nothing, in terms of information carried by the data x_1, \dots, x_n , provided the data are condensed in the right way. This is precisely the concept of sufficiency to be introduced below. For a motivation of the definition, consider the following example.

EXAMPLE 11

In Example 1, each x_i , $i = 1, \dots, 10$ takes on the value either 0 or 1, and we are given that $\bar{x} = 0.6$. There are $2^{10} = 1,024$ arrangements of 10 0's or 1's with respective probabilities given by $\theta^{x_i}(1 - \theta)^{1-x_i}$ for each one of the 1,024 arrangements of 0's and 1's. These probabilities, of course, depend on θ . Now, restrict attention to the $\binom{10}{6} = 210$ arrangements of 0's and 1's only, which produce a sum of 6 or an average of 0.6; their probability is $210\theta^6(1 - \theta)^4$. Finally, calculate the conditional probability of each one of these arrangements, given that the average is 0.6 or that the sum is 6. In other words, calculate

$$P_\theta(X_i = x_i, i = 1, \dots, 10 \mid T = 6), \quad T = \sum_{i=1}^{10} X_i. \quad (5)$$

Suppose that all these conditional probabilities have the same value, which, in addition, is independent of θ . This would imply two things: First, given that the sum is 6, all possible arrangements, summing up to 6, have the same probability independent of the location of occurrences of 1's; and second, this probability has the same numerical value for all values of θ in $(0, 1)$. So, from a probabilistic viewpoint, given the information that the sum is 6, it does not really matter either what the arrangement is or what the value of θ is; we can reconstruct each one of all those arrangements giving sum 6 by choosing each one of the 210 possible arrangements, with probability $1/210$ each. It is in this sense that, restricting ourselves to the sum and ignoring or not knowing the individual values, we deprive ourselves of no information about θ .

We proceed now with the calculation of the probabilities in (5). Although we can refer to existing results, let us derive the probabilities here.

$$\begin{aligned} P_\theta(X_i = x_i, i = 1, \dots, 10 \mid T = 6) &= P_\theta(X_i = x_i, i = 1, \dots, 10, T = 6) / P_\theta(T = 6) \\ &= P_\theta(X_i = x_i, i = 1, \dots, 10) / P_\theta(T = 6) \\ &\quad (\text{since } X_i = x_i, i = 1, \dots, 10 \text{ implies } T = 6) \\ &= \theta^6(1 - \theta)^4 / \binom{10}{6} \theta^6(1 - \theta)^4 \quad (\text{since } T \sim B(10, \theta)) \\ &= 1 / \binom{10}{6} = 1/210 (\simeq 0.005). \end{aligned}$$

Thus, what was supposed above is, actually, true.

This example and the elaboration associated with it lead to the following definition of sufficiency.

DEFINITION 1

Let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and let $T = T(X_1, \dots, X_n)$ be a statistic (i.e., a known function of the X_i 's). Then, if the conditional distribution of the X_i 's, given $T = t$, does not depend on θ , we say that T is a *sufficient* statistic for θ .

REMARK 5 If g is a real-valued one-to-one function defined on the range of T , it is clear that knowing T is equivalent to knowing $T^* = g(T)$, and vice versa. Thus, if T is a sufficient statistic for θ , so is T^* . In particular, if $T = \sum_{i=1}^n X_i$ or $\frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2$ (or $\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$) is a sufficient statistic for θ so is \bar{X} or $\sum_{i=1}^n (X_i - \mu)^2$ (or $\sum_{i=1}^n (X_i - \bar{X})^2$).

REMARK 6 The definition given for one parameter also applies for more than one parameter, but then we also need a multidimensional sufficient statistic, usually, with dimensionality equal to the number of the parameters. In all cases, we often use simply the term “*sufficient*” instead of “*sufficient statistic(s) for θ* ,” if no confusion is possible.

As is often the case, definitions do not lend themselves easily to identifying the quantity defined. This is also the case in Definition 1. A sufficient statistic is, actually, found by way of the theorem stated below.

THEOREM 3

(Fisher-Neyman Factorization Theorem) Let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and let $T = T(X_1, \dots, X_n)$ be a statistic. Then T is a sufficient statistic for θ , if and only if the joint p.d.f. of the X_i 's may be written as follows:

$$f_{X_1, \dots, X_n}(x_1, \dots, x_n; \theta) = g[T(x_1, \dots, x_n); \theta]h(x_1, \dots, x_n). \quad (6)$$

The way this theorem applies is the following: One writes out the joint p.d.f. of the X_i 's and then one tries to rewrite it as the product of two factors, one factor, $g[T(x_1, \dots, x_n); \theta]$, which contains the x_i 's only through the function $T(x_1, \dots, x_n)$ and the parameter θ , and another factor, $h(x_1, \dots, x_n)$, which involves the x_i 's in whatever form but not θ in any form.

REMARK 7 The theorem just stated also holds for multidimensional parameters θ , but then the statistic T is also multidimensional, usually of the same dimension as that of θ . A rigorous proof of the theorem can be given, at least for the case of discrete X_i 's, but we choose to omit it.

In all of the Examples 2–10, the MLE's are, actually, sufficient statistics or functions thereof, as demonstrated below. This fact should certainly reinforce our appreciation for these MLE's.

APPLICATION In Example 2, the p.d.f. is written as follows in a compact form $f(x_i; \theta) = \theta^{x_i}(1 - \theta)^{1-x_i} I_{(0,1)}(x_i)$, so that the joint p.d.f. becomes:

$$L(\theta | \mathbf{x}) = \theta^t(1 - \theta)^{n-t} \times \prod_{i=1}^n I_{(0,1)}(x_i), \quad t = \sum_{i=1}^n x_i.$$

Then $g[T(x_1, \dots, x_n); \theta] = \theta^t(1 - \theta)^{n-t}$, and $h(x_1, \dots, x_n) = \prod_{i=1}^n I_{(0,1)}(x_i)$. It follows that $T = \sum_{i=1}^n X_i$ is sufficient and so is $\frac{T}{n} = \bar{X}$.

Examples 3 and 4 are treated similarly.

In Example 5(i),

$$L(\theta | \mathbf{x}) = \exp \left[\frac{n\mu(2\bar{x} - \mu)}{2\sigma^2} \right] \times \left(\frac{1}{\sqrt{2\pi\sigma}} \right)^n \exp \left(-\frac{1}{2\sigma^2} \sum_{i=1}^n x_i^2 \right),$$

so that \bar{X} is sufficient for μ . Likewise, in Example 5(ii),

$$L(\sigma^2 | \mathbf{x}) = \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^n \exp \left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right] \times 1,$$

so that $\sum_{i=1}^n (X_i - \mu)^2$ is sufficient for σ^2 and so is $\frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2$.

Example 6 is treated similarly.

In Example 7,

$$L(\mu, \sigma^2 | \mathbf{x}) = \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^n \exp \left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \bar{x})^2 - \frac{1}{2\sigma^2} n(\bar{x} - \mu)^2 \right] \times 1$$

because $\sum_{i=1}^n (x_i - \mu)^2 = \sum_{i=1}^n [(x_i - \bar{x}) + (\bar{x} - \mu)]^2 = \sum_{i=1}^n (x_i - \bar{x})^2 + n(\bar{x} - \mu)^2$. It follows that the pair of statistics $(\bar{X}, \sum_{i=1}^n (X_i - \bar{X})^2)$ is sufficient for the pair of parameters (μ, σ^2) .

Examples 8, 9, and 10 are treated similarly.

REMARK 8 Under certain regularity conditions, it is always the case that a MLE is only a function of a sufficient statistic.

Here is another example, in four parts, where a sufficient statistic is determined by way of Theorem 3.

EXAMPLE 12

On the basis of a random sample of size n , X_1, \dots, X_n , from each one of the p.d.f.'s given below with observed values x_1, \dots, x_n , determine a sufficient statistic for θ .

- (i) $f(x; \theta) = \frac{\theta}{x^{\theta+1}}, \quad x \geq 1, \quad \theta \in \Omega = (0, \infty)$.
- (ii) $f(x; \theta) = \frac{x}{\theta} e^{-x^2/2\theta}, \quad x > 0, \quad \theta \in \Omega = (0, \infty)$.
- (iii) $f(x; \theta) = (1 + \theta)x^\theta, \quad 0 < x < 1, \quad \theta \in \Omega = (-1, \infty)$.
- (iv) $f(x; \theta) = \frac{\theta}{x^2}, \quad x \geq \theta, \quad \theta \in \Omega = (0, \infty)$.

DISCUSSION In the first place, the functions given above are, indeed, p.d.f.'s (see Exercise 2.12). Next, rewriting each one of the p.d.f.'s by using the indicator function, we have:

(i) $f(x; \theta) = \frac{\theta}{x^{\theta+1}} I_{[1, \infty)}(x)$, so that:

$$\prod_{i=1}^n f(x_i; \theta) = \frac{\theta^n}{\left(\prod_{i=1}^n x_i\right)^{\theta+1}} \times \prod_{i=1}^n I_{[1, \infty)}(x_i) = \frac{\theta^n}{\left(\prod_{i=1}^n x_i\right)^{\theta+1}} \times I_{[1, \infty)}(x_{(1)}),$$

and therefore $\prod_{i=1}^n X_i$ is sufficient for θ .

(ii) $f(x; \theta) = \frac{x}{\theta} e^{-x^2/2\theta} I_{(0, \infty)}(x)$, so that:

$$\begin{aligned} \prod_{i=1}^n f(x_i; \theta) &= \frac{1}{\theta^n} \prod_{i=1}^n x_i e^{-\frac{1}{2\theta} \sum_{i=1}^n x_i^2} \prod_{i=1}^n I_{(0, \infty)}(x_i) \\ &= \frac{1}{\theta^n} e^{-\frac{1}{2\theta} \sum_{i=1}^n x_i^2} \times \left(\prod_{i=1}^n x_i \right) I_{(0, \infty)}(x_{(1)}), \end{aligned}$$

and therefore $\sum_{i=1}^n X_i^2$ is sufficient for θ .

(iii) $f(x; \theta) = (1 + \theta)x^\theta I_{(0, 1)}(x)$, so that

$$\begin{aligned} \prod_{i=1}^n f(x_i; \theta) &= (1 + \theta)^n \left(\prod_{i=1}^n x_i \right)^\theta \prod_{i=1}^n I_{(0, 1)}(x_i) \\ &= (1 + \theta)^n \left(\prod_{i=1}^n x_i \right)^\theta \times I_{(0, 1)}(x_{(1)}) I_{(0, 1)}(x_{(n)}), \end{aligned}$$

and therefore $\prod_{i=1}^n X_i$ is sufficient for θ .

(iv) $f(x; \theta) = \frac{\theta}{x^2} I_{[\theta, \infty)}(x)$, so that:

$$\prod_{i=1}^n f(x_i; \theta) = \frac{\theta^n}{\left(\prod_{i=1}^n x_i^2\right)} \prod_{i=1}^n I_{[\theta, \infty)}(x_i) = \theta^n I_{[\theta, \infty)}(x_{(1)}) \times \frac{1}{\left(\prod_{i=1}^n x_i^2\right)},$$

and therefore $X_{(1)}$ is sufficient for θ .

This section is concluded with two desirable asymptotic properties of a MLE. The first is consistency (in the probability sense), and the other is asymptotic normality. However, we will not bother either to list the conditions needed or to justify the results stated.

THEOREM 4

Let $\hat{\theta}_n = \hat{\theta}_n(X_1, \dots, X_n)$ be the MLE of $\theta \in \Omega \subseteq \Re$ based on the random sample X_1, \dots, X_n with p.d.f. $f(\cdot, \theta)$. Then, under certain regularity conditions, $\{\hat{\theta}_n\}$ is consistent in the probability sense; that is, $\hat{\theta}_n \rightarrow \theta$ in P_θ -probability as $n \rightarrow \infty$.

The usefulness of this result is, of course, that, for sufficiently large n , $\hat{\theta}_n$ is as close to (the unknown) θ as we please with probability as close to 1 as we desire. The tool usually employed in establishing Theorem 4 is either the Weak Law of Large Numbers (WLLN) or the Tchebichev inequality. In exercises at the end of this section, the validity of Theorem 4 is illustrated in some of the examples of the previous section.

THEOREM 5

In the notation of Theorem 4, and under suitable regularity conditions, the MLE $\hat{\theta}_n$ is asymptotically normal. More precisely, under P_θ -probability,

$$\sqrt{n}(\hat{\theta}_n - \theta) \xrightarrow{d} N(0, \sigma_\theta^2), \quad \text{as } n \rightarrow \infty,$$

$$\text{where } \sigma_\theta^2 = 1/I(\theta) \text{ and } I(\theta) = E_\theta \left[\frac{\partial}{\partial \theta} \log f(X; \theta) \right]^2, \quad X \sim f(\cdot; \theta). \quad (7)$$

To state it loosely, $\hat{\theta}_n \simeq N(\theta, \sigma_\theta^2/n)$ for sufficiently large n . That is, the MLE $\hat{\theta}_n$ is approximately Normally distributed around θ , and therefore various probabilities related to it may be approximately calculated in principle. The justification of the theorem is done by using a Taylor expansion of the derivative $\frac{\partial}{\partial \theta} \log L(\theta | \mathbf{X})$ up to terms of third-order, employing the fact that $\frac{\partial}{\partial \theta} \log L(\theta | \mathbf{X}) |_{\theta=\hat{\theta}_n} = 0$, and suitably utilizing the WLLN and the Central Limit Theorem (CLT). For some applications of this theorem, see, e.g., Example 21, page 324, of the book *A Course in Mathematical Statistics*, 2nd edition, Academic Press (1977), by G. G. Roussas.

REMARK 9 The quantity $I(\theta)$ is referred to as the *Fisher information* carried by the random sample X_1, \dots, X_n about the parameter θ . A justification for the “information” stems by the fact that $\sigma_\theta^2 = 1/I(\theta)$, so that the larger $I(\theta)$ is the smaller the variance σ_θ^2 is, and therefore the more concentrated $\hat{\theta}$ is about θ . The opposite happens for small values of $I(\theta)$.

Exercises

2.1 Let X_1, \dots, X_n be i.i.d. r.v.’s with the Negative Exponential p.d.f. $f(x; \theta) = \theta e^{-\theta x}$, $x > 0$, $\theta \in \Omega = (0, \infty)$. Then:

(i) Show that $1/\bar{X}$ is the MLE of θ .

(ii) Use Theorem 1 in order to conclude that the MLE of θ^* in the parameterized form $f(x; \theta^*) = \frac{1}{\theta^*} e^{-x/\theta^*}$, $x > 0$, is \bar{X} .

2.2 Let X be a r.v. denoting the life span of an equipment. Then the *reliability* of the equipment at time x , $R(x)$, is defined as the probability that $X > x$;

i.e., $R(x) = P(X > x)$. Now, suppose that X has the Negative Exponential p.d.f. $f(x; \theta) = \frac{1}{\theta}e^{-x/\theta}$, $x > 0$, $\theta \in \Omega = (0, \infty)$. Then:

- (i) Calculate the reliability $R(x; \theta)$ based on this r.v. X .
- (ii) Use Theorem 1 in order to determine the MLE of $R(x; \theta)$, on the basis of a random sample X_1, \dots, X_n from the underlying p.d.f.

2.3 Let X be a r.v. describing the lifetime of a certain equipment, and suppose that the p.d.f. of X is $f(x; \theta) = \theta e^{-\theta x}$, $x > 0$, $\theta \in \Omega = (0, \infty)$.

- (i) Show that the probability that X is greater than or equal to t time units is $g(\theta) = e^{-t\theta}$.
- (ii) We know (see Exercise 2.1) that the MLE of θ , based on a random sample of size n from the above p.d.f., is $\hat{\theta} = 1/\bar{X}$. Then determine the MLE of $g(\theta)$.

2.4 Consider the independent r.v.'s X_1, \dots, X_n with the Weibull p.d.f. $f(x; \theta) = \frac{\gamma}{\theta} x^{\gamma-1} \exp(-x^\gamma/\theta)$, $x > 0$, $\theta \in \Omega = (0, \infty)$, $\gamma > 0$ known, and:

- (i) Show that $\hat{\theta} = (\sum_{i=1}^n X_i^\gamma)/n$ is the MLE of θ .
- (ii) Take $\gamma = 1$ and relate the result in part (i) to the result in Exercise 2.1(ii).

2.5 Let X_1, \dots, X_n be a random sample of size n from the $N(\mu, \sigma^2)$ distribution, where both μ and σ^2 are unknown. Set $\theta = (\mu, \sigma^2)$ and let p be a (known) number with $0 < p < 1$. Then:

- (i) Show that the point c for which $P_\theta(\bar{X} \leq c) = p$ is given by: $c = \mu + \frac{\sigma}{\sqrt{n}} \Phi^{-1}(p)$.
- (ii) Given that the MLE's of μ and σ^2 are, respectively, $\hat{\mu} = \bar{X}$ and $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$, determine the MLE of c , call it \hat{c} .
- (iii) Express \hat{c} in terms of the X_i 's, if $n = 25$ and $p = 0.95$.

2.6 (i) Show that the function $f(x; \theta) = \theta x^{-(\theta+1)}$, $x \geq 1$, $\theta \in \Omega = (0, \infty)$ is a p.d.f.

- (ii) On the basis of a random sample of size n from this p.d.f., show that the statistic $X_1 \cdots X_n$ is sufficient for θ , and so is the statistic $\sum_{i=1}^n \log X_i$.

2.7 Let X be a r.v. having the Geometric p.d.f. $f(x; \theta) = \theta(1 - \theta)^{x-1}$, $x = 1, 2, \dots$, $\theta \in \Omega = (0, 1)$. Then show that X is sufficient for θ .

2.8 In reference to Exercise 1.9, show that $\sum_{i=1}^n |X_i|$ is a sufficient statistic for θ .

2.9 (i) In reference to Example 3, use Theorem 3 in order to find a sufficient statistic for θ .

- (ii) Do the same in reference to Example 4.
- (iii) Do the same in reference to Example 6(i), (ii).

2.10 Same as in Exercise 2.9 in reference to Examples 8 and 9.

2.11 Refer to Exercise 1.11, and determine:

- (i) A sufficient statistic for α when β is known.

- (ii) Also, a sufficient statistic for β when α is known.
- (iii) A set of sufficient statistics for α and β when they are both unknown.

2.12 Show that the functions (i)–(iv) given in Example 12 are, indeed, p.d.f.'s.

9.3 Uniformly Minimum Variance Unbiased Estimates

Perhaps the second most popular method of estimating a parameter is that based on the concepts of unbiasedness and variance. This method will be discussed here to a certain extent and will also be illustrated by specific examples.

To start with, let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and let us introduce the notation $U = U(X_1, \dots, X_n)$ for an estimate of θ .

DEFINITION 2

The estimate U is said to be *unbiased* if $E_\theta U = \theta$ for all $\theta \in \Omega$.

Some examples of unbiased estimates follow.

EXAMPLE 13

Let X_1, \dots, X_n be having any one of the following distributions:

- (i) $B(1, \theta)$, $\theta \in (0, 1)$. Then the sample mean \bar{X} is an unbiased estimate of θ .
- (ii) $P(\theta)$, $\theta > 0$. Then again \bar{X} is an unbiased estimate of θ . Here, since $E_\theta X_1 = \text{Var}_\theta(X_1) = \theta$, \bar{X} is also an unbiased estimate of the variance.
- (iii) $N(\theta, \sigma^2)$, $\theta \in \mathfrak{R}$, σ known. Then, once again, \bar{X} is an unbiased estimate of θ .
- (iv) $N(\mu, \theta)$, μ known, $\theta > 0$. Then the sample variance $\frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2$ is an unbiased estimate of θ . This is so because $\sum_{i=1}^n \left(\frac{X_i - \mu}{\sqrt{\theta}} \right)^2 \sim \chi_n^2$, so that

$$E_\theta \left[\sum_{i=1}^n \left(\frac{X_i - \mu}{\sqrt{\theta}} \right)^2 \right] = n \quad \text{or} \quad E_\theta \left[\frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2 \right] = \theta.$$

- (v) Gamma with $\alpha = \theta$ and $\beta = 1$. Then \bar{X} is an unbiased estimate of θ . This is so because, in the Gamma distribution, the expectation is $\alpha\beta$, so that for $\alpha = \theta$ and $\beta = 1$, $E_\theta X_1 = \theta$ and hence $E_\theta \bar{X} = \theta$.
- (vi) Gamma with $\alpha = 1$ and $\beta = \theta$, $\theta > 0$ (which gives the reparameterized Negative Exponential distribution). Then \bar{X} is an unbiased estimate of θ as explained in part (v).

EXAMPLE 14

Let X_1, \dots, X_n be a random sample from the $U(0, \theta)$ ($\theta > 0$). Determine an unbiased estimate of θ .

DISCUSSION Let $Y_n = \max(X_1, \dots, X_n)$; i.e., the largest order statistic of the X_i 's. Then, by (29) in Chapter 6, the p.d.f. of Y_n is given by:

$$g(y) = n[F(y)]^{n-1} f(y), \quad \text{for } 0 < y < \theta \text{ (and 0 otherwise),}$$

where:

$$f(y) = \frac{1}{\theta}, \quad 0 < y < \theta, \quad F(y) = \begin{cases} 0, & y \leq 0 \\ \frac{y}{\theta}, & 0 < y < \theta \\ 1, & y \geq \theta. \end{cases}$$

Then, for $0 < y < \theta$, $g(y) = n \left(\frac{y}{\theta}\right)^{n-1} \left(\frac{1}{\theta}\right) = \frac{n}{\theta^n} y^{n-1}$, so that


$$E_{\theta} Y_n = \int_0^{\theta} y \cdot \frac{n}{\theta^n} y^{n-1} dy = \frac{n}{\theta^n} \int_0^{\theta} y^n dy = \frac{n}{(n+1)\theta^n} \cdot y^{n+1} \Big|_0^{\theta} = \frac{n}{n+1} \theta.$$

It follows that $E_{\theta} \left(\frac{n+1}{n} Y_n\right) = \theta$, so that $\frac{n+1}{n} Y_n$ is an unbiased estimate of θ .

The desirability of *unbiasedness* of an estimate stems from the interpretation of the expectation as an average value. Typically, one may construct many unbiased estimates for the same parameter θ . This fact then raises the question of selecting one such estimate from the class of all unbiased estimates. Here is where the concept of the variance enters the picture. From two unbiased estimates $U_1 = U_1(X_1, \dots, X_n)$ and $U_2 = U_2(X_1, \dots, X_n)$ of θ , one would select the one with the smaller variance. This estimate will be more concentrated around θ than the other. Pictorially, this is illustrated by Figure 9.2.

Figure 9.2

(a) p.d.f. of U_1 (for a Fixed θ); (b) p.d.f. of U_2 (for a Fixed θ)



The next natural step is to look for an unbiased estimate which has the smallest variance in the class of all unbiased estimates, and this should happen for all $\theta \in \Omega$. Thus, we are led to the following concept.

DEFINITION 3

The unbiased estimate $U = U(X_1, \dots, X_n)$ of θ is said to be *Uniformly Minimum Variance Unbiased (UMVU)*, if for any other unbiased estimate $V = V(X_1, \dots, X_n)$, it holds that:

$$\text{Var}_{\theta}(U) \leq \text{Var}_{\theta}(V) \quad \text{for all } \theta \in \Omega.$$

That a UMVU estimate is desirable is more or less indisputable (see, however, Exercise 3.18). The practical question which then arises is how one goes about finding such an estimate. The process of seeking a UMVU estimate is facilitated by the Cramér–Rao inequality stated next. First, this inequality is stated, and then we describe how it is used.

THEOREM 6

(Cramér–Rao inequality) Let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \Re$, and suppose certain regularity conditions are met. Then, for any unbiased estimate $U = U(X_1, \dots, X_n)$ of θ , it holds that:

$$\text{Var}_\theta(U) \geq 1/nI(\theta), \quad \text{for all } \theta \in \Omega, \quad (8)$$

where

$$I(\theta) = E_\theta \left[\frac{\partial}{\partial \theta} \log f(X; \theta) \right]^2, \quad X \sim f(\cdot; \theta). \quad (9)$$

REMARK 10 The unspecified conditions mentioned in the formulation of the theorem include the assumption that the domain of x in the p.d.f. $f(x; \theta)$ does not depend on θ ; thus, the $U(0, \theta)$ distribution, for example, is left out. Also, the conditions include the validity of interchanging the operations of differentiation and integration in certain expressions. The proof of Theorem 6 is relatively long and involves an extensive list of regularity conditions. It may be found in considerable detail in Subsection 12.4.1 of Chapter 12 of the book *A Course in Mathematical Statistics*, 2nd edition, Academic Press (1997), by G. G. Roussas. Actually, in the reference just cited what is proved is a generalized version of Theorem 6, where the estimated function is a real-valued function of θ , $g(\theta)$, rather than θ itself.

REMARK 11 It can be shown that, under suitable conditions, the quantity $I(\theta)$ in (9) may also be calculated as follows:

$$I(\theta) = -E_\theta \left[\frac{\partial^2}{\partial \theta^2} \log f(X; \theta) \right]. \quad (10)$$

This expression is often easier to calculate.

The Cramér–Rao inequality is used in the following way.

- (i) Calculate the Fisher information either through (9) or by way of (10).
- (ii) Form the Cramér–Rao (C-R) lower bound figuring in inequality (8).
- (iii) Try to identify an unbiased estimate whose variance is equal to the C-R lower bound (for all $\theta \in \Theta$). If such an estimate is found,
- (iv) Declare the estimate described in (iii) as the UMVU estimate of θ .

In connection with steps (i)–(iv), it should be noted that it is possible that a UMVU estimate exists and yet such an estimate is not located through this process. The reason for such a failure is that the C-R inequality provides, simply, a lower bound for the variances of unbiased estimates, which may be strictly smaller than the variance of a UMVU estimate. It is, nevertheless, a good try!

The use of the inequality will be illustrated by two examples; other cases are left as exercises.

EXAMPLE 15

Refer to Example 2 and seek a UMVU estimate of θ through the C-R inequality.

DISCUSSION Here $f(x; \theta) = \theta^x(1 - \theta)^{1-x}$, $x = 0, 1$, so that:

$$(i) \log f(X; \theta) = X \log \theta + (1 - X) \log(1 - \theta),$$

$$\frac{\partial}{\partial \theta} \log f(X; \theta) = \frac{X}{\theta} - \frac{1 - X}{1 - \theta} \quad \text{and} \quad \frac{\partial^2}{\partial \theta^2} \log f(X; \theta) = -\frac{X}{\theta^2} - \frac{1 - X}{(1 - \theta)^2},$$

$$I(\theta) = -E_{\theta} \left[\frac{\partial^2}{\partial \theta^2} \log f(X; \theta) \right] = \frac{\theta}{\theta^2} + \frac{1 - \theta}{(1 - \theta)^2} = \frac{1}{\theta(1 - \theta)}, \quad \text{since } E_{\theta} X = \theta.$$

$$(ii) \text{ The C-R lower bound} = \frac{1}{nI(\theta)} = \theta(1 - \theta)/n.$$

(iii) Consider $U = \bar{X}$. Then $E_{\theta} \bar{X} = \theta$, so that \bar{X} is unbiased, and next $\sigma_{\theta}^2(\bar{X}) = \theta(1 - \theta)/n = 1/nI(\theta)$, since $\sigma_{\theta}^2(X) = \theta(1 - \theta)$.

(iv) The estimate \bar{X} is UMVU.

EXAMPLE 16

Refer to Example 4 and use the following parameterization:

$$X \sim f(x; \theta) = \frac{1}{\theta} e^{-\frac{x}{\theta}}, \quad x > 0, \quad \text{so that } E_{\theta} X = \theta, \quad \sigma_{\theta}^2(X) = \theta^2.$$

Then seek a UMVU estimate for θ through the C-R inequality.

DISCUSSION Here

$$(i) \log f(X; \theta) = -\log \theta - \frac{X}{\theta},$$

$$\frac{\partial}{\partial \theta} \log f(X; \theta) = -\frac{1}{\theta} + \frac{X}{\theta^2} \quad \text{and} \quad \frac{\partial^2}{\partial \theta^2} \log f(X; \theta) = \frac{1}{\theta^2} - \frac{2X}{\theta^3},$$

$$I(\theta) = -E_{\theta} \left[\frac{\partial^2}{\partial \theta^2} \log f(X; \theta) \right] = -\frac{1}{\theta^2} + \frac{2\theta}{\theta^3} = \frac{1}{\theta^2}.$$

$$(ii) \text{ The C-R lower bound} = \frac{1}{nI(\theta)} = \theta^2/n.$$

(iii) Consider $U = \bar{X}$. Then $E_{\theta} U = \theta$, so that \bar{X} is unbiased, and $\sigma_{\theta}^2(\bar{X}) = \frac{\theta^2}{n} = 1/nI(\theta)$.

(iv) The estimate \bar{X} is UMVU.

There is an alternative way of looking for UMVU estimates, in particular, when the approach by means of the Cramér–Rao inequality fails to produce such an estimate. This approach hinges heavily on the concept of sufficiency already introduced and also an additional technical concept, so-called *completeness*. The concept of completeness is a technical concept, and it says, in effect, that the only unbiased estimate of 0 is essentially the 0 statistics. More technically, if T is a r.v. with p.d.f. $f_T(\cdot; \theta)$, $\theta \in \Omega \subseteq \Re$, the family $\{f_T(\cdot; \theta); \theta \in \Omega\}$ (or the r.v. T) is said to be *complete* if, for $h: \Re \rightarrow \Re$, $E_{\theta} h(T) = 0$ for all $\theta \in \Omega$ implies $h(t)$ is, essentially, equal to 0. For the precise definition and a number of illustrative examples, the reader is referred to Section 11.2 of Chapter 11 in the reference cited in Remark 10 here. The concepts of sufficiency and completeness combined lead constructively to a UMVU estimate by means

of two theorems, the Rao–Blackwell and the Lehmann–Scheffé theorem. The procedure is summarized in the following result.

THEOREM 7

(Rao–Blackwell, Lehmann–Scheffé) Let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and let $T = T(X_1, \dots, X_n)$ be a sufficient statistic for θ (which is *complete*). Let $U = U(X_1, \dots, X_n)$ be any unbiased estimate of θ , and define the statistic $\varphi(T)$ by the relation:

$$\varphi(T) = E_{\theta}(U | T). \quad (11)$$

Then $\varphi(T)$ is unbiased, $\text{Var}_{\theta}[\varphi(T)] \leq \text{Var}_{\theta}(U)$ for all $\theta \in \Omega$, and, indeed, $\varphi(T)$ is a UMVU estimate of θ . If U is already a function of T only, then the conditioning in (11) is superfluous.

PROOF (ROUGH OUTLINE) That $\varphi(T)$ is independent of θ (and hence a statistic), despite the fact that we use quantities depending on θ in forming the conditional expectation, is due to the sufficiency of T . Recall that sufficiency of T means that the conditional distribution of U , given T , is independent of θ , and hence so is the expectation of U formed by using this conditional distribution. That $\varphi(T)$ is unbiased is due to a property of the conditional expectation (namely, for two r.v.'s X and Y : $E[E(X | Y)] = EX$), and the inequality involving the variances is also due to a property of the variance for conditional expectations (namely, $\text{Var}[E(X | Y)] \leq \text{Var}(X)$). The concept of completeness guarantees, through the Lehmann–Scheffé theorem, that no matter which unbiased estimate U we start out with, we end up (essentially) with the same UMVU estimate $\varphi(T)$ through the procedure (11), which is known as *Rao–Blackwellization*. ▲

REMARK 12 This theorem also applies suitably to multidimensional parameters θ , although, it must be stated here that a version of the Cramér–Rao inequality also exists for such parameters.

The following examples illustrate how one goes about applying Theorem 7 in concrete cases.

EXAMPLE 17

Determine the UMVU estimate of θ on the basis of the random sample X_1, \dots, X_n from the distribution $P(\theta)$.

DISCUSSION Perhaps, the simplest unbiased estimate of θ is X_1 , and we already know that $T = X_1 + \dots + X_n$ is sufficient for θ . For the Rao–Blackwellization of X_1 , we need the conditional distribution of X_1 , given $T = t$. It is known, however (see Exercise 2.10(ii), in Chapter 5), that this conditional distribution is $B(t, \frac{1}{n})$; that is, $P_{\theta}(X_1 = x | T = t) = \binom{t}{x} \left(\frac{1}{n}\right)^x \left(1 - \frac{1}{n}\right)^{t-x}$. It follows that $E_{\theta}(X_1 | T = t) = \frac{t}{n}$, so that $\varphi(T) = E_{\theta}(X_1 | T) = \frac{T}{n} = \bar{X}$. It so happens that the conditions of Theorem 7 hold (see Exercise 3.17) and therefore \bar{X} is the UMVU estimate of θ .

A well-known case where Theorem 7 works whereas Theorem 6 does not (or more properly, their suitable versions for two parameters do or do not) is illustrated by the following example.

EXAMPLE 18

Let X_1, \dots, X_n be a random sample from the $N(\mu, \sigma^2)$ distribution, where both μ and σ are unknown.

By an application of either theorem, it is seen that \bar{X} is the UMVU estimate of μ . Working with the Cramér–Rao inequality regarding σ^2 as the estimated parameter, it is seen that the C-R lower bound is equal to $2\sigma^4/n$. Next, Theorem 7 leads to the UMVU estimate of σ^2 , $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$. Furthermore, it has been seen (Exercise 1.4) that $\text{Var}_{\sigma^2}(S^2) = \frac{2\sigma^4}{n-1}$, which is strictly larger than $\frac{2\sigma^4}{n}$. This is the reason the Cramér–Rao inequality approach fails. For a little more extensive discussion on this example, see, e.g., Example 9, pages 299–301, in the reference cited in Remark 10.

Exercises

- 3.1** If X is a r.v. distributed as $B(n, \theta)$, $\theta \in \Omega = (0, 1)$, show that there is no unbiased estimate of $1/\theta$.

Hint: If $h(X)$ were such an estimate, then $E_\theta h(X) = \frac{1}{\theta}$ for all $\theta \in (0, 1)$. Write out the expectation, set $\frac{\theta}{1-\theta} = t$, and by expanding the right-hand side, conclude that $\binom{n+1}{n+1} (= 1) = 0$, which, of course, is a contradiction.

- 3.2** Let X_1, \dots, X_n be independent r.v.'s with p.d.f. $f(x; \theta) = \theta e^{-\theta x}$, $x > 0$, $\theta \in \Omega = (0, \infty)$, and let Y_1 be the smallest order statistic of the X_i 's. Then, by Example 11 in Chapter 6, the p.d.f. of Y_1 is $g_1(y) = (n\theta)e^{-(n\theta)y}$, $y > 0$.
- Show that both \bar{X} and nY_1 are unbiased estimates of $1/\theta$.
 - On the basis of variance considerations, which of these two estimates would you prefer?

- 3.3** Let X_1, \dots, X_n be a random sample of size n from the $U(0, \theta)$ distribution, $\theta \in \Omega = (0, \infty)$, and let Y_n be the largest order statistic of the X_i 's. Then:
- Employ formula (29) in Chapter 6 in order to obtain the p.d.f. of Y_n .
 - Use part (i) in order to construct an unbiased estimate of θ depending only on Y_n .
 - By Example 6 here (with $\alpha = 0$ and $\beta = \theta$) in conjunction with Theorem 3, show that the unbiased estimate in part (ii) depends only on a sufficient statistic for θ .

- 3.4** Let X_1, \dots, X_n be a random sample of size n from the $U(\theta_1, \theta_2)$ distribution, $\theta_1 < \theta_2$, and let Y_1 and Y_n be the smallest and the largest order statistics of the X_i 's.
- Use formulas (28) and (29) in Chapter 6 to obtain the p.d.f.'s of Y_1 and Y_n , and then, by calculating the $E_\theta Y_1$ and $E_\theta Y_n$, construct unbiased

estimates of the mean $(\theta_1 + \theta_2)/2$ and of the range $\theta_2 - \theta_1$ depending only on Y_1 and Y_n .

- (ii) Employ Example 6 here (with $\alpha = \theta_1$ and $\beta = \theta_2$) in conjunction with Theorem 3 and Remark 7 in order to show that the unbiased estimates in part (i) depend only on a set of sufficient statistics for (θ_1, θ_2) .

3.5 Let X_1, \dots, X_n be a random sample of size n from the $U(\theta, 2\theta)$ distribution, $\theta \in \Omega = (0, \infty)$, and set:

$$U_1 = \frac{n+1}{2n+1}Y_n \quad \text{and} \quad U_2 = \frac{n+1}{5n+4}(2Y_n + Y_1),$$

where Y_1 and Y_n are the smallest and the largest order statistics, respectively, of the X_i 's.

- (i) Use relations (28) and (29) in Chapter 6 in order to obtain the p.d.f.'s g_1 and g_n of Y_1 and Y_n , respectively.
(ii) By using part (i), show that:

$$E_\theta Y_1 = \frac{n+2}{n+1}\theta, \quad E_\theta Y_n = \frac{2n+1}{n+1}\theta.$$

- (iii) By means of part (ii), conclude that both U_1 and U_2 are unbiased estimates of θ .

3.6 Refer to Exercise 3.5, and show that:

$$\begin{aligned} \text{(i)} \quad E_\theta Y_1^2 &= \frac{n^2+5n+8}{(n+1)(n+2)}\theta^2, & \text{Var}_\theta(Y_1) &= \frac{n}{(n+1)^2(n+2)}\theta^2. \\ \text{(ii)} \quad E_\theta Y_n^2 &= \frac{2(2n^2+4n+1)}{(n+1)(n+2)}\theta^2, & \text{Var}_\theta(Y_n) &= \frac{n}{(n+1)^2(n+2)}\theta^2. \end{aligned}$$

3.7 Refer to Exercise 3.5, and:

- (i) Use Exercise 5.3 (ii) in Chapter 6 in order to show that the joint p.d.f. g_{1n} of Y_1 and Y_n is given by:

$$g_{1n}(y_1, y_n) = \frac{n(n-1)}{\theta^n}(y_n - y_1)^{n-2}, \quad \theta \leq y_1 < y_n \leq 2\theta.$$

- (ii) Employ part (i) here and also part (ii) of Exercise 3.5 in order to show that:

$$E_\theta(Y_1 Y_n) = \frac{2n^2 + 7n + 5}{(n+1)(n+2)}\theta^2, \quad \text{Cov}_\theta(Y_1, Y_n) = \frac{\theta^2}{(n+1)^2(n+2)}.$$

3.8 Refer to Exercise 3.5, and:

- (i) Use Exercises 3.6 and 3.7 (ii) in order to show that:

$$\text{Var}_\theta(U_1) = \frac{n}{(2n+1)^2(n+2)}\theta^2, \quad \text{Var}_\theta(U_2) = \frac{1}{(5n+4)(n+2)}\theta^2.$$

- (ii) From part (i), conclude that $\text{Var}_\theta(U_2) \leq \text{Var}_\theta(U_1)$ for all θ (with equality holding only for $n = 1$), so that the (unbiased) estimate U_2 is uniformly better (in terms of variance) than the (unbiased) estimate U_1 .

3.9 Let X_1, \dots, X_m and Y_1, \dots, Y_n be independent random samples with the same mean θ and known variances σ_1^2 and σ_2^2 , respectively. For any fixed

c with $0 \leq c \leq 1$, set $U_c = c\bar{X} + (1 - c)\bar{Y}$, where \bar{X} and \bar{Y} are the sample means of the X_i 's and of the Y_i 's, respectively. Then:

- (i) Show that U_c is an unbiased estimate of θ for every c as specified above.
- (ii) Calculate the variance of U_c , and show that it is minimized for $c = c_0 = m\sigma_2^2 / (n\sigma_1^2 + m\sigma_2^2)$.

3.10 Let X_1, \dots, X_n be i.i.d. r.v.'s with mean μ and variance σ^2 , both unknown. Then for any known constants c_1, \dots, c_n , consider the *linear* estimate of μ defined by: $U_c = \sum_{i=1}^n c_i X_i$.

- (i) Identify the condition that the c_i 's must satisfy, so that U_c is an unbiased estimate of μ .
- (ii) Show that the sample mean \bar{X} is the unbiased linear estimate of μ with the smallest variance (among all unbiased linear estimates of μ).

Hint: For part (ii), one has to minimize the expression $\sum_{i=1}^n c_i^2$ subject to the side restriction that $\sum_{i=1}^n c_i = 1$. For this minimization, use the Lagrange multipliers method, which calls for the minimization of the function $\phi(c_1, \dots, c_n) = \sum_{i=1}^n c_i^2 + \lambda(\sum_{i=1}^n c_i - 1)$ with respect to c_1, \dots, c_n , where λ is a constant (Lagrange multiplier). Alternatively, one may employ a geometric argument to the same effect.

In all of the following Exercises 3.11–3.15, employ steps (i)–(iv) listed after Remark 11 in an attempt to determine UMVU estimates. Use relation (10) whenever possible.

3.11 If X_1, \dots, X_n is a random sample from the Poisson distribution $P(\theta)$, show that the sample mean \bar{X} is the UMVU estimate of θ .

3.12 Let X_1, \dots, X_n be i.i.d. r.v.'s distributed as $N(\mu, \sigma^2)$.

- (i) If $\mu = \theta \in \Omega = \Re$ and σ is known, show that \bar{X} is the UMVU estimate of θ .
- (ii) If $\sigma^2 = \theta \in \Omega = (0, \infty)$ and μ is known show that $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2$ is the UMVU estimate of σ^2 .

Hint: For part (ii), recall that $Y = \sum_{i=1}^n \left(\frac{X_i - \mu}{\sqrt{\theta}} \right)^2 \sim \chi_n^2$, and hence $E_\theta Y = n$, $Var_\theta(Y) = 2n$.

3.13 Let X_1, \dots, X_n be i.i.d. r.v.'s from the Gamma distribution with parameters α known and $\beta = \theta \in \Omega = (0, \infty)$ unknown.

- (i) Determine the Fisher information $I(\theta)$.
- (ii) Show that the estimate $U = U(X_1, \dots, X_n) = \frac{1}{n\alpha} \sum_{i=1}^n X_i$ is unbiased and calculate its variance.
- (iii) Show that $Var_\theta(U) = 1/nI(\theta)$, so that U is the UMVU estimate of θ .

3.14 Let X_1, \dots, X_n be i.i.d. r.v.'s from the Negative Exponential p.d.f. in the following parametric form: $f(x; \theta) = \frac{1}{\theta} e^{-x/\theta}$, $x > 0$, $\theta \in \Omega = (0, \infty)$, so

that $E_\theta X_1 = \theta$. Use Exercise 3.13 in order to show that \bar{X} is the UMVU estimate of θ .

Hint: Recall that the Negative Exponential distribution is a special case of the Gamma distribution.

3.15 Let X be a r.v. with p.d.f. $f(x; \theta) = \frac{1}{2\theta} e^{-|x|/\theta}$, $x \in \mathfrak{R}$, $\theta \in \Omega = (0, \infty)$. Then:

- (i) Show that $E_\theta |X| = \theta$, and $E_\theta X^2 = 2\theta^2$.
- (ii) Show that the statistic $U = U(X_1, \dots, X_n) = \frac{1}{n} \sum_{i=1}^n |X_i|$ is an unbiased estimate of θ , and calculate its variance.
- (iii) Show that the Fisher information number $I(\theta) = -E_\theta \left[\frac{\partial^2}{\partial \theta^2} \log f(X; \theta) \right] = 1/\theta^2$.
- (iv) Conclude that U is the UMVU estimate of θ .

In Exercises 3.16–3.20, the purpose is to construct the UMVU estimates of the parameters involved by invoking Theorem 7. Sufficiency can always be established through Theorem 3; completeness sometimes will be established, but it will always be assumed when appropriate.

3.16 If X_1, \dots, X_n are independent r.v.'s distributed as $B(1, \theta)$, $\theta \in \Omega = (0, 1)$, then:

- (i) Show that $T = \sum_{i=1}^n X_i$ is sufficient for θ .
- (ii) Also, show that T is complete.
- (iii) From parts (i) and (ii), conclude that \bar{X} is the UMVU estimate of θ .

3.17 Let X_1, \dots, X_n be a random sample of size n from the $P(\theta)$ distribution, $\theta \in \Omega = (0, \infty)$. With $T = \sum_{i=1}^n X_i$, it has been seen that the conditional distribution $P_\theta(X_1 = x | T = t)$ is $B(t, \frac{1}{n})$ (see Exercise 2.10(ii) in Chapter 5) and that T is sufficient for θ (see Exercise 2.9 here). Show that T is complete, so that the conclusion reached in Example 17 will be fully justified.

3.18 Let the independent r.v.'s X_1, \dots, X_n have the Geometric p.d.f. $f(x; \theta) = \theta(1 - \theta)^{x-1}$, $x = 1, 2, \dots$, $\theta \in \Omega = (0, 1)$.

- (i) Show that X is both sufficient and complete.
- (ii) Show that the estimate U defined by: $U(X) = 1$ if $X = 1$, and $U(X) = 0$ if $X = 0$, is an unbiased estimate of θ .
- (iii) Conclude that U is the UMVU estimate of θ and also an entirely unreasonable estimate.
- (iv) Prove that the variance of U is uniformly bigger than the Cramér–Rao lower bound. (So, on account of this, the Cramér–Rao inequality could not produce the UMVU estimate.)

Remark: We have stipulated that an estimate always takes values in the appropriate parameter Ω . In order to be consistent with this stipulation, we take $\Omega = [0, 1]$ in part (ii).

3.19 Let X_1, \dots, X_n be independent r.v.'s with the Negative Exponential p.d.f. $f(x; \theta) = \frac{1}{\theta} e^{-\frac{1}{\theta}x}$, $x > 0$, $\theta \in \Omega = (0, \infty)$. Then:

- (i) \bar{X} is sufficient (and complete, although completeness will not be established here).
- (ii) \bar{X} is the UMVU estimate of θ .

3.20 Let X_1, \dots, X_n be independent r.v.'s distributed as $U(0, \theta)$, $\theta \in \Omega = (0, \infty)$. Then:

- (i) The largest order statistic Y_n of the X_i 's is sufficient (and complete, although completeness will not be established here).
- (ii) The unbiased estimate $U = \frac{n+1}{n} Y_n$ is the UMVU estimate of θ .
- (iii) Explain why the Cramér–Rao inequality approach is not applicable here.

9.4 Decision-Theoretic Approach to Estimation

In this section, a brief discussion is presented of still another approach to parameter estimation, which, unlike the previous two approaches, is kind of penalty driven. In order to introduce the relevant concepts and notation, let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and let δ be a function defined on \mathfrak{R}^n into Ω ; i.e., $\delta : \mathfrak{R}^n \rightarrow \Omega$. If x_1, \dots, x_n are the observed values of X_1, \dots, X_n , then the value $\delta(x_1, \dots, x_n)$ is the proposed estimate of θ . The quality of this estimate is usually measured by its squared distance from the estimated quantity θ ; that is, $[\theta - \delta(x_1, \dots, x_n)]^2$. Denote it by $L[\theta; \delta(x_1, \dots, x_n)]$ and call it a *loss function*. So $L[\theta; \delta(x_1, \dots, x_n)] = [\theta - \delta(x_1, \dots, x_n)]^2$. The closer the estimate $\delta(x_1, \dots, x_n)$ is to θ (on either side of it) the smaller is the loss we suffer, and vice versa. The objective here is to select δ in some optimal way to be discussed below. The first step to this effect is that δ be selected so that it minimizes the average loss we suffer by using this estimate. For this purpose, consider the r.v. $L[\theta; \delta(X_1, \dots, X_n)] = [\theta - \delta(X_1, \dots, X_n)]^2$ and take its expectation to be denoted by $R(\theta; \delta)$; namely,

$$\begin{aligned}
 R(\theta; \delta) &= E_\theta[\theta - \delta(X_1, \dots, X_n)]^2 \\
 &= \begin{cases} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} [\theta - \delta(x_1, \dots, x_n)]^2 f(x_1; \theta) \cdots f(x_n; \theta) dx_1 \cdots dx_n, & \text{for the continuous case,} \\ \sum_{x_1} \cdots \sum_{x_n} [\theta - \delta(x_1, \dots, x_n)]^2 f(x_1; \theta) \cdots f(x_n; \theta), & \text{for the discrete case.} \end{cases} \quad (12)
 \end{aligned}$$

The average loss $R(\theta; \delta)$ is called the *risk function*, corresponding to δ . The value $R(\theta; \delta)$ is the average loss suffered corresponding to the point θ , when δ is used. At this point, there are two options available to us in pursuing the issue of selecting δ . One is to choose δ , so as to minimize the worst which can happen to us. More formally, choose δ so that, for any other estimate δ^* , it holds that:

$$\sup[R(\theta; \delta); \theta \in \Omega] \leq \sup[R(\theta; \delta^*); \theta \in \Omega].$$

Such an estimate, if it exists, is called *minimax* (by the fact that it minimizes the maximum risk). The second option would be to average $R(\theta; \delta)$, with respect to θ , and then choose δ to minimize this average. The implementation of this

plan goes as follows. Let $\lambda(\theta)$ be a p.d.f. on Ω and average $R(\theta; \delta)$ by using this p.d.f.; let $r(\delta)$ be the resulting average. Thus,

$$r(\delta) = E_{\lambda} R(\theta; \delta) = \begin{cases} \int_{\Omega} R(\theta; \delta) \lambda(\theta) d\theta, & \text{for the continuous case,} \\ \sum_{\theta \in \Omega} R(\theta; \delta) \lambda(\theta), & \text{for the discrete case.} \end{cases} \quad (13)$$

Then select δ to minimize $r(\delta)$. Such an estimate is called a *Bayes estimate*, corresponding to the p.d.f. λ , and it may be denoted by δ_{λ} . In this context, the parameter θ is interpreted as a r.v. taking values in Ω according to the p.d.f. $\lambda(\theta)$, which is called a *prior* or a *priori* p.d.f. It so happens that, under minimal assumptions, a Bayes estimate always exists and is given by an explicit formula.

THEOREM 8

Suppose θ is a r.v. of the continuous type with prior p.d.f. $\lambda(\theta)$, and that the three quantities $\int_{\Omega} f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta$, $\int_{\Omega} \theta f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta$, and $\int_{\Omega} \theta^2 f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta$ are finite. Then the Bayes estimate corresponding to $\lambda(\theta)$, δ_{λ} , is given by the expression

$$\delta_{\lambda}(x_1, \dots, x_n) = \frac{\int_{\Omega} \theta f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta}{\int_{\Omega} f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta}. \quad (14)$$

If θ is a discrete r.v., all integrals above are to be replaced by summation signs.

This theorem has the following corollary.

COROLLARY The Bayes estimate $\delta_{\lambda}(x_1, \dots, x_n)$ defined in relation (14) can also be calculated thus:

$$\delta_{\lambda}(x_1, \dots, x_n) = \int_{\Omega} \theta h(\theta | x_1, \dots, x_n) d\theta, \quad (15)$$

where $h(\theta | x_1, \dots, x_n)$ is the conditional p.d.f. of θ , given $X_i = x_i$, $i = 1, \dots, n$, which is also called the *posterior* p.d.f. The integral is to be replaced by a summation sign in the discrete case.

PROOF Observe that $f(x_1; \theta) \cdots f(x_n; \theta)$ is, actually, the joint conditional p.d.f. of X_1, \dots, X_n , given θ , so that the product $f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta)$ is the joint p.d.f. of X_1, \dots, X_n and θ . Then its integral over Ω is the marginal (joint) p.d.f. of X_1, \dots, X_n , and therefore

$$f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) / \int_{\Omega} f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta$$

is $h(\theta | x_1, \dots, x_n)$ as described above. Then expression (14) completes the proof. \blacktriangle

So, in computing $\delta_{\lambda}(x_1, \dots, x_n)$, one may use relation (14) or, alternatively, first calculate $h(\theta | x_1, \dots, x_n)$ and then apply formula (15).

We now proceed with the justification of (14).

PROOF OF THEOREM 8 All operations below are valid without any further explanation. The derivations are carried out for the continuous case; in the discrete case, the integrals are to be replaced by summation signs. By relation (13),

$$\begin{aligned} r(\delta) &= \int_{\Omega} R(\theta; \delta) \lambda(\theta) d\theta = \int_{\Omega} \left\{ \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} [\theta - \delta(x_1, \dots, x_n)]^2 \times \right. \\ &\quad \left. f(x_1; \theta) \cdots f(x_n; \theta) dx_1 \cdots dx_n \right\} \lambda(\theta) d\theta \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \left\{ \int_{\Omega} [\theta - \delta(x_1, \dots, x_n)]^2 \lambda(\theta) f(x_1; \theta) \cdots f(x_n; \theta) d\theta \right\} \\ &\quad \times dx_1 \cdots dx_n. \end{aligned}$$

Then, in order to minimize $r(\delta)$, it suffices to minimize the inner integral for each x_1, \dots, x_n . However,

$$\begin{aligned} &\int_{\Omega} [\theta - \delta(x_1, \dots, x_n)]^2 \lambda(\theta) f(x_1; \theta) \cdots f(x_n; \theta) d\theta \\ &= \delta^2(x_1, \dots, x_n) \left[\int_{\Omega} f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta \right] \\ &\quad - 2\delta(x_1, \dots, x_n) \left[\int_{\Omega} \theta f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta \right] \\ &\quad + \left[\int_{\Omega} \theta^2 f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta \right], \end{aligned}$$

and this is of the form: $g(t) = at^2 - 2bt + c$, where

$$\begin{aligned} a &= \int_{\Omega} f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta, \\ b &= \int_{\Omega} \theta f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta, \\ c &= \int_{\Omega} \theta^2 f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta, \end{aligned}$$

and

$$t = \delta(x_1, \dots, x_n).$$

The quadratic expression $g(t) = at^2 - 2bt + c$ is minimized for $t = \frac{b}{a}$, since $g'(t) = 2at - 2b = 0$ gives $t = \frac{b}{a}$ and $g''(t) = 2a > 0$. But $\frac{b}{a}$ is equal to the right-hand side in expression (14). The proof is complete. \blacktriangle

REMARK 13 In the context of the present section, the function δ and the estimate $\delta(x_1, \dots, x_n)$ are also referred to as a *decision function* and a *decision* (associated with the specific outcome x_1, \dots, x_n). Hence the title of the section.

REMARK 14 The Bayes approach presents us with both advantages and disadvantages. An issue which often arises is how the prior p.d.f. $\lambda(\theta)$ is to

be chosen. People have given various considerations in selecting $\lambda(\theta)$, including mathematical convenience. Perhaps the most significant advantage of this approach is that, in selecting the prior $\lambda(\theta)$, we have flexibility in incorporating whatever information we may have about the parameter θ .

The theorem is illustrated with one example.

EXAMPLE 19

Let X_1, \dots, X_n be a random sample from the $B(1, \theta)$ distribution, $\theta \in \Omega = (0, 1)$, and choose $\lambda(\theta)$ to be the so-called *Beta* density with parameters α and β ; that is,

$$\lambda(\theta) = \begin{cases} \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \theta^{\alpha-1} (1-\theta)^{\beta-1}, & \text{if } \theta \in (0, 1) \\ 0, & \text{otherwise.} \end{cases} \quad (16)$$

(For a proof that λ is, indeed, p.d.f., see, e.g., pages 70–71 in the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G. G. Roussas.) Then the Bayes estimate is given by relation (20) below.

DISCUSSION Now, from the definition of the p.d.f. of a Beta distribution with parameters α and β , we have

$$\int_0^1 x^{\alpha-1} (1-x)^{\beta-1} dx = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \quad (17)$$

and, of course, $\Gamma(\gamma) = (\gamma-1)\Gamma(\gamma-1)$. Then, for simplicity, writing $\sum_j x_j$ rather than $\sum_{j=1}^n x_j$ when this last expression appears as an exponent, we have

$$\begin{aligned} I_1 &= \int_{\Omega} f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 \theta^{\sum_j x_j} (1-\theta)^{n-\sum_j x_j} \theta^{\alpha-1} (1-\theta)^{\beta-1} d\theta \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 \theta^{(\alpha+\sum_j x_j)-1} (1-\theta)^{(\beta+n-\sum_j x_j)-1} d\theta, \end{aligned}$$

which, by means of (17), becomes as follows:

$$I_1 = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \frac{\Gamma(\alpha+\sum_{j=1}^n x_j) \Gamma(\beta+n-\sum_{j=1}^n x_j)}{\Gamma(\alpha+\beta+n)}. \quad (18)$$

Next,

$$\begin{aligned} I_2 &= \int_{\Omega} \theta f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 \theta \theta^{\sum_j x_j} (1-\theta)^{n-\sum_j x_j} \theta^{\alpha-1} (1-\theta)^{\beta-1} d\theta \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 \theta^{(\alpha+\sum_j x_j+1)-1} (1-\theta)^{(\beta+n-\sum_j x_j)-1} d\theta. \end{aligned}$$

Once more, relation (17) gives

$$I_2 = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \times \frac{\Gamma(\alpha+\sum_{j=1}^n x_j+1) \Gamma(\beta+n-\sum_{j=1}^n x_j)}{\Gamma(\alpha+\beta+n+1)}. \quad (19)$$

Relations (18) and (19) imply, by virtue of (14),

$$\delta(x_1, \dots, x_n) = \frac{\Gamma(\alpha + \beta + n)\Gamma(\alpha + \sum_{j=1}^n x_j + 1)}{\Gamma(\alpha + \beta + n + 1)\Gamma(\alpha + \sum_{j=1}^n x_j)} = \frac{\alpha + \sum_{j=1}^n x_j}{\alpha + \beta + n};$$

that is,

$$\delta(x_1, \dots, x_n) = \frac{\sum_{j=1}^n x_j + \alpha}{n + \alpha + \beta}. \quad (20)$$

REMARK 15 When $\alpha = \beta = 1$, the Beta distribution becomes $U(0, 1)$, as follows from (16), since $\Gamma(2) = 1 \times \Gamma(1) = 1$. In this case, the corresponding Bayes estimate is $\delta(x_1, \dots, x_n) = (\sum_{i=1}^n x_i + 1)/(n + 2)$.

A minimax estimate is usually found indirectly, by showing that a Bayes estimate is also minimax. The following theorem tells the story.

THEOREM 9

Let $\delta_\lambda(x_1, \dots, x_n)$ be the Bayes estimate corresponding to the prior p.d.f. $\lambda(\theta)$, and suppose its risk $R(\theta; \delta_\lambda)$, as given in (12), is independent of $\theta \in \Omega$. Then $\delta_\lambda(x_1, \dots, x_n)$ is minimax.

PROOF The justification is straightforward and goes like this. Set $R(\theta; \delta_\lambda) = c$, and let $\delta^* = \delta^*(x_1, \dots, x_n)$ be any other estimate. Then

$$\begin{aligned} \sup[R(\theta; \delta_\lambda); \theta \in \Omega] &= c = \int_{\Omega} c\lambda(\theta) d\theta \\ &= \int_{\Omega} R(\theta; \delta_\lambda)\lambda(\theta) d\theta \leq \int_{\Omega} R(\theta; \delta^*)\lambda(\theta) d\theta \quad (\text{since } \delta_\lambda \text{ is Bayes}) \\ &\leq \sup[R(\theta; \delta^*); \theta \in \Omega] \quad (\text{since } \lambda \text{ is a p.d.f. on } \Omega). \end{aligned}$$

This completes the proof. ▲

The following example illustrates this theorem.

EXAMPLE 20

Let X_1, \dots, X_n and $\lambda(\theta)$ be as in Example 19. Then the corresponding Bayes estimate δ is given by (20), and the estimate δ^* given in (21) is minimax.

DISCUSSION By setting $X = \sum_{j=1}^n X_j$ and taking into consideration that $E_\theta X = n\theta$ and $E_\theta X^2 = n\theta(1 - \theta + n\theta)$, we obtain

$$\begin{aligned} R(\theta; \delta) &= E_\theta \left(\theta - \frac{X + \alpha}{n + \alpha + \beta} \right)^2 \\ &= \frac{1}{(n + \alpha + \beta)^2} \{[(\alpha + \beta)^2 - n]\theta^2 - (2\alpha^2 + 2\alpha\beta - n)\theta + \alpha^2\}. \end{aligned}$$

By taking $\alpha = \beta = \frac{1}{2}\sqrt{n}$ and denoting by δ^* the resulting estimate, we have

$$(\alpha + \beta)^2 - n = 0, \quad 2\alpha^2 + 2\alpha\beta - n = 0,$$

so that

$$R(\theta; \delta^*) = \frac{\alpha^2}{(n + \alpha + \beta)^2} = \frac{n}{4(n + \sqrt{n})^2} = \frac{1}{4(1 + \sqrt{n})^2}.$$

Since $R(\theta; \delta^*)$ is independent of θ , Theorem 9 implies that

$$\delta^*(x_1, \dots, x_n) = \frac{\sum_{j=1}^n x_j + \frac{1}{2}\sqrt{n}}{n + \sqrt{n}} = \frac{2\sqrt{n}\bar{x} + 1}{2(1 + \sqrt{n})} \quad (21)$$

is minimax.

Exercises

4.1 Consider one observation from the p.d.f. $f(x; \theta) = (1 - \theta)\theta^{x-1}$, $x = 1, 2, \dots$, $\theta \in \Omega = (0, 1)$, and let the prior p.d.f. λ on $(0, 1)$ be the $U(0, 1)$ distribution. Then, determine:

- (i) The posterior p.d.f. of θ , given $X = x$.
- (ii) The Bayes estimate of θ , by using relation (15).

4.2 If the r.v. X has the Beta distribution with parameters α and β ; i.e., its p.d.f. is given by expression (16), then without integration and by using the recursive property of the Gamma function ($\Gamma(\gamma) = (\gamma - 1)\Gamma(\gamma - 1)$, $\gamma > 1$), show that $EX = \alpha/(\alpha + \beta)$.

4.3 In reference to Example 19:

- (i) Show that the marginal p.d.f., $h(x_1, \dots, x_n)$, defined by $h(x_1, \dots, x_n) = \int_0^1 f(x_1; \theta) \cdots f(x_n; \theta) \lambda(\theta) d\theta$ with $f(x; \theta) = \theta^x(1 - \theta)^{1-x}$, $x = 0, 1$, and $\lambda(\theta)$ as in relation (16), is given by:

$$h(x_1, \dots, x_n) = \frac{\Gamma(\alpha + \beta)\Gamma(\alpha + t)\Gamma(\beta + n - t)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\alpha + \beta + t)},$$

where $t = x_1 + \dots + x_n$. Do it without, actually, carrying out any integrations, by taking notice of the form of a Beta p.d.f.

- (ii) Show that the posterior p.d.f. of θ , given $X_1 = x_1, \dots, X_n = x_n$, $h(\theta | x_1, \dots, x_n)$, is the Beta p.d.f. with parameters $\alpha + t$ and $\beta + n - t$.
- (iii) Use the posterior p.d.f. obtained in part (ii) in order to rederive the Bayes estimate $\delta(x_1, \dots, x_n)$ given in (20) by utilizing relation (15). Do it without carrying out any integrations, by using Exercise 4.2.
- (iv) Construct a $100(1 - \alpha)\%$ Bayes confidence interval for θ ; that is, determine a set $\{\theta \in (0, 1); h(\theta | x_1, \dots, x_n) \geq c(x_1, \dots, x_n)\}$, where $c(x_1, \dots, x_n)$ is determined by the requirement that the P_λ -probability of this set is equal to $1 - \alpha$.

4.4 Let X_1, \dots, X_n be independent r.v.'s from the $N(\theta, 1)$ distribution, $\theta \in \Omega = \Re$, and on \Re , consider the p.d.f. λ to be that of $N(\mu, 1)$ with μ known. Then show that the Bayes estimate of θ , $\delta_\lambda(x_1, \dots, x_n)$, is given by: $\delta(x_1, \dots, x_n) = \frac{n\bar{x} + \mu}{n + 1}$.

Hint: By (14), we have to find suitable expressions for the integrals:

$$I_1 = \int_{-\infty}^{\infty} \frac{1}{(\sqrt{2\pi})^n} \exp\left[-\frac{1}{2} \sum_{i=1}^n (x_i - \theta)^2\right] \times \frac{1}{\sqrt{2\pi}} \exp\left[-\frac{(\theta - \mu)^2}{2}\right] d\theta,$$

$$I_2 = \int_{-\infty}^{\infty} \theta \cdot \frac{1}{(\sqrt{2\pi})^n} \exp\left[-\frac{1}{2} \sum_{i=1}^n (x_i - \theta)^2\right] \times \frac{1}{\sqrt{2\pi}} \exp\left[-\frac{(\theta - \mu)^2}{2}\right] d\theta.$$

The integrand of I_1 is equal to (the constant for the integration):

$$\begin{aligned} & \frac{1}{(\sqrt{2\pi})^n} \exp\left[-\frac{1}{2} \left(\sum_{i=1}^n x_i^2 + \mu^2\right)\right] \\ & \times \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{1}{2}[(n+1)\theta^2 - 2(n\bar{x} + \mu)\theta]\right\}. \end{aligned}$$

However,

$$\begin{aligned} (n+1)\theta^2 - 2(n\bar{x} + \mu)\theta &= (n+1) \left(\theta^2 - 2\frac{n\bar{x} + \mu}{n+1}\theta\right) \\ &= (n+1) \left[\left(\theta - \frac{n\bar{x} + \mu}{n+1}\right)^2 - \left(\frac{n\bar{x} + \mu}{n+1}\right)^2\right] \\ &= \frac{(\theta - \frac{n\bar{x} + \mu}{n+1})^2}{1/(\sqrt{n+1})^2} - \frac{(n\bar{x} + \mu)^2}{n+1}, \end{aligned}$$

so that

$$\begin{aligned} & \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{1}{2}[(n+1)\theta^2 - 2(n\bar{x} + \mu)\theta]\right\} \\ &= \frac{1}{\sqrt{n+1}} \exp\left[\frac{1}{2} \times \frac{(n\bar{x} + \mu)^2}{n+1}\right] \times \frac{1}{\sqrt{2\pi}(1/\sqrt{n+1})} \exp\left[-\frac{(\theta - \frac{n\bar{x} + \mu}{n+1})^2}{2/(\sqrt{n+1})^2}\right], \end{aligned}$$

and the second factor is the p.d.f. of $N(\frac{n\bar{x} + \mu}{n+1}, \frac{1}{n+1})$. Therefore the integration produces the constant:

$$\frac{1}{\sqrt{n+1}} \times \frac{1}{(\sqrt{2\pi})^n} \exp\left\{-\frac{1}{2} \left[\sum_{i=1}^n x_i^2 + \mu^2 - \frac{(n\bar{x} + \mu)^2}{n+1}\right]\right\}.$$

Likewise, the integrand in I_2 is rewritten thus:

$$\begin{aligned} & \frac{1}{\sqrt{n+1}} \times \frac{1}{(\sqrt{2\pi})^n} \exp\left\{-\frac{1}{2} \left[\sum_{i=1}^n x_i^2 + \mu^2 - \frac{(n\bar{x} + \mu)^2}{n+1}\right]\right\} \\ & \times \frac{1}{\sqrt{2\pi}(1/\sqrt{n+1})} \theta \exp\left[-\frac{(\theta - \frac{n\bar{x} + \mu}{n+1})^2}{2/(\sqrt{n+1})^2}\right], \end{aligned}$$

and the second factor, when integrated with respect to θ , is the mean of $N(\frac{n\bar{x} + \mu}{n+1}, \frac{1}{n+1})$ distribution, which is $\frac{n\bar{x} + \mu}{n+1}$. Dividing then I_2 by I_1 , we obtain the desired result.

4.5 Refer to Exercise 4.4, and:

- (i) By utilizing the derivations in the hint, derive the posterior p.d.f. $h(\theta \mid x_1, \dots, x_n)$.
- (ii) Construct a $100(1 - \alpha)\%$ Bayes confidence interval for θ as in Exercise 4.3(iv).

4.6 Let X_1, \dots, X_n be independent r.v.'s distributed as $P(\theta)$, $\theta \in \Omega = (0, \infty)$, and consider the estimate $\delta(x_1, \dots, x_n) = \bar{x}$ and the loss function $L(\theta; \delta) = [\theta - \delta(x_1, \dots, x_n)]^2/\theta$.

Calculate the risk $R(\theta; \delta) = \frac{1}{\theta} E_\theta L[\theta - \delta(X_1, \dots, X_n)]^2$, and use Theorem 9 in order to conclude that the estimate $\delta(x_1, \dots, x_n) = \bar{x}$ is, actually, minimax.

9.5 Other Methods of Estimation

In addition to the methods of estimation discussed so far, there are also other methods and approaches, such as the so-called *minimum Chi-Square method*, the method of *least squares*, and the *method of moments*. The method of least squares is usually associated with the so-called linear models, and therefore we defer its discussion to a later chapter (see Chapter 13). Here, we are going to present only a brief outline of the method of moments, and illustrate it with three examples.

To this end, let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and suppose that $E_\theta X_1 = m_1(\theta)$ is finite. The objective is to estimate θ by means of the random sample at hand. By the WLLN,

$$\frac{1}{n} \sum_{i=1}^n X_i = \bar{X}_n \xrightarrow[n \rightarrow \infty]{P_\theta} m_1(\theta). \quad (22)$$

Therefore, for large n , it would make sense to set $\bar{X}_n = m_1(\theta)$ (since it will be approximately so with probability as close to 1 as one desires), and make an attempt to solve for θ . Assuming that this can be done and that there is a unique solution, we declare that solution as the *moment estimate* of θ .

This methodology applies in principle also in the case that there are r parameters involved, $\theta_1, \dots, \theta_r$, or, as we say, when θ has r coordinates, $r \geq 1$. In such a case, we have to assume that the r first moments of the X_i 's are finite; that is,

$$E_\theta X_1^k = m_k(\theta_1, \dots, \theta_r) \in \mathfrak{R}, \quad k = 1, \dots, r, \quad \theta = (\theta_1, \dots, \theta_r).$$

Then form the first r sample moments $\frac{1}{n} \sum_{i=1}^n X_i^k$, $k = 1, \dots, r$, and equate them to the corresponding (population) moments; that is,

$$\frac{1}{n} \sum_{i=1}^n X_i^k = m_k(\theta_1, \dots, \theta_r), \quad k = 1, \dots, r. \quad (23)$$

The reasoning for doing this is the same as the one explained above in conjunction with (22). Assuming that we can solve for $\theta_1, \dots, \theta_r$ in (23), and that the solutions are unique, we arrive at what we call the *moment estimates* of the parameters $\theta_1, \dots, \theta_r$.

The following examples should help shed some light on the above exposition.

EXAMPLE 21

On the basis of the random sample X_1, \dots, X_n from the $B(1, \theta)$ distribution, find the moment estimate of θ .

DISCUSSION Here $E_\theta X_1 = \theta$ and there is only one parameter. Thus, it suffices to set $\bar{X} = \theta$, so that the moment estimate of θ is the same as the MLE and the UMVU estimate, but (slightly) different from the Bayes (and the minimax estimate).

EXAMPLE 22

On the basis of the random sample X_1, \dots, X_n from the $N(\mu, \sigma^2)$ distribution with both μ and σ^2 unknown, determine the moment estimates of μ and σ^2 .

DISCUSSION The conditions referred to above are satisfied here, and, specifically, $E_\theta X_1 = \mu$, $E_\theta X_1^2 = \sigma^2 + \mu^2$, $\theta = (\mu, \sigma^2)$. Here we need the first two sample moments, \bar{X} and $\frac{1}{n} \sum_{i=1}^n X_i^2$. We have then: $\bar{X} = \mu$ and $\frac{1}{n} \sum_{i=1}^n X_i^2 = \sigma^2 + \mu^2$. Hence $\mu = \bar{X}$ and $\sigma^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2 = \frac{1}{n} (\sum_{i=1}^n X_i^2 - n\bar{X}^2) = \frac{1}{n} \times \sum_{i=1}^n (X_i - \bar{X})^2$. Thus, the moment estimates are $\tilde{\mu} = \bar{X}$ and $\tilde{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$. The estimate $\tilde{\mu}$ is identical with the MLE and the UMVU estimate, whereas $\tilde{\sigma}^2$ is the same as the MLE, but (slightly) different from the UMVU estimate.

EXAMPLE 23

Let the random sample X_1, \dots, X_n be from the $U(\alpha, \beta)$ distribution, where both α and β are unknown. Determine their moment estimates.

DISCUSSION Recall that $E_\theta X_1 = \frac{\alpha + \beta}{2}$ and $\sigma_\theta^2(X_1) = \frac{(\alpha - \beta)^2}{12}$, $\theta = (\alpha, \beta)$, so that:

$$\bar{X} = \frac{\alpha + \beta}{2}, \quad \frac{1}{n} \sum_{i=1}^n X_i^2 = \frac{(\alpha - \beta)^2}{12} + \left(\frac{\alpha + \beta}{2} \right)^2.$$

Hence $\frac{(\alpha - \beta)^2}{12} = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$, call it S^2 . Thus, $\beta + \alpha = 2\bar{X}$ and $(\alpha - \beta)^2 = 12S^2$, or $\beta - \alpha = 2S\sqrt{3}$, so that the moment estimates of α and β are: $\tilde{\alpha} = \bar{X} - S\sqrt{3}$, $\tilde{\beta} = \bar{X} + S\sqrt{3}$. These estimates are entirely different from the MLE's of these parameters.

Exercises

- 5.1** Refer to Exercise 1.6, and derive the moment estimate of θ . Also, compare it with the MLE $\hat{\theta} = 1/\bar{X}$.
- 5.2** (i) Refer to Exercise 1.7, and derive the moment estimate of θ , $\tilde{\theta}$.
 (ii) Find the numerical values of $\tilde{\theta}$ and of the MLE $\hat{\theta}$ (see Exercise 1.7), if: $n = 10$ and:

$$\begin{aligned} x_1 = 0.92, \quad x_2 = 0.79, \quad x_3 = 0.90, \quad x_4 = 0.65, \quad x_5 = 0.86, \\ x_6 = 0.47, \quad x_7 = 0.73, \quad x_8 = 0.97, \quad x_9 = 0.94, \quad \text{and} \quad x_{10} = 0.77. \end{aligned}$$

- 5.3** Refer to Exercise 1.8, and derive the moment estimate of θ .
- 5.4** Refer to Exercise 1.9 and show that $E_{\theta}|X| = \theta$, and therefore the moment estimate of θ is $\tilde{\theta} = \frac{1}{n} \sum_{i=1}^n |X_i|$.
- 5.5** Refer to Exercise 1.10 and find the expectation of the given p.d.f. by recalling that, if $X \sim \text{Gamma}$ with parameters α and β , then $EX = \alpha\beta$ (and $\text{Var}(X) = \alpha\beta^2$). Then derive the moment estimate of θ , and compare it with the MLE found in Exercise 1.10(ii).
- 5.6** Refer to Exercise 1.11, and:
- Show that $EX = \alpha + \beta$ and $EX^2 = \alpha^2 + 2\alpha\beta + 2\beta^2$, where X is a r.v. with the p.d.f. given in the exercise cited. Also, calculate the $\text{Var}(X)$.
 - Derive the moment estimates of α and β .
- 5.7** Let X_1, \dots, X_n be independent r.v.'s from the $U(\theta - a, \theta + b)$ distribution, where a and b are positive constants and $\theta \in \Omega = \Re$. Determine the moment estimate $\tilde{\theta}$ of θ , and compute its expectation and variance.
- 5.8** If the independent r.v.'s X_1, \dots, X_n have the $U(-\theta, \theta)$ distribution, $\theta \in \Omega = (0, \infty)$, how can one construct a moment estimate of θ ?
- 5.9** If the independent r.v.'s X_1, \dots, X_n have the Gamma distribution with parameters α and β , show that the moment estimates of α and β are: $\tilde{\alpha} = \bar{X}^2/S^2$ and $\tilde{\beta} = S^2/\bar{X}$, where $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$.
- Hint:** Recall that, if $X \sim \text{Gamma}$ with parameters α and β , then $EX = \alpha\beta$, $\text{Var}(X) = \alpha\beta^2$.
- 5.10** Let X be a r.v. with p.d.f. $f(x; \theta) = \frac{2}{\theta^2}(\theta - x)$, $0 < x < \theta$, $\theta \in \Omega = (0, \infty)$. Then:
- Show that $f(\cdot; \theta)$ is, indeed, a p.d.f.
 - Show that $E_{\theta}X = \frac{\theta}{3}$ and $\text{Var}_{\theta}(X) = \frac{\theta^2}{18}$.
 - On the basis of a random sample of size n from $f(\cdot; \theta)$, find the moment estimate of θ , $\tilde{\theta}$, and show that it is unbiased. Also, calculate the variance of $\tilde{\theta}$.
- 5.11** Let X be a r.v. having the Beta p.d.f. with parameters α and β ; i.e., $f(x; \alpha, \beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1}(1-x)^{\beta-1}$, $0 < x < 1$ ($\alpha, \beta > 0$). Then, by Exercise 4.2, $EX = \alpha/(\alpha + \beta)$.
- Follow the same approach used in proving Exercise 4.2 in order to establish that $EX^2 = \alpha(\alpha + 1)/(\alpha + \beta)(\alpha + \beta + 1)$.
 - On the basis of a random sample of size n from the underlying p.d.f., determine the moment estimates of α and β .
- 5.12** Let X and Y be *any* two r.v.'s with finite second moments, so that their correlation coefficient, $\rho(X, Y)$, is given by $\rho(X, Y) = \text{Cov}(X, Y)/\sigma(X)\sigma(Y)$. Let X_i and Y_i , $i = 1, \dots, n$ be i.i.d. r.v.'s distributed as the r.v.'s X and

Y , respectively. From the expression, $Cov(X, Y) = E[(X - EX) \times (Y - EY)]$, it makes sense to estimate $\rho(X, Y)$ by $\hat{\rho}_n(X, Y)$ give by:

$$\hat{\rho}(X, Y) = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y}) / \hat{\sigma}(X)\hat{\sigma}(Y),$$

where

$$\hat{\sigma}(x) = \sqrt{\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2} \quad \text{and} \quad \hat{\sigma}(Y) = \sqrt{\frac{1}{n} \sum_{i=1}^n (Y_i - \bar{Y})^2}.$$

Then set $EX = \mu_1$, $EY = \mu_2$, $\text{Var}(X) = \sigma_1^2$, $\text{Var}(Y) = \sigma_2^2$, $\rho(X, Y) = \rho$, and show that:

- (i) $\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y}) = \frac{1}{n} \sum_{i=1}^n (X_i Y_i) - \bar{X}\bar{Y}$.
- (ii) $E(XY) = \sigma_1\sigma_2 \text{Cov}(X, Y) + \mu_1\mu_2 = \rho\sigma_1\sigma_2 + \mu_1\mu_2$.
- (iii) Use the WLLN (Theorem 3 in Chapter 7) in conjunction with the Corollary to Theorem 5 in Chapter 7 in order to show that $\hat{\rho}_n(X, Y) \xrightarrow{P} \rho(X, Y) = \rho$, so that $\hat{\rho}_n(X, Y)$ is consistent (in the probability sense) estimate of ρ .

(Notice that $\hat{\rho}_n(X, Y)$ is the same as the MLE of ρ for the case that the pair (X, Y) has the Bivariate Normal distribution; see Exercise 1.14 in this chapter.)

- 5.13** (i) For any n pairs of real numbers (α_i, β_i) , $i = 1, \dots, n$, show that:
 $(\sum_{i=1}^n \alpha_i \beta_i)^2 \leq (\sum_{i=1}^n \alpha_i^2)(\sum_{i=1}^n \beta_i^2)$.

Hint: One way of proving it is to consider the function in λ , $g(\lambda) = \sum_{i=1}^n (\alpha_i - \lambda \beta_i)^2$, and observe that $g(\lambda) \geq 0$ for all real λ , and, in particular, for $\lambda = (\sum_{i=1}^n \alpha_i \beta_i) / (\sum_{i=1}^n \beta_i^2)$, which is actually the minimizing value for $g(\lambda)$.

- (ii) Use part (i) in order to show that $[\hat{\rho}_n(X, Y)]^2 \leq 1$.

- 5.14** In reference to Example 25 in Chapter 1, denote by x_i and y_i , $i = 1, \dots, 15$, respectively, the observed measurements for the cross-fertilized and the self-fertilized pairs. Then calculate the (observed) sample means \bar{x} , \bar{y} , sample variances s_x^2 , s_y^2 , and the sample s.d.'s s_x , s_y .

Confidence Intervals and Confidence Regions

In Section 2 of Chapter 8, the basic concepts about confidence intervals etc. were introduced; the detailed discussion was deferred to the present chapter. The point estimation problem, in its simplest form, discussed extensively in the previous chapter, is as follows: On the basis of a random sample X_1, \dots, X_n with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and its observed values x_1, \dots, x_n , construct a point estimate of θ , call it $\hat{\theta} = \hat{\theta}(x_1, \dots, x_n)$. Thus, for example, in the $N(\theta, 1)$ case, we are invited to pinpoint a value of $\theta \in \mathfrak{R}$ as the (unknown to us but) true value of θ . Such estimates were, actually, constructed by way of at least three methods. Also, certain desirable properties of estimates (fixed sample size properties, as well as asymptotic properties) were established or stated.

Now, declaring that (the unknown value of) θ is, actually, \bar{x} may look quite unreasonable. How is it possible to single out one value out of \mathfrak{R} , \bar{x} , and identify it as the true value of θ ? The concept of a confidence interval with a given confidence coefficient mitigates this seemingly unreasonable situation. It makes much more sense to declare that θ lies within an interval in \mathfrak{R} with high confidence. This is, in effect, what we are doing in this chapter by formulating the questions and problems in rigorous probabilistic/statistical terms.

The chapter consists of four sections. The first section concerns itself with confidence intervals of one real-valued parameter. The following section considers the same kind of a problem when *nuisance* (unknown but of no interest to us) parameters are present. In the third section, an example is discussed, where a confidence region of two parameters is constructed; no general theory is developed. (See, however, Theorem 4 in Chapter 12.) In the final section, some confidence intervals are constructed with given approximate confidence coefficient.

10.1 Confidence Intervals

We formalize in the form of a definition some concepts already introduced in the second section of Chapter 8.

DEFINITION 1

Let X_1, \dots, X_n be a random sample with p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \Re$. Then:

- (i) A *random interval* is an interval whose end-points are r.v.'s.
- (ii) A *confidence interval* for θ with *confidence coefficient* $1 - \alpha$ ($0 < \alpha < 1$, α small) is a random interval whose end-points are statistics $L(X_1, \dots, X_n)$ and $U(X_1, \dots, X_n)$, say, such that $L(X_1, \dots, X_n) \leq U(X_1, \dots, X_n)$ and

$$P_\theta[L(X_1, \dots, X_n) \leq \theta \leq U(X_1, \dots, X_n)] \geq 1 - \alpha, \quad \text{for all } \theta \in \Omega.$$

(1)

- (iii) The statistic $L(X_1, \dots, X_n)$ is called a *lower confidence limit* for θ with *confidence coefficient* $1 - \alpha$, if the interval $[L(X_1, \dots, X_n), \infty)$ is a confidence interval for θ with confidence coefficient $1 - \alpha$. Likewise, $U(X_1, \dots, X_n)$ is said to be an *upper confidence limit* for θ with *confidence coefficient* $1 - \alpha$, if the interval $(-\infty, U(X_1, \dots, X_n)]$ is a confidence interval for θ with confidence coefficient $1 - \alpha$.
-

REMARK 1 The significance of a confidence interval stems from the relative frequency interpretation of probability. Thus, on the basis of the observed values x_1, \dots, x_n of X_1, \dots, X_n , construct the interval with end-points $L(x_1, \dots, x_n)$ and $U(x_1, \dots, x_n)$, and denote it by $[L_1, U_1]$. Repeat the underlying random experiment independently another n times and likewise form the interval $[L_2, U_2]$. Repeat this process a large number of times N independently each time, and let $[L_N, U_N]$ be the corresponding interval. Then the fact that $[L(X_1, \dots, X_n), U(X_1, \dots, X_n)]$ is a confidence interval for θ with confidence coefficient $1 - \alpha$ means that approximately $100(1 - \alpha)\%$ of the above N intervals will cover θ , no matter what its value is.

REMARK 2 When the underlying r.v.'s are of the continuous type, the inequalities in the above definition, regarding the confidence coefficient $1 - \alpha$, become equalities.

REMARK 3 If $L(X_1, \dots, X_n)$ is a lower confidence limit for θ with confidence coefficient $1 - \frac{\alpha}{2}$, and $U(X_1, \dots, X_n)$ is an upper confidence limit for θ with confidence coefficient $1 - \frac{\alpha}{2}$, then $[L(X_1, \dots, X_n), U(X_1, \dots, X_n)]$ is a confidence interval for θ with confidence coefficient $1 - \alpha$.

Indeed, writing L and U instead of $L(X_1, \dots, X_n)$ and $U(X_1, \dots, X_n)$, and keeping in mind that $L \leq U$, we have:

$$\begin{aligned} P_\theta(L \leq \theta) &= P_\theta(L \leq \theta, U \geq \theta) + P_\theta(L \leq \theta, U < \theta) \\ &= P_\theta(L \leq \theta \leq U) + P_\theta(U < \theta), \quad \text{since } (U < \theta) \subseteq (L \leq \theta), \end{aligned}$$

and

$$\begin{aligned} P_\theta(\theta \leq U) &= P_\theta(U \geq \theta, L \leq \theta) + P_\theta(U \geq \theta, L > \theta) \\ &= P_\theta(L \leq \theta \leq U) + P_\theta(L > \theta), \quad \text{since } (L > \theta) \subseteq (U \geq \theta). \end{aligned}$$

Summing them up, we have then

$$P_\theta(L \leq \theta) + P_\theta(U \geq \theta) = 2P_\theta(L \leq \theta \leq U) + P_\theta(U < \theta) + P_\theta(L > \theta),$$

or

$$\begin{aligned} 2P_\theta(L \leq \theta \leq U) &= P_\theta(L \leq \theta) + P_\theta(U \geq \theta) - P_\theta(U < \theta) - P_\theta(L > \theta) \\ &= P_\theta(L \leq \theta) + P_\theta(U \geq \theta) - 1 + P_\theta(U \geq \theta) - 1 + P_\theta(L \leq \theta) \\ &= 2[P_\theta(L \leq \theta) + P_\theta(U \geq \theta) - 1], \end{aligned}$$

or

$$\begin{aligned} P_\theta(L \leq \theta \leq U) &= P_\theta(L \leq \theta) + P_\theta(U \geq \theta) - 1 \\ &\geq 1 - \frac{\alpha}{2} + 1 - \frac{\alpha}{2} - 1 = 1 - \alpha, \end{aligned}$$

as was to be seen.

This section is concluded with the construction of confidence intervals in some concrete examples. In so doing, we draw heavily on distribution theory and point estimates. It would be, perhaps, helpful to outline the steps we usually follow in constructing a confidence interval.

- Think of a r.v. which contains the parameter θ , the r.v.'s X_1, \dots, X_n , preferably in the form of a sufficient statistic, and whose distribution is (exactly or at least approximately) known.
- Determine suitable points $a < b$ such that the r.v. in step (a) lies in $[a, b]$ with P_θ -probability $\geq 1 - \alpha$.
- In the expression of step (b), rearrange the terms to arrive at an interval with the end-points being statistics and containing θ .
- The interval in step (c) is the required confidence interval.

EXAMPLE 1

Let X_1, \dots, X_n be a random interval from the $N(\mu, \sigma^2)$ distribution, where only one of μ or σ^2 is unknown. Construct a confidence interval for it with confidence coefficient $1 - \alpha$.

DISCUSSION

- Let μ be unknown. The natural r.v. to think of is $\sqrt{n}(\bar{X} - \mu)/\sigma$, which satisfies the requirements in step (a). Next, determine any two points $a < b$

from the Normal tables for which $P(a \leq Z \leq b) = 1 - \alpha$ where $Z \sim N(0, 1)$. (See, however, Exercise 1.1 for the best choice of a and b .) Since $\frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \sim N(0, 1)$, it follows that

$$P_{\mu} \left[a \leq \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \leq b \right] = 1 - \alpha, \quad \text{for all } \mu,$$

so that step (b) is satisfied. Rearranging the terms inside the square brackets, we obtain:

$$P_{\mu} \left(\bar{X} - \frac{b\sigma}{\sqrt{n}} \leq \mu \leq \bar{X} - \frac{a\sigma}{\sqrt{n}} \right) = 1 - \alpha, \quad \text{for all } \mu,$$

so that step (c) is fulfilled. In particular, for $b = z_{\alpha/2}$ (recall $P(Z \geq z_{\alpha/2}) = \frac{\alpha}{2}$) and $a = -z_{\alpha/2}$, we have

$$P_{\mu} \left(\bar{X} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{X} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right) = 1 - \alpha, \quad \text{for all } \mu.$$

It follows that

$$\left[\bar{X} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}, \bar{X} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right] = \bar{X} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \quad (\text{for brevity}) \quad (2)$$

is the required confidence interval.

(ii) Let σ^2 be unknown. Set $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2$ and recall that $\frac{nS^2}{\sigma^2} = \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 \sim \chi_n^2$. The r.v. $\frac{nS^2}{\sigma^2}$ satisfies the requirements of step (a). From the Chi-Square tables, determine any pair $0 < a < b$ for which $P(a \leq X \leq b) = 1 - \alpha$, where $X \sim \chi_n^2$. Then

$$\begin{aligned} P_{\sigma^2} \left(a \leq \frac{nS^2}{\sigma^2} \leq b \right) &= 1 - \alpha, \quad \text{for all } \sigma^2, \quad \text{or } P_{\sigma^2} \left(\frac{nS^2}{b} \leq \sigma^2 \leq \frac{nS^2}{a} \right) \\ &= 1 - \alpha, \quad \text{for all } \sigma^2 \text{ and steps (b) and (c) are satisfied.} \end{aligned}$$

In particular,

$$P_{\sigma^2} \left(\frac{nS^2}{\chi_{n, \alpha/2}^2} \leq \sigma^2 \leq \frac{nS^2}{\chi_{n, 1-\alpha/2}^2} \right) = 1 - \alpha, \quad \text{for all } \sigma^2,$$

where $P(X \leq \chi_{n, 1-\alpha/2}^2) = P(X \geq \chi_{n, \alpha/2}^2) = \frac{\alpha}{2}$. It follows that

$$\left[\frac{nS^2}{\chi_{n, \alpha/2}^2}, \frac{nS^2}{\chi_{n, 1-\alpha/2}^2} \right], \quad S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2 \quad (3)$$

is the required confidence interval.

Numerical Example Let $n = 25$ and $1 - \alpha = 0.95$. For part (i), we have $z_{\alpha/2} = z_{0.025} = 1.96$, so that $\bar{X} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}} = \bar{X} \pm 1.96 \times \frac{\sigma}{5} = \bar{X} \pm 0.392\sigma$. For $\sigma = 1$, for example, the required interval is then: $\bar{X} \pm 0.392$. For the second part, we have $\chi_{n, \alpha/2}^2 = \chi_{25, 0.025}^2 = 40.646$, and $\chi_{n, 1-\alpha/2}^2 = \chi_{25, 0.975}^2 = 13.120$. The required interval is then:

$$\left[\frac{25S^2}{40.646}, \frac{25S^2}{13.120} \right] \simeq [0.615S^2, 1.905S^2].$$

EXAMPLE 2

On the basis of the random sample X_1, \dots, X_n from the $U(0, \theta)$ ($\theta > 0$) distribution, construct a confidence interval for θ with confidence coefficient $1 - \alpha$.

DISCUSSION It has been seen (just apply Example 6(ii) in Chapter 9 with $\alpha = 0$ and $\beta = \theta$) that $X = X_{(n)}$ is a sufficient statistic for θ . Also, the p.d.f. of X is given by (see Example 14 in Chapter 9) $f_X(x; \theta) = \frac{n}{\theta^n} x^{n-1}$, $0 \leq x \leq \theta$. Setting $Y = X/\theta$, it is easily seen that the p.d.f. of Y is: $f_Y(y) = ny^{n-1}$, $0 \leq y \leq 1$. The r.v. Y satisfies the requirements of step (a). Next, determine any $0 \leq a < b < 1$ such that $\int_a^b f_Y(y) dy = \int_a^b ny^{n-1} dy = \int_a^b dy^n = y^n|_a^b = b^n - a^n = 1 - \alpha$. Then

$$P_\theta(a \leq Y \leq b) = P_\theta\left(a \leq \frac{X}{\theta} \leq b\right) = P_\theta\left(\frac{X}{b} \leq \theta \leq \frac{X}{a}\right) = 1 - \alpha, \quad \text{for all } \theta,$$

so that steps (b) and (c) are satisfied. It follows that $[\frac{X}{b}, \frac{X}{a}] = [\frac{X_{(n)}}{b}, \frac{X_{(n)}}{a}]$ is the required confidence interval.

Looking at the length of this interval, $X_{(n)}(\frac{1}{a} - \frac{1}{b})$, setting $a = a(b)$ and minimizing with respect to b , we find that the shortest interval is taken for $b = 1$ and $a = \alpha^{1/n}$. That is, $[X_{(n)}, \frac{X_{(n)}}{\alpha^{1/n}}]$. (See also Exercise 1.5.)

Numerical Example For $n = 32$ and $1 - \alpha = 0.95$, we get (approximately) $[X_{(32)}, 1.098X_{(32)}]$.

Exercises

- 1.1 Let Φ be the d.f. of the $N(0, 1)$ distribution, and let $a < b$ be such that $\Phi(b) - \Phi(a) = \gamma$, some fixed number with $0 < \gamma < 1$. Show that the length $b - a$ of the interval (a, b) is minimum, if, for some $c > 0$, $b = c$ and $a = -c$.
- 1.2 If X_1, \dots, X_n are independent r.v.'s distributed as $N(\mu, \sigma^2)$ with μ unknown and σ known, then a $100(1 - \alpha)\%$ confidence interval for μ is given by $\bar{X}_n \pm z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}$ (see Example 1(i)). Suppose that the length of this interval is 7.5 and we wish to halve it. What sample size $m = m(n)$ will be needed?

Hint: Set $m = cn$ and determine c .
- 1.3 The stray-load loss (in watts) for a certain type of induction motor, when the line current is held at 10 amps for a speed of 1,500 rpm, is a r.v. $X \sim N(\mu, 9)$.
(i) Compute a 99% confidence interval for μ when $n = 100$ and $\bar{x} = 58.3$.
(ii) Determine the sample size n , if the length of the 99% confidence interval is required to be 1.
- 1.4 If the independent r.v.'s X_1, \dots, X_n are distributed as $N(\theta, \sigma^2)$ with σ known, the $100(1 - \alpha)\%$ confidence interval for θ is given by $\bar{X}_n \pm z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}$ (see Example 1(i)).

- (i) If the length of the confidence interval is to be equal to a preassigned number l , determine the sample size n as a function of l , σ , and α .
- (ii) Compute the numerical value of n , if $l = 0.1$, $\sigma = 1$, and $\alpha = 0.05$.

1.5 Refer to Example 2 and show that the shortest length of the confidence interval is, indeed, $[X_{(n)}, X_{(n)}/\alpha^{1/n}]$ as asserted.

Hint: Set $a = a(b)$, differentiate $g(b) = \frac{1}{a} - \frac{1}{b}$, with respect to b , and use the derivative of $b^n - a^n = 1 - \alpha$ in order to show that $\frac{dg(b)}{db} < 0$, so that $g(b)$ is decreasing. Conclude that $g(b)$ is minimized for $b = 1$.

1.6 Let X_1, \dots, X_n be independent r.v.'s with the Negative Exponential p.d.f. given in the form $f(x; \theta) = \frac{1}{\theta}e^{-x/\theta}$, $x > 0$, $\theta \in \Omega = (0, \infty)$. Then:

- (i) By using the m.g.f. approach, show that the r.v. $U = \sum_{i=1}^n X_i$ has the Gamma distribution with parameters $\alpha = n$ and $\beta = \theta$.
- (ii) Also, show that the r.v. $V = \frac{2U}{\theta}$ is distributed as χ_{2n}^2 .
- (iii) By means of part (ii), construct a confidence interval for θ with confidence coefficient $1 - \alpha$.

1.7 If X is a r.v. with the Negative Exponential p.d.f. $f(x; \theta) = \frac{1}{\theta}e^{-x/\theta}$, $x > 0$, $\theta \in \Omega = (0, \infty)$, then, by Exercise 2.2 in Chapter 9, the reliability $R(x; \theta) = P_\theta(X > x) = e^{-x/\theta}$. If X_1, \dots, X_n is a random sample of size n from this p.d.f., use Exercise 1.6(iii) in order to construct a confidence interval for $R(x; \theta)$ with confidence coefficient $1 - \alpha$.

1.8 Let X_1, \dots, X_n be a random sample of size n from the p.d.f. $f(x; \theta) = e^{-(x-\theta)}$, $x > \theta$, $\theta \in \Omega = \Re$, and let Y_1 be the smallest order statistic of the X_i 's.

- (i) Use formula (28) in order to show that the p.d.f. of Y_1 , call it g , is given by: $g(y) = ne^{-n(y-\theta)}$, $y > \theta$.
- (ii) Set $T(\theta) = 2n(Y_1 - \theta)$ and show that $T \sim \chi_2^2$.
- (iii) Use part (ii) in order to show that a $100(1 - \alpha)\%$ confidence interval for θ , based on $T(\theta)$, is given by: $[Y_1 - \frac{b}{2n}, Y_1 - \frac{a}{2n}]$, for suitable $0 < a < b$; a special choice of a and b is: $a = \chi_{2, 1-\frac{\alpha}{2}}^2$ and $b = \chi_{2, \frac{\alpha}{2}}^2$.

1.9 Let the independent r.v.'s X_1, \dots, X_n have the Weibull distribution with parameters γ and θ with $\theta \in \Omega = (0, \infty)$ and $\gamma > 0$ known; i.e., their p.d.f. $f(\cdot; \theta)$ is given by:

$$f(x; \theta) = \frac{\gamma}{\theta} x^{\gamma-1} e^{-x^\gamma/\theta}, \quad x > 0.$$

- (i) For $i = 1, \dots, n$, set $Y_i = X_i^\gamma$ and show that the p.d.f. of Y_i , $g(\cdot; \theta)$, is Negative Exponential parameterized as follows: $g(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$, $y > 0$.
- (ii) For $i = 1, \dots, n$, set $T_i(\theta) = \frac{2Y_i}{\theta}$ and show that the p.d.f. of $T_i(\theta)$, $g_T(\cdot; \theta)$, is that of a χ_2^2 distributed r.v., and conclude that the r.v. $T(\theta) = \sum_{i=1}^n T_i(\theta) \sim \chi_{2n}^2$.

(iii) Show that a $100(1 - \alpha)\%$ confidence interval for θ , based on $T(\theta)$, is of the form $[\frac{2Y}{b}, \frac{2Y}{a}]$, for suitable $0 < a < b$, where $Y = \sum_{i=1}^n X_i^y$. In particular, a and b may be chosen to be $\chi_{2n; 1-\frac{\alpha}{2}}^2$ and $\chi_{2n; \frac{\alpha}{2}}^2$, respectively.

1.10 If the independent r.v.'s X_1, \dots, X_n have p.d.f. $f(x; \theta) = \frac{1}{2\theta} e^{-|x|/\theta}$, $x \in \mathfrak{R}$, $\theta \in \Omega = (0, \infty)$, then:

- (i) The independent r.v.'s $Y_i = |X_i|$, $i = 1, \dots, n$ have the Negative Exponential p.d.f. $g(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$, $y > 0$.
- (ii) The independent r.v.'s $T_i(\theta) = \frac{2Y_i}{\theta}$, $i = 1, \dots, n$ are χ_2^2 -distributed, so that the r.v. $T(\theta) = \sum_{i=1}^n T_i(\theta) = \frac{2}{\theta} \sum_{i=1}^n Y_i = \frac{2Y}{\theta} \sim \chi_{2n}^2$, where $Y = \sum_{i=1}^n Y_i = \sum_{i=1}^n |X_i|$.
- (iii) A $100(1 - \alpha)\%$ confidence interval for θ , based on $T(\theta)$, is given by $[\frac{2Y}{b}, \frac{2Y}{a}]$, for suitable $0 < a < b$. In particular, a and b may be chosen to be $a = \chi_{2n; 1-\frac{\alpha}{2}}^2$, $b = \chi_{2n; \frac{\alpha}{2}}^2$.

1.11 Consider the p.d.f. $f(x; \alpha, \beta) = \frac{1}{\beta} e^{-(x-\alpha)/\beta}$, $x \geq \alpha$, $\alpha \in \mathfrak{R}$, $\beta > 0$ (see Exercise 1.11 in Chapter 9), and suppose that β is known and α is unknown, and denote it by θ . Thus, we have here:

$$f(x; \theta) = \frac{1}{\beta} e^{-(x-\theta)/\beta}, \quad x \geq \theta, \quad \theta \in \Omega = \mathfrak{R}.$$

- (i) Show that the corresponding d.f., $F(\cdot; \theta)$, is given by: $F(x; \theta) = 1 - e^{-(x-\theta)/\beta}$, $x \geq \theta$, so that $1 - F(x; \theta) = e^{-(x-\theta)/\beta}$, $x \geq \theta$.
- (ii) Let X_1, \dots, X_n be independent r.v.'s drawn from the p.d.f. $f(\cdot; \theta)$, and let Y_1 be the smallest order statistic. Use relation (28) in Chapter 6 in order to show that the p.d.f. of Y_1 is given by:

$$f_{Y_1}(y; \theta) = \frac{n}{\beta} e^{-n(y-\theta)/\beta}, \quad y \geq \theta.$$

- (iii) Consider the r.v. $T = T_n(\theta)$ defined by: $T = n(Y_1 - \theta)/\beta$, and show that its p.d.f. is given by: $f_T(t) = e^{-t}$, $t \geq 0$.

1.12 In reference to Exercise 1.11:

- (i) Determine $0 \leq a < b$, so that $P(a \leq T \leq b) = 1 - \alpha$, for some $0 < \alpha < 1$.
- (ii) By part (i), $P_\theta[a \leq \frac{n(Y_1 - \theta)}{\beta} \leq b] = 1 - \alpha$, since T has the p.d.f. $f_T(t) = e^{-t}$, $t \geq 0$. Use this relation to conclude that $[Y_1 - \frac{b\beta}{n}, Y_1 - \frac{a\beta}{n}]$ is a $100(1 - \alpha)\%$ confidence interval of θ .
- (iii) The length l of the confidence interval in part (ii) is $l = \frac{\beta}{n}(b - a)$. Set $b = b(a)$ and show that the shortest confidence interval is given by: $[Y_1 + \frac{b \log \alpha}{n}, Y_1]$.

Hint: For part (iii), set $b = b(a)$, and from $e^{-a} - e^{-b} = 1 - \alpha$, obtain $\frac{db}{da} = e^{b-a}$ by differentiation. Then replace $\frac{db}{da}$ in $\frac{dl}{da}$ and observe that it is > 0 . This implies that l obtain its minimum at $a = 0$.

1.13 Let X_1, \dots, X_n be independent r.v.'s with d.f. F and p.d.f. f with $f(x) > 0$ for $-\infty \leq a < x < b \leq \infty$, and let Y_1 and Y_n be, respectively, the smallest and the largest order statistics of the X_i 's.

(i) By using the hint given below, show that the joint p.d.f., f_{Y_1, Y_n} , of Y_1 and Y_n is given by:

$$f_{Y_1, Y_n}(y_1, y_n) = n(n-1)[F(y_n) - F(y_1)]^{n-2} f(y_1) f(y_n),$$

$$a < y_1 < y_n < b.$$

Hint: $P(Y_n \leq y_n) = P(Y_1 \leq y_1, Y_n \leq y_n) + P(Y_1 > y_1, Y_n \leq y_n) = F_{Y_1, Y_n}(y_1, y_n) + P(y_1 < Y_1 < Y_n \leq y_n)$. But $P(Y_n \leq y_n) = P(\text{all } X_i\text{'s} \leq y_n) = P(X_1 \leq y_n, \dots, X_n \leq y_n) = P(X_1 \leq y_n) \dots P(X_n \leq y_n) = [F(y_n)]^n$, and: $P(y_1 < Y_1 < Y_n \leq y_n) = P(\text{all } X_i\text{'s are } > y_1 \text{ and also } \leq y_n) = P(y_1 < X_1 \leq y_n, \dots, y_1 < X_n \leq y_n) = P(y_1 < X_1 \leq y_n) \dots P(y_1 < X_n \leq y_n) = [P(y_1 < X_1 \leq y_n)]^n = [F(y_n) - F(y_1)]^n$. Thus,

$$[F(y_n)]^n = F_{Y_1, Y_n}(y_1, y_n) + [F(y_n) - F(y_1)]^n, \quad a < y_1 < y_n < b.$$

Solving for $F_{Y_1, Y_n}(y_1, y_n)$ and taking the partial derivatives with respect to y_1 and y_n , we get the desired result.

(ii) Find the p.d.f. f_{Y_1, Y_n} when the X_i 's are distributed as $U(0, \theta)$, $\theta \in \Omega = (0, \infty)$.

(iii) Do the same for the case the X_i 's have the Negative Exponential p.d.f. $f(x) = \frac{1}{\theta} e^{-\theta x}$, $x > 0$, $\theta \in \Omega = (0, \infty)$.

1.14 Refer to Exercise 1.13(ii), and show that the p.d.f. of the range $R = Y_n - Y_1$ is given by:

$$f_R(r; \theta) = \frac{n(n-1)}{\theta^n} r^{n-2} (\theta - r), \quad 0 < r < \theta.$$

1.15 Refer to Exercise 1.13(iii), and show that the p.d.f. of the range $R = Y_n - Y_1$ is given by:

$$f_R(r; \theta) = \frac{n-1}{\theta} e^{-\frac{r}{\theta}} (1 - e^{-\frac{r}{\theta}})^{n-2}, \quad r > 0.$$

1.16 In reference to Exercise 1.14:

(i) Set $T = \frac{R}{\theta}$ and show that $f_T(t) = n(n-1)t^{n-2}(1-t)$, $0 < t < 1$.

(ii) Take $0 < c < 1$ such that $P_\theta(c \leq T \leq 1) = 1 - \alpha$, and construct a confidence interval for θ , based on the range R , with confidence coefficient $1 - \alpha$. Also, show that c is a root of the equation $c^{n-1} \times [n - (n-1)c] = \alpha$.

1.17 Consider the independent random samples X_1, \dots, X_m from the $N(\mu_1, \sigma_1^2)$ distribution and Y_1, \dots, Y_n from the $N(\mu_2, \sigma_2^2)$ distribution, where μ_1, μ_2 are unknown and σ_1^2, σ_2^2 are known, and define the r.v. $T = T_{m,n}(\mu_1 - \mu_2)$ by: $T_{m,n}(\mu_1 - \mu_2) = \frac{(\bar{X}_m - \bar{Y}_n) - (\mu_1 - \mu_2)}{\sqrt{(\sigma_1^2/m) + (\sigma_2^2/n)}}$.

Then show that:

- (i) A $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$, based on T , is given by: $[(\bar{X}_m - \bar{Y}_n) - b\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}, (\bar{X}_m - \bar{Y}_n) - a\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}]$ for suitable constants a and b .
- (ii) The confidence interval in part (i) with the shortest length is taken for $b = z_{\alpha/2}$ and $a = -z_{\alpha/2}$.

1.18 Refer to Exercise 1.17, and suppose that μ_1, μ_2 are known and σ_1^2, σ_2^2 are unknown. Then define the r.v. $\bar{T} = \bar{T}_{m,n}(\sigma_1^2/\sigma_2^2) = \frac{\sigma_1^2 S_X^2}{\sigma_2^2 S_Y^2}$, where $S_X^2 = \frac{1}{m} \sum_{i=1}^m (X_i - \mu_1)^2$ and $S_Y^2 = \frac{1}{n} \sum_{j=1}^n (Y_j - \mu_2)^2$, and show that a $100(1 - \alpha)\%$ confidence interval for σ_1^2/σ_2^2 , based on \bar{T} , is given by $[a \frac{S_X^2}{S_Y^2}, b \frac{S_X^2}{S_Y^2}]$ for $0 < a < b$ with $P(a \leq X \leq b) = 1 - \alpha$, $X \sim F_{n,m}$. In particular, we may choose $a = F_{n,m;1-\frac{\alpha}{2}}$ and $b = F_{n,m;\frac{\alpha}{2}}$.

1.19 Consider the independent random samples X_1, \dots, X_m and Y_1, \dots, Y_n from the Negative Exponential distributions $f(x; \theta_1) = \frac{1}{\theta_1} e^{-x/\theta_1}$, $x > 0$, $\theta_1 \in \Omega = (0, \infty)$, and $f(y; \theta_2) = \frac{1}{\theta_2} e^{-y/\theta_2}$, $y > 0$, $\theta_2 \in \Omega = (0, \infty)$, and set $U = \sum_{i=1}^m X_i$, $V = \sum_{j=1}^n Y_j$. Then, by Exercise 1.6(ii), $\frac{2U}{\theta_1} \sim \chi_{2m}^2$, $\frac{2V}{\theta_2} \sim \chi_{2n}^2$ and they are independent. It follows that $\frac{\frac{2V}{\theta_2}}{\frac{2U}{\theta_1}/2m} = \frac{\theta_1}{\theta_2} \times \frac{mV}{nU} = \frac{\theta_1}{\theta_2} \times \frac{m \sum_{j=1}^n Y_j}{n \sum_{i=1}^m X_i} \sim F_{2n,2m}$. Use this result in order to construct a $100(1 - \alpha)\%$ confidence interval for θ_1/θ_2 .

10.2 Confidence Intervals in the Presence of Nuisance Parameters

In Example 1, the position was adopted that only one of the parameters in the $N(\mu, \sigma^2)$ distribution was unknown. This is a rather artificial assumption as, in practice, both μ and σ^2 are most often unknown. What was done in that example did, however, pave the way to solving the problem here in its natural setting.

EXAMPLE 3

Let X_1, \dots, X_n be a random sample from the $N(\mu, \sigma^2)$ distribution, where both μ and σ^2 are unknown. Construct confidence intervals for μ and σ^2 , each with confidence coefficient $1 - \alpha$.

DISCUSSION We have that:

$$\frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \sim N(0, 1) \quad \text{and} \quad \frac{(n-1)S^2}{\sigma^2} = \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 \sim \chi_{n-1}^2,$$

where $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$, and these two r.v.'s are independent. It follows that $\frac{\sqrt{n}(\bar{X} - \mu)/\sigma}{\sqrt{(n-1)S^2/\sigma^2(n-1)}} = \frac{\sqrt{n}(\bar{X} - \mu)}{S} \sim t_{n-1}$. From the t -tables, determine any pair

(a, b) with $a < b$ such that $P(a \leq X \leq b) = 1 - \alpha$, where $X \sim t_{n-1}$. It follows that:

$$P_{\theta} \left[a \leq \frac{\sqrt{n}(\bar{X} - \mu)}{S} \leq b \right] = 1 - \alpha, \quad \text{for all } \theta = (\mu, \sigma^2),$$

or

$$P_{\theta} \left(\bar{X} - b \frac{S}{\sqrt{n}} \leq \mu \leq \bar{X} - a \frac{S}{\sqrt{n}} \right) = 1 - \alpha, \quad \text{for all } \theta.$$

In particular,

$$P_{\theta} \left(\bar{X} - t_{n-1;\alpha/2} \frac{S}{\sqrt{n}} \leq \mu \leq \bar{X} + t_{n-1;\alpha/2} \frac{S}{\sqrt{n}} \right) = 1 - \alpha, \quad \text{for all } \theta,$$

where $P(X \geq t_{n-1;\alpha/2}) = \frac{\alpha}{2}$ (and $X \sim t_{n-1}$). It follows that the required confidence interval for μ is:

$$\left[\bar{X} - t_{n-1;\alpha/2} \frac{S}{\sqrt{n}}, \bar{X} + t_{n-1;\alpha/2} \frac{S}{\sqrt{n}} \right] = \bar{X} \pm t_{n-1;\alpha/2} \frac{S}{\sqrt{n}} \quad (\text{for brevity}). \quad (4)$$

The construction of a confidence interval for σ^2 in the presence of (an unknown) μ is easier. We have already mentioned that $\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$. Then repeat the process in Example 1(ii), replacing χ_n^2 by χ_{n-1}^2 , to obtain the confidence interval.

$$\left[\frac{(n-1)S^2}{\chi_{n-1;\alpha/2}^2}, \frac{(n-1)S^2}{\chi_{n-1;1-\alpha/2}^2} \right], \quad S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2. \quad (5)$$

REMARK 4 Observe that the confidence interval in (4) differs from that in (2) in that σ in (2) is replaced by an estimate S , and then the constant $z_{\alpha/2}$ in (2) is adjusted to $t_{n-1;\alpha/2}$. Likewise, the confidence intervals in (3) and (5) are of the same form, with the only difference that (the unknown) μ in (3) is replaced by its estimate \bar{X} in (5). The constants n , $\chi_{n;\alpha/2}^2$, and $\chi_{n;1-\alpha/2}^2$ are also adjusted as indicated in (5).

Numerical Example Let $n = 25$ and $1 - \alpha = 0.95$. Then $t_{n-1;\alpha/2} = t_{24;0.025} = 2.0639$, and the interval in (4) becomes $\bar{X} \pm 0.41278S$. Also, $\chi_{n-1;\alpha/2}^2 = \chi_{24;0.025}^2 = 39.364$, $\chi_{n-1;1-\alpha/2}^2 = \chi_{24;0.975}^2 = 12.401$, so that the interval in (5) is $\left[\frac{24S^2}{39.364}, \frac{24S^2}{12.401} \right] \simeq [0.610S^2, 1.935S^2]$.

Actually, a somewhat more important problem from a practical viewpoint is that of constructing confidence intervals for the difference of the means of two normal populations and the ratio of their variances. This is a way of comparing two normal populations. The precise formulation of the problem is given below.

EXAMPLE 4

Let X_1, \dots, X_m and Y_1, \dots, Y_n be two independent random samples from the $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$ distributions, respectively, with all μ_1, μ_2, σ_1^2 , and σ_2^2 unknown. We wish to construct confidence intervals for $\mu_1 - \mu_2$ and σ_1^2/σ_2^2 .

DISCUSSION

- (i) *Confidence interval for $\mu_1 - \mu_2$.* In order to be able to resolve this problem, we have to *assume* that the variances, although unknown, are equal; i.e., $\sigma_1^2 = \sigma_2^2 = \sigma^2$, say.

Let us review briefly some distribution results. Recall that $\bar{X} - \mu_1 \sim N(0, \frac{\sigma^2}{m})$, $\bar{Y} - \mu_2 \sim N(0, \frac{\sigma^2}{n})$, and by independence,

$$[(\bar{X} - \bar{Y}) - (\mu_1 - \mu_2)]/\sigma\sqrt{\frac{1}{m} + \frac{1}{n}} \sim N(0, 1). \quad (6)$$

Also, if

$$S_X^2 = \frac{1}{m-1} \sum_{i=1}^m (X_i - \bar{X})^2, \quad S_Y^2 = \frac{1}{n-1} \sum_{j=1}^n (Y_j - \bar{Y})^2,$$

then $\frac{(m-1)S_X^2}{\sigma^2} \sim \chi_{m-1}^2$, $\frac{(n-1)S_Y^2}{\sigma^2} \sim \chi_{n-1}^2$, and by independence,

$$\frac{(m-1)S_X^2 + (n-1)S_Y^2}{\sigma^2} \sim \chi_{m+n-2}^2. \quad (7)$$

From (6) and (7), we obtain then:

$$\frac{(\bar{X} - \bar{Y}) - (\mu_1 - \mu_2)}{\sqrt{\frac{(m-1)S_X^2 + (n-1)S_Y^2}{m+n-2} \left(\frac{1}{m} + \frac{1}{n}\right)}} \sim t_{m+n-2}. \quad (8)$$

Then working with (8) as in Example 1(i), we arrive at the following confidence interval

$$\begin{aligned} & \left[(\bar{X} - \bar{Y}) - t_{m+n-2, \alpha/2} \sqrt{\frac{(m-1)S_X^2 + (n-1)S_Y^2}{m+n-2} \left(\frac{1}{m} + \frac{1}{n}\right)}, \right. \\ & \left. (\bar{X} - \bar{Y}) + t_{m+n-2, \alpha/2} \sqrt{\frac{(m-1)S_X^2 + (n-1)S_Y^2}{m+n-2} \left(\frac{1}{m} + \frac{1}{n}\right)} \right] \\ & = (\bar{X} - \bar{Y}) \pm t_{m+n-2, \alpha/2} \sqrt{\frac{(m-1)S_X^2 + (n-1)S_Y^2}{m+n-2} \left(\frac{1}{m} + \frac{1}{n}\right)}. \quad (9) \end{aligned}$$

- (ii) *Confidence interval for σ_1^2/σ_2^2 .* By the fact that $\frac{(m-1)S_X^2}{\sigma_1^2} \sim \chi_{m-1}^2$, $\frac{(n-1)S_Y^2}{\sigma_2^2} \sim \chi_{n-1}^2$, and independence, we have $\frac{S_Y^2/\sigma_2^2}{S_X^2/\sigma_1^2} = \frac{\sigma_1^2}{\sigma_2^2} \times \frac{S_Y^2/\sigma_1^2}{S_X^2/\sigma_1^2} \sim F_{n-1, m-1}$. From the F -tables, determine any pair (a, b) with $0 < a < b$ such that $P(a \leq X \leq b) = 1 - \alpha$, where $X \sim F_{n-1, m-1}$. Then, for all $\theta = (\mu_1, \mu_2, \sigma_1^2, \sigma_2^2)$,

$$P_\theta \left(a \leq \frac{\sigma_1^2}{\sigma_2^2} \times \frac{S_Y^2}{S_X^2} \leq b \right) = 1 - \alpha, \quad \text{or} \quad P_\theta \left(a \frac{S_X^2}{S_Y^2} \leq \frac{\sigma_1^2}{\sigma_2^2} \leq b \frac{S_X^2}{S_Y^2} \right) = 1 - \alpha.$$

In particular, for all θ ,

$$P_\theta \left(\frac{S_X^2}{S_Y^2} F_{n-1, m-1; 1-\alpha/2} \leq \frac{\sigma_1^2}{\sigma_2^2} \leq \frac{S_X^2}{S_Y^2} F_{n-1, m-1; \alpha/2} \right) = 1 - \alpha,$$

where $P(X \leq F_{n-1, m-1; 1-\alpha/2}) = P(X \geq F_{n-1, m-1; \alpha/2}) = \frac{\alpha}{2}$ (and $X \sim F_{n-1, m-1}$). The required confidence interval is then

$$\left[\frac{S_X^2}{S_Y^2} F_{n-1, m-1; 1-\alpha/2}, \frac{S_X^2}{S_Y^2} F_{n-1, m-1; \alpha/2} \right]. \quad (10)$$

Numerical Example Let $m = 13$, $n = 14$, and $1 - \alpha = 0.95$. Then $t_{m+n-2, \alpha/2} = t_{25; 0.025} = 2.0595$, so that the interval in (9) becomes

$$(\bar{X} - \bar{Y}) \pm 2.0595 \sqrt{\frac{12S_X^2 + 13S_Y^2}{25} \left(\frac{1}{13} + \frac{1}{14} \right)} \simeq (\bar{X} - \bar{Y}) \pm 0.1586 \sqrt{12S_X^2 + 13S_Y^2}.$$

Next, $F_{n-1, m-1; \alpha/2} = F_{13, 12; 0.025} = 3.2388$, $F_{n-1, m-1; 1-\alpha/2} = F_{13, 12; 0.975} = \frac{1}{F_{12, 13; 0.025}} = \frac{1}{3.1532} \simeq 0.3171$. Therefore the interval in (10) is $[0.3171 \frac{S_X^2}{S_Y^2}, 3.2388 \frac{S_X^2}{S_Y^2}]$.

Exercises

- 2.1 If the independent r.v.'s X_1, \dots, X_n are $N(\mu, \sigma^2)$ distributed with both μ and σ^2 unknown, construct a $100(1 - \alpha)\%$ confidence interval for σ .
- 2.2 Refer to Exercise 1.18 and suppose that all μ_1, μ_2 , and σ_1^2, σ_2^2 are unknown. Then construct a $100(1 - \alpha)\%$ confidence interval for σ_1^2/σ_2^2 .

10.3 A Confidence Region for (μ, σ^2) in the $N(\mu, \sigma^2)$ Distribution

Refer again to Example 1 and suppose that both μ and σ^2 are unknown, as is most often the case. In this section, we wish to construct a *confidence region* for the pair (μ, σ^2) ; i.e., a subset of the plane determined in terms of statistics and containing (μ, σ^2) with probability $1 - \alpha$. This problem is resolved in the following example.

EXAMPLE 5

On the basis of the random sample X_1, \dots, X_n from the $N(\mu, \sigma^2)$ distribution, construct a confidence region for the pair (μ, σ^2) with confidence coefficient $1 - \alpha$.

DISCUSSION In solving this problem, we draw heavily on what we have done in the previous example. Let \bar{X} be the sample mean and define S^2 by $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$. Then

$$\frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \sim N(0, 1), \quad \frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2 \quad (11)$$

and the two r.v.'s involved here are independent. From the Normal tables, define $c > 0$ so that $P(-c \leq Z \leq c) = \sqrt{1 - \alpha}$, $Z \sim N(0, 1)$; c is uniquely determined. From the χ^2 -tables, determine a pair (a, b) with $0 < a < b$ and

$P(a \leq X \leq b) = \sqrt{1 - \alpha}$, where $X \sim \chi_{n-1}^2$. Then, by means of (11), and with $\theta = (\mu, \sigma^2)$, we have:

$$P_{\theta} \left[-c \leq \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \leq c \right] = \sqrt{1 - \alpha}, \quad (12)$$

$$P_{\theta} \left[a \leq \frac{(n-1)S^2}{\sigma^2} \leq b \right] = \sqrt{1 - \alpha}.$$

These relations are rewritten thus:

$$P_{\theta} \left[-c \leq \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \leq c \right] = P_{\theta} \left[(\mu - \bar{X})^2 \leq \frac{c^2 \sigma^2}{n} \right] (= \sqrt{1 - \alpha}), \quad (13)$$

$$P_{\theta} \left[a \leq \frac{(n-1)S^2}{\sigma^2} \leq b \right] = P_{\theta} \left[\frac{(n-1)S^2}{b} \leq \sigma^2 \leq \frac{(n-1)S^2}{a} \right] (= \sqrt{1 - \alpha}), \quad (14)$$

so that, by means of (12)–(14) and independence, we have:

$$\begin{aligned} & P_{\theta} \left[-c \leq \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \leq c, a \leq \frac{(n-1)S^2}{\sigma^2} \leq b \right] \\ &= P_{\theta} \left[-c \leq \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \leq c \right] P_{\theta} \left[a \leq \frac{(n-1)S^2}{\sigma^2} \leq b \right] \\ &= P_{\theta} \left[(\mu - \bar{X})^2 \leq \frac{c^2 \sigma^2}{n} \right] P_{\theta} \left[\frac{(n-1)S^2}{b} \leq \sigma^2 \leq \frac{(n-1)S^2}{a} \right] = 1 - \alpha. \end{aligned} \quad (15)$$

Let \bar{x} and s^2 be the observed values of \bar{X} and S^2 . Then in a system of orthogonal (μ, σ^2) -axis, the equation $(\mu - \bar{x})^2 = \frac{c^2 \sigma^2}{n}$ is the equation of a parabola with vertex V located at the point $(\bar{x}, 0)$, with focus F with coordinates $(\bar{x}, \frac{c^2}{4n})$, and with directrix L with equation $\sigma^2 = -\frac{c^2}{4n}$ (see Figure 10.1). Then the part of the plane for which $(\mu - \bar{x})^2 \leq \frac{c^2 \sigma^2}{n}$ is the inner part of the parabola along with the points on the parabola. Since

$$\sigma^2 = \frac{(n-1)s^2}{b} = \frac{1}{b} \sum_{i=1}^n (x_i - \bar{x})^2 \quad \text{and} \quad \sigma^2 = \frac{(n-1)s^2}{a} = \frac{1}{a} \sum_{i=1}^n (x_i - \bar{x})^2$$

are straight lines parallel to the μ -axis, the set of points (μ, σ^2) in the plane, which satisfy simultaneously all inequalities:


$$(\mu - \bar{x})^2 \leq \frac{c^2 \sigma^2}{n}, \quad \frac{1}{b} \sum_{i=1}^n (x_i - \bar{x})^2 \leq \sigma^2 \leq \frac{1}{a} \sum_{i=1}^n (x_i - \bar{x})^2$$

is the part of the plane between the straight lines mentioned above and the inner part of the parabola (along with the points on the parabola) (see shaded area in Figure 10.1).

From relation (15), it follows then that, when replacing \bar{x} by \bar{X} and s^2 by S^2 , the shaded region with random boundary (determined completely as described above) becomes the required confidence region for (μ, σ^2) . What is depicted

Figure 10.1

Confidence Region for (μ, σ^2) with Confidence Coefficient $1 - \alpha$



in Figure 10.1 is a realization of such a confidence region, evaluated for the observed values of the X_i 's.

Actually, the point c above is z_γ , where $\gamma = (1 - \sqrt{1 - \alpha})/2$, and for definiteness, we may choose to split the probability $1 - \sqrt{1 - \alpha}$ equally among the two tails of the Chi-Square distribution. Thus, we take $b = \chi_{n-1; \gamma}^2$ and $a = \chi_{n-1; 1-\gamma}^2$. Then the confidence region is:

$$(\mu - \bar{X})^2 \leq \frac{z_\gamma^2}{n} \sigma^2, \quad \frac{1}{\chi_{n-1; \gamma}^2} \sum_{i=1}^n (X_i - \bar{X})^2 \leq \sigma^2 \leq \frac{1}{\chi_{n-1; 1-\gamma}^2} \sum_{i=1}^n (X_i - \bar{X})^2, \quad (16)$$

$$\gamma = (1 - \sqrt{1 - \alpha})/2.$$

Numerical Example As a numerical example, take $n = 25$ and $\alpha = 0.05$, so that $\gamma \simeq 0.012661$, and (by linear interpolation) $z_\gamma \simeq 2.236$, $\chi_{24; \gamma}^2 \simeq 42.338$, $\chi_{24; 1-\gamma}^2 \simeq 11.130$, and the confidence region becomes:

$$(\mu - \bar{X})^2 \leq 0.199988\sigma^2, \quad 0.023619 \sum_{i=1}^{25} (X_i - \bar{X})^2 \leq \sigma^2 \leq 0.089847 \sum_{i=1}^{25} (X_i - \bar{X})^2$$

or, approximately,

$$(\mu - \bar{X})^2 \leq 0.2\sigma^2, \quad 0.024 \sum_{i=1}^{25} (X_i - \bar{X})^2 \leq \sigma^2 \leq 0.09 \sum_{i=1}^{25} (X_i - \bar{X})^2.$$

REMARK 5 A somewhat general theory for constructing confidence regions is discussed in Chapter 12 (see Theorem 4 there and the examples following it).

10.4 Confidence Intervals with Approximate Confidence Coefficient

It is somewhat conspicuous that in this chapter we have not yet dealt with examples, such as the Binomial, the Poisson, and the Negative Exponential. There is a reason, however, behind it, and that is that the expressions which would serve as the basis for constructing confidence intervals do not have

a known exact distribution. They do have, however, an approximate Normal distribution, and this fact leads to the construction of confidence intervals with *approximate* (rather than exact) confidence coefficient $1 - \alpha$. The remainder of this section is devoted to constructing such intervals.

EXAMPLE 6

On the basis of the random sample X_1, \dots, X_n from the $B(1, \theta)$ distribution, construct a confidence interval for θ with confidence coefficient approximately $1 - \alpha$.

DISCUSSION The tools employed here, as well as in the following two examples, are the CLT and the WLLN in conjunction with either Theorem 7(ii) or Theorem 6(iii) in Chapter 7. It will be assumed throughout that n is large enough, so that these theorems apply.

Recall that $E_\theta X_1 = \theta$ and $\sigma_\theta^2(X_1) = \theta(1 - \theta)$, so that, by the CLT,

$$\frac{\sqrt{n}(\bar{X}_n - \theta)}{\sqrt{\theta(1 - \theta)}} \simeq N(0, 1). \quad (17)$$

In the denominator in (17), replace $\theta(1 - \theta)$ by S_n^2 , where $S_n^2 = \frac{1}{n} \times \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{1}{n}(\sum_{i=1}^n X_i^2 - n\bar{X}_n^2) = \frac{1}{n}(\sum_{i=1}^n X_i - n\bar{X}_n^2) = \bar{X} - \bar{X}^2 = \bar{X}(1 - \bar{X})$, in order to obtain (by Theorem 7(ii) in Chapter 7),

$$\frac{\sqrt{n}(\bar{X}_n - \theta)}{\sqrt{\bar{X}_n(1 - \bar{X}_n)}} \simeq N(0, 1). \quad (18)$$

It follows from (18) that

$$P_\theta \left[-z_{\alpha/2} \leq \frac{\sqrt{n}(\bar{X}_n - \theta)}{\sqrt{\bar{X}_n(1 - \bar{X}_n)}} \leq z_{\alpha/2} \right] \simeq 1 - \alpha, \quad \text{for all } \theta.$$

This expression is equivalent to:

$$P_\theta \left[\bar{X}_n - z_{\alpha/2} \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{n}} \leq \theta \leq \bar{X}_n + z_{\alpha/2} \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{n}} \right] \simeq 1 - \alpha, \quad \text{for all } \theta,$$

which leads to the confidence interval

$$\left[\bar{X}_n - z_{\alpha/2} \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{n}}, \bar{X}_n + z_{\alpha/2} \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{n}} \right] = \bar{X}_n \pm z_{\alpha/2} \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{n}} \quad (19)$$

with confidence coefficient approximately $1 - \alpha$.

Numerical Example For $n = 100$ and $1 - \alpha = 0.95$, the confidence interval in (19) becomes: $\bar{X}_n \pm 1.96 \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{100}} = \bar{X}_n \pm 0.196 \sqrt{\bar{X}_n(1 - \bar{X}_n)}$.

EXAMPLE 7

Construct a confidence interval for θ with confidence coefficient approximately $1 - \alpha$ on the basis of the random sample X_1, \dots, X_n from the $P(\theta)$ distribution.

DISCUSSION Here $E_\theta X_1 = \sigma_\theta^2(X_1) = \theta$, so that, working as in the previous example, and employing Theorem 6(iii) in Chapter 7, we have

$$\frac{\sqrt{n}(\bar{X}_n - \theta)}{\sqrt{\theta}} \simeq N(0, 1), \quad \text{or} \quad \frac{\sqrt{n}(\bar{X}_n - \theta)}{\sqrt{\bar{X}_n}} \simeq N(0, 1).$$

Hence $P_\theta[-z_{\alpha/2} \leq \frac{\sqrt{n}(\bar{X}_n - \theta)}{\sqrt{\bar{X}_n}} \leq z_{\alpha/2}] \simeq 1 - \alpha$, for all θ , which leads to the required confidence interval

$$\left[\bar{X}_n - z_{\alpha/2} \sqrt{\frac{\bar{X}_n}{n}}, \bar{X}_n + z_{\alpha/2} \sqrt{\frac{\bar{X}_n}{n}} \right] = \bar{X}_n \pm z_{\alpha/2} \sqrt{\frac{\bar{X}_n}{n}}. \quad (20)$$

Numerical Example For $n = 100$ and $1 - \alpha = 0.95$, the confidence interval in (20) becomes: $\bar{X}_n \pm 0.196\sqrt{\bar{X}_n}$.

EXAMPLE 8

Let X_1, \dots, X_n be a random sample from the Negative Exponential distribution in the following parameterization: $f(x; \theta) = \frac{1}{\theta} e^{-x/\theta}$, $x > 0$. Construct a confidence interval for θ with confidence coefficient approximately $1 - \alpha$.

DISCUSSION In the adopted parameterization above, $E_\theta X_1 = \theta$ and $\sigma_\theta^2(X_1) = \theta^2$. Then working as in the previous example, we have that

$$\frac{\sqrt{n}(\bar{X}_n - \theta)}{\theta} \simeq N(0, 1), \quad \text{or} \quad \frac{\sqrt{n}(\bar{X}_n - \theta)}{\bar{X}_n} \simeq N(0, 1).$$

It follows that the required confidence interval is given by:

$$\left[\bar{X}_n - z_{\alpha/2} \frac{\bar{X}_n}{\sqrt{n}}, \bar{X}_n + z_{\alpha/2} \frac{\bar{X}_n}{\sqrt{n}} \right] = \bar{X}_n \pm z_{\alpha/2} \frac{\bar{X}_n}{\sqrt{n}}. \quad (21)$$

Numerical Example For $n = 100$ and $1 - \alpha = 0.95$, the confidence interval in (21) becomes: $\bar{X}_n \pm 0.196\bar{X}_n$.

Exercises

4.1 Let the independent r.v.'s X_1, \dots, X_n have unknown (finite) mean μ and known (finite) variance σ^2 , and suppose that n is large. Then:

- (i) Use the CLT in order to construct a confidence interval for μ with approximate confidence coefficient $1 - \alpha$.
- (ii) Provide the form of the interval in part (i) for $n = 100$, $\sigma = 1$, and $\alpha = 0.05$.

- (iii) Refer to part (i) and suppose that $\sigma = 1$ and $\alpha = 0.05$. Then determine the sample size n , so that the length of the confidence interval is 0.1.
- (iv) Observe that the length of the confidence interval in part (i) tends to 0 as $n \rightarrow \infty$, for any σ and any α .

4.2 Refer to Exercise 4.1, and suppose that both μ and σ^2 are unknown. Then:

- (i) Construct a confidence interval for μ with approximate confidence coefficient $1 - \alpha$.
- (ii) Provide the form of the interval in part (i) for $n = 100$ and $\alpha = 0.05$.
- (iii) Show that the length of the interval in part (i) tends to 0 in probability as $n \rightarrow \infty$.

Hint: For part (i), refer to Theorem 7(ii) in Chapter 7, and for part (iii), refer to Theorem 7(i) and Theorem 6(ii) in the same chapter.

- 4.3** (i) Let $X \sim N(\mu, \sigma^2)$, and for $0 < \alpha < 1$, let x_α and $x_{1-\alpha}$ be the α th and $(1 - \alpha)$ th quantiles, respectively, of X ; i.e., $P(X \leq x_\alpha) = P(X \geq x_{1-\alpha}) = \alpha$, so that $P(x_\alpha \leq X \leq x_{1-\alpha}) = 2\alpha$. Show that $x_\alpha = \mu + \sigma \Phi^{-1}(\alpha)$, $x_{1-\alpha} = \mu + \sigma \Phi^{-1}(1 - \alpha)$, so that $[x_\alpha, x_{1-\alpha}] = [\mu + \sigma \Phi^{-1}(\alpha), \mu + \sigma \Phi^{-1}(1 - \alpha)]$.
- (ii) Refer to Exercise 4.5(i) of Chapter 9 (see also Exercise 4.4 there), where it is found that the posterior p.d.f. of θ , given $X_1 = x_1, \dots, X_n = x_n$, $h(\cdot | x_1, \dots, x_n)$, is $N(\frac{n\bar{x} + \mu}{n+1}, \frac{1}{n+1})$. Use part (i) in order to find the expression of the interval $[x_\alpha, x_{1-\alpha}]$ here.

Remark: In the present context, the interval $[x_\alpha, x_{1-\alpha}]$ is called a *prediction interval* for θ with confidence coefficient $1 - 2\alpha$.

- (iii) Compute the prediction interval in part (ii) when $n = 9$, $\mu = 1$, $\bar{x} = 1.5$, and $\alpha = 0.025$.

4.4 Let X_1, \dots, X_n be independent r.v.'s with strictly increasing d.f. F , and let Y_i be the i th order statistic of the X_i 's, $1 \leq i \leq n$. For $0 < p < 1$, let x_p be the (unique) p th quantile of F . Then:

- (i) Show that for any i and j with $1 \leq i < j \leq n - 1$,

$$P(Y_i \leq x_p \leq Y_j) = \sum_{k=i}^{j-1} \binom{n}{k} p^k q^{n-k} \quad (q = 1 - p).$$

Thus, $[Y_i, Y_j]$ is a confidence interval for x_p with confidence coefficient $\sum_{k=i}^{j-1} \binom{n}{k} p^k q^{n-k}$. This probability is often referred to as *probability of coverage* of x_p .

- (ii) For $n = 10$ and $p = 0.25$, identify the respective coverage probabilities for the pairs (Y_1, Y_3) , (Y_1, Y_4) , (Y_2, Y_4) , (Y_2, Y_5) .
- (iii) For $p = 0.50$, do the same as in part (ii) for the pairs (Y_3, Y_9) , (Y_4, Y_7) , (Y_4, Y_8) , (Y_5, Y_7) .


- (iv) For $p = 0.75$, do the same as in part (ii) for the pairs (Y_8, Y_{10}) , (Y_7, Y_{10}) , (Y_7, Y_9) , (Y_6, Y_9) .

Hint: For part (i), observe that: $P(Y_i \leq x_p) = P(\text{at least } i \text{ of } X_1, \dots, X_n \leq x_p) = \sum_{k=i}^n \binom{n}{k} p^k q^{n-k}$, since $P(X_k \leq x_p) = p$ and $q = 1 - p$. Also, $P(Y_i \leq x_p) = P(Y_i \leq x_p, Y_j \geq x_p) + P(Y_i \leq x_p, Y_j < x_p) = P(Y_i \leq x_p \leq Y_j) + P(Y_j < x_p)$, so that $P(Y_i \leq x_p \leq Y_j) = P(Y_i \leq x_p) - P(X_j \leq x_p)$.

For part (iv), observe that $\binom{n}{k} p^k q^{n-k} = \binom{n}{n-r} q^r p^{n-r} = \binom{n}{r} q^r p^{n-r}$ (by setting $n - k = r$ and recalling that $\binom{n}{n-r} = \binom{n}{r}$).

- 4.5** Let X be a r.v. with a strictly increasing d.f. F , and let p be a number with $0 < p < 1$. Consider the event: $A_p = \{F(X) \leq p\} = \{s \in \mathcal{S}; F(X(s)) \leq p\} = \{s \in \mathcal{S}; X(s) \leq F^{-1}(p)\}$. So, A_p is the event in the underlying sample space \mathcal{S} for the sample points s of which $F(X(s)) \leq p$. Since for each fixed x , $F(x)$ represents the proportion of the (unit) distribution mass of F which is covered (or carried) by the interval $(-\infty, x]$, it follows that the random interval $(-\infty, X]$ covers (carries) the (random) proportion $F(X)$ of the distribution mass of F , and on the event A_p , the random interval $(-\infty, X]$ covers (carries) at most 100

p % of the mass of F . Equivalently, the random interval (X, ∞) covers (carries) at least 100(1 - p)% of the distribution mass of F .



Use Theorem 10 in Chapter 6 in order to show that $P(A_p) = p$; i.e., $(-\infty, X]$ covers at most 100

p % of the distribution mass of F with probability p . Equivalently, the random interval (X, ∞) covers at least 100(1 - p)% of the distribution mass of F with probability p .

Testing Hypotheses

In this chapter, the problem of testing hypotheses is considered to some extent. Additional topics are discussed in Chapter 12. The chapter consists of four sections, the first of which is devoted to some general concepts and the formulation of a null hypothesis and its alternative. A number of examples discussed provide sufficient motivation for what is done in this section.

Section 2 is somewhat long and enters into the essence of the testing hypotheses issue. Specifically, the Neyman–Pearson Fundamental Lemma is stated, and the main points of its proof are presented for the case that the underlying r.v.'s are of the continuous type. It is stated that this result by itself is of limited use; nevertheless, it does serve as the stepping stone in establishing other more complicated and truly useful results. This is obtained when the underlying family of distributions is the so-called family of distributions of the exponential type. Thus, the definition of an exponential type p.d.f. follows, and it is next illustrated by means of examples that such families occur fairly often. In an exponential type p.d.f. (in the real-valued parameter θ), uniformly most powerful (UMP) tests are presented for one-sided and two-sided hypotheses, which arise in practice in a natural way. This is done in Theorems 2 and 3.

In the following section, Theorems 2 and 3 are applied to concrete cases, such as the Binomial distribution, the Poisson distribution, and the Normal distribution. All applications are accompanied by numerical examples.

The last section of this chapter, Section 4, is also rather extensive and deals with Likelihood Ratio (LR) tests. General concepts, the necessary notation, and some motivation for the tests used are given. The better part of the section is devoted to deriving LR tests in Normal distributions. The problem is divided into two parts. The first part considers the case where we are dealing with one sample from an underlying Normal distribution, and LR tests are derived for the mean and the variance of the distribution. In the second part, two independent random samples are available coming from two underlying Normal populations. Then LR tests are derived in comparing the means and the variances of the distributions. In all cases, the results produced are illustrated by means of numerical examples.

11.1 General Concepts, Formulation of Some Testing Hypotheses

In order to motivate the formulation of a null hypothesis and its alternative, consider some specific examples. Most of them are taken from Chapter 1.

EXAMPLE 1

In reference to Example 6 in Chapter 1, let θ be the unknown proportion of unemployed workers, and let θ_0 be an acceptable level of unemployment, e.g., $\theta_0 = 6.25\%$. Then the parameter space is split into the sets $(0.0625, 1)$ and $(0, 0.0625]$, and one of them will be associated with the (null) hypothesis. It is proposed that that set be $(0.0625, 1)$; i.e., $H_0: \theta > 0.0625$ (and therefore $H_A: \theta \leq 0.0625$). The rule of thumb for selecting H_0 is this: "Select as null hypothesis that hypothesis whose false rejection has the most serious consequences." Indeed, if θ is, actually, greater than 6.25% and is (falsely) rejected, then human suffering may occur, due to the fact that the authorities in charge had no incentives to take the necessary measures. On the other hand, if $\theta \leq 0.0625$ was selected as the null hypothesis and was falsely rejected, then the most likely consequence would be for the authorities to undertake some unnecessary measures and, perhaps, waste some money. However, the former consequence is definitely more serious than the latter. Another way of looking at the problem of determining the null hypothesis is to formulate as such a position, which we wish to challenge, and which we are willing to accept only in the face of convincing evidence, provided by the interested party. To summarize then, if X is the r.v. denoting the number of unemployed workers among n sampled, then $X \sim B(n, \theta)$ and the hypothesis to be tested is $H_0: \theta > 0.0625$ against the alternative $H_A: \theta \leq 0.0625$ at (some given) level of significance α .

EXAMPLE 2

In reference to Example 8 in Chapter 1, if X is the r.v. denoting those young adults, among the n sampled, who listen to this particular weekend music program, then $X \sim B(n, \theta)$. Then, arguing as in the previous example, we have that the hypothesis to be tested is $H_0: \theta \leq \theta_0 (=100p\%)$ against the alternative $H_A: \theta > \theta_0$ at level of significance α .

EXAMPLE 3

Refer to Example 12 of Chapter 1, and let X be the r.v. denoting the mean bacteria count per unit volume of water at a lake beach. Then $X \sim P(\theta)$ and the hypothesis to be tested is $H_0: \theta > 200$ against $H_A: \theta \leq 200$ at level of significance α .

EXAMPLE 4

Suppose that the mean θ of a r.v. X represents the dosage of a drug which is used for the treatment of a certain disease. For this medication to be both safe and effective, θ must satisfy the requirements $\theta_1 < \theta < \theta_2$, for two specified values θ_1 and θ_2 . Then, on the basis of previous discussions, the hypothesis to be tested here is $H_0: \theta \leq \theta_1$ or $\theta \geq \theta_2$ against the alternative $H_A: \theta_1 < \theta < \theta_2$ at the level of significance α . Of course, we have to assume a certain

distribution for the r.v. X , which for good reasons is taken to be $N(\theta, \sigma^2)$, σ known.

EXAMPLE 5

Refer to Example 16 in Chapter 1, and suppose that the survival time for a terminal cancer patient treated with the standard treatment is a r.v. $X \sim N(\theta_1, \sigma_1^2)$. Likewise, let the r.v. Y stand for the survival time for such a patient subject to the new treatment, and let $Y \sim N(\theta_2, \sigma_2^2)$. Then the hypothesis to be tested here is $H_0: \theta_2 = \theta_1$ against the alternative $H_A: \theta_2 > \theta_1$ at level of significance α .

REMARK 1 The hypothesis to be tested could also be $\theta_2 \leq \theta_1$, but the possibility that $\theta_2 < \theta_1$ may be excluded; it can be assumed that the new treatment cannot be inferior to the existing one. The supposition that $\theta_2 = \theta_1$, there is no difference between the two treatments, leads to the term “null” for the hypothesis $H_0: \theta_2 = \theta_1$.

Examples 1–4 have the following common characteristics. A r.v. X is distributed according to the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, and we are interested in testing one of the following hypotheses, each one at some specified level of significance α : $H_0: \theta > \theta_0$ against $H_A: \theta \leq \theta_0$; $H_0: \theta \leq \theta_0$ against $H_A: \theta > \theta_0$; $H_0: \theta \leq \theta_1$ or $\theta \geq \theta_2$ against $H_A: \theta_1 < \theta < \theta_2$. It is understood that in all cases θ remains in Ω . In Example 5, two Normally distributed populations are compared in terms of their means, and the hypothesis tested is $H_0: \theta_2 = \theta_1$ against $H_A: \theta_2 > \theta_1$. An example of a different nature would lead to testing the hypothesis $H_0: \theta_2 < \theta_1$ against $H_A: \theta_2 = \theta_1$.

In the first four examples, the hypotheses stated are to be tested by means of a random sample X_1, \dots, X_n from the underlying distribution. In the case of Example 5, the hypothesis is to be tested by utilizing two independent random samples X_1, \dots, X_m and Y_1, \dots, Y_n from the underlying distributions.

Observe that in all cases the hypotheses tested are composite, and so are the alternatives. We wish, of course, for the proposed tests to be optimal in some satisfactory sense. If the tests were to be UMP (uniformly most powerful), then they would certainly be highly desirable. In the following section, a somewhat general theory will be provided, which, when applied to the examples under consideration, will produce UMP tests.

Exercises

1.1 In the following examples, indicate which statements constitute a simple and which a composite hypothesis:

- (i) X is a r.v. whose p.d.f. f is given by $f(x) = 2e^{-2x}$, $x > 0$.
- (ii) When tossing a coin, let X be the r.v. taking the value 1 if the head appears and 0 if the tail appears. Then the statement is: The coin is biased.
- (iii) X is a r.v. whose expectation is equal to 5.

1.2 Let X_1, \dots, X_n be i.i.d. r.v.'s with p.d.f. f which may be either Normal, $N(\mu, \sigma^2)$, to be denoted by f_N , or Cauchy with parameters μ and σ^2 , to be

denoted by f_C , where, we recall that:

$$f_N(x; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}, \quad x \in \mathfrak{R}, \mu \in \mathfrak{R}, \sigma > 0,$$

$$f_C(x; \mu, \sigma^2) = \frac{\sigma}{\pi} \times \frac{1}{(x-\mu)^2 + \sigma^2}, \quad x \in \mathfrak{R}, \mu \in \mathfrak{R}, \sigma > 0.$$

Consider the following null hypotheses and the corresponding alternatives:

- (i) H_{01} : f is Normal, H_{A1} : f is Cauchy.
- (ii) H_{02} : f is Normal with $\mu \leq \mu_0$, H_{A2} : f is Cauchy with $\mu \leq \mu_0$.
- (iii) H_{03} : f is Normal with $\mu = \mu_0$, H_{A3} : f is Cauchy with $\mu = \mu_0$.
- (iv) H_{04} : f is Normal with $\mu = \mu_0$, $\sigma \geq \sigma_0$, H_{A4} : f is Cauchy with $\mu = \mu_0$, $\sigma \geq \sigma_0$.
- (v) H_{05} : f is Normal with $\mu = \mu_0$, $\sigma < \sigma_0$, H_{A5} : f is Cauchy with $\mu = \mu_0$, $\sigma = \sigma_0$.
- (vi) H_{06} : f is Normal with $\mu = \mu_0$, $\sigma = \sigma_0$, H_{A6} : f is Cauchy with $\mu = \mu_0$, $\sigma = \sigma_0$.

State which of the H_{0i} and which of the H_{Ai} , $i = 1, \dots, 6$, are simple and which are composite.

11.2 Neyman–Pearson Fundamental Lemma, Exponential Type Families, Uniformly Most Powerful Tests for Some Composite Hypotheses

In reference to Example 1, one could certainly consider testing the simple hypothesis $H_0: \theta = \theta_0$ (e.g., 0.05) against the simple alternative $H_A: \theta = \theta_1$, for some fixed θ_1 either $>\theta_0$ or $<\theta_0$. However, such a testing framework would be highly unrealistic. It is simply not reasonable to isolate two single values from the continuum of values $(0, 1)$ and test one against the other. What is meaningful is the way we actually formulated H_0 in this example. Nevertheless, it is still true that a long journey begins with the first step, and this applies here as well. Accordingly, we are going to start out with the problem of testing a simple hypothesis against a simple alternative, which is what the celebrated Neyman–Pearson Fundamental Lemma is all about.

THEOREM 1

(Neyman–Pearson Fundamental Lemma) Let X_1, \dots, X_n be a random sample with p.d.f. f unknown. We are interested in testing the simple hypothesis $H_0: f = f_0$ (specified) against the simple alternative $H_A: f = f_1$ (specified) at level of significance α ($0 < \alpha < 1$). To this end, define the test φ as follows:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } f_1(x_1) \dots f_1(x_n) > C f_0(x_1) \dots f_0(x_n) \\ \gamma & \text{if } f_1(x_1) \dots f_1(x_n) = C f_0(x_1) \dots f_0(x_n) \\ 0 & \text{if } f_1(x_1) \dots f_1(x_n) < C f_0(x_1) \dots f_0(x_n), \end{cases} \quad (1)$$

where the constants C and γ ($C > 0$, $0 \leq \gamma \leq 1$) are defined through the relationship:

$$\begin{aligned} E_{f_0} \varphi(X_1, \dots, X_n) &= P_{f_0}[f_1(X_1) \cdots f_1(X_n) > C f_0(X_1) \cdots f_0(X_n)] \\ &\quad + \gamma P_{f_0}[f_1(X_1) \cdots f_1(X_n) = C f_0(X_1) \cdots f_0(X_n)] = \alpha. \end{aligned} \quad (2)$$

Then the test φ is MP among all tests with level of significance $\leq \alpha$.

REMARK 2 The test φ is a *randomized* test, if $0 < \gamma < 1$. The necessity for a randomized test stems from relation (2), where the left-hand side has to be equal to α . If the X_i 's are discrete, the presence of γ ($0 < \gamma < 1$) is indispensable. In case, however, the X_i 's are of the continuous type, then $\gamma = 0$ and the test is *nonrandomized*.

The appearance of f_0 as a subscript indicates, of course, that expectations and probabilities are calculated by using the p.d.f. f_0 for the X_i 's.

PROOF OF THEOREM 1 (Outline for X_i 's of the Continuous Type) To simplify the notation, write 0 (or 1) rather than f_0 (or f_1) when f_0 (or f_1) occurs as a subscript. Also, it would be convenient to use the vector notation $\mathbf{X} = (X_1, \dots, X_n)$ and $\mathbf{x} = (x_1, \dots, x_n)$. First, we show that the test φ is of level α . Indeed, let $T = \{\mathbf{x} \in \mathfrak{R}^n; L_0(\mathbf{x}) > 0\}$, where $L_0(\mathbf{x}) = f_0(x_1) \cdots f_0(x_n)$, and likewise $L_1(\mathbf{x}) = f_1(x_1) \cdots f_1(x_n)$. Then, if $D = \mathbf{X}^{-1}(T)$; i.e., $D = \{s \in \mathcal{S}; \mathbf{X}(s) \in T\}$, so that $D^c = \{s \in \mathcal{S}; \mathbf{X}(s) \in T^c\}$, it follows that $P_0(D^c) = P_0(\mathbf{X} \in T^c) = \int_{T^c} L_0(\mathbf{x}) d\mathbf{x} = 0$. Therefore, in calculating probabilities by using the p.d.f. L_0 , it suffices to restrict ourselves to the set D . Then, by means of (2),

$$\begin{aligned} E_0 \varphi(\mathbf{X}) &= P_0[L_1(\mathbf{X}) > C L_0(\mathbf{X})] \\ &= P_0\{[L_1(\mathbf{X}) > C L_0(\mathbf{X})] \cap D\} \\ &= P_0\left\{\left[\frac{L_1(\mathbf{X})}{L_0(\mathbf{X})} > C\right] \cap D\right\} \quad (\text{since } L_0(\mathbf{X}) > 0 \text{ on } D) \\ &= P_0(Y > C) = 1 - P_0(Y \leq C) = g(C), \text{ say,} \end{aligned}$$


where $Y = \frac{L_1(\mathbf{X})}{L_0(\mathbf{X})}$ on D , and arbitrary on D^c . The picture of $1 - P_0(Y \leq C)$ is depicted in Figure 11.1, and it follows that, for each α ($0 < \alpha < 1$), there is (essentially) a unique C such that $1 - P_0(Y \leq C) = \alpha$. That is, $E_0 \varphi(\mathbf{X}) = \alpha$, which shows that the test φ is of level α .

Next, it is shown that φ is MP as described by showing that, if φ^* is any other test with $E_0 \varphi^*(\mathbf{X}) = \alpha^* \leq \alpha$, then $\pi_\varphi(1) = E_1 \varphi(\mathbf{X}) \geq E_1 \varphi^*(\mathbf{X}) = \pi_{\varphi^*}(1)$ (i.e., the power of φ is not smaller than the power of any such test φ^* of level of significance $\leq \alpha$). Indeed, define B^+ and B^- by:

$$\begin{aligned} B^+ &= \{\mathbf{x} \in \mathfrak{R}^n; \varphi(\mathbf{x}) - \varphi^*(\mathbf{x}) > 0\} = (\varphi > \varphi^*), \\ B^- &= \{\mathbf{x} \in \mathfrak{R}^n; \varphi(\mathbf{x}) - \varphi^*(\mathbf{x}) < 0\} = (\varphi < \varphi^*). \end{aligned}$$

Figure 11.1

The Graph of the
Function of $(C) =$
 $1 - P_0(\gamma \leq C)$



Then, clearly, $B^+ \cap B^- = \emptyset$, and, by means of (1),

$$\begin{aligned} B^+ &= (\varphi > \varphi^*) \subseteq (\varphi = 1) \subseteq (L_1 > CL_0), \\ B^- &= (\varphi < \varphi^*) \subseteq (\varphi = 0) \subseteq (L_1 < CL_0). \end{aligned} \quad (3)$$

Therefore

$$\begin{aligned} &\int_{\mathfrak{R}^n} [\varphi(\mathbf{x}) - \varphi^*(\mathbf{x})][L_1(\mathbf{x}) - CL_0(\mathbf{x})]d\mathbf{x} \\ &= \int_{B^+} [\varphi(\mathbf{x}) - \varphi^*(\mathbf{x})][L_1(\mathbf{x}) - CL_0(\mathbf{x})]d\mathbf{x} \\ &\quad + \int_{B^-} [\varphi(\mathbf{x}) - \varphi^*(\mathbf{x})][L_1(\mathbf{x}) - CL_0(\mathbf{x})]d\mathbf{x} \geq 0 \text{ by (3)}. \end{aligned}$$

Hence

$$\begin{aligned} &\int_{\mathfrak{R}^n} \varphi(\mathbf{x})L_1(\mathbf{x})d\mathbf{x} - \int_{\mathfrak{R}^n} \varphi^*(\mathbf{x})L_1(\mathbf{x})d\mathbf{x} \\ &\geq C \left[\int_{\mathfrak{R}^n} \varphi(\mathbf{x})L_0(\mathbf{x})d\mathbf{x} - \int_{\mathfrak{R}^n} \varphi^*(\mathbf{x})L_0(\mathbf{x})d\mathbf{x} \right] \\ &= C(\alpha - \alpha^*) \geq 0 \quad (\text{since } \alpha^* \leq \alpha). \end{aligned}$$

Hence $\int_{\mathfrak{R}^n} \varphi(\mathbf{x})L_1(\mathbf{x})d\mathbf{x} = E_1\varphi(\mathbf{X}) \geq E_1\varphi^*(\mathbf{X}) = \int_{\mathfrak{R}^n} \varphi^*(\mathbf{x})L_1(\mathbf{x})d\mathbf{x}$. \blacktriangle

This theorem has the following corollary, according to which the power of the MP test φ cannot be $< \alpha$; not very much to be sure, but yet somewhat reassuring.

COROLLARY For the MP test φ , $\pi_\varphi(1) = E_1\varphi(\mathbf{x}) \geq \alpha$.

PROOF Just compare the power of φ with that of $\varphi^* \equiv \alpha$ whose level of significance and power are both equal to α . \blacktriangle

REMARK 3 The theorem was formulated in terms of any two p.d.f.'s f_0 and f_1 as the two possible options for f . In a parametric setting, where f is of the form $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}^r$, $r \geq 1$, the p.d.f.'s f_0 and f_1 will correspond to two specified values of θ ; θ_0 and θ_1 , say. That is, $f_0 = f(\cdot; \theta_0)$ and $f_1 = f(\cdot; \theta_1)$.

The following examples will help illustrate how the Neyman–Pearson Fundamental Lemma actually applies in concrete cases.

EXAMPLE 6

On the basis of a random sample of size 1 from the p.d.f. $f(x; \theta) = \theta x^{\theta-1}$, $0 < x < 1$ ($\theta > 1$):

- (i) Use the Neyman–Pearson Fundamental Lemma to derive the MP test for testing the hypothesis $H_0: \theta = \theta_0$ against the alternative $H_A: \theta = \theta_1$ at level of significance α .
- (ii) Derive the formula for the power $\pi(\theta_1)$.
- (iii) Give numerical values for parts (i) and (ii) when $\theta_0 = 4$ and $\theta_1 = 6$, $\theta_1 = 2$; take $\alpha = 0.05$.

DISCUSSION In the first place, the given function is, indeed, a p.d.f., since $\int_0^1 \theta x^{\theta-1} dx = x^\theta |_0^1 = 1$. Next:

- (i) H_0 is rejected, if for some positive constant C^* :

$$\frac{\theta_1 x^{\theta_1-1}}{\theta_0 x^{\theta_0-1}} > C^*, \text{ or } x^{\theta_1-\theta_0} > \frac{\theta_0 C^*}{\theta_1}, \text{ or } (\theta_1 - \theta_0) \log x > \log \left(\frac{\theta_0 C^*}{\theta_1} \right).$$

Now, if $\theta_1 > \theta_0$, this last inequality is equivalent to:

$$\log x > \log \left(\frac{\theta_0 C^*}{\theta_1} \right)^{1/(\theta_1-\theta_0)} = \log C \quad \left(C = \left(\frac{\theta_0 C^*}{\theta_1} \right)^{1/(\theta_1-\theta_0)} \right),$$

or $x > C$. If $\theta_1 < \theta_0$, the final form of the inequality becomes $x < C$. For $\theta_1 > \theta_0$, the cutoff point is calculated by:

$$P_{\theta_0}(X > C) = \int_C^1 \theta_0 x^{\theta_0-1} dx = x^{\theta_0} |_C^1 = 1 - C^{\theta_0} = \alpha, \quad \text{or} \quad C = (1 - \alpha)^{1/\theta_0}.$$

For $\theta_1 < \theta_0$, we have:

$$P_{\theta_0}(X < C) = x^{\theta_0} |_0^C = C^{\theta_0} = \alpha, \quad \text{or} \quad C = \alpha^{1/\theta_0}.$$

Then, for $\theta_1 > \theta_0$, reject H_0 when $x > (1 - \alpha)^{1/\theta_0}$; and, for $\theta_1 < \theta_0$, reject H_0 when $x < \alpha^{1/\theta_0}$.

- (ii) For $\theta_1 > \theta_0$, the power of the test is given by:

$$\pi(\theta_1) = P_{\theta_1}(X > C) = \int_C^1 \theta_1 x^{\theta_1-1} dx = x^{\theta_1} |_C^1 = 1 - C^{\theta_1}, \text{ or}$$

$\pi(\theta_1) = 1 - (1 - \alpha)^{\theta_1/\theta_0}$. For $\theta_1 < \theta_0$, we have:

$$\pi(\theta_1) = P_{\theta_1}(X < C) = \int_0^C \theta_1 x^{\theta_1-1} dx = x^{\theta_1} |_0^C = C^{\theta_1} = \alpha^{\theta_1/\theta_0}.$$

That is,

$$\pi(\theta_1) = 1 - (1 - \alpha)^{\theta_1/\theta_0} \quad \text{for } \theta_1 > \theta_0; \quad \pi(\theta_1) = \alpha^{\theta_1/\theta_0} \quad \text{for } \theta_1 < \theta_0.$$

- (iii) For $\theta_1 = 6$, the cutoff point is:

$$(1 - 0.05)^{1/4} = 0.95^{0.25} \simeq 0.987,$$

and the power is: $\pi(6) = 1 - (0.95)^{1.5} \simeq 1 - 0.926 = 0.074$. For $\theta_1 = 2$, the cutoff point is: $(0.05)^{1/4} = (0.05)^{0.25} \simeq 0.473$, and the power is: $\pi(2) = (0.05)^{1/2} \simeq 0.224$.

EXAMPLE 7

Refer to Example 6 and:

- (i) Show that the Neyman–Pearson test which rejects the (simple) hypothesis $H_0: \theta = \theta_0$ when tested against the (simple) alternative $H_{A,\theta_1}: \theta = \theta_1$, for some fixed $\theta_1 > \theta_0$, at level of significance α , is, actually, UMP for testing H_0 against the composite alternative $H_A: \theta > \theta_0$ at level of significance α .
- (ii) Also, show that the Neyman–Pearson test which rejects the hypothesis $H_0: \theta = \theta_0$ when tested against the (simple) alternative $H'_{A,\theta_1}: \theta = \theta_1$, for some fixed $\theta_1 < \theta_0$, at level of significance α , is, actually, UMP for testing H_0 against the composite alternative $H'_A: \theta < \theta_0$ at level of significance α .
- (iii) Show that there is no UMP test for testing the hypothesis $H_0: \theta = \theta_0$ against the (double-sided) composite alternative $H''_A: \theta \neq \theta_0$ at level of significance α .

DISCUSSION

- (i) Indeed, by part (i) of Example 6, the MP test for testing $H_0: \theta = \theta_0$ against $H_{A,\theta_1}: \theta = \theta_1$ rejects H_0 when $x > (1 - \alpha)^{1/\theta_0}$, regardless of the *specific* value of θ_1 , provided $\theta_1 > \theta_0$. Thus, this test becomes a UMP test when H_{A,θ_1} is replaced by $H_A: \theta > \theta_0$.
- (ii) Likewise, by Example 6(i), the MP test for testing $H_0: \theta = \theta_0$ against $H'_{A,\theta_1}: \theta = \theta_1$ rejects H_0 when $x < \alpha^{1/\theta_0}$, regardless of the *specific* value θ_1 , provided $\theta_1 < \theta_0$. Thus, this test becomes a UMP test when H'_{A,θ_1} is replaced by $H'_A: \theta < \theta_0$.
- (iii) The rejection region for testing the hypotheses $H_0: \theta = \theta_0$ against the alternative $H_A: \theta > \theta_0$ is $R_1 = ((1 - \alpha)^{1/\theta_0}, 1)$, and the rejection region for testing H_0 against $H'_A: \theta < \theta_0$ is $R_2 = (0, \alpha^{1/\theta_0})$. Since these MP regions depend on which side of θ_0 lie the alternative θ 's and are different, there cannot exist a UMP test for testing H_0 against $H''_A: \theta \neq \theta_0$.

EXAMPLE 8

On the basis of a random sample of size 1 from the p.d.f. $f(x; \theta) = 1 + \theta^2(\frac{1}{2} - x)$, $0 < x < 1$, $-1 \leq \theta \leq 1$:

- (i) Use the Neyman–Pearson Fundamental Lemma to derive the MP test for testing the hypothesis $H_0: \theta = 0$ (i.e., the p.d.f. is $U(0, 1)$) against the alternative $H_A: \theta = \theta_1$ at level of significance α .
- (ii) Investigate whether or not the test derived in part (i) is a UMP test for testing $H_0: \theta = 0$ against the alternative $H'_A: \theta \neq 0$.
- (iii) Determine the test in part (i) for $\alpha = 0.05$.
- (iv) Determine the power of the test in part (i).

DISCUSSION First, the function given is a p.d.f., because it is nonnegative and $\int_0^1 [1 + \theta^2(\frac{1}{2} - x)] dx = 1 + \theta^2(\frac{1}{2} - \frac{1}{2}) = 1$. Next:

- (i) H_0 is rejected whenever $1 + \theta_1^2(\frac{1}{2} - x) > C^*$, or $x < C$, where $C = \frac{1}{2} - (C^* - 1)/\theta_1^2$, and C is determined by $P_0(X < C) = \alpha$, so that $C = \alpha$, since $X \sim U(0, 1)$ under H_0 . Thus, H_0 is rejected when $x < \alpha$.
- (ii) Observe that the test is independent of θ_1 , and since it is MP against each fixed θ_1 , it follows that it is UMP for testing H_0 against $H_A: \theta \neq 0$.
- (iii) For $\alpha = 0.05$, the test in part (i) rejects H_0 whenever $x < 0.05$.
- (iv) For $\theta \neq 0$, the power of the test is:

$$\pi(\theta) = P_\theta(X < \alpha) = \int_0^\alpha \left[1 + \theta^2 \left(\frac{1}{2} - x \right) \right] dx = \frac{1}{2}\alpha(1 - \alpha)\theta^2 + \alpha.$$

Thus, e.g., $\pi(\pm 1) = \frac{1}{2}\alpha(1 - \alpha) + \alpha$, $\pi(\pm \frac{1}{2}) = \frac{1}{8}\alpha(1 - \alpha) + \alpha$, which for $\alpha = 0.05$ become: $\pi(\pm 1) \simeq 0.074$, $\pi(\pm \frac{1}{2}) \simeq 0.056$.

11.2.1 Exponential Type Families of p.d.f.'s

The remarkable thing here is that, if the p.d.f. $f(\cdot; \theta)$ is of a certain general form to be discussed below, then the apparently simple-minded Theorem 1 leads to UMP tests; it is the stepping stone for getting to those tests.

DEFINITION 1

The p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$, is said to be of the *exponential type*, if

$$f(x; \theta) = C(\theta)e^{Q(\theta)T(x)} \times h(x), \quad x \in \mathfrak{R}, \quad (4)$$

where Q is strictly monotone and h does not involve θ in any way; $C(\theta)$ is simply a normalizing constant.

Most of the p.d.f.'s we have encountered so far are of the form (4). Here are some examples.

EXAMPLE 9

The $B(n, \theta)$ p.d.f. is of the exponential type.

DISCUSSION Indeed,

$$f(x; \theta) = \binom{n}{x} \theta^x (1 - \theta)^{n-x} I_A(x), \quad A = \{0, 1, \dots, n\},$$

where, we recall that I_A is the indicator of A ; i.e., $I_A(x) = 1$ if $x \in A$, and $I_A(x) = 0$ if $x \in A^c$.

Hence

$$f(x; \theta) = (1 - \theta)^n \cdot e^{[\log(\frac{\theta}{1-\theta})]x} \times \binom{n}{x} I_A(x),$$

so that $f(x; \theta)$ is of the form (4) with $C(\theta) = (1 - \theta)^n$, $Q(\theta) = \log(\frac{\theta}{1-\theta})$ strictly increasing (since $\frac{d}{d\theta}(\frac{\theta}{1-\theta}) = \frac{1}{(1-\theta)^2} > 0$ and $\log(\cdot)$ is strictly increasing), $T(x) = x$, and $h(x) = \binom{n}{x} I_A(x)$.

EXAMPLE 10

The $P(\theta)$ p.d.f. is of the exponential type.

DISCUSSION Here

$$f(x; \theta) = \frac{e^{-\theta} \theta^x}{x!} I_A(x), \quad A = \{0, 1, \dots\}.$$

Hence

$$f(x; \theta) = e^{-\theta} \times e^{(\log \theta)x} \times \frac{1}{x!} I_A(x),$$

so that $f(x; \theta)$ is of the form (4) with $C(\theta) = e^{-\theta}$, $Q(\theta) = \log \theta$ strictly increasing, $T(x) = x$, and $h(x) = \frac{1}{x!} I_A(x)$.

EXAMPLE 11

The $N(\theta, \sigma^2)$ (σ known) p.d.f. is of the exponential type.

DISCUSSION In fact,

$$f(x; \theta) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\theta)^2}{2\sigma^2}} = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{\theta^2}{2\sigma^2}} \times e^{\frac{\theta}{\sigma^2}x} \times e^{-\frac{x^2}{2\sigma^2}},$$

and this is of the form (4) with $C(\theta) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{\theta^2}{2\sigma^2}}$, $Q(\theta) = \frac{\theta}{\sigma^2}$ strictly increasing, $T(x) = x$, and $h(x) = e^{-x^2/2\sigma^2}$.

EXAMPLE 12

The $N(\mu, \theta)$ (μ known) p.d.f. is of the exponential type.

DISCUSSION Here

$$f(x; \theta) = \frac{1}{\sqrt{2\pi\theta}} e^{-\frac{1}{2\theta}(x-\mu)^2},$$

and this is of the form (4) with $C(\theta) = \frac{1}{\sqrt{2\pi\theta}}$, $Q(\theta) = -\frac{1}{2\theta}$ strictly increasing (since $\frac{d}{d\theta}(-\frac{1}{2\theta}) = \frac{1}{2\theta^2} > 0$), $T(x) = (x - \mu)^2$, and $h(x) = 1$.

11.2.2 Uniformly Most Powerful Tests for Some Composite Hypotheses

We may now proceed with the formulation of the following important results.

THEOREM 2

Let X_1, \dots, X_n be a random sample with exponential type p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}$; i.e.,

$$f(x; \theta) = C(\theta) e^{Q(\theta)T(x)} \times h(x), \quad x \in \mathfrak{R},$$

and set $V(x_1, \dots, x_n) = \sum_{i=1}^n T(x_i)$. Then each one of the tests defined below is UMP of level α for testing the hypothesis specified against the respective alternative among all tests of level $\leq \alpha$. Specifically:

(i) Let Q be strictly increasing.

Then for testing $H_0: \theta \leq \theta_0$ against $H_A: \theta > \theta_0$, the UMP test is given by:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } V(x_1, \dots, x_n) > C \\ \gamma & \text{if } V(x_1, \dots, x_n) = C \\ 0 & \text{if } V(x_1, \dots, x_n) < C, \end{cases} \quad (5)$$

where the constants C and γ ($C > 0$, $0 \leq \gamma \leq 1$) are determined by:

$$E_{\theta_0} \varphi(X_1, \dots, X_n) = P_{\theta_0}[V(X_1, \dots, X_n) > C] + \gamma P_{\theta_0}[V(X_1, \dots, X_n) = C] = \alpha. \quad (6)$$

The power of the test is given by:

$$\pi_\varphi(\theta) = P_\theta[V(X_1, \dots, X_n) > C] + \gamma P_\theta[V(X_1, \dots, X_n) = C] \quad (\theta > \theta_0). \quad (7)$$

If the hypothesis to be tested is $H_0: \theta \geq \theta_0$, so that the alternative is $H_A: \theta < \theta_0$, then the UMP test is given by (5) and (6) with reversed inequalities; i.e.,

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } V(x_1, \dots, x_n) < C \\ \gamma & \text{if } V(x_1, \dots, x_n) = C \\ 0 & \text{if } V(x_1, \dots, x_n) > C, \end{cases} \quad (8)$$

where the constants C and γ ($C > 0$, $0 \leq \gamma \leq 1$) are determined by:

$$E_{\theta_0} \varphi(X_1, \dots, X_n) = P_{\theta_0}[V(X_1, \dots, X_n) < C] + \gamma P_{\theta_0}[V(X_1, \dots, X_n) = C] = \alpha. \quad (9)$$

The power of the test is given by:

$$\pi_\varphi(\theta) = P_\theta[V(X_1, \dots, X_n) < C] + \gamma P_\theta[V(X_1, \dots, X_n) = C] \quad (\theta < \theta_0). \quad (10)$$

(ii) Let Q be strictly decreasing.

Then for testing $H_0: \theta \leq \theta_0$ against $H_A: \theta > \theta_0$, the UMP test is given by (8) and (9), and the power is given by (10).

For testing $H_0: \theta \geq \theta_0$ against $H_A: \theta < \theta_0$, the UMP test is given by (5) and (6), and the power is given by (7).

PROOF (Just Pointing Out the Main Points) The proof of this theorem is based on Theorem 1 and also the specific form assumed for the p.d.f. $f(\cdot; \theta)$. As a rough illustration, consider the case that Q is strictly increasing and the hypothesis to be tested is $H_0: \theta \leq \theta_0$. For an arbitrary $\theta_1 < \theta_0$, it is shown that $E_{\theta_1} \varphi(X_1, \dots, X_n) < \alpha$. This establishes that $E_{\theta} \varphi(X_1, \dots, X_n) \leq \alpha$ for all $\theta \leq \theta_0$, so that φ is of level α . Next, take an arbitrary $\theta_1 > \theta_0$ and consider the problem

of testing the simple hypothesis $H_0: \theta = \theta_0$ against the simple alternative $H_{A,1}: \theta = \theta_1$. It is shown that the MP test, provided by Theorem 1, actually, coincides with the test φ given by (5) and (6). This shows that the test φ is UMP. The same reasoning applies for the remaining cases. ▲

Figures 11.2 and 11.3 depict the form of the power of the UMP tests for the one-sided hypotheses $H_0: \theta \leq \theta_0$ and $H_0: \theta \geq \theta_0$.

Figure 11.2

$H_0: \theta \leq \theta_0, H_A:$
 $\theta > \theta_0$: The Power
Curve





Figure 11.3

$H_0: \theta \geq \theta_0, H_A:$
 $\theta < \theta_0$: The Power
Curve



THEOREM 3

Let X_1, \dots, X_n be a random sample with p.d.f. as in Theorem 2, and let $V(x_1, \dots, x_n)$ be as in the same theorem. Consider the problem of testing the hypothesis $H_0: \theta \leq \theta_1$ or $\theta \geq \theta_2$ against the alternative $H_A: \theta_1 < \theta < \theta_2$ at level of significance α . Then the tests defined below are UMP of level α among all tests of level $\leq \alpha$. Specifically:

(i) If Q is strictly increasing, the UMP test is given by:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } C_1 < V(x_1, \dots, x_n) < C_2 \\ \gamma_1 & \text{if } V(x_1, \dots, x_n) = C_1 \\ \gamma_2 & \text{if } V(x_1, \dots, x_n) = C_2 \\ 0 & \text{otherwise,} \end{cases} \quad (11)$$

where the constants C_1, C_2 and γ_1, γ_2 ($C_1 > 0, C_2 > 0, 0 \leq \gamma_1 \leq 1, 0 \leq \gamma_2 \leq 1$) are determined through the relationships:

$$\begin{aligned} E_{\theta_1} \varphi(X_1, \dots, X_n) &= P_{\theta_1}[C_1 < V(X_1, \dots, X_n) < C_2] \\ &\quad + \gamma_1 P_{\theta_1}[V(X_1, \dots, X_n) = C_1] \\ &\quad + \gamma_2 P_{\theta_1}[V(X_1, \dots, X_n) = C_2] = \alpha, \end{aligned} \quad (12)$$

$$\begin{aligned} E_{\theta_2} \varphi(X_1, \dots, X_n) &= P_{\theta_2}[C_1 < V(X_1, \dots, X_n) < C_2] \\ &\quad + \gamma_1 P_{\theta_2}[V(X_1, \dots, X_n) = C_1] \\ &\quad + \gamma_2 P_{\theta_2}[V(X_1, \dots, X_n) = C_2] = \alpha. \end{aligned} \quad (13)$$

The power of the test is given by:

$$\begin{aligned} \pi_\varphi(\theta) &= P_\theta[C_1 < V(X_1, \dots, X_n) < C_2] + \gamma_1 P_\theta[V(X_1, \dots, X_n) = C_1] \\ &\quad + \gamma_2 P_\theta[V(X_1, \dots, X_n) = C_2] \quad (\theta_1 < \theta < \theta_2). \end{aligned} \quad (14)$$

(ii) If Q is strictly decreasing, then the UMP test is given by (11) and (12)–(13) with reversed inequalities; i.e.,

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } V(x_1, \dots, x_n) < C_1 \text{ or } V(x_1, \dots, x_n) > C_2 \\ \gamma_1 & \text{if } V(x_1, \dots, x_n) = C_1 \\ \gamma_2 & \text{if } V(x_1, \dots, x_n) = C_2 \\ 0 & \text{otherwise,} \end{cases} \quad (15)$$

and the constants C_1, C_2 and γ_1, γ_2 are determined by:

$$\begin{aligned} E_{\theta_1} \varphi(X_1, \dots, X_n) &= P_{\theta_1}[V(X_1, \dots, X_n) < C_1 \text{ or } V(X_1, \dots, X_n) > C_2] \\ &\quad + \gamma_1 P_{\theta_1}[V(X_1, \dots, X_n) = C_1] \\ &\quad + \gamma_2 P_{\theta_1}[V(X_1, \dots, X_n) = C_2] = \alpha, \end{aligned} \quad (16)$$

$$\begin{aligned}
 E_{\theta_2} \varphi(X_1, \dots, X_n) &= P_{\theta_2}[V(X_1, \dots, X_n) < C_1 \text{ or } V(X_1, \dots, X_n) > C_2] \\
 &\quad + \gamma_1 P_{\theta_2}[V(X_1, \dots, X_n) = C_1] \\
 &\quad + \gamma_2 P_{\theta_2}[V(X_1, \dots, X_n) = C_2] = \alpha. \quad (17)
 \end{aligned}$$


The power of the test is given by:

$$\begin{aligned}
 \pi_{\varphi}(\theta) &= P_{\theta}[V(X_1, \dots, X_n) < C_1 \text{ or } V(X_1, \dots, X_n) > C_2] \\
 &\quad + \gamma_1 P_{\theta}[V(X_1, \dots, X_n) = C_1] + \gamma_2 P_{\theta}[V(X_1, \dots, X_n) = C_2] \\
 &\quad (\theta_1 < \theta < \theta_2). \quad (18)
 \end{aligned}$$

The power of the UMP test is depicted in Figure 11.4.

Figure 11.4

$H_0: \theta \leq \theta_1 \text{ or } \theta \geq \theta_2$, H_A :
 $\theta_1 < \theta < \theta_2$: The
 Power Curve



Exercises

2.1 If X_1, \dots, X_{16} are independent r.v.'s:

- (i) Construct the MP test of the hypothesis H_0 : the common distribution of the X_i 's is $N(0, 9)$ against the alternative H_A : the common distribution of the X_i 's is $N(1, 9)$; take $\alpha = 0.05$.
- (ii) Also, determine the power of the test.

2.2 Let X_1, \dots, X_n be independent r.v.'s distributed as $N(\mu, \sigma^2)$, where μ is unknown and σ is known.

- (i) For testing the hypothesis $H_0: \mu = 0$ against the alternative $H_A: \mu = 1$, show that the sample size n can be determined to achieve a given level of significance α and given power $\pi(1)$.
- (ii) What is the numerical value of n for $\alpha = 0.05$, $\pi(1) = 0.9$ when $\sigma = 1$?

- 2.3** (i) Let X_1, \dots, X_n be independent r.v.'s distributed as $N(\mu, \sigma^2)$, where μ is unknown and σ is known. Derive the MP test for testing the hypothesis $H_0: \mu = \mu_1$ against the alternative $H_A: \mu = \mu_2$ ($\mu_2 > \mu_1$) at level of significance α .
- (ii) Find an expression for computing the power of the test.
- (iii) Carry out the testing hypothesis and compute the power for $n = 100$, $\sigma^2 = 4$, $\mu_1 = 3$, $\mu_2 = 3.5$, $\bar{x} = 3.2$, and $\alpha = 0.01$.
- 2.4** Let X_1, \dots, X_n be independent r.v.'s distributed as $N(\mu, \sigma^2)$ with μ unknown and σ known. Suppose we wish to test the hypothesis $H_0: \mu = \mu_0$ against the alternative $H_A: \mu = \mu_1$ ($\mu_1 > \mu_0$).
- (i) Derive the MP test for testing H_0 against H_A .
- (ii) For a given level of significance $\alpha_n (< 0.5)$ and given power $\pi_n (> 0.5)$, determine the cutoff point C_n and the sample size for which both α_n and π_n are attained.
- (iii) Show that $\alpha_n \rightarrow 0$ and $\pi_n \rightarrow 1$ as $n \rightarrow \infty$.
- (iv) Determine the sample size n and the cutoff point C_n for $\mu_0 = 0$, $\mu_1 = 1$, $\sigma = 1$, $\alpha_n = 0.001$, and $\pi_n = 0.995$.
- 2.5** Let X_1, \dots, X_n be independent r.v.'s having the Gamma distribution with α known and β unknown.
- (i) Construct the MP test for testing the hypothesis $H_0: \beta = \beta_1$ against the alternative $H_A: \beta = \beta_2$ ($\beta_2 > \beta_1$) at level of significance α .
- (ii) By using the m.g.f. approach, show that, if $X \sim \text{Gamma}(\alpha, \beta)$, then $X_1 + \dots + X_n \sim \text{Gamma}(n\alpha, \beta)$, where the X_i 's are independent and distributed as X .
- (iii) Use the CLT to carry out the test when $n = 30$, $\alpha = 10$, $\beta_1 = 2$, $\beta_2 = 3$, and $\alpha = 0.05$.
- (iv) Compute the power of the test, also by using the CLT.
- 2.6** Let X be a r.v. with p.d.f. $f(x; \theta) = \frac{1}{\theta} e^{-x/\theta}$, $x > 0$, $\theta \in \Omega = (0, \infty)$.
- (i) Refer to Definition 1 in order to show that $f(\cdot; \theta)$ is of the exponential type.
- (ii) Use Theorem 2 in order to derive the UMP test for testing the hypothesis $H_0: \theta \geq \theta_0$ against the alternative $H_A: \theta < \theta_0$ at level of significance α , on the basis of the random sample X_1, \dots, X_n from the above p.d.f.
- (iii) Use the m.g.f. approach in order to show that the r.v. $Y = 2 \times (\sum_{i=1}^n X_i)/\theta$ is distributed as χ_{2n}^2 .
- (iv) Use parts (ii) and (iii) in order to find an expression for the cutoff point C and the power function of the test.
- (v) If $\theta_0 = 1,000$ and $\alpha = 0.05$, determine the sample size n , so that the power of the test at $\theta_1 = 500$ is at least 0.95.
- 2.7** The life of an electronic equipment is a r.v. X whose p.d.f. is $f(x; \theta) = \theta e^{-\theta x}$, $x > 0$, $\theta \in \Omega = (0, \infty)$, and let ℓ be its expected lifetime. On the basis of the random sample X_1, \dots, X_n from this distribution:

- (i) Derive the MP test for testing the hypothesis $H_0: \ell = \ell_0$ against the alternative $H_A: \ell = \ell_1$ ($\ell_1 > \ell_0$) at level of significance α , and write the expression giving the power of the test.
- (ii) Use the m.g.f. approach in order to show that the r.v. $Y = 2\theta \times (\sum_{i=1}^n X_i)$ is distributed as χ_{2n}^2 .
- (iii) Use part (ii) in order to relate the cutoff point and the power of the test to χ^2 -percentiles.
- (iv) Employ the CLT (assuming that n is sufficiently large) in order to find (approximate) values for the cutoff point and the power of the test.
- (v) Use parts (iii) and (iv) in order to carry out the test and also calculate the power when $n = 22$, $\ell_0 = 10$, $\ell_1 = 12.5$, and $\alpha = 0.05$.

2.8 Let X be a r.v. whose p.d.f. f is either the $U(0, 1)$, to be denoted by f_0 , or the Triangular over the interval $[0, 1]$, to be denoted by f_1 (that is, $f_1(x) = 4x$ for $0 \leq x < \frac{1}{2}$; $f_1(x) = 4 - 4x$ for $\frac{1}{2} \leq x \leq 1$, and 0 otherwise).

- (i) Test the hypothesis $H_0: f = f_0$ against the alternative $H_A: f = f_1$ at level of significance $\alpha = 0.05$.
- (ii) Compute the power of the test.
- (iii) Draw the picture of f_1 and compute the power by means of geometric consideration.

2.9 The number of times that an electric light switch can be turned on and off until failure occurs is a r.v. X , which may be assumed to have the Geometric p.d.f. with parameter θ ; i.e., $f(x; \theta) = \theta(1 - \theta)^{x-1}$, $x = 1, 2, \dots$, $\theta \in \Omega = (0, 1)$.

- (i) Refer to Definition 1 in order to show that $f(\cdot; \theta)$ is of the exponential type.
- (ii) Use Theorem 2 in order to derive the UMP test for testing the hypothesis $H_0: \theta = \theta_0$ against the alternative $H_A: \theta > \theta_0$ at level of significance α , on the basis of a random sample of size n from the p.d.f. $f(\cdot; \theta)$.
- (iii) Use the CLT to find an approximate value for the cutoff point C .
- (iv) Carry out the test if $n = 15$, the observed sample mean $\bar{x} = 15,150$, $\theta_0 = 10^{-4}$, and $\alpha = 0.05$.

2.10 Let X be a r.v. with p.d.f. f which is either the $P(1)$ (Poisson with $\lambda = 1$), to be denoted by f_0 , or the $f_1(x) = 1/2^{x+1}$, $x = 0, 1, \dots$. For testing the hypothesis $H_0: f = f_0$ against the alternative $H_A: f = f_1$ on the basis of one observation X :

- (i) Show that the rejection region is defined by: $\{x \geq 0 \text{ integer}; 1.36 \times \frac{x!}{2^x} \geq C\}$ for some positive number C .
- (ii) Determine the level of significance α of the test when $C = 2$.

Hint: Observe that the function $g(x) = \frac{x!}{2^x}$ is nondecreasing for x integer ≥ 1 .

11.3 Some Applications of Theorems 2 and 3

APPLICATION 1 The Binomial Case In reference to Example 9 with $n = 1$, we have $T(x) = x$ where $x = 0, 1$, and then in Theorem 2, $V(x_1, \dots, x_n) = \sum_{i=1}^n x_i$ and $V(X_1, \dots, X_n) = \sum_{i=1}^n X_i \sim B(n, \theta)$. Since $Q(\theta) = \log\left(\frac{\theta}{1-\theta}\right)$ is strictly increasing, consider relations (5) and (6), which, for testing the hypothesis $H_0: \theta \leq \theta_0$ against the alternative $H_A: \theta > \theta_0$, become here:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n x_i > C \\ \gamma & \text{if } \sum_{i=1}^n x_i = C \\ 0 & \text{if } \sum_{i=1}^n x_i < C, \end{cases} \quad (19)$$

$$E_{\theta_0} \varphi(X_1, \dots, X_n) = P_{\theta_0}(X > C) + \gamma P_{\theta_0}(X = C) = \alpha, \quad X \sim B(n, \theta_0). \quad (20)$$

Relation (20) is rewritten below to allow the usage of the Binomial tables for the determination of C and γ ; namely,

$$P_{\theta_0}(X \leq C) - \gamma P_{\theta_0}(X = C) = 1 - \alpha, \quad X \sim B(n, \theta_0). \quad (21)$$

The power of the test is:

$$\begin{aligned} \pi_\varphi(\theta) &= P_\theta(X > C) + \gamma P_\theta(X = C) = 1 - P_\theta(X \leq C) + \gamma P_\theta(X = C), \\ &(\theta > \theta_0), \quad X \sim B(n, \theta). \end{aligned} \quad (22)$$

Numerical Example Refer to Example 2 and suppose that $n = 25$, $\theta_0 = 100p\% = 0.125$, and $\alpha = 0.05$.

DISCUSSION Here, for $\theta = \theta_0$, $X \sim B(25, 0.125)$, and (21) becomes

$$P_{0.125}(X \leq C) - \gamma P_{0.125}(X = C) = 0.95.$$

From the Binomial tables, the value of C which renders $P_{0.125}(X \leq C)$ just above 0.95 is 6 and $P_{0.125}(X \leq 6) = 0.9703$. Also, $P_{0.125}(X = 6) = 0.9703 - 0.9169 = 0.0534$, so that $\gamma = \frac{0.9703 - 0.95}{0.0534} = \frac{0.0203}{0.0534} \simeq 0.38$. Thus, the test in (19) is:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } x > 6 \\ 0.38 & \text{if } x = 6 \\ 0 & \text{if } x < 6. \end{cases}$$

Reject outright the hypothesis that $100p\% = 12.5\%$ if the number of listeners among the sample of 25 is 7 or more, reject the hypothesis with probability 0.38 if this number is 6, and accept the hypothesis if this number is 5 or smaller.

The power of the test is calculated to be as follows, by relation (22):

$$\pi_\varphi(0.1875) \simeq 1 - 0.8261 + 0.38 \times 0.1489 \simeq 0.230,$$

$$\pi_\varphi(0.25) \simeq 1 - 0.5611 + 0.38 \times 0.1828 \simeq 0.508,$$

$$\pi_\varphi(0.375) \simeq 1 - 0.1156 + 0.38 \times 0.0652 \simeq 0.909.$$

If we suppose that the observed value of X is 7, then the P -value is:

$$1 - P_{0.125}(X \leq 7) + 0.38P_{0.125}(X = 7) = 1 - 0.9910 + 0.38 \times 0.0207 \simeq 0.017,$$

so that the result is statistically significant.

Next, for testing the hypothesis $H_0: \theta \geq \theta_0$ against the alternative $H_A: \theta < \theta_0$, relations (8) and (9) become:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n x_i < C \\ \gamma & \text{if } \sum_{i=1}^n x_i = C \\ 0 & \text{if } \sum_{i=1}^n x_i > C, \end{cases} \quad (23)$$

and

$$P_{\theta_0}(X \leq C - 1) + \gamma P_{\theta_0}(X = C) = \alpha, \quad X = \sum_{i=1}^n X_i \sim B(n, \theta_0). \quad (24)$$

The power of the test is:

$$\pi_\varphi(\theta) = P_\theta(X \leq C - 1) + \gamma P_\theta(X = C) \quad (\theta < \theta_0), \quad X \sim B(n, \theta). \quad (25)$$

Numerical Example Refer to Example 1 and suppose that $n = 25$, $\theta_0 = 0.0625$, and $\alpha = 0.05$.

DISCUSSION Here, under θ_0 , $X \sim B(25, 0.0625)$, and (24) becomes

$$P_{0.0625}(X \leq C - 1) + \gamma P_{0.0625}(X = C) = 0.05,$$

so that $C = 0$, and $\gamma P_{0.0625}(X = 0) = 0.1992\gamma = 0.05$. It follows that $\gamma \simeq 0.251$. Therefore the hypothesis is rejected with probability 0.251, if $x = 0$, and is accepted otherwise.

APPLICATION 2 The Poisson Case In reference to Example 10, we have $T(x) = x$, $x = 0, 1, \dots$, and then in Theorem 2, $V(x_1, \dots, x_n) = \sum_{i=1}^n x_i$ and $V(X_1, \dots, X_n) = \sum_{i=1}^n X_i \sim P(n\theta)$. Since $Q(\theta) = \log \theta$ is strictly increasing, consider relations (8) and (9) for testing $H_0: \theta > \theta_0$ against $H_A: \theta \leq \theta_0$. They become here:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n x_i < C \\ \gamma & \text{if } \sum_{i=1}^n x_i = C \\ 0 & \text{if } \sum_{i=1}^n x_i > C, \end{cases} \quad (26)$$

$$E_{\theta_0} \varphi(X_1, \dots, X_n) = P_{\theta_0}(X < C) + \gamma P_{\theta_0}(X = C) = \alpha, \quad X = \sum_{i=1}^n X_i \sim P(n\theta_0),$$

or

$$P_{\theta_0}(X \leq C - 1) + \gamma P_{\theta_0}(X = C) = \alpha, \quad X \sim P(n\theta_0). \quad (27)$$

The power of the test is:

$$\pi_\varphi(\theta) = P_\theta(X \leq C - 1) + \gamma P_\theta(X = C) \quad (\theta \leq \theta_0), \quad X \sim P(n\theta). \quad (28)$$

Unfortunately, no numerical application for Example 3 can be given as the Poisson tables do not provide entries for $\theta_0 = 200$. In order to be able to apply the test defined by (26) and (27), consider the following example.

EXAMPLE 13

Let X_1, \dots, X_{20} be i.i.d. r.v.'s denoting the number of typographical errors in 20 pages of a book. We may assume that the X_i 's are independently distributed as $P(\theta)$, and let us test the hypothesis $H_0: \theta > 0.5$ (the average number of errors is more than 1 per couple of pages) against the alternative $H_A: \theta \leq 0.5$ at level $\alpha = 0.05$.

DISCUSSION In (27), $X \sim P(10)$, so that

$$P_{0.5}(X \leq C - 1) + \gamma P_{0.5}(X = C) = 0.05,$$

and hence $C - 1 = 4$ and $P_{0.5}(X \leq 4) = 0.0293$, $P_{0.5}(X = 5) = 0.0671 - 0.0293 = 0.0378$. It follows that $\gamma = \frac{0.05 - 0.0293}{0.0378} \simeq 0.548$. Therefore by (26), reject the hypothesis outright if $x \leq 4$, reject it with probability 0.548 if $x = 5$, and accept it otherwise. The power of the test is: For $\theta = 0.2$, $X \sim P(4)$ and:

$$\pi_\varphi(0.2) = P_{0.2}(X \leq 4) + 0.548 P_{0.2}(X = 5) = 0.6288 + 0.548 \times 0.1563 \simeq 0.714.$$

If the observed value x is 6, then the P -value is (for $X \sim P_{0.5}(10)$): $P(X \leq 5) + 0.548 P_{0.5}(X = 6) \simeq 0.102$.

APPLICATION 3 The Normal Case: Testing Hypotheses About the Mean Refer to Example 11 and observe that $T(x) = x$ and $Q(\theta)$ is strictly increasing. Therefore the appropriate test for testing $H_0: \theta \leq \theta_0$ against $H_A: \theta > \theta_0$ at level of significance α is given by (5) and (6) with $\gamma = 0$. That is,

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n x_i > C \\ 0 & \text{otherwise,} \end{cases} \quad (29)$$

or

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \frac{\sqrt{n}(\bar{x} - \theta_0)}{\sigma} > z_\alpha \\ 0 & \text{otherwise,} \end{cases} \quad (29')$$

because

$$\begin{aligned} \alpha &= E_{\theta_0} \varphi(X_1, \dots, X_n) = P_{\theta_0} \left(\sum_{i=1}^n X_i > C \right) \\ &= P_{\theta_0} \left[\frac{\sqrt{n}(\bar{X} - \theta_0)}{\sigma} > \frac{C - n\theta_0}{\sigma\sqrt{n}} \right], \end{aligned}$$


so that $\frac{C - n\theta_0}{\sigma\sqrt{n}} = z_\alpha$ and therefore

$$C = n\theta_0 + z_\alpha \sigma \sqrt{n}; \quad (30)$$

this is so, because $\frac{\sqrt{n}(\bar{X} - \theta_0)}{\sigma} = Z \sim N(0, 1)$, and recall that $P(Z > z_\alpha) = \alpha$.

Figure 11.5

Rejection Region of the Hypothesis H_0 : $\theta \leq \theta_0$ (the Shaded Area) in the Form (29') and the Respective Probability



The power of the test is given by:

$$\pi_{\varphi}(\theta) = 1 - \Phi \left[z_{\alpha} + \frac{\sqrt{n}(\theta_0 - \theta)}{\sigma} \right], \quad \theta > \theta_0, \quad (31)$$

because, on account of (30):

$$\begin{aligned} \pi_{\varphi}(\theta) &= P_{\theta} \left(\sum_{i=1}^n X_i > C \right) = P_{\theta} \left[\sum_{i=1}^n X_i - n\theta > n(\theta_0 - \theta) + z_{\alpha} \sigma \sqrt{n} \right] \\ &= P_{\theta} \left[\frac{\sum_{i=1}^n X_i - n\theta}{\sigma \sqrt{n}} > z_{\alpha} + \frac{\sqrt{n}(\theta_0 - \theta)}{\sigma} \right] \\ &= P \left[Z > z_{\alpha} + \frac{\sqrt{n}(\theta_0 - \theta)}{\sigma} \right] = 1 - \Phi \left[z_{\alpha} + \frac{\sqrt{n}(\theta_0 - \theta)}{\sigma} \right], \end{aligned}$$

since $\frac{\sum_{i=1}^n X_i - n\theta}{\sigma \sqrt{n}} = Z \sim N(0, 1)$.

Numerical Example In reference to Example 5, focus on patients treated with the new treatment, and call $Y \sim N(\theta, \sigma^2)$ (σ known) the survival time. On the basis of observations on $n = 25$ such patients, we wish to test the hypothesis $H_0: \theta \leq 5$ (in years) against $H_A: \theta > 5$ at level of significance $\alpha = 0.01$. For simplicity, take $\sigma = 1$.

DISCUSSION Here $z_{\alpha} = z_{0.01} = 2.33$, so that $C = 25 \times 5 + 2.33 \times 1 \times 5 = 136.65$. Thus, reject H_0 if the total of survival years is > 136.65 , and accept H_0 otherwise.

The power of the test is given by (31) and is:

$$\begin{aligned} \text{For } \theta = 5.5, \pi_{\varphi}(5.5) &= 1 - \Phi[2.33 + 5(5 - 5.5)] = 1 - \Phi(-0.17) \\ &= \Phi(0.17) = 0.567495; \end{aligned}$$

$$\begin{aligned} \text{and for } \theta = 6, \pi_{\varphi}(6) &= 1 - \Phi[2.33 + 5(6 - 5.5)] = 1 - \Phi(-2.67) \\ &= \Phi(2.67) = 0.996207. \end{aligned}$$

If we suppose that the observed value of $\sum_{i=1}^{25} x_i$ is equal to 138, then the P -value is $P(\sum_{i=1}^{25} X_i > 138) = 1 - \Phi(\frac{138-125}{5}) = 1 - \Phi(2.6) = 1 - 0.995339 = 0.004661$, so that the result is highly statistically significant.

APPLICATION 4 The Normal Case (continued) Testing Hypotheses About the Variance Refer to Example 12, where $T(x) = (x - \mu)^2$ (μ known) and Q is strictly increasing. Then, for testing the hypothesis $H_0: \sigma^2 \geq \sigma_0^2$ against the alternative $H_A: \sigma^2 < \sigma_0^2$ (or $\theta \geq \theta_0$ against $\theta < \theta_0$ with $\theta = \sigma^2$ and $\theta_0 = \sigma_0^2$) at level of significance α , the appropriate test is given by (8) and (9) (with $\gamma = 0$), and it is here:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n (x_i - \mu)^2 < C \\ 0 & \text{otherwise,} \end{cases} \quad (32)$$

or

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma_0}\right)^2 < \chi_{n;1-\alpha}^2 \\ 0 & \text{otherwise,} \end{cases} \quad (32')$$

because

$$\begin{aligned} \alpha &= E_{\sigma_0^2} \varphi(X_1, \dots, X_n) = P_{\sigma_0^2} \left[\sum_{i=1}^n (X_i - \mu)^2 < C \right] \\ &= P_{\sigma_0^2} \left[\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma_0}\right)^2 < \frac{C}{\sigma_0^2} \right], \end{aligned}$$


so that $\frac{C}{\sigma_0^2} = \chi_{n;1-\alpha}^2$ and therefore

$$C = \sigma_0^2 \chi_{n;1-\alpha}^2; \quad (33)$$

this is so, because $\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma_0}\right)^2 \sim \chi_n^2$.

Figure 11.6

Rejection Region of the Hypothesis $H_0: \sigma^2 \geq \sigma_0^2$ (the Shaded Area) in the Form (32') and the Respective Probability



By slightly abusing the notation and denoting by χ_n^2 also a r.v. which has the χ_n^2 distribution, the power of the test is given by:

$$\pi_\varphi(\sigma^2) = P\left(\chi_n^2 < \frac{\sigma_0^2}{\sigma^2} \chi_{n;1-\alpha}^2\right), \quad \sigma^2 < \sigma_0^2, \quad (34)$$

because, on account of (33):

$$\begin{aligned}\pi_\varphi(\sigma^2) &= P_{\sigma^2} \left[\sum_{i=1}^n (X_i - \mu)^2 < C \right] = P_{\sigma^2} \left[\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 < \frac{C}{\sigma^2} \right] \\ &= P_{\sigma^2} \left[\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 < \frac{\sigma_0^2}{\sigma^2} \chi_{n;1-\alpha}^2 \right] = P \left(\chi_n^2 < \frac{\sigma_0^2}{\sigma^2} \chi_{n;1-\alpha}^2 \right),\end{aligned}$$

since $\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 \sim \chi_n^2$.

Numerical Example Suppose $n = 40$, $\sigma_0 = 2$, and $\alpha = 0.025$. For simplicity, take $\mu = 0$.

DISCUSSION Here $\chi_{n;1-\alpha}^2 = \chi_{40;0.975}^2 = 24.433$ and $C = 4 \times 24.433 = 97.732$. Thus, by means of (32), the hypothesis is rejected if $\sum_{i=1}^{40} x_i^2 < 97.732$, and it is accepted otherwise.

For $\sigma = 1.25$, for example, the power of the test is, by means of (34),

$$\begin{aligned}\pi_\varphi(\sigma^2) &= \pi_\varphi(2.25) = P \left(\chi_{40}^2 < \frac{97.732}{1.5625} \right) \\ &= P(\chi_{40}^2 < 62.548) = 0.986\end{aligned}$$

(by linear interpolation).

If we suppose that the observed value of $\sum_{i=1}^{40} x_i^2$ is 82.828, then the P -value is

$$P_4 \left(\sum_{i=1}^{40} X_i^2 < 82.828 \right) = P_4 \left[\sum_{i=1}^{40} \left(\frac{X_i}{2} \right)^2 < 20.707 \right] = 0.05,$$

which indicates strong rejection.

APPLICATION 5 The Normal Case (continued) Testing Further Hypotheses About the Mean In reference to Example 11, $T(x) = x$ and $Q(\theta)$ is strictly increasing. Therefore, for testing $H_0: \theta \leq \theta_1$ or $\theta \geq \theta_2$ against $H_A: \theta_1 < \theta < \theta_2$ at level of significance α , the test to be employed is the one given by (11) and (12)–(13), which here becomes ($\gamma_1 = \gamma_2 = 0$):

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } C_1 \leq \sum_{i=1}^n x_i \leq C_2 \\ 0 & \text{otherwise,} \end{cases} \quad (35)$$

$$\begin{aligned}E_{\theta_1} \varphi(X_1, \dots, X_n) &= P_{\theta_1} \left(C_1 \leq \sum_{i=1}^n X_i \leq C_2 \right) = \alpha, \\ E_{\theta_2} \varphi(X_1, \dots, X_n) &= P_{\theta_2} \left(C_1 \leq \sum_{i=1}^n X_i \leq C_2 \right) = \alpha,\end{aligned} \quad (36)$$

and $\sum_{i=1}^n X_i \sim N(n\theta_i, n\sigma^2)$, $i = 1, 2$.

For the purpose of utilizing the Normal tables, (36) are rewritten thus:

$$\Phi\left(\frac{C_2 - n\theta_i}{\sigma\sqrt{n}}\right) - \Phi\left(\frac{C_1 - n\theta_i}{\sigma\sqrt{n}}\right) = \alpha, \quad i = 1, 2. \quad (37)$$

The power of the test is calculated as follows:

$$\pi_\varphi(\theta) = \Phi\left(\frac{C_2 - n\theta}{\sigma\sqrt{n}}\right) - \Phi\left(\frac{C_1 - n\theta}{\sigma\sqrt{n}}\right), \quad (\theta_1 < \theta < \theta_2). \quad (38)$$

Numerical Example In reference to Example 4, suppose $n = 25$, $\theta_1 = 1$, $\theta_2 = 3$, and $\alpha = 0.01$. For simplicity, let us take $\sigma = 1$.

DISCUSSION Here $n\theta_1 = 25$, $n\theta_2 = 75$, and (37) yields:

$$\Phi\left(\frac{C_2 - 25}{5}\right) - \Phi\left(\frac{C_1 - 25}{5}\right) = \Phi\left(\frac{C_2 - 75}{5}\right) - \Phi\left(\frac{C_1 - 75}{5}\right) = 0.01. \quad (39)$$

Placing the four quantities $\frac{C_1 - 75}{5}$, $\frac{C_2 - 75}{5}$, $\frac{C_1 - 25}{5}$, and $\frac{C_2 - 25}{5}$ under the $N(0, 1)$ curve, we observe that relation (39) obtains only for:

$$\frac{C_1 - 25}{5} = -\frac{C_2 - 75}{5} \quad \text{and} \quad \frac{C_2 - 25}{5} = -\frac{C_1 - 75}{5},$$

which imply that $C_1 + C_2 = 100$. Setting $C_1 = C$, we have then that $C_2 = 100 - C$, and (39) gives:

$$\Phi\left(\frac{75 - C}{5}\right) - \Phi\left(\frac{C - 25}{5}\right) = 0.01. \quad (40)$$

From the Normal tables, we find that (40) is closely satisfied for $C = 36.5$. So $C_1 = 36.5$ and hence $C_2 = 63.5$, and the test rejects the hypothesis H_0 whenever $\sum_{i=1}^{25} x_i$ is between 36.5 and 63.5 and accepts it otherwise.

The power of the test, calculated through (38), is, for example, for $\theta = 2.5$ and $\theta = 2$:

$$\pi_\varphi(1.5) = \pi_\varphi(2.5) = 0.57926 \quad \text{and} \quad \beta_\varphi(2) = 0.99307.$$

Exercises

- 3.1** (i) In reference to Example 8 in Chapter 1, the appropriate model is the Binomial model with $X_i = 1$ if the i th young adult listens to the program, and $X_i = 0$ otherwise, where $P(X_i = 1) = p$, and the X_i 's are independent, so that $X = \sum_{i=1}^n X_i \sim B(n, p)$.
- (ii) The claim is that $p > p_0$ some specified number $0 < p_0 < 1$, and the claim is checked by testing the hypothesis $H_0: p \leq p_0$ against the alternative $H_A: p > p_0$ at level of significance α .
- (iii) For $p_0 = 5\%$, $n = 100$, and $\alpha = 0.02$, use the CLT to carry out the test.
- 3.2** (i) In reference to Example 9 in Chapter 1, the appropriate model is the Binomial model with $X_i = 1$ if the i th item is defective, and 0 otherwise, where $P(X_i = 1) = p$, and the X_i 's are independent, so that $X = \sum_{i=1}^n X_i \sim B(n, p)$.

- (ii) The process is under control if $p \leq p_0$, where p_0 is a specified number with $0 < p_0 < 1$, and the hypothesis to be checked is $H_0: p > p_0$ against the alternative $H_A: p \leq p_0$ at level of significance α .
- (iii) For $p_0 = 0.0625$, $n = 100$, and $\alpha = 0.10$, use the CLT to carry out the test.
- 3.3** (i) In reference to Example 10 in Chapter 1, the appropriate model is the Binomial model with $X_i = 1$ if the i th flawed specimen is identified as such, and $X_i = 0$, otherwise, where $P(X_i = 1) = p$, and the X_i 's are independent, so that $X = \sum_{i=1}^n X_i \sim B(n, p)$.
- (ii) The electronic scanner is superior to the mechanical testing if $p > p_0$, some specified p_0 with $0 < p_0 < 1$, and this is checked by testing the hypothesis $H_0: p \leq p_0$ against the alternative $H_A: p > p_0$ at level of significance α .
- (iii) For $p_0 = 90\%$, $n = 100$, and $\alpha = 0.05$, use the CLT to carry out the test.
- 3.4** (i) In a certain university, 400 students were chosen at random and it was found that 95 of them were women. On the basis of this, test the hypothesis H_0 : the proportion of women is 25% against the alternative H_A : the proportion of women is less than 25% at level of significance $\alpha = 0.05$.
- (ii) Use the CLT in order to determine the cutoff point.
- 3.5** Let X_1, \dots, X_n be independent r.v.'s distributed as $B(1, p)$. For testing the hypothesis $H_0: p \leq \frac{1}{2}$ against the alternative $H_A: p > \frac{1}{2}$, use the CLT in order to determine the sample size n for which the level of significance and power are, respectively, $\alpha = 0.05$ and $\pi(7/8) = 0.95$.
- 3.6** Let X be a r.v. distributed as $B(n, \theta)$, $\theta \in \Omega = (0, 1)$.
- (i) Use relations (19) and (20) to set up the UMP test for testing the hypothesis $H_0: \theta \leq \theta_0$ against the alternative $H_A: \theta > \theta_0$ at level of significance α .
- (ii) Specify the test in part (i) for $n = 10$, $\theta_0 = 0.25$, and $\alpha = 0.05$.
- (iii) Compute the power of the test for $\theta_1 = 0.375, 0.500$.
- (iv) For $\theta > 0.5$, show that: $P_\theta(X \leq C) = 1 - P_{1-\theta}(X \leq n - C - 1)$ and hence $P_\theta(X = C) = P_{1-\theta}(X \leq n - C) - P_{1-\theta}(X \leq n - C - 1)$.
- (v) Use part (iv) to compute the power of the test for $\theta_1 = 0.625, 0.875$.
- (vi) Use the CLT in order to determine the sample size n if $\theta_0 = 0.125$, $\alpha = 0.1$, and $\pi(0.25) = 0.9$.
- 3.7** (i) In reference to Example 12 in Chapter 1, the appropriate model to be used is the Poisson model; i.e., $X \sim P(\lambda)$.
- (ii) The safety level is specified by $\lambda \leq 200$, and this is checked by testing the hypothesis $H_0: \lambda > 200$ against the alternative $H_A: \lambda \leq 200$ at level of significance α .
- (iii) On the basis of a random sample of size $n = 100$, use the CLT in order to carry out the test for $\alpha = 0.05$.

3.8 The number of total traffic accidents in a certain city during a year is a r.v. X , which may be assumed to be distributed as $P(\lambda)$. For the last year, the observed value of X was $x = 4$, whereas for the past several years, the average was 10.

- (i) Formulate the hypothesis that the average remains the same against the alternative that there is an improvement.
- (ii) Refer to Application 2 in order to derive the UMP test for testing the hypothesis of part (i) at level $\alpha = 0.01$.

3.9 (i) In reference to Example 16 in Chapter 1, a suitable model would be to assume that $X \sim N(\mu_1, \sigma^2)$, $Y \sim N(\mu_2, \sigma^2)$ and that they are independent.

- (ii) Let μ_0 be the known mean survival period (in years) for the existing treatment. Then the claim is that $\mu_2 > \mu_0$, and this is to be checked by testing the hypothesis $H_0: \mu_2 \leq \mu_0$ against the alternative $H_A: \mu_2 > \mu_0$ at level of significance α .
- (iii) Carry out the test if $n = 100$, $\mu_0 = 5$, and $\alpha = 0.05$.

3.10 The life length of a 50-watt light bulb of a certain brand is a r.v. X , which may be assumed to be distributed as $N(\mu, \sigma^2)$ with unknown μ and σ known. Let X_1, \dots, X_n be a random sample from this distribution and suppose that we are interested in testing the hypothesis $H_0: \mu = \mu_0$ against the alternative $H_A: \mu < \mu_0$ at level of significance α .

- (i) Derive the UMP test.
- (ii) Derive the formula for the power of the test.
- (iii) Carry out the testing hypothesis problem when $n = 25$, $\mu_0 = 1,800$, $\sigma = 150$ (in hours), $\alpha = 0.01$, and $\bar{x} = 1,730$. Also, calculate the power at $\mu = 1,700$.

3.11 The rainfall at a certain station during a year is a r.v. X , which may be assumed to be distributed as $N(\mu, \sigma^2)$ with μ unknown and $\sigma = 3$ inches. For the past 10 years, the record provides the following rainfalls:

$$x_1 = 30.5, \quad x_2 = 34.1, \quad x_3 = 27.9, \quad x_4 = 29.4, \quad x_5 = 35.0,$$

$$x_6 = 26.9, \quad x_7 = 30.2, \quad x_8 = 28.3, \quad x_9 = 31.7, \quad x_{10} = 25.8.$$

Test the hypothesis $H_0: \mu = 30$ against the alternative $H_A: \mu < 30$ at level of significance $\alpha = 0.05$.

3.12 Let $X_i, i = 1, \dots, 4$ and $Y_j, j = 1, \dots, 4$ be two independent random samples from the distributions $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$, respectively. Suppose that the observed values of the X_i 's and the Y_j 's are as follows:

$$x_1 = 10.1, \quad x_2 = 8.4, \quad x_3 = 14.3, \quad x_4 = 11.7,$$

$$y_1 = 9.0, \quad y_2 = 8.2, \quad y_3 = 12.1, \quad y_4 = 10.3.$$

Suppose that $\sigma_1 = 4$ and $\sigma_2 = 3$.

Then test the hypothesis that the two means differ in absolute value by at least 1 unit. That is, if $\theta = \mu_1 - \mu_2$, then the hypothesis to be tested

is $H_0: |\theta| \leq 1$, or, equivalently, $H_0: \theta \leq -1$ or $\theta \geq 1$. The alternative is $H_A: -1 < \theta < 1$. Take $\alpha = 0.05$.

Hint: Set $Z_i = X_i - Y_i$, so that the Z_i 's are independent and distributed as $N(\mu, 25)$. Then use appropriately Theorem 3.

3.13 (i) On the basis of the independent r.v.'s X_1, \dots, X_{25} , distributed as $N(0, \sigma^2)$, test the hypothesis $H_0: \sigma \leq 2$ against the alternative $H_A: \sigma > 2$ at level of significance $\alpha = 0.05$.

(ii) Specify the test when the observed values x_i 's of the X_i 's are such that $\sum_{i=1}^{25} x_i^2 = 120$.

3.14 The diameters of bolts produced by a certain machine are independent r.v.'s distributed as $N(\mu, \sigma^2)$ with μ known. In order for the bolts to be usable for the intended purpose, the s.d. σ must not exceed 0.04 inch. A random sample of size $n = 16$ is taken and it is found that $s = 0.05$ inch. Formulate the appropriate testing hypothesis problem and carry out the test at level of significance $\alpha = 0.05$.

11.4 Likelihood Ratio Tests

In the previous sections, UMP tests were constructed for several important hypotheses and were illustrated by specific examples. Those tests have the UMP property, provided the underlying p.d.f. is of the exponential type given in (4). What happens if either the p.d.f. is not of this form and/or the hypotheses to be tested are not of the type for which UMP tests exist? One answer is for sure that the testing activities will not be terminated here; other procedures are to be invented and investigated. Such a procedure is one based on the Likelihood Ratio, which gives rise to the so-called *Likelihood Ratio* (LR) tests. The rationale behind this procedure was given in Section 3 of Chapter 8. What we are doing in this section is to apply it to some specific cases and produce the respective LR tests in a usable form.

As already explained, LR tests do have a motivation which is, at least intuitively, satisfactory, although they do not possess, in general, a property such as the UMP property. The LR approach also applies to multidimensional parameters and leads to manageable tests. In addition, much of the work needed to set up a LR test has already been done in Section 1 of Chapter 9 about MLE's. In our discussions below, we restrict ourselves to the Normal case, where exact tests do exist. In the next chapter, we proceed with the Multinomial distribution, where we have to be satisfied with approximations.

The basics here, as we recall from Chapter 8, Section 8.3, are as follows: X_1, \dots, X_n is a random sample from the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}^r$, $r \geq 1$, and ω is a (proper) subset of Ω . On the basis of this random sample, test the hypothesis $H_0: \theta \in \omega$ at level of significance α . (In the present framework, the alternative is $H_A: \theta \notin \omega$, but is not explicitly stated.) Then, by relation (8) in Chapter 8 and the discussion following it, reject H_0 whenever

$$\lambda < \lambda_0, \text{ where } \lambda_0 \text{ is a constant to be specified,} \quad (41)$$

or

$$g(\lambda) > g(\lambda_0), \text{ or } g(\lambda) < g(\lambda_0), \text{ for some strictly monotone function } g; \quad (42)$$

$g(\lambda) = -2 \log \lambda$ is such a function, and H_0 may be rejected whenever

$$-2 \log \lambda > C, \text{ a constant to be determined} \quad (43)$$

(see relation (10) in Chapter 8).

Recall that

$$\lambda = \lambda(x_1, \dots, x_n) = \frac{L(\hat{\omega})}{L(\hat{\Omega})}, \quad (44)$$

where, with $\mathbf{x} = (x_1, \dots, x_n)$, the observed value of (X_1, \dots, X_n) , $L(\hat{\Omega})$ is the maximum of the likelihood function $L(\theta | \mathbf{x})$, which obtains if θ is replaced by its MLE, and $L(\hat{\omega})$ is again the maximum of the likelihood function under the restriction that θ lies in ω . Clearly, $L(\hat{\omega}) = L(\hat{\theta}_\omega)$, where $\hat{\theta}_\omega$ is the MLE of θ under the restriction that θ lies in ω . Actually, much of the difficulty associated with the present method stems from the fact that, in practice, obtaining $\hat{\theta}_\omega$ is far from a trivial problem.

The following two examples shed some light on how a LR test is actually constructed. These examples are followed by a series of applications to normal populations.

EXAMPLE 14

Determine the LR test for testing the hypothesis $H_0: \theta = 0$ (against the alternative $H_A: \theta \neq 0$) at level of significance α on the basis of one observation from the p.d.f. $f(x; \theta) = \frac{1}{\pi} \times \frac{1}{1+(x-\theta)^2}$, $x \in \mathfrak{R}$, $\theta > 0$ (the Cauchy p.d.f.)

DISCUSSION First, $f(\cdot; \theta)$ is a p.d.f., since

$$\begin{aligned} \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{dx}{1+(x-\theta)^2} &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{dy}{1+y^2} \quad (\text{by setting } x-\theta = y) \\ &= \frac{1}{\pi} \int_{-\pi/2}^{\pi/2} dt = 1 \\ &\left(\text{by setting } y = \tan t, \text{ so that } 1+y^2 = 1 + \frac{\sin^2 t}{\cos^2 t} = \frac{1}{\cos^2 t}, \right. \\ &\quad \left. \frac{dy}{dt} = \frac{d}{dt} \left(\frac{\sin t}{\cos t} \right) = \frac{1}{\cos^2 t}, \text{ and } -\frac{\pi}{2} < t < \frac{\pi}{2} \right). \end{aligned}$$

Next, clearly, $L(\theta | x) (= f(x; \theta))$ is maximized for $\theta = x$, so that $\lambda = \frac{1}{\pi} \times \frac{1}{1+x^2} / \frac{1}{\pi} = \frac{1}{1+x^2}$, and $\lambda < \lambda_0$, if and only if $x^2 > \frac{1}{\lambda_0} - 1 = C$, or $x < -C$ or $x > C$, where C is determined through the relation: $P_0(X < -C \text{ or } X > C) = \alpha$, or $P(X > C) = \frac{\alpha}{2}$ due to the symmetry (around 0) of the p.d.f. $f(x; 0)$. But

$$P(X > C) = \int_C^{\infty} \frac{1}{\pi} \times \frac{dx}{1+x^2} = \frac{1}{\pi} \int_{\tan^{-1} C}^{\pi/2} dt = \frac{1}{\pi} \left(\frac{\pi}{2} - \tan^{-1} C \right) = \frac{\alpha}{2},$$

or $\tan^{-1} C = \frac{(1-\alpha)\pi}{2}$, and hence $C = \tan\left(\frac{(1-\alpha)\pi}{2}\right)$. So, H_0 is rejected whenever $x < -\tan\left(\frac{(1-\alpha)\pi}{2}\right)$ or $x > \tan\left(\frac{(1-\alpha)\pi}{2}\right)$. For example, for $\alpha = 0.05$, $C = \tan(0.475\pi) \simeq 12.706$, and H_0 is rejected when $x < -12.706$ or $x > 12.706$.

EXAMPLE 15

Let X_1, \dots, X_n be a random sample of size n from the Negative Exponential p.d.f. $f(x; \theta) = \theta e^{-\theta x}$, $x > 0$ ($\theta > 0$). Derive the LR test for testing the hypothesis $H_0: \theta = \theta_0$ (against the alternative $H_A: \theta \neq \theta_0$) at level of significance α .

DISCUSSION Here

$$L(\theta | \mathbf{x}) = \theta^n e^{-\theta t}, \quad \text{where } \mathbf{x} = (x_1, \dots, x_n) \text{ and } t = \sum_{i=1}^n x_i.$$

We also know that the MLE of θ is $\hat{\theta} = 1/\bar{x} = n/t$. Therefore the LR λ is given by:


$$\lambda = \theta_0^n e^{-\theta_0 t} / \left(\frac{n}{t}\right)^n e^{-n} = \left(\frac{e\theta_0}{n}\right)^n t^n e^{-\theta_0 t},$$

and hence $\lambda < \lambda_0$, if and only if $te^{-\frac{\theta_0}{n}t} < C_0 (= \frac{n\lambda_0^{1/n}}{e\theta_0})$. We wish to determine the cutoff point C_0 . To this end, set $g(t) = te^{-dt}$ ($d = \theta_0/n$) and observe that $g(t)$ is increasing for $0 < t < \frac{1}{d} = \frac{n}{\theta_0}$, decreasing for $t > \frac{n}{\theta_0}$, and attains its maximum at $t = n/\theta_0$ (see Figure 11.7). It follows that $te^{-dt} < C_0$, if and only if $t < C_1$ or $t > C_2$. Therefore, by setting $T = \sum_{i=1}^n X_i$, we have: $P_{\theta_0}(Te^{-\frac{\theta_0}{n}T} < C_0) = \alpha$, if and only if $P_{\theta_0}(T < C_1 \text{ or } T > C_2) = \alpha$. For simplicity, let us take the two-tail probabilities equal. Thus,

$$P_{\theta_0}(T < C_1) = P(T > C_2) = \frac{\alpha}{2}.$$

Figure 11.7

Graphical Determination of the Rejection Region



By the fact that the independent X_i 's have the $f(x; \theta_0) = \theta_0 e^{-\theta_0 x}$ p.d.f. (under H_0), it follows that T is distributed as Gamma with $\alpha = n$ and $\beta = \frac{1}{\theta_0}$. Therefore its p.d.f. is given by:

$$f_T(t) = \frac{\theta_0^n}{\Gamma(n)} t^{n-1} e^{-\theta_0 t}, \quad t > 0.$$

Then C_1 and C_2 are determined by:

$$\int_0^{C_1} f_T(t) dt = \int_{C_2}^{\infty} f_T(t) dt = \frac{\alpha}{2}.$$

In order to be able to proceed further, take, e.g., $n = 2$. Then

$$f_T(t) = \theta_0^2 t e^{-\theta_0 t}, \quad t > 0,$$

and

$$\int_0^{C_1} \theta_0^2 t e^{-\theta_0 t} dt = 1 - e^{-\theta_0 C_1} - \theta_0 C_1 e^{-\theta_0 C_1},$$

$$\int_{C_2}^{\infty} \theta_0^2 t e^{-\theta_0 t} dt = e^{-\theta_0 C_2} + \theta_0 C_2 e^{-\theta_0 C_2}.$$

Thus, the relations $P_{\theta_0}(T < C_1) = P_{\theta_0}(T > C_2) = \frac{\alpha}{2}$ become, equivalently, for $\alpha = 0.05$:

$$0.975e^p - p = 1, \quad p = \theta_0 C_1; \quad 0.025e^q - q = 1, \quad q = \theta_0 C_2.$$

By trial and error, we find: $p = 0.242$ and $q = 5.568$, so that $C_1 = 0.242/\theta_0$ and $C_2 = 5.568/\theta_0$. Thus, for $n = 2$ and by splitting the error $\alpha = 0.05$ equally between the two tails, the LR test rejects H_0 when $t(=x_1 + x_2) < 0.242/\theta_0$ or $t > 5.568/\theta_0$. For example, for $\theta_0 = 1$, the test rejects H_0 when $t < 0.242$ or $t > 5.568$.

APPLICATIONS TO THE NORMAL CASE The applications to be discussed here are organized as follows: First, we consider the one-sample case and test a hypothesis about the mean, regardless of whether the variance is known or unknown. Next, a hypothesis is tested about the variance, regardless of whether the mean is known or not. Second, we consider the two-sample problem and make the realistic assumption that all parameters are unknown. Then the hypothesis is tested about the equality of the means, and, finally, the variances are compared through their ratio.

11.4.1 Testing Hypotheses for the Parameters in a Single Normal Population

Here X_1, \dots, X_n is a random sample from the $N(\mu, \sigma^2)$, and we are interested in testing: (i) $H_0: \mu = \mu_0, \sigma$ known; (ii) $H_0: \mu = \mu_0, \sigma$ unknown; (iii) $H_0: \sigma = \sigma_0$ (or $\sigma^2 = \sigma_0^2$), μ known; (iv) $H_0: \sigma = \sigma_0$ (or $\sigma^2 = \sigma_0^2$), μ unknown.

DISCUSSION

(i) $H_0: \mu = \mu_0, \sigma$ known. Under H_0 ,

$$L(\hat{\omega}) = (2\pi\sigma^2)^{-n/2} \exp\left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu_0)^2\right],$$

and

$$L(\hat{\Omega}) = (2\pi\sigma^2)^{-n/2} \exp\left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \bar{x})^2\right], \quad \text{since } \hat{\mu}_\Omega = \bar{x}.$$

Forming the likelihood ratio λ and taking $-2 \log \lambda$, we have:

$$-2 \log \lambda = \frac{1}{\sigma^2} \sum_{i=1}^n [(x_i - \mu_0)^2 - (x_i - \bar{x})^2] = \left[\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} \right]^2. \quad (45)$$

Then

$$-2 \log \lambda > \lambda_0, \quad \text{if and only if } \left[\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} \right]^2 > C^2 \quad (\text{some } C = \lambda_0^{1/2} > 0),$$

and this happens, if and only if

$$\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} < -C, \quad \text{or} \quad \frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} > C.$$


Under H_0 , $\frac{\sqrt{n}(\bar{X} - \mu_0)}{\sigma} \sim N(0, 1)$, so that the relation

$$P_{\mu_0} \left[\frac{\sqrt{n}(\bar{X} - \mu_0)}{\sigma} < -C, \text{ or } \frac{\sqrt{n}(\bar{X} - \mu_0)}{\sigma} > C \right] = \alpha, \quad \text{gives } C = z_{\alpha/2}.$$

Thus, the likelihood ratio test is:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} < -z_{\alpha/2}, \text{ or } \frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} > z_{\alpha/2} \\ 0 & \text{otherwise.} \end{cases} \quad (46)$$

Figure 11.8
Rejection Region of the Hypothesis H_0 in (i) (the Shaded Areas), and the Respective Probabilities of Rejection



Since, for any μ ,

$$\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} > z_{\alpha/2} \text{ is equivalent to } \frac{\sqrt{n}(\bar{x} - \mu)}{\sigma} > \frac{\sqrt{n}(\mu_0 - \mu)}{\sigma} + z_{\alpha/2}, \quad (47)$$

and likewise

$$\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} < -z_{\alpha/2} \text{ is equivalent to } \frac{\sqrt{n}(\bar{x} - \mu)}{\sigma} < \frac{\sqrt{n}(\mu_0 - \mu)}{\sigma} - z_{\alpha/2}, \quad (48)$$

it follows that the power of the test is given by:

$$\pi_{\varphi}(\mu) = 1 - \Phi \left[\frac{\sqrt{n}(\mu_0 - \mu)}{\sigma} + z_{\alpha/2} \right] + \Phi \left[\frac{\sqrt{n}(\mu_0 - \mu)}{\sigma} - z_{\alpha/2} \right]. \quad (49)$$

Numerical Example Suppose $n = 36$, $\mu_0 = 10$, and let $\alpha = 0.01$.

DISCUSSION Here $z_{\alpha/2} = z_{0.005} = 2.58$, and, if $\sigma = 4$, the power of the test is:

For $\mu = 12$, $\frac{\sqrt{n}(\mu_0 - \mu)}{\sigma} = \frac{6(10-12)}{4} = -3$, so that

$$\pi_\varphi(12) = 1 - \Phi(-0.42) + \Phi(-5.58) \simeq \Phi(0.42) = 0.662757;$$

and for $\mu = 6$, $\pi_\varphi(6) = 1 - \Phi(8.58) + \Phi(3.42) \simeq 0.999687$.

(ii) $H_0: \mu = \mu_0$, σ unknown. Under H_0 ,

$$L(\hat{\omega}) = (2\pi\hat{\sigma}_\omega^2)^{-n/2} \exp\left[-\frac{1}{2\hat{\sigma}_\omega^2} \sum_{i=1}^n (x_i - \mu_0)^2\right] = (2\pi\hat{\sigma}_\omega^2)^{-n/2} \exp\left(-\frac{n}{2}\right),$$

and

$$L(\hat{\Omega}) = (2\pi\hat{\sigma}_\Omega^2)^{-n/2} \exp\left[-\frac{1}{2\hat{\sigma}_\Omega^2} \sum_{i=1}^n (x_i - \bar{x})^2\right] = (2\pi\hat{\sigma}_\Omega^2)^{-n/2} \exp\left(-\frac{n}{2}\right),$$

since $\hat{\sigma}_\omega^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu_0)^2$ and $\hat{\sigma}_\Omega^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$. Then

$$\lambda = \left(\frac{\hat{\sigma}_\Omega^2}{\hat{\sigma}_\omega^2}\right)^{n/2}, \quad \text{or} \quad \lambda^{2/n} = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{\sum_{i=1}^n (x_i - \mu_0)^2}.$$

Observe that

$$\sum_{i=1}^n (x_i - \mu_0)^2 = \sum_{i=1}^n [(x_i - \bar{x}) + (\bar{x} - \mu_0)]^2 = \sum_{i=1}^n (x_i - \bar{x})^2 + n(\bar{x} - \mu_0)^2,$$

and set $t = \sqrt{n}(\bar{x} - \mu_0) / \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$. Then

$$\begin{aligned} \lambda^{2/n} &= \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{\sum_{i=1}^n (x_i - \bar{x})^2 + n(\bar{x} - \mu_0)^2} = \frac{1}{1 + \frac{n(\bar{x} - \mu_0)^2}{\sum_{i=1}^n (x_i - \bar{x})^2}} \\ &= \frac{1}{1 + \frac{1}{n-1} \times \frac{n(\bar{x} - \mu_0)^2}{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}} = \frac{1}{1 + \frac{1}{n-1} \left[\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}} \right]^2} \\ &= \frac{1}{1 + \frac{t^2}{n-1}}. \end{aligned}$$

Since $g(\lambda) = \lambda^{2/n}$ is a strictly increasing function of λ , the LR test rejects H_0 when $\lambda^{2/n} > C_1$ or $\frac{1}{1 + \frac{t^2}{n-1}} > C_1$ or $1 + \frac{t^2}{n-1} < C_2$ or $t^2 < C_3$ or, finally, $t < -C$ or $t > C$. Under H_0 , the distribution of

$$t(\mathbf{X}) = \frac{\sqrt{n}(\bar{X} - \mu_0)}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}}$$

is t_{n-1} . Since $P_{\mu_0}[t(\mathbf{X}) < -C, \text{ or } t(\mathbf{X}) > C] = \alpha$, it follows that $C = t_{n-1; \alpha/2}$. Therefore, the LR test is:


$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } t < -t_{n-1; \alpha/2}, \text{ or } t > t_{n-1; \alpha/2} \\ 0 & \text{otherwise,} \end{cases} \quad (50)$$

where

$$t = t(\mathbf{x}) = \sqrt{n}(\bar{x} - \mu_0) / \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}. \quad (51)$$

Figure 11.9

Rejection Region of the Hypothesis H_0 in (ii) (the Shaded Areas), and the Respective Probabilities; Here $C = t_{n-1; \alpha/2}$



Numerical Example If $n = 85$ and $\alpha = 0.01$, we find that $t_{n-1; \alpha/2} = t_{84; 0.005} = 2.6356$. Thus, the test rejects H_0 whenever t is < -2.6356 or $t > 2.6356$.

(iii) $H_0: \sigma = \sigma_0$ (or $\sigma^2 = \sigma_0^2$), μ known. Under H_0 ,

$$L(\hat{\omega}) = (2\pi\sigma_0^2)^{-n/2} \exp\left[-\frac{1}{2\sigma_0^2} \sum_{i=1}^n (x_i - \mu)^2\right],$$

and

$$L(\hat{\Omega}) = (2\pi\hat{\sigma}_\Omega^2)^{-n/2} \exp\left[-\frac{1}{2\hat{\sigma}_\Omega^2} \sum_{i=1}^n (x_i - \mu)^2\right] = (2\pi\hat{\sigma}_\Omega^2)^{-n/2} \exp\left(-\frac{n}{2}\right),$$

since $\hat{\sigma}_\Omega^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2$. Therefore

$$\begin{aligned} \lambda &= \left(\frac{\hat{\sigma}_\Omega^2}{\sigma_0^2}\right)^{n/2} e^{n/2} \exp\left[-\frac{1}{2\sigma_0^2} \sum_{i=1}^n (x_i - \mu)^2\right] \\ &= e^{n/2} \left[\frac{1}{n} \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma_0}\right)^2\right]^{n/2} \exp\left[-\frac{1}{2} \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma_0}\right)^2\right] \\ &= e^{n/2} u^{n/2} \exp\left(-\frac{nu}{2}\right), \text{ where } u = \frac{1}{n} \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma_0}\right)^2. \end{aligned}$$


The function $\lambda = \lambda(u)$, $u \geq 0$, has the following properties:

$$\left. \begin{aligned} \lambda(u) \text{ is strictly increasing for } 0 \leq u \leq 1, \\ \lambda(u) \text{ is strictly decreasing for } u > 1, \\ \max\{\lambda(u); 0 \leq u < \infty\} = \lambda(1) = 1, \text{ and} \\ \lambda(u) \rightarrow 0, \text{ as } u \rightarrow \infty, \text{ and, of course,} \\ \lambda(0) = 0. \end{aligned} \right\} \quad (52)$$

On the basis of these observations, the picture of $\lambda(u)$ is as in Figure 11.10.

Figure 11.10

**Graphical
Determination of the
Rejection Region**



Therefore $\lambda(u) \leq \lambda_0$ if and only if $u \leq C_1$, or $u \geq C_2$, where C_1 and C_2 are determined by the requirement that


$$P_{\mu_0}(U \leq C_1, \text{ or } U \geq C_2) = \alpha, \quad U = \frac{1}{n} \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma_0} \right)^2.$$

However, under H_0 , $\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma_0} \right)^2 \sim \chi_n^2$, so that $U = \frac{X}{n}$ with $X \sim \chi_n^2$. Then $P_{\mu_0}(U \leq C_1, \text{ or } U \geq C_2) = P_{\mu_0}(X \leq nC_1, \text{ or } X \geq nC_2) = \alpha$, and, for convenience, we may take the two-tail probabilities equal to $\frac{\alpha}{2}$. Then $nC_1 = \chi_{n;1-\alpha/2}^2$, $nC_2 = \chi_{n;\alpha/2}^2$. Summarizing what we have done so far, we have:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma_0} \right)^2 \leq \chi_{n;1-\alpha/2}^2, \text{ or } \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma_0} \right)^2 > \chi_{n;\alpha/2}^2 \\ 0 & \text{otherwise.} \end{cases} \quad (53)$$

Figure 11.11

**Rejection Region of
the Hypothesis H_0 in
(iii) (the Shaded
Areas), and the
Respective
Probabilities; Here
 $C_1 = \chi_{n;1-\frac{\alpha}{2}}^2$, $C_2 =$
 $\chi_{n;\frac{\alpha}{2}}^2$**



Numerical Example For $n = 40$ and $\alpha = 0.01$, we find $\chi_{n;1-\alpha/2}^2 = \chi_{40;0.995}^2 = 20.707$ and $\chi_{n;\alpha/2}^2 = \chi_{40;0.005}^2 = 66.766$. Therefore the test rejects H_0 whenever $\sum_{i=1}^{40} \left(\frac{x_i - \mu}{\sigma_0} \right)^2$ is either ≤ 20.707 or ≥ 66.766 .

(iv) $H_0: \sigma = \sigma_0$ (or $\sigma^2 = \sigma_0^2$), μ unknown. Under ω ,

$$L(\hat{\omega}) = (2\pi\sigma_0^2)^{-n/2} \exp \left[-\frac{1}{2\sigma_0^2} \sum_{i=1}^n (x_i - \bar{x})^2 \right], \quad \text{since } \hat{\mu}_\omega = \bar{x},$$

and

$$L(\hat{\Omega}) = (2\pi\hat{\sigma}_{\Omega}^2)^{-n/2} \exp\left[-\frac{1}{2\hat{\sigma}_{\Omega}^2} \sum_{i=1}^n (x_i - \bar{x})^2\right] = (2\pi\hat{\sigma}_{\Omega}^2)^{-n/2} \exp\left(-\frac{n}{2}\right),$$

since $\hat{\sigma}_{\Omega}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$. Therefore

$$\lambda = \left(\frac{\hat{\sigma}_{\Omega}^2}{\sigma_0^2}\right)^{n/2} e^{n/2} \exp\left[-\frac{1}{2\sigma_0^2} \sum_{i=1}^n (x_i - \bar{x})^2\right],$$

and then proceed exactly as in the previous case with $u = \frac{1}{n} \sum_{i=1}^n \left(\frac{x_i - \bar{x}}{\sigma_0}\right)^2$, in order to arrive at the following modified test; namely,


$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n \left(\frac{x_i - \bar{x}}{\sigma_0}\right)^2 \leq \chi_{n-1; 1-\alpha/2}^2, \text{ or } \sum_{i=1}^n \left(\frac{x_i - \bar{x}}{\sigma_0}\right)^2 \geq \chi_{n-1; \alpha/2}^2 \\ 0 & \text{otherwise.} \end{cases} \quad (54)$$

Figure 11.12

Rejection Region of the Hypothesis H_0 in (iv) (Shaded Areas), and the Respective Probabilities; Here

$$C_1 = \chi_{n-1; 1-\frac{\alpha}{2}}^2,$$

$$C_2 = \chi_{n-1; \frac{\alpha}{2}}^2$$



Numerical Example With the values of n and α as in the previous case ($n = 40$, $\alpha = 0.01$), we find $\chi_{n-1; 1-\alpha/2}^2 = \chi_{39; 0.995}^2 = 19.996$ and $\chi_{n-1; \alpha/2}^2 = \chi_{39; 0.005}^2 = 65.476$, so that the test rejects H_0 whenever $\sum_{i=1}^n \left(\frac{x_i - \bar{x}}{\sigma_0}\right)^2$ is ≤ 19.996 or ≥ 65.476 .

11.4.2 Comparing the Parameters of Two Normal Populations

Here, we have two independent random samples $X_1, \dots, X_m \sim N(\mu_1, \sigma_1^2)$ and $Y_1, \dots, Y_n \sim N(\mu_2, \sigma_2^2)$ with all parameters unknown. The two populations are compared, first, by way of their means, and second, through their variances. When comparing these populations through their means, it is necessary from a mathematical viewpoint (i.e., in order to be able to derive an exact distribution for the test statistic) that the variances, although unknown, be equal.

(i) $H_0: \mu_1 = \mu_2 = \mu$, say, unknown, $\sigma_1 = \sigma_2 = \sigma$, say, unknown.

The (joint) likelihood function of the X_i 's and the Y_j 's here is

$$(2\pi\sigma^2)^{-(m+n)/2} \exp\left\{-\frac{1}{2\sigma^2} \left[\sum_{i=1}^m (x_i - \mu_1)^2 + \sum_{j=1}^n (y_j - \mu_2)^2 \right]\right\}. \quad (55)$$

Maximizing (55) with respect to μ_1, μ_2 , and σ^2 , we find for their MLE's:

$$\hat{\mu}_1 = \bar{x}, \quad \hat{\mu}_2 = \bar{y}, \quad \hat{\sigma}_\Omega^2 = \frac{1}{m+n} \left[\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2 \right]. \quad (56)$$

Hence

$$L(\hat{\Omega}) = (2\pi \hat{\sigma}_\Omega^2)^{-\frac{m+n}{2}} \exp\left(-\frac{m+n}{2}\right). \quad (57)$$

Next, under H_0 , the (joint) likelihood function becomes

$$(2\pi\sigma^2)^{-(m+n)/2} \exp\left\{-\frac{1}{2\sigma^2} \left[\sum_{i=1}^m (x_i - \mu)^2 + \sum_{j=1}^n (y_j - \mu)^2 \right]\right\}, \quad (58)$$

from the maximization (with respect to μ and σ^2) of which we obtain the MLE's

$$\hat{\mu}_\omega = \frac{1}{m+n} \left(\sum_{i=1}^m x_i + \sum_{j=1}^n y_j \right) = \frac{m\bar{x} + n\bar{y}}{m+n}, \quad (59)$$

$$\hat{\sigma}_\omega^2 = \frac{1}{m+n} \left[\sum_{i=1}^m (x_i - \hat{\mu}_\omega)^2 + \sum_{j=1}^n (y_j - \hat{\mu}_\omega)^2 \right].$$

Inserting these expressions in (58), we then have

$$L(\hat{\omega}) = (2\pi \hat{\sigma}_\omega^2)^{-\frac{m+n}{2}} \exp\left(-\frac{m+n}{2}\right). \quad (60)$$

Thus, the likelihood function becomes, on account of (57) and (60),

$$\lambda = \left(\frac{\hat{\sigma}_\Omega^2}{\hat{\sigma}_\omega^2} \right)^{\frac{m+n}{2}}, \quad \text{or} \quad \lambda^{2/(m+n)} = \frac{\hat{\sigma}_\Omega^2}{\hat{\sigma}_\omega^2}. \quad (61)$$

Next,

$$\begin{aligned} \sum_{i=1}^m (x_i - \hat{\mu}_\omega)^2 &= \sum_{i=1}^m [(x_i - \bar{x}) + (\bar{x} - \hat{\mu}_\omega)]^2 = \sum_{i=1}^m (x_i - \bar{x})^2 + m(\bar{x} - \hat{\mu}_\omega)^2 \\ &= \sum_{i=1}^m (x_i - \bar{x})^2 + m \left(\bar{x} - \frac{m\bar{x} + n\bar{y}}{m+n} \right)^2 = \sum_{i=1}^m (x_i - \bar{x})^2 + \frac{mn^2(\bar{x} - \bar{y})^2}{(m+n)^2}, \end{aligned}$$

and likewise,

$$\sum_{j=1}^n (y_j - \hat{\mu}_\omega)^2 = \sum_{j=1}^n (y_j - \bar{y})^2 + \frac{m^2n(\bar{x} - \bar{y})^2}{(m+n)^2}.$$

Then, by means of (56) and (59), $\hat{\sigma}_\omega^2$ is written as follows:

$$\begin{aligned} \hat{\sigma}_\omega^2 &= \frac{1}{m+n} \left[\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2 \right] + \frac{mn^2(\bar{x} - \bar{y})^2 + m^2n(\bar{x} - \bar{y})^2}{(m+n)^3} \\ &= \hat{\sigma}_\Omega^2 + \frac{mn(\bar{x} - \bar{y})^2(m+n)}{(m+n)^3} = \hat{\sigma}_\Omega^2 + \frac{mn(\bar{x} - \bar{y})^2}{(m+n)^2}. \end{aligned} \quad (62)$$

Therefore (61) yields, by way of (62) and (56),

$$\begin{aligned}
 \lambda^{2/(m+n)} &= \frac{\hat{\sigma}_\Omega^2}{\hat{\sigma}_\Omega^2 + \frac{mn(\bar{x}-\bar{y})^2}{(m+n)^2}} = \frac{1}{1 + \frac{mn(\bar{x}-\bar{y})^2}{(m+n)^2} \times \frac{1}{\hat{\sigma}_\Omega^2}} \\
 &= \left[1 + \frac{mn(\bar{x}-\bar{y})^2}{(m+n)^2} \Big/ \hat{\sigma}_\Omega^2 \right]^{-1} \\
 &= \left\{ 1 + \frac{mn}{m+n} (\bar{x}-\bar{y})^2 \Big/ \left[\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2 \right] \right\}^{-1} \\
 &= \left\{ 1 + \frac{mn}{m+n} (\bar{x}-\bar{y})^2 / (m+n-2) \times \frac{1}{m+n-2} \right. \\
 &\quad \left. \times \left[\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2 \right] \right\}^{-1} \\
 &= \left[1 + \left\{ \frac{\sqrt{\frac{mn}{m+n}} (\bar{x}-\bar{y})}{\sqrt{\frac{1}{m+n-2} \left[\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2 \right]}} \right\}^2 / (m+n-2) \right]^{-1} \\
 &= \left(1 + \frac{t^2}{m+n-2} \right)^{-1}, \text{ where} \\
 t &= t(\mathbf{x}, \mathbf{y}) = \sqrt{\frac{mn}{m+n}} (\bar{x}-\bar{y}) \Big/ \sqrt{\frac{1}{m+n-2} \left[\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2 \right]}. \tag{63}
 \end{aligned}$$

So, $\lambda^{2/(m+n)} = (1 + \frac{t^2}{m+n-2})^{-1}$ and hence $\lambda = (1 + \frac{t^2}{m+n-2})^{-\frac{m+n}{2}}$. Since λ is strictly decreasing in t^2 , the LR test rejects H_0 whenever $t^2 > C_0$ or, equivalently, $t \leq -C$ or $t \geq C$. The constant C is to be determined by

$$P_{H_0}[t(\mathbf{X}, \mathbf{Y}) \leq -C, \text{ or } t(\mathbf{X}, \mathbf{Y}) \geq C] = \alpha, \tag{64}$$

where $t(\mathbf{X}, \mathbf{Y})$ is taken from (63) with the x_i 's and the y_j 's being replaced by the r.v.'s X_i 's and Y_j 's. However, under H_0 ,

$$t(\mathbf{X}, \mathbf{Y}) \sim t_{m+n-2}, \tag{65}$$


so that (64) yields $C = t_{m+n-2; \alpha/2}$. In conclusion, then,

$$\varphi(\mathbf{x}, \mathbf{y}) = \begin{cases} 1 & \text{if } t(\mathbf{x}, \mathbf{y}) < -t_{m+n-2; \alpha/2}, \text{ or } t(\mathbf{x}, \mathbf{y}) > t_{m+n-2; \alpha/2} \\ 0 & \text{otherwise,} \end{cases} \tag{66}$$

where $t(\mathbf{x}, \mathbf{y})$ is given by (63).

Figure 11.13

Rejection Region of the Hypothesis H_0 in (i) (Shaded Areas), and the Respective Probabilities; Here $C = t_{m+n-2; \frac{\alpha}{2}}$



Numerical Example For $m = 40, n = 50$, and $\alpha = 0.01$, we get $t_{m+n-2; \alpha/2} = t_{88; 0.005} = 2.6329$, and hence the hypothesis about equality of means is rejected whenever $|t(\mathbf{x}, \mathbf{y})| \geq 2.6329$.

(ii) $H_0: \sigma_1 = \sigma_2 = \sigma$, say (or $\sigma_1^2 = \sigma_2^2 = \sigma^2$, say), μ_1, μ_2 unknown. The (joint) likelihood function of the X_i 's and Y_j 's is here

$$(2\pi)^{-\frac{m+n}{2}} (\sigma_1^2)^{-m/2} (\sigma_2^2)^{-n/2} \exp \left[-\frac{1}{2\sigma_1^2} \sum_{i=1}^m (x_i - \mu_1)^2 - \frac{1}{2\sigma_2^2} \sum_{j=1}^n (y_j - \mu_2)^2 \right]. \quad (67)$$

Maximizing (67) with respect to all four parameters, we find the following MLE's:

$$\hat{\mu}_{1,\Omega} = \bar{x}, \quad \hat{\mu}_{2,\Omega} = \bar{y}, \quad \hat{\sigma}_{1,\Omega}^2 = \frac{1}{m} \sum_{i=1}^m (x_i - \bar{x})^2, \quad \hat{\sigma}_{2,\Omega}^2 = \frac{1}{n} \sum_{j=1}^n (y_j - \bar{y})^2. \quad (68)$$

Then

$$L(\hat{\Omega}) = (2\pi)^{-\frac{m+n}{2}} (\hat{\sigma}_{1,\Omega}^2)^{-m/2} (\hat{\sigma}_{2,\Omega}^2)^{-n/2} \exp \left(-\frac{m+n}{2} \right). \quad (69)$$

Under H_0 , the likelihood function has the form (55), and the MLE's are already available and given by (56). That is,

$$\hat{\mu}_{1,\omega} = \bar{x}, \quad \hat{\mu}_{2,\omega} = \bar{y}, \quad \hat{\sigma}_{1,\omega}^2 = \frac{1}{m+n} \left[\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2 \right]. \quad (70)$$

Therefore,

$$L(\hat{\omega}) = (2\pi)^{-\frac{m+n}{2}} (\hat{\sigma}_{1,\omega}^2)^{-\frac{m+n}{2}} \exp \left(-\frac{m+n}{2} \right). \quad (71)$$

For simplicity, set $\sum_{i=1}^m (x_i - \bar{x})^2 = a$, $\sum_{j=1}^n (y_j - \bar{y})^2 = b$. Then the LR is, by means of (68) through (71),

$$\begin{aligned} \lambda &= \frac{(\hat{\sigma}_{1,\Omega}^2)^{m/2} (\hat{\sigma}_{2,\Omega}^2)^{n/2}}{(\hat{\sigma}_{1,\omega}^2)^{(m+n)/2}} = \frac{m^{-m/2} n^{-n/2} a^{m/2} b^{n/2}}{(m+n)^{-(m+n)/2} (a+b)^{(m+n)/2}} \\ &= \frac{(m+n)^{(m+n)/2}}{m^{m/2} n^{n/2}} \times \frac{(a/b)^{m/2}}{\left(1 + \frac{a}{b}\right)^{(m+n)/2}} \quad (\text{dividing by } b^{(m+n)/2}) \\ &= \frac{(m+n)^{(m+n)/2}}{m^{m/2} n^{n/2}} \times \frac{\left[\left(\frac{m-1}{n-1}\right) \left(\frac{a}{m-1} / \frac{b}{n-1}\right)\right]^{m/2}}{\left[1 + \left(\frac{m-1}{n-1}\right) \left(\frac{a}{m-1} / \frac{b}{n-1}\right)\right]^{(m+n)/2}} \\ &= \frac{(m+n)^{(m+n)/2}}{m^{m/2} n^{n/2}} \times \frac{\left(\frac{m-1}{n-1} u\right)^{m/2}}{\left(1 + \frac{m-1}{n-1} u\right)^{(m+n)/2}}, \quad \text{where } u = \frac{a}{m-1} / \frac{b}{n-1}. \end{aligned}$$

So

$$\lambda = \lambda(u) = \frac{(m+n)^{(m+n)/2}}{m^{m/2} n^{n/2}} \times \frac{\left(\frac{m-1}{n-1} u\right)^{m/2}}{\left(1 + \frac{m-1}{n-1} u\right)^{(m+n)/2}}, \quad u \geq 0. \quad (72)$$


The function $\lambda(u)$ has the following properties:

$$\left. \begin{aligned} \lambda(0) &= 0 \text{ and } \lambda(u) \rightarrow 0, \text{ as } u \rightarrow \infty, \\ \frac{d}{du} \lambda(u) &= 0 \text{ for } u = u_0 = \frac{m(n-1)}{n(m-1)}, \quad \frac{d}{du} \lambda(u) > 0 \text{ for } u < u_0, \text{ and} \\ \frac{d}{du} \lambda(u) &< 0 \text{ for } u > u_0, \text{ so that } \lambda(u) \text{ is} \\ &\text{maximized for } u = u_0, \text{ and } \lambda(u_0) = 1. \end{aligned} \right\} \quad (73)$$

On the basis of these properties, the picture of $\lambda(u)$ is as in Figure 11.14.

Figure 11.14

**The Graph of the
Function $\lambda = \lambda(u)$
Given in Relation
(72)**




Therefore $\lambda(u) \leq \lambda_0$, if and only if $u \leq C_1$ or $u > C_2$, where C_1 and C_2 are determined by the requirement that

$$P_{H_0}(U \leq C_1, \text{ or } U \geq C_2) = \alpha, \quad \text{where } U = \frac{\sum_{i=1}^m (X_i - \bar{X})^2 / (m-1)}{\sum_{j=1}^n (Y_j - \bar{Y})^2 / (n-1)}. \quad (74)$$

Figure 11.15

Rejection Region of the Hypothesis H_0 in (ii) (Shaded Areas), and the Respective Probabilities; Here $C_1 = F_{m-1, n-1; 1-\frac{\alpha}{2}}$, $C_2 = F_{m-1, n-1; \frac{\alpha}{2}}$



Under H_0 , $U \sim F_{m-1, n-1}$, and this allows the determination of C_1, C_2 . For simplicity, we may split the probability α equally among the two tails, in which case

$$C_1 = F_{m-1, n-1; 1-\alpha/2}, \quad C_2 = F_{m-1, n-1; \alpha/2}.$$

To summarize then, the LR test is as follows:

$$\varphi(\mathbf{x}, \mathbf{y}) = \begin{cases} 1 & \text{if } u(\mathbf{x}, \mathbf{y}) \leq F_{m-1, n-1; 1-\alpha/2}, \text{ or } u(\mathbf{x}, \mathbf{y}) \geq F_{m-1, n-1; \alpha/2} \\ 0 & \text{otherwise,} \end{cases} \quad (75)$$

where

$$u(\mathbf{x}, \mathbf{y}) = \frac{\sum_{i=1}^m (x_i - \bar{x})^2 / (m-1)}{\sum_{j=1}^n (y_j - \bar{y})^2 / (n-1)}. \quad (76)$$

Numerical Example Let $m = 13$, $n = 19$, and take $\alpha = 0.05$.

DISCUSSION If $X \sim F_{12, 18}$, then we get from the F -tables: $P(X > F_{12, 18; 0.025}) = 0.025$, or $P(X \leq F_{12, 18; 0.025}) = 0.975$, and hence $F_{12, 18; 0.025} = 2.7689$. Also, $P(X \leq F_{12, 18; 0.975}) = 0.025$, or $P(X > F_{12, 18; 0.975}) = 0.975$, or $P(\frac{1}{X} < \frac{1}{F_{12, 18; 0.975}}) = 0.975$. But then $\frac{1}{X} \sim F_{18, 12}$, and therefore $\frac{1}{F_{12, 18; 0.975}} = 3.1076$, and hence $F_{12, 18; 0.975} \simeq 0.3218$. Thus, the hypothesis H_0 is rejected whenever $u(\mathbf{x}, \mathbf{y}) \leq 0.3218$, or $u(\mathbf{X}, \mathbf{Y}) \geq 2.7689$, and it is accepted otherwise.

Exercises

4.1 A coin, with probability θ of falling heads, is tossed independently 100 times and 60 heads are observed. At level of significance $\alpha = 0.1$:

- (i) Use the LR test in order to test the hypothesis $H_0: \theta = 1/2$ (against the alternative $H_A: \theta \neq 1/2$).

- (ii) Employ the appropriate approximation (see relation (10) in Chapter 8) to determine the cutoff point.
- 4.2** Let X_1, X_2, X_3 be independent r.v.'s distributed as $B(1, \theta)$, $\theta \in \Omega = (0, 1)$, and let $t = x_1 + x_2 + x_3$, where the x_i 's are the observed values of the X_i 's.
- (i) Derive the LR test λ for testing the hypothesis $H_0: \theta = 0.25$ (against the alternative $H_A: \theta \neq 0.25$) at level of significance $\alpha = 0.02$.
- (ii) Calculate the distribution of $\lambda(T)$ and carry out the test, where $T = X_1 + X_2 + X_3$.
- 4.3** (i) In reference to Example 15 in Chapter 1, the appropriate model to be employed is the Normal distribution $N(\mu, \sigma^2)$ (with $\mu > 0$, of course).
- (ii) If μ_0 is the stipulated average growth, then this will be checked by testing the hypothesis $H_0: \mu = \mu_0$ (against the alternative $H_A: \mu \neq \mu_0$) at level of significance α .
- (iii) On the basis of a random sample of size n , use the likelihood ratio test to test H_0 when $n = 25$, $\mu_0 = 6$ inch, and $\alpha = 0.05$.
- 4.4** (i) In reference to Example 17 in Chapter 1, an appropriate model would be the following. Let X_i and Y_i be the blood pressure of the i th individual before and after the use of the pill, and set $Z_i = Y_i - X_i$, $i = 1, \dots, n$. Furthermore, it is reasonable to assume that the X_i 's and the Y_i 's are independent and Normally distributed, so that the Z_i 's are independently distributed as $N(\mu, \sigma^2)$.
- (ii) With μ denoting the difference of blood pressure after the usage of the pill and before it, the claim is that $\mu < 0$. This claim is checked by testing the hypothesis $H_0: \mu = 0$ (against the alternative $H_A: \mu \neq 0$, with the only viable part of it here being $\mu > 0$) at level of significance α , by using the likelihood ratio test.
- (iii) Carry out the test if $n = 90$ and $\alpha = 0.05$.
- 4.5** In reference to Example 25 in Chapter 1:
- (i) For $i = 1, \dots, 15$, let X_i and Y_i be the heights of the cross-fertilized plants and self-fertilized plants, respectively. It is reasonable to assume that the X_i 's and the Y_i 's are independent random samples with respective distributions $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$ (the estimates of σ_1^2 and σ_2^2 do not justify the possible assumption of a common variance). Setting $Z_i = X_i - Y_i$, we have that the Z_i 's are independent and distributed as $N(\mu, \sigma^2)$, where $\mu = \mu_1 - \mu_2$, $\sigma^2 = \sigma_1^2 + \sigma_2^2$.
- (ii) The claim is that $\mu > 0$, and is to be checked by testing the hypothesis $H_0: \mu = 0$ (against the alternative $H_A: \mu \neq 0$, with the only viable part of it being that $\mu > 0$) at level of significance α , by using the likelihood ratio test.
- (iii) Carry out the test when $\alpha = 0.05$ and $\alpha = 0.10$.
- 4.6** The diameters of certain cylindrical items produced by a machine are r.v.'s distributed as $N(\mu, 0.01)$. A sample of size 16 is taken and it is found

that $\bar{x} = 2.48$ inches. If the desired value for μ is 2.5 inches, formulate the appropriate testing hypothesis problem and carry out the test if $\alpha = 0.05$.

4.7 A manufacturer claims that packages of certain goods contain 18 ounces. In order to check his claim, 100 packages are chosen at random from a large lot and it is found that $\sum_{i=1}^{100} x_i = 1,752$ and $\sum_{i=1}^{100} x_i^2 = 31,157$. Assume that the observations are Normally distributed, and formulate the manufacturer's claim as a testing hypothesis problem. Carry out the test at level of significance $\alpha = 0.01$.

4.8 The breaking powers of certain steel bars produced by processes A and B are r.v.'s distributed as Normal with possibly different means but the same variance. A random sample of size 25 is taken from bars produced by each one of the processes, and it is found that $\bar{x} = 60$, $s_x = 6$, $\bar{y} = 65$, $s_y = 7$. Test whether there is a difference between the two processes at the level of significance $\alpha = 0.05$.

4.9 (i) Let X_i , $i = 1, \dots, 9$ and Y_j , $j = 1, \dots, 10$ be independent r.v.'s from the distributions $N(\mu_1, \sigma_1^2)$ and $N(\mu_2, \sigma_2^2)$, respectively. Suppose that the observed values of the sample s.d.'s are $s_x = 2$, $s_y = 3$. At level of significance $\alpha = 0.05$, test the hypothesis $H_0: \sigma_1 = \sigma_2$ (against the alternative $H_A: \sigma_1 \neq \sigma_2$.)

(ii) Find an expression for the computation of the power of the test for $\sigma_1 = 2$ and $\sigma_2 = 3$.

4.10 Refer to Exercise 3.12, and suppose that the variances σ_1^2 and σ_2^2 are unknown. Then test the hypothesis $H_0: \sigma_1 = \sigma_2$ (against the alternative $H_A: \sigma_1 \neq \sigma_2$) at level of significance $\alpha = 0.05$.

4.11 The independent random samples X_i and Y_i , $i = 1, \dots, 5$ represent resistance measurements taken on two test pieces, and the observed values (in ohms) are as follows:

$$\begin{aligned} x_1 &= 0.118, & x_2 &= 0.125, & x_3 &= 0.121, & x_4 &= 0.117, & x_5 &= 0.120, \\ y_1 &= 0.114, & y_2 &= 0.115, & y_3 &= 0.119, & y_4 &= 0.120, & y_5 &= 0.110. \end{aligned}$$

Assume that the X_i 's and the Y_i 's are Normally distributed, and test the hypothesis $H_0: \sigma_1 = \sigma_2$ (against the alternative $H_A: \sigma_1 \neq \sigma_2$) at level of significance $\alpha = 0.05$.

4.12 Refer to Exercise 4.11, and assume now that $\sigma_1 = \sigma_2 = \sigma$, say, unknown (which is supported by the fact that the hypothesis $H_0: \sigma_1 = \sigma_2$ was not rejected). Then test the hypothesis $H_0: \mu_1 = \mu_2$ (against the alternative $H_A: \mu_1 \neq \mu_2$) at level of significance $\alpha = 0.05$.

4.13 Consider the independent random samples X_1, \dots, X_m and Y_1, \dots, Y_n from the respective distributions $N(\mu_1, \sigma^2)$ and $N(\mu_2, \sigma^2)$ where σ is known, and suppose we are interested in testing the hypothesis $H_0: \mu_1 = \mu_2 = \mu$, say, unknown (against the alternative $H_A: \mu_1 \neq \mu_2$) at level of

significance α , by means of the likelihood ratio test. Set $\mathbf{x} = (x_1, \dots, x_m)$ and $\mathbf{y} = (y_1, \dots, y_n)$ for the observed values of the X_i 's and the Y_i 's.

- (i) Form the joint likelihood function $L(\mu_1, \mu_2 | \mathbf{x}, \mathbf{y})$ of the X_i 's and the Y_i 's, as well as the likelihood function $L(\mu | \mathbf{x}, \mathbf{y})$.
- (ii) From part (i), conclude immediately that the MLE's of μ_1 and μ_2 are $\hat{\mu}_1 = \bar{x}$ and $\hat{\mu}_2 = \bar{y}$. Also, show that the MLE of μ is given by $\hat{\mu}_\omega = \frac{m\bar{x} + n\bar{y}}{m+n}$.
- (iii) Show that $-2 \log \lambda = mn(\bar{x} - \bar{y})^2 / \sigma^2(m+n)$, where $\lambda = L(\hat{\omega}) / L(\hat{\Omega})$.
- (iv) From part (iii), conclude that the likelihood ratio test $-2 \log \lambda > C_0$ is equivalent to $|\bar{x} - \bar{y}| > C (= \sigma \sqrt{(m+n)C_0 / mn})$.
- (v) Show that $C = z_{\alpha/2} \sigma \sqrt{\frac{1}{m} + \frac{1}{n}}$.
- (vi) For any μ_1 and μ_2 , show that the power of the test depends on μ_1 and μ_2 through their difference $\mu_1 - \mu_2 = \Delta$, say, and is given by the formula:

$$\pi(\Delta) = 2 - \Phi\left(\frac{C - \Delta}{\sigma \sqrt{\frac{1}{m} + \frac{1}{n}}}\right) - \Phi\left(\frac{C + \Delta}{\sigma \sqrt{\frac{1}{m} + \frac{1}{n}}}\right).$$

- (vii) Determine the cutoff point when $m = 10$, $n = 15$, $\sigma = 1$, and $\alpha = 0.05$.
- (viii) Determine the power of the test when $\Delta = 1$ and $\Delta = 2$.

4.14 In reference to Example 15, verify the results:

$$\int_0^{C_1} \theta_0^2 t e^{-\theta_0 t} dt = 1 - e^{-\theta_0 C_1} - \theta_0 C_1 e^{-\theta_0 C_1},$$

$$\int_{C_2}^{\infty} \theta_0^2 t e^{-\theta_0 t} dt = e^{-\theta_0 C_2} + \theta_0 C_2 e^{-\theta_0 C_2}.$$

4.15 Verify expression (49) for the power of the test.

4.16 Verify the assertions made in expressions (52) about the function $\lambda = \lambda(u)$, $u \geq 0$.

4.17 Verify the assertion made in relation (56) that $\hat{\mu}_1$, $\hat{\mu}_2$, and $\hat{\sigma}_\Omega^2$ are the MLE's of μ_1 , μ_2 , and σ^2 , respectively.

4.18 Show that the expressions in relation (59) are, indeed, the MLE's of $(\mu_1 = \mu_2 = \mu)$ and σ^2 , respectively.

4.19 Show that $\lambda = \lambda(t^2) = (1 + \frac{t^2}{m+n-2})^{-\frac{m+n}{2}}$ is, indeed, strictly increasing in t^2 as asserted right after relation (63).

4.20 Justify the statement made in relation (65) that $t(\mathbf{X}, \mathbf{Y}) \sim t_{m+n-2}$.

4.21 Show that the expressions in relation (68) are, indeed, the MLE's of μ_1 , μ_2 , σ_1^2 , and σ_2^2 , respectively.

4.22 Verify the assertions made in expression (73) about the function $\lambda = \lambda(u)$, $u \geq 0$.

4.23 Refer to the Bivariate Normal distribution discussed in Chapter 4, Section 5, whose p.d.f. is given by:

$$f_{X,Y}(x, y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} e^{-q/2}, \quad x, y \in \Re,$$

where $q = \frac{1}{1-\rho^2} \left[\left(\frac{x-\mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x-\mu_1}{\sigma_1} \right) \left(\frac{y-\mu_2}{\sigma_2} \right) + \left(\frac{y-\mu_2}{\sigma_2} \right)^2 \right]$, $\mu_1, \mu_2 \in \Re$, $\sigma_1^2, \sigma_2^2 > 0$, and $-1 \leq \rho \leq 1$ are the parameters of the distribution. Also, recall that independence between X and Y is equivalent to their being uncorrelated; i.e., $\rho = 0$. In this exercise, a test is derived for testing the hypothesis $H_0: \rho = 0$ (against the alternative $H_A: \rho \neq 0$, the X and Y are not independent). The test statistic is based on the likelihood ratio statistic.

(i) On the basis of a random sample of size n from a Bivariate Normal distribution, (X_i, Y_i) , $i = 1, \dots, n$, the MLE's of the parameters involved are given by:

$$\hat{\mu}_1 = \bar{x}, \quad \hat{\mu}_2 = \bar{y}, \quad \hat{\sigma}_1^2 = S_x, \quad \hat{\sigma}_2^2 = S_y, \quad \hat{\rho} = S_{xy}/\sqrt{S_x S_y},$$

where $S_x = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$, $S_y = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2$, $S_{xy} = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})$, and the x_i 's and y_i 's are the observed values of the X_i 's and Y_i 's. (See Exercise 1.14 (vii) in Chapter 9.)

(ii) Under the hypothesis $H_0: \rho = 0$, the MLE's of μ_1, μ_2, σ_1^2 , and σ_2^2 are the same as in part (i).

Hint: It follows immediately, because the joint p.d.f. of the pairs factorizes to the joint p.d.f. of the X_i 's times the joint p.d.f. of the Y_i 's.

(iii) When replacing the parameters by their MLE's, the likelihood function, call it $L(\mathbf{x}, \mathbf{y})$, is given by:

$$L(\mathbf{x}, \mathbf{y}) = [2\pi (S_x S_y - S_{xy}^2)]^{-\frac{n}{2}} e^{-n},$$

where $\mathbf{x} = (x_1, \dots, x_n)$, $\mathbf{y} = (y_1, \dots, y_n)$.

(iv) Under the hypothesis $H_0(\rho = 0)$, when the parameters are replaced by their MLE's, the likelihood function, call it $L_0(\mathbf{x}, \mathbf{y})$, is given by:

$$L_0(\mathbf{x}, \mathbf{y}) = (2\pi S_x S_y)^{-\frac{n}{2}} e^{-n}.$$

(v) From parts (iii) and (iv), it follows that the likelihood ratio statistic λ is given by:

$$\lambda = (1 - \hat{\rho}^2)^{n/2}, \quad \hat{\rho} = S_{xy}/\sqrt{S_x S_y}.$$

4.24 (i) By differentiation, show that the function $f(r) = (1 - r)^{n/2}$ is decreasing in r . Therefore, in reference to Exercise 4.23(v), $\lambda < \lambda_0$ is equivalent to $\hat{\rho}^2 > C_1$, some constant C_1 (actually, $C_1 = 1 - \lambda_0^{2/n}$); equivalently, $\hat{\rho} < -C_2$ or $\hat{\rho} > C_2$ ($C_2 = \sqrt{C_1}$).

(ii) Since the LR test rejects the hypothesis H_0 when $\lambda < \lambda_0$, part (i) states that the LR test is equivalent to rejecting H_0 wherever $\hat{\rho} < -C_2$ or $\hat{\rho} > C_2$.

- (iii) In $\hat{\rho}$, replace the x_i 's and the y_i 's by the respective r.v.'s X_i and Y_i , and set R for the resulting r.v. Then, in part (ii), carrying out the test based on $\hat{\rho}$, requires knowledge of the cutoff point C_2 , which in turn, presupposes knowledge of the distribution of R (under H_0). Although the distribution of R can be determined (see, e.g., Corollary to Theorem 7, page 474, in the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G. G. Roussas), it is not of any of the known forms, and hence no tables can be used.
- (iv) Set $W = W(R) = \frac{\sqrt{n-2}R}{\sqrt{1-R^2}}$, and show that W is an increasing function of R by showing that $\frac{d}{dr}W(r)$ is positive.
- (v) By parts (ii) and (iv), it follows that the likelihood ratio test is equivalent to rejecting H_0 whenever $W(r) < -C$ or $W(r) > C$, where C is determined by the requirement that $P_{H_0}[W(R) < C \text{ or } W(R) > C] = \alpha$ (the given level of significance).
- (vi) Under H_0 , it can be shown (see, e.g., pages 472–474, in the book cited in part (iii) above) that $W(R)$ has the t_{n-2} distribution. It follows that $C = t_{n-2; \frac{\alpha}{2}}$.

To summarize then, for testing $H_0: \rho = 0$ at level of significance α , reject H_0 whenever $W(r) < -t_{n-2; \frac{\alpha}{2}}$ or $W(r) > t_{n-2; \frac{\alpha}{2}}$, where $W(r) = \frac{\sqrt{n-2}r}{\sqrt{1-r^2}}$, $r = \hat{\rho} = S_{xy}/\sqrt{S_x S_y}$; this test is equivalent to the likelihood ratio test.

Chapter 12

More About Testing Hypotheses

In this chapter, a few more topics are discussed on testing hypotheses problems. More specifically, LR tests are presented for the Multinomial distribution with further applications to contingency tables. A brief section is devoted to the so-called (Chi-Square) goodness-of-fit tests, and another also brief section discusses the decision-theoretic approach to testing hypotheses. The chapter is concluded with a result connecting testing hypotheses and construction of confidence regions.

12.1 Likelihood Ratio Tests in the Multinomial Case and Contingency Tables

It was stated in Section 3 of Chapter 8 that the statistic $-2 \log \lambda$ is distributed approximately as χ_f^2 with certain degrees of freedom f , provided some regularity conditions are met. In this section, this result is stated in a more formal way, although the required conditions will not be spelled out.

THEOREM 1

On the basis of the random sample X_1, \dots, X_n from the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}^r$, $r \geq 1$, we wish to test the hypothesis $H_0: \theta \in \omega \subset \Omega$ at level of significance α and on the basis of the Likelihood Ratio statistic $\lambda = \lambda(X_1, \dots, X_n)$. Then, provided certain conditions are met, it holds that:

$$-2 \log \lambda \simeq \chi_{r-m}^2, \text{ for all sufficiently large } n \text{ and } \theta \in \omega;$$

more formally,

$$P_\theta(-2 \log \lambda \leq x) \rightarrow G(x), \quad x \geq 0, \quad \text{as } n \rightarrow \infty, \quad (1)$$

where G is the d.f. of the χ_{r-m}^2 distribution; r is the dimensionality of Ω , m is the dimensionality of ω , and $\theta \in \omega$.

The practical use of (1) is that (for sufficiently large n) we can use the χ_{r-m}^2 distribution in order to determine the cutoff point C of the test, which rejects H_0 when $-2 \log \lambda \geq C$. Specifically, $C \simeq \chi_{r-m; \alpha}^2$. Thus, for testing the hypothesis H_0 at level of significance α , H_0 is to be rejected whenever $-2 \log \lambda$ is $\geq \chi_{r-m; \alpha}^2$ (always provided n is sufficiently large).

EXAMPLE 1

The Multinomial Case A multinomial experiment, with k possible outcomes O_1, \dots, O_k and respective unknown probabilities p_1, \dots, p_k , is carried out independently n times, and let X_1, \dots, X_k be the r.v.'s denoting the number of times outcomes O_1, \dots, O_k occur, respectively. Then the joint p.d.f. of the X_i 's is:

$$f(x_1, \dots, x_k; \theta) = \frac{n!}{x_1! \cdots x_k!} p_1^{x_1} \cdots p_k^{x_k}, \quad (2)$$

for $x_1, \dots, x_k \geq 0$ integers with $x_1 + \cdots + x_k = n$, and $\theta = (p_1, \dots, p_k)$. The parameter space Ω is $(k-1)$ -dimensional and is defined by:

$$\Omega = \{(p_1, \dots, p_k) \in \mathfrak{N}^k; \quad p_i > 0, \quad i = 1, \dots, k, \quad p_1 + \cdots + p_k = 1\}.$$

DISCUSSION Suppose we wish to test the hypothesis $H_0: p_i = p_{i0}, i = 1, \dots, k$ (specified) at level of significance α . Under H_0 ,

$$L(\hat{\omega}) = \frac{n!}{x_1! \cdots x_k!} p_{10}^{x_1} \cdots p_{k0}^{x_k}$$

and we know that the MLE's of the p_i 's are: $\hat{p}_i = \frac{x_i}{n}, i = 1, \dots, k$. Therefore

$$\begin{aligned} L(\hat{\Omega}) &= \frac{n!}{x_1! \cdots x_k!} \hat{p}_1^{x_1} \cdots \hat{p}_k^{x_k} = \frac{n!}{x_1! \cdots x_k!} \left(\frac{x_1}{n}\right)^{x_1} \cdots \left(\frac{x_k}{n}\right)^{x_k} \\ &= n^{-n} \frac{n!}{x_1! \cdots x_k!} x_1^{x_1} \cdots x_k^{x_k}. \end{aligned}$$

Therefore

$$\lambda = n^n \left(\frac{p_{10}}{x_1}\right)^{x_1} \cdots \left(\frac{p_{k0}}{x_k}\right)^{x_k}, \text{ and } H_0 \text{ is rejected when } -2 \log \lambda \geq \chi_{k-1; \alpha}^2$$

since here $r = k - 1$ and $m = 0$.

Numerical Example The fairness of a die is to be tested on the basis of the following outcomes of 30 independent rollings: $x_1 = 4, x_2 = 7, x_3 = 3, x_4 = 8, x_5 = 4, x_6 = 4$. Take $\alpha = 0.05$.


DISCUSSION Here $H_0: p_i = \frac{1}{6}, i = 1, \dots, 6$ and the LR λ is given by:

$$\begin{aligned}\lambda &= 30^{30} \left(\frac{1}{6 \times 4}\right)^4 \times \left(\frac{1}{6 \times 7}\right)^7 \times \left(\frac{1}{6 \times 3}\right)^3 \times \left(\frac{1}{6 \times 8}\right)^8 \times \left(\frac{1}{6 \times 4}\right)^4 \times \left(\frac{1}{6 \times 4}\right)^4 \\ &= 30^{30} \times 6^{-30} \times 4^{-4} \times 7^{-7} \times 3^{-3} \times 8^{-8} \times 4^{-4} \times 4^{-4}.\end{aligned}$$

It follows that $-2 \log \lambda \simeq 3.826$, whereas $\chi_{5,0.05}^2 = 11.071$. Thus, the hypothesis H_0 is not rejected.

Figure 12.1

**An $r \times s$
Contingency Table**



Application to Contingency Tables Consider a multinomial experiment with $r \times s$ possible outcomes arranged in a rectangular array with r rows and s columns. Such a rectangular array is referred to as an $r \times s$ contingency table. The r rows and s columns generate $r \times s$ cells. (See Figure 12.1.) Denote by p_{ij} the probability that an outcome will fall into the (i, j) th cell. Carry out the multinomial experiment under consideration n independent times, and let X_{ij} be the r.v. denoting the number of outcomes falling into the (i, j) th cell. Define p_i and p_j by the formulas:

$$p_i = \sum_{j=1}^s p_{ij}, \quad i = 1, \dots, r, \quad p_j = \sum_{i=1}^r p_{ij}, \quad j = 1, \dots, s. \quad (3)$$

Then, clearly, p_i is the probability that an outcome falls in the i th row regardless of column, and p_j is the probability that an outcome falls in the j th column regardless of row. Of course, $\sum_{i=1}^r p_i = \sum_{j=1}^s p_j = \sum_{i=1}^r \sum_{j=1}^s p_{ij} = 1$. Also, define the r.v.'s X_i and X_j as follows:

$$X_i = \sum_{j=1}^s X_{ij}, \quad i = 1, \dots, r, \quad X_j = \sum_{i=1}^r X_{ij}, \quad j = 1, \dots, s. \quad (4)$$

Thus, clearly, X_i denotes the number of outcomes falling in the i th row regardless of column, and X_j denotes the number of outcomes falling in the

j th column regardless of row. It is also clear, that

$$\sum_{i=1}^r X_{i.} = \sum_{j=1}^s X_{.j} = \sum_{i=1}^r \sum_{j=1}^s X_{ij} = n.$$

The parameters p_{ij} , $i = 1, \dots, r$, $j = 1, \dots, s$ are, in practice, unknown and are estimated by the MLE's $\hat{p}_{ij} = \frac{x_{ij}}{n}$. In the testing hypotheses framework, one could test the hypothesis $H_0: p_{ij} = p_{ij0}$, $i = 1, \dots, r$, $j = 1, \dots, s$, specified. However, from a practical viewpoint, this is not an interesting hypothesis. What is of true interest here is to test independence of rows and columns. In order to provide some motivation, suppose that some subjects (e.g., human beings) are classified according to two *characteristics* to be denoted by A and B (e.g., human beings are classified according to gender, characteristic A , and whether or not they are cigarette smokers, characteristic B). Suppose that characteristic A has r levels and characteristic B has s levels. (In the concrete example at hand, $r = 2$ (Male, Female) and $s = 2$ (Smoker, Nonsmoker).) We agree to have the r rows in an $r \times s$ contingency table represent the r levels of characteristic A and the s columns of the contingency table represent the s levels of characteristic B . Then independence of rows and columns, as mentioned earlier, is restated as independence of characteristics A and B or, more precisely, independence of the r levels of characteristic A and the s levels of characteristic B . (In the concrete example this would mean that gender and smoking/nonsmoking are independent events.) The probabilistic formulation of the independence stated is as follows:

Observe that $P(A_i \cap B_j) = p_{ij}$, $P(A_i) = p_{i.}$, and $P(B_j) = p_{.j}$. Independence of A_i and B_j for all i and j means then that

$$P(A_i \cap B_j) = P(A_i)P(B_j), \quad \text{all } i \text{ and } j, \quad \text{or} \quad p_{ij} = p_{i.}p_{.j}, \quad \text{all } i \text{ and } j.$$

To put it differently, we wish to test the hypothesis that there exist (probabilities) $p_i > 0$, $i = 1, \dots, r$, $p_1 + \dots + p_r = 1$ and $q_j > 0$, $j = 1, \dots, s$, $q_1 + \dots + q_s = 1$, such that

$$H_0: p_{ij} = p_i q_j, \quad i = 1, \dots, r, \quad j = 1, \dots, s. \quad (5)$$

(Of course, then $p_i = p_{i.}$ and $q_j = p_{.j}$, all i and j .) The MLE of p_{ij} is $\hat{p}_{ij} = \frac{x_{ij}}{n}$, $i = 1, \dots, r$, $j = 1, \dots, s$. Therefore, writing $\prod_{i,j}$ for $\prod_{i=1}^r \prod_{j=1}^s$ and setting θ for $(p_{ij}, i = 1, \dots, r, j = 1, \dots, s)$, we have, for the likelihood function

$$L(\theta | x_{ij}, i = 1, \dots, r, j = 1, \dots, s) = \frac{n!}{\prod_{i,j} x_{ij}!} \prod_{i,j} p_{ij}^{x_{ij}}, \quad (6)$$

and

$$L(\hat{\Omega}) = \frac{n!}{\prod_{i,j} x_{ij}!} \prod_{i,j} \left(\frac{x_{ij}}{n} \right)^{x_{ij}} = \frac{n!}{n^n \prod_{i,j} x_{ij}!} \prod_{i,j} x_{ij}^{x_{ij}}. \quad (7)$$

Under H_0 , the likelihood function becomes

$$\begin{aligned} \frac{n!}{\prod_{i,j} x_{ij}!} \prod_{i,j} (p_i q_j)^{x_{ij}} &= \frac{n!}{\prod_{i,j} x_{ij}!} \prod_i \prod_j p_i^{x_{ij}} q_j^{x_{ij}} \\ &= \frac{n!}{\prod_{i,j} x_{ij}!} \prod_i p_i^{x_i} q_1^{x_{i1}} \cdots q_s^{x_{is}} = \frac{n!}{\prod_{i,j} x_{ij}!} \left(\prod_i p_i^{x_i} \right) \left(\prod_j q_j^{x_j} \right), \end{aligned} \quad (8)$$

because

$$\begin{aligned} \prod_i p_i^{x_i} q_1^{x_{i1}} \cdots q_s^{x_{is}} &= (p_1^{x_1} q_1^{x_{11}} \cdots q_s^{x_{1s}}) \cdots (p_r^{x_r} q_1^{x_{r1}} \cdots q_s^{x_{rs}}) \\ &= (p_1^{x_1} \cdots p_r^{x_r}) (q_1^{x_{11}} \cdots q_1^{x_{r1}}) \cdots (q_s^{x_{1s}} \cdots q_s^{x_{rs}}) \\ &= \left(\prod_i p_i^{x_i} \right) (q_1^{x_{11}} \cdots q_s^{x_{1s}}) \cdots (q_1^{x_{r1}} \cdots q_s^{x_{rs}}) \\ &= \left(\prod_i p_i^{x_i} \right) \left(\prod_j q_j^{x_j} \right). \end{aligned}$$

The MLE's of p_i and q_j are given by

$$\hat{p}_i = \frac{x_i}{n}, \quad i = 1, \dots, r, \quad \hat{q}_j = \frac{x_j}{n}, \quad j = 1, \dots, s, \quad (9)$$

so that

$$L(\hat{\omega}) = \frac{n!}{\prod_{i,j} x_{ij}!} \prod_i \binom{x_i}{n} \prod_j \binom{x_j}{n} = \frac{n!}{n^{2n} \prod_{i,j} x_{ij}!} \left(\prod_i x_i^{x_i} \right) \left(\prod_j x_j^{x_j} \right). \quad (10)$$

By (7) and (10), we have then

$$\lambda = \frac{(\prod_i x_i^{x_i}) (\prod_j x_j^{x_j})}{n^n \prod_{i,j} x_{ij}^{x_{ij}}}, \quad (11)$$

and

$$-2 \log \lambda = 2 \left[\left(n \log n + \sum_{i=1}^r \sum_{j=1}^s x_{ij} \log x_{ij} \right) - \left(\sum_{i=1}^r x_i \log x_i + \sum_{j=1}^s x_j \log x_j \right) \right]. \quad (12)$$

Here the dimension of Ω is $rs - 1$ because we have rs p_{ij} , $i = 1, \dots, r$, $j = 1, \dots, s$, which, however, satisfy the relationship $\sum_{i=1}^r \sum_{j=1}^s p_{ij} = 1$. In order to determine the dimension of ω , observe that we have $r + s$ parameters p_i , $i = 1, \dots, r$ and q_j , $j = 1, \dots, s$, which, however, satisfy two relationships; namely, $\sum_{i=1}^r p_i = 1$ and $\sum_{j=1}^s q_j = 1$. Therefore the dimension of ω is $r + s - 2$ and

$$\dim \Omega - \dim \omega = (rs - 1) - (r + s - 2) = (r - 1)(s - 1).$$

Furthermore, it so happens that the (unspecified) conditions of Theorem 1 are satisfied here, so that, under H_0 , $-2 \log \lambda$ is distributed approximately (for all sufficiently large n) as $\chi_{(r-1)(s-1)}^2$. It follows that the hypothesis (5) about independence is rejected, at level of significance α , whenever

$$-2 \log \lambda \geq \chi_{(r-1)(s-1); \alpha}^2. \quad (13)$$

Numerical Example A population consisting of $n = 100$ males (M) and females (F) is classified according to their smoking (S) or nonsmoking (NS) cigarettes habit. Suppose the resulting 2×2 contingency table is as given below. Then test independence of gender and smoking/nonsmoking habit at the level of significance $\alpha = 0.05$.

	S	NS	
M	20	35	55
F	15	30	45
	35	65	100

DISCUSSION The values x_{ij} are shown in the cells and the $x_{i\cdot}$, $x_{\cdot j}$ are shown in the margins, and they are: $x_{11} = 20$, $x_{12} = 35$, $x_{21} = 15$, $x_{22} = 30$, $x_{1\cdot} = 55$, $x_{2\cdot} = 45$, $x_{\cdot 1} = 35$, $x_{\cdot 2} = 65$. Replacing these values in the expression of $-2 \log \lambda$ given by (12), we find $-2 \log \lambda = 0.061$. Here $r = s = 2$, so that $\chi_{(r-1)(s-1); \alpha}^2 = \chi_{1; 0.05}^2 = 3.841$. Therefore the hypothesis is not rejected.

Exercises

- 1.1** (i) In reference to Example 18 in Chapter 1, the appropriate probability model is the Multinomial distribution with parameters n and p_A , p_B , p_{AB} , p_O , where p_A through p_O are the probabilities that an individual, chosen at random from among the n persons has blood type either A or B or AB or O , respectively.
- (ii) Let p_{A0} , p_{B0} , p_{AB0} , and p_{O0} be a priori stipulated numbers. Then, checking agreement of the actual probabilities with the stipulated values amounts to testing the hypothesis

$$H_0: p_A = p_{A0}, \quad p_B = p_{B0}, \quad p_{AB} = p_{AB0}, \quad p_O = p_{O0}.$$

- (iii) The hypothesis H_0 is tested by means of either the log-LR test (see Example 1 here) or the χ^2 goodness-of-fit test. (See also Exercise 2.1.)
- 1.2** (i) In reference to Example 19 in Chapter 1, the appropriate probability model is the Multinomial distribution with parameters $n = 41,208$ and p_i , $i = 1, \dots, 12$, where $p_i = P(\text{a birth chosen at random from among the } n \text{ births falls in the } i\text{th month})$.
- (ii) Checking uniform distribution of the n births over the 12 months amounts to testing the hypothesis

$$H_0: p_i = p_{i0} = \frac{1}{12}, \quad i = 1, \dots, 12.$$

- (iii) The hypothesis H_0 is tested by means of either the log-LR test (see Example 1 here) or the χ^2 goodness-of-fit test. (See also Exercise 2.2.) The hypothesis H_0 is rejected when $-2 \log \lambda > \chi_{11; \alpha}^2$.

- 1.3** (i) In reference to Example 20 in Chapter 1, the appropriate probability model is the 2×3 contingency table setup in the example.
- (ii) If p_{ij} is the probability that a randomly chosen subject from among the 150 falls into the (i, j) th cell, then independence between the factors health and diet is checked by testing the hypothesis

$$H_0: p_{ij} = p_i q_j, \quad i = 1, 2 \text{ and } j = 1, 2, 3.$$

An appropriate test statistic for testing the hypothesis H_0 is either the log-LR test or the χ^2 goodness-of-fit-statistic. (See also Exercise 2.3.)

- 1.4** (i) In reference to Example 21 in Chapter 1, the appropriate probability model is the 3×4 contingency table setup in the example.
- (ii) If p_{ij} is the probability that a randomly chosen subject from among the 200 falls into the (i, j) th cell, then checking the stipulation that change of bone minerals does not vary for different groups amounts to testing the hypothesis

$$H_0: p_{ij} = p_i q_j, \quad i = 1, 2, 3 \text{ and } j = 1, 2, 3.$$

The hypothesis H_0 may be checked by means of either the log-LR test or the χ^2 goodness-of-fit test. (See also Exercise 2.4.)

- 1.5** In reference to Example 1 of Chapter 1, the n landfills are classified according to two levels of concentration (High and Low) and three levels of hazardous chemicals (Arsenic, Barium, and Mercury) to produce the following 2×3 contingency table:

		HAZARDOUS CHEMICALS			
		Arsenic	Barium	Mercury	Totals
Level of	High	x_{11}	x_{12}	x_{13}	$x_{1.}$
Concentration	Low	x_{21}	x_{22}	x_{23}	$x_{2.}$
	Totals	$x_{.1}$	$x_{.2}$	$x_{.3}$	$x_{..} = n$

Then, if p_{ij} is the probability that a landfill chosen at random from among the n landfills falls into the (i, j) th cell, part (ii) of the example becomes that of testing the hypothesis $H_0: p_{ij} = p_{ij0}$, where p_{ij0} , $i = 1, 2$ and $j = 1, 2, 3$ are a priori stipulated numbers. The hypothesis H_0 is tested by means of either the log-LR test or the χ^2 goodness-of-fit test. (See also Exercise 2.5.)

12.2 A Goodness-of-Fit Test

This test applies primarily to the Multinomial distribution, although other distributions can also be suitably reduced to a multinomial framework. In the notation of the previous section, we have that, for each fixed $i = 1, \dots, k$, $X_i \sim B(n, p_i)$, so that $E_{\theta} X_i = np_i$, $i = 1, \dots, k$, $\theta = (p_1, \dots, p_k)$. Thus, the i th outcome would be expected to appear np_i times, whereas the actual number of times it appears is X_i . It then makes sense to compare what we expect

and what we, actually, observe, and do this simultaneously for all $i = 1, \dots, k$. One way of doing this is to look at the quantity $\sum_{i=1}^k (X_i - np_i)^2$. Small values of this quantity would indicate agreement between expected and observed values, and large values would indicate the opposite. For distributional reasons, the above expression is modified as indicated below, and in this form it is denoted by χ^2 ; namely,

$$\chi^2 = \sum_{i=1}^k \frac{(X_i - np_i)^2}{np_i}. \quad (14)$$

Expression (14) is the basis for constructing test statistics for testing various hypotheses. In this setting, we will consider the hypothesis $H_0: p_i = p_{i0}$, $i = 1, \dots, k$, specified as we did in the previous section. Under H_0 , (14) is denoted by χ_ω^2 and is equal to:

$$\chi_\omega^2 = \sum_{i=1}^k \frac{(X_i - np_{i0})^2}{np_{i0}}. \quad (15)$$

This is a statistic and is used for testing H_0 . Accordingly, H_0 is rejected, at level of significance α , if $\chi_\omega^2 \geq C$, where C is determined by the requirement $P_{H_0}(\chi_\omega^2 \geq C) = \alpha$. It can be seen that, under H_0 , $\chi_\omega^2 \simeq \chi_{k-1}^2$ for all sufficiently large n . Consequently, $C \simeq \chi_{k-1, \alpha}^2$. The test used here is called a test of *goodness-of-fit* for obvious reasons. It is also referred to as *chi-square* (or χ^2) *goodness-of-fit* test, because of the symbol used in relation (15), and because its asymptotic distribution (under the null hypothesis) is chi-square with certain degrees of freedom. Thus, the (Chi-Square) goodness-of-fit test rejects H_0 whenever $\chi_\omega^2 \geq \chi_{k-1, \alpha}^2$.

For illustrative and also comparison purposes, let us consider the first numerical example in the previous section.

Numerical Example Here $np_{10} = \dots = np_{60} = \frac{30}{6} = 5$, and then the observed value of χ_ω^2 is:

$$\chi_\omega^2 = \frac{1}{5}[(4-5)^2 + (7-5)^2 + (3-5)^2 + (8-5)^2 + (4-5)^2 + (4-5)^2] = 4.$$

For $\alpha = 0.05$, $\chi_{k-1, \alpha}^2 = \chi_{5, 0.05}^2 = 11.071$, and since $\chi_\omega^2 = 4 < 11.071$, the hypothesis H_0 is not rejected, as was also the case with the LR test.

In the framework of a contingency table, expression (14) becomes

$$\chi^2 = \sum_{i=1}^r \sum_{j=1}^s \frac{(X_{ij} - np_{ij})^2}{np_{ij}}. \quad (16)$$

Under the hypothesis of independence stated in (5), expression (16) takes the form

$$\chi_\omega^2 = \sum_{i=1}^r \sum_{j=1}^s \frac{(X_{ij} - np_i q_j)^2}{np_i q_j}. \quad (17)$$

From (17), we form the test statistic $\chi_{\hat{\omega}}^2$ defined below:

$$\chi_{\hat{\omega}}^2 = \sum_{i=1}^r \sum_{j=1}^s \frac{(X_{ij} - n\hat{p}_i\hat{q}_j)^2}{n\hat{p}_i\hat{q}_j}, \quad (18)$$

where \hat{p}_i , $i = 1, \dots, r$ and \hat{q}_j , $j = 1, \dots, s$ are given in (9). Once again, it may be seen that, under H_0 , $\chi_{\hat{\omega}}^2 \simeq \chi_{(r-1)(s-1)}^2$ for all sufficiently large n . Thus, the hypothesis H_0 is rejected, at level of significance α , whenever $\chi_{\hat{\omega}}^2 \geq \chi_{(r-1)(s-1); \alpha}^2$. The contingency table numerical example of the previous section is as below.

Numerical Example Here $\hat{p}_1 = 0.55$, $\hat{p}_2 = 0.45$, $\hat{q}_1 = 0.35$, $\hat{q}_2 = 0.65$, so that

$$n\hat{p}_1\hat{q}_1 = 19.25, \quad n\hat{p}_1\hat{q}_2 = 35.75, \quad n\hat{p}_2\hat{q}_1 = 15.75, \quad n\hat{p}_2\hat{q}_2 = 29.25.$$

DISCUSSION Therefore

$$\chi_{\hat{\omega}}^2 = \frac{(20 - 19.25)^2}{19.25} + \frac{(35 - 35.75)^2}{35.75} + \frac{(15 - 15.75)^2}{15.75} + \frac{(30 - 29.25)^2}{29.25} \simeq 0.0998.$$

Since $\chi_{(r-1)(s-1); \alpha}^2 = \chi_{1; 0.05}^2 = 3.841$, the hypothesis H_0 is not rejected, as was also the case with the LR test.

Exercises

- 2.1 Same as Exercise 1.1.
- 2.2 Same as Exercise 1.2.
- 2.3 Same as Exercise 1.3.
- 2.4 Same as Exercise 1.4.
- 2.5 Same as Exercise 1.5.
- 2.6 A coin, with probability p of falling heads, is tossed independently 100 times, and 60 heads are observed.
 - (i) Test the hypothesis $H_0: p = 1/2$ (against the alternative $H_A: p \neq 1/2$) at level of significance $\alpha = 0.1$, by using the appropriate χ^2 goodness-of-fit test.
 - (ii) Determine the P -value of the test (use linear interpolation).
- 2.7 A die is cast independently 600 times, and the numbers 1 through 6 appear with the frequencies recorded below.

1	2	3	4	5	6
100	94	103	89	110	104

Use the appropriate χ^2 goodness-of-fit test to test fairness for the die at level of significance $\alpha = 0.1$.

- 2.8** In a certain genetic experiment, two different varieties of a certain species are crossed and a specific characteristic of the offspring can occur at only three levels A , B , and C , say. According to a proposed model, the probabilities for A , B , and C are $\frac{1}{12}$, $\frac{3}{12}$, and $\frac{8}{12}$, respectively. Out of 60 offspring, 6, 18, and 36 fall into levels A , B , and C , respectively. Test the validity of the proposed model at the level of significance $\alpha = 0.05$. Use the appropriate χ^2 goodness-of-fit test.
- 2.9** Course work grades are often assumed to be Normally distributed. In a certain class, suppose that letter grades are given in the following manner: A for grades in the range from 90 to 100 inclusive, B for grades in the range from 75 to 89 inclusive, C for grades in the range from 60 to 74 inclusive, D for grades in the range from 50 to 59 inclusive, and F for grades in the range from 0 to 49. Use the data given below to check the assumption that the data are coming from an $N(75, 9^2)$ distribution. For this purpose, employ the appropriate χ^2 goodness-of-fit test, and take $\alpha = 0.05$.

A	B	C	D	F
3	12	10	4	1

Hint: Assuming that the grade of a student chosen at random is a r.v. $X \sim N(75, 81)$, compute the probabilities of an A , B , C , D , and F . Then use these probabilities in applying the χ^2 goodness-of-fit test.

- 2.10** It is often assumed that the I.Q. scores of human beings are Normally distributed. On the basis of the following data, test this claim at level of significance $\alpha = 0.05$ by using the appropriate χ^2 goodness-of-fit test. Specifically, if X is the r.v. denoting the I.Q. score of an individual chosen at random, then:
- Set $p_1 = P(X \leq 90)$, $p_2 = P(90 < X \leq 100)$, $p_3 = P(100 < X \leq 110)$, $p_4 = P(110 < X \leq 120)$, $p_5 = P(120 < X \leq 130)$, $p_6 = P(X > 130)$.
 - Calculate the probabilities p_i , $i = 1, \dots, 6$ under the assumption that $X \sim N(100, 15^2)$ and call them p_{i0} , $i = 1, \dots, 6$. Then set up the hypothesis $H_0: p_i = p_{i0}$, $i = 1, \dots, 6$.
 - Use the appropriate χ^2 goodness-of-fit test to test the hypothesis at level of significance $\alpha = 0.05$.

The available data are given below, where x denotes the observed number of individuals lying in a given interval.

$x \leq 90$	$90 < x \leq 100$	$100 < x \leq 110$	$110 < x \leq 120$	$120 < x \leq 130$	$x > 130$
10	18	23	22	18	9

- 2.11** Consider a group of 100 people living and working under very similar conditions. Half of them are given a preventive shot against a certain disease and the other half serve as controls. Of those who received the treatment, 40 did not contract the disease whereas the remaining 10 did so. Of those not treated, 30 did contract the disease and the remaining 20 did not. Test effectiveness of the vaccine at the level of significance $\alpha = 0.05$, by using the appropriate χ^2 goodness-of-fit test.

Hint: For an individual chosen at random from the target population of 100 individuals, denote by T_1, T_2 and D_1, D_2 the following events: $T_1 =$ “treated,” $T_2 =$ “not treated,” $D_1 =$ “diseased,” $D_2 =$ “not diseased,” and set up the appropriate 2×2 contingency table.

2.12 On the basis of the following scores, appropriately taken, test whether there are gender-associated differences in mathematical ability (as is often claimed!). Take $\alpha = 0.05$, and use the appropriate χ^2 goodness-of-fit test.

Boys:	80	96	98	87	75	83	70	92	97	82
Girls:	82	90	84	70	80	97	76	90	88	86

Hint: Group the grades into the following intervals: $[70, 75)$, $[75, 80)$, $[80, 85)$, $[85, 90)$, $[90, 95)$, $[95, 100)$, and count the grades of boys and girls falling into each one of these intervals. Then form a 2×6 contingency table with rows the two levels of gender (Boy, Girl), and columns the six levels of grades. Finally, with p_{ij} standing for the probability that an individual, chosen at random from the target population, falls into the (i, j) th cell, stipulate the hypothesis $H_0: p_{ij} = p_i q_j$, $i = 1, 2$ and $j = 1, \dots, 6$, and proceed to test it as suggested.

2.13 From each of four political wards of a city with approximately the same number of voters, 100 voters were chosen at random and their opinions were asked regarding a certain legislative proposal. On the basis of the data given below, test whether the fractions of voters favoring the legislative proposal under consideration differ in the four wards. Take $\alpha = 0.05$, and use the appropriate χ^2 goodness-of-fit test.

	WARD				
	1	2	3	4	Totals
Favor proposal	37	29	32	21	119
Do not favor proposal	63	71	68	79	281
Totals	100	100	100	100	400

12.3 Decision-Theoretic Approach to Testing Hypotheses

There are chapters and books written on this subject. What we plan to do in this section is to deal with the simplest possible case of a testing hypothesis problem in order to illustrate the underlying concepts.

To this end, let X_1, \dots, X_n be a random sample with an unknown p.d.f. f . We adopt the (somewhat unrealistic) position that f can be one of two possible specified p.d.f.'s, f_0 or f_1 . On the basis of the observed values x_1, \dots, x_n of X_1, \dots, X_n , we are invited to decide which is the true p.d.f. This decision will be made on the basis of a (nonrandomized) *decision function* $\delta = \delta(x_1, \dots, x_n)$ defined on \mathfrak{R}^n into \mathfrak{R} . More specifically, let R be a subset of \mathfrak{R}^n , and suppose that if $\mathbf{x} = (x_1, \dots, x_n)$ lies in R , we decide that f_1 is the true p.d.f., and if \mathbf{x}

lies in R^c (the complement of R with respect to \mathfrak{R}^n), we decide in favor of f_0 . In terms of a decision function, we reach the same conclusion by taking $\delta(\mathbf{x}) = I_R(\mathbf{x})$ (the indicator function of R) and deciding in favor of f_1 if $\delta(\mathbf{x}) = 1$ and in favor of f_0 if $\delta(\mathbf{x}) = 0$. Or

$$\delta(\mathbf{x}) = \begin{cases} 1 & \text{(which happens when } \mathbf{x} \in R \text{) leads to selection} \\ & \text{of } f_1, \text{ and hence rejection of } f_0, \\ 0 & \text{(which happens when } \mathbf{x} \in R^c \text{) leads to selection} \\ & \text{of } f_0, \text{ and hence rejection of } f_1. \end{cases} \quad (19)$$

At this point, we introduce monetary penalties for making wrong decisions, which are expressed in terms of a *loss function*. Specifically, let $L(f; \delta)$ be a function in two arguments, the p.d.f. f and the decision function $\delta = \delta(\mathbf{x})$. Then it makes sense to define $L(f, \delta)$ in the following way.

$$L(f; \delta) = \begin{cases} 0 & \text{if } f = f_0 \text{ and } \delta(\mathbf{x}) = 0 \text{ or } f = f_1 \text{ and } \delta(\mathbf{x}) = 1, \\ L_1 & \text{if } f = f_0 \text{ and } \delta(\mathbf{x}) = 1, \\ L_2 & \text{if } f = f_1 \text{ and } \delta(\mathbf{x}) = 0, \end{cases} \quad (20)$$

where L_1 and L_2 are positive quantities.

Next, consider the average (expected) loss when the decision function δ is used, which is denoted by $R(f; \delta)$ and is called the *risk function*. In order to find the expression of $R(f; \delta)$, let us suppose that $P_{f_0}(\mathbf{X} \in R) = P_{f_0}[\delta(\mathbf{X}) = 1] = \alpha$ and $P_{f_1}(\mathbf{X} \in R) = P_{f_1}[\delta(\mathbf{X}) = 1] = \pi$. Then α is the probability of deciding in favor of f_1 if, actually, f_0 is true, and π is the probability of deciding in favor of f_1 when f_1 is, actually, true. Then:

$$R(f; \delta) = \begin{cases} L_1 P_{f_0}(\mathbf{X} \in R) = L_1 P_{f_0}[\delta(\mathbf{X}) = 1] = L_1 \alpha, & \text{if } f = f_0 \\ L_2 P_{f_1}(\mathbf{X} \in R^c) = L_2 P_{f_1}[\delta(\mathbf{X}) = 0] = L_2(1 - \pi), & \text{if } f = f_1, \end{cases} \quad (21)$$

or,

$$\begin{aligned} R(f_0; \delta) &= L_1 P_{f_0}(\mathbf{X} \in R) = L_1 \alpha, \\ R(f_1; \delta) &= L_2 P_{f_1}(\mathbf{X} \in R^c) = L_2(1 - \pi). \end{aligned} \quad (22)$$

Let us recall that our purpose is to construct an optimal decision function $\delta = \delta(\mathbf{x})$, where optimality is defined below on the basis of two different criteria. From relation (22), we know which is the bigger among the risk values $R(f_0; \delta) = L_1 \alpha$ and $R(f_1; \delta) = L_2(1 - \pi)$. That is, we have the quantity $\max\{R(f_0; \delta), R(f_1; \delta)\}$. For any other (nonrandomized) decision function δ^* the corresponding quantity is $\max\{R(f_0; \delta^*), R(f_1; \delta^*)\}$. Then it makes sense to choose δ so that

$$\max\{R(f_0; \delta), R(f_1; \delta)\} \leq \max\{R(f_0; \delta^*), R(f_1; \delta^*)\} \quad (23)$$

for any other decision function δ^* as described above. A decision function δ , if it exists, which satisfies inequality (23) is called *minimax* (since it minimizes the

maximum risk). The result below, Theorem 2, provides conditions under which the decision function δ defined by (19) is, actually, minimax. The problem is stated as a testing problem of a simple hypothesis against a simple alternative.

THEOREM 2

Let X_1, \dots, X_n be a random sample with p.d.f. f which is either f_0 or f_1 , both completely specified. For testing the hypothesis $H_0: f = f_0$ against the alternative $H_A: f = f_1$ at level of significance α , define the rejection region R by:

$$R = \{(x_1, \dots, x_n) \in \mathfrak{N}^n; f_1(x_1) \cdots f_1(x_n) > C f_0(x_1) \cdots f_0(x_n)\},$$

and let the test function $\delta = \delta(\mathbf{x})$ ($\mathbf{x} = (x_1, \dots, x_n)$) be defined by (19); i.e.,

$$\delta(\mathbf{x}) = \begin{cases} 1 & \text{if } \mathbf{x} \in R, \\ 0 & \text{if } \mathbf{x} \in R^c. \end{cases}$$

The constant C is defined by the requirement that $E_{f_0} \delta(\mathbf{X}) = P_{f_0}(\mathbf{X} \in R) = \alpha$ ($\mathbf{X} = (X_1, \dots, X_n)$), and it is assumed that the level of significance α , the power π of the test δ , and the quantities L_1 and L_2 satisfy the relationship

$$(R(f_0; \delta) =) L_1 \alpha = L_2 (1 - \pi) (=R(f_1; \delta)). \quad (24)$$

Then the decision function $\delta = \delta(\mathbf{x})$ is minimax.

REMARK 1 In connection with relation (24), observe that, if we determine the level of significance α , then the power π is also determined, and therefore relation (24) simply specifies a relationship between the losses L_1 and L_2 ; they cannot be determined independently but rather one will be a function of the other. In the present context, however, we wish to have the option of specifying the losses L_1 and L_2 , and then see what is a possible determination of the constant C , which will produce a test of level of significance α (and of power π) satisfying relation (24).

PROOF OF THEOREM 2 For simplicity, let us write P_0 and P_1 instead of P_{f_0} and P_{f_1} , respectively, and likewise, $R(0; \delta)$ and $R(1; \delta)$ instead of $R(f_0; \delta)$ and $R(f_1; \delta)$, respectively. Then assumption (24) is rewritten thus: $R(0; \delta) = L_1 \alpha = L_2 (1 - \pi) = R(1; \delta)$. Recall that we are considering only nonrandomized decision functions. With this in mind, let T be any (other than R) subset of \mathfrak{N}^n , and let δ^* be its indicator function, $\delta^*(\mathbf{x}) = I_T(\mathbf{x})$, so that δ^* is the decision function associated with T . Then, in analogy with (22),

$$R(0; \delta^*) = L_1 P_0(\mathbf{X} \in T), \quad R(1; \delta^*) = L_2 P_1(\mathbf{X} \in T^c). \quad (25)$$

Look at $R(0; \delta)$ and $R(0; \delta^*)$ and suppose that $R(0; \delta^*) \leq R(0; \delta)$. This is equivalent to $L_1 P_0(\mathbf{X} \in T) \leq L_1 P_0(\mathbf{X} \in R) = L_1 \alpha$, or $P_0(\mathbf{X} \in T) \leq \alpha$. So δ^* , being looked upon as a test, is of level of significance $\leq \alpha$. Then by Theorem 1

in Chapter 11, the power of the test δ^* , which is $P_1(\mathbf{X} \in T)$, is less than or equal to $P_1(\mathbf{X} \in R)$, which is the power of the test δ . This is so because δ is of level of significance α , MP among all tests of level of significance $\leq \alpha$. From $P_1(\mathbf{X} \in T) \leq P_1(\mathbf{X} \in R)$ we have, equivalently, $P_1(\mathbf{X} \in T^c) \geq P_1(\mathbf{X} \in R^c)$ or $L_2 P_1(\mathbf{X} \in T^c) \geq L_2 P_1(\mathbf{X} \in R^c)$ or $R(1; \delta^*) \geq R(1; \delta)$. To summarize, the assumption $R(0; \delta^*) \leq R(0; \delta)$ leads to $R(1; \delta) \leq R(1; \delta^*)$. Hence

$$R(0; \delta^*) \leq R(0; \delta) = R(1; \delta) \text{ (by (24))} \leq R(1; \delta^*),$$

and therefore

$$\max\{R(0; \delta^*), R(1; \delta^*)\} = R(1; \delta^*) \geq R(1; \delta) = \max\{R(0; \delta), R(1; \delta)\}, \quad (26)$$

as desired. Next, the assumption,

$$R(0; \delta) < R(0; \delta^*) \quad (27)$$

leads likewise to the inequalities

$$R(1; \delta^*) \leq R(1; \delta) = R(0; \delta) \text{ (by (24))} < R(0; \delta^*), \quad (28)$$

so that

$$\max\{R(0; \delta^*), R(1; \delta^*)\} = R(0; \delta^*) > R(0; \delta) = \max\{R(0; \delta), R(1; \delta)\}. \quad (29)$$

Relations (26) and (29) yield

$$\max\{R(0; \delta), R(1; \delta)\} \leq \max\{R(0; \delta^*), R(1; \delta^*)\},$$

so that δ is, indeed, minimax. \blacktriangle

REMARK 2 It is to be pointed out that the minimax decision function $\delta = \delta(\mathbf{x})$ above is the MP test of level of significance $P_{f_0}(\mathbf{X} \in R)$ for testing the (simple) hypothesis $H_0: f = f_0$ against the (simple) alternative $H_A: f = f_1$.

REMARK 3 If the underlying p.d.f. f depends on a parameter $\theta \in \Omega$, then the two possible options f_0 and f_1 for f will correspond to two values of the parameter θ , θ_0 , and θ_1 , say.

The theorem of this section is illustrated now by two examples.

EXAMPLE 2

On the basis of the random sample X_1, \dots, X_n from the $N(\theta, 1)$ distribution, determine the minimax decision function $\delta = \delta(\mathbf{x})$ for testing the hypothesis $H_0: \theta = \theta_0$ against the alternative $H_A: \theta = \theta_1$.

DISCUSSION Here the joint p.d.f. of the X_i 's is

$$L(\mathbf{x}; \theta) = (2\pi)^{-n/2} \exp\left[-\frac{1}{2} \sum_{i=1}^n (x_i - \theta)^2\right],$$

so that the rejection region R is defined by $L(\mathbf{x}; \theta_1) > CL(\mathbf{x}; \theta_0)$ or, equivalently, by

$$\exp[n(\theta_1 - \theta_0)\bar{x}] > C \exp\left[n(\theta_1^2 - \theta_0^2)\right],$$

or

$$\begin{aligned} \bar{x} > C_0 \quad \text{for } \theta_1 > \theta_0, \quad \text{and} \quad \bar{x} < C_0 \quad \text{for } \theta_1 < \theta_0, \\ \text{where } C_0 = \frac{1}{2}(\theta_1 + \theta_0) + \frac{\log C}{n(\theta_1 - \theta_0)}. \end{aligned} \quad (30)$$

Then the requirement in (24) becomes, accordingly,

$$L_1 P_{\theta_0}(\bar{X} > C_0) = L_2 P_{\theta_1}(\bar{X} \leq C_0) \quad \text{for } \theta_1 > \theta_0,$$

and

$$L_1 P_{\theta_0}(\bar{X} < C_0) = L_2 P_{\theta_1}(\bar{X} \geq C_0) \quad \text{for } \theta_1 < \theta_0,$$

or

$$L_1 \{1 - \Phi[\sqrt{n}(C_0 - \theta_0)]\} = L_2 \Phi[\sqrt{n}(C_0 - \theta_1)] \quad \text{for } \theta_1 > \theta_0,$$

and

$$L_1 \Phi[\sqrt{n}(C_0 - \theta_0)] = L_2 \{1 - \Phi[\sqrt{n}(C_0 - \theta_1)]\} \quad \text{for } \theta_1 < \theta_0. \quad (31)$$

Consider the following numerical application.

Numerical Example Suppose $n = 25$ and let $\theta_0 = 0$ and $\theta_1 = 1$. In the spirit of Remark 1, take, e.g., $L_1 = 5$ and $L_2 = 2.5$.

DISCUSSION Then the first relation in (31), which is applicable here, becomes

$$\Phi[5(C_0 - 1)] = 2[1 - \Phi(5C_0)] \quad \text{or} \quad 2\Phi(5C_0) - \Phi(5 - 5C_0) = 1.$$

From the Normal tables, we find $C_0 = 0.53$, so that the minimax decision function is given by:

$$\delta(\mathbf{x}) = 1 \quad \text{if } \bar{x} > 0.53, \quad \text{and} \quad \delta(\mathbf{x}) = 0 \quad \text{if } \bar{x} \leq 0.53.$$

Let us now calculate the level of significance and the power of this test. We have

$$P_0(\bar{X} > 0.53) = 1 - \Phi(5 \times 0.53) = 1 - \Phi(2.65) = 1 - 0.995975 \simeq 0.004,$$

and

$$\pi(1) = P_1(\bar{X} > 0.53) = 1 - \Phi[5(0.53 - 1)] = \Phi(2.35) = 0.990613 \simeq 0.991.$$

EXAMPLE 3

In terms of the random sample X_1, \dots, X_n from the $B(1, \theta)$ distribution, determine the minimax function $\delta = \delta(\mathbf{x})$ for testing the hypothesis $H_0: \theta = \theta_0$ against the alternative $H_A: \theta = \theta_1$.

DISCUSSION The joint p.d.f. of the X_i 's is here

$$L(\mathbf{x}; \theta) = \theta^t (1 - \theta)^{n-t}, \quad t = x_1 + \dots + x_n,$$

so that the rejection region R is determined by

$$L(\mathbf{x}; \theta_1) > CL(\mathbf{x}; \theta_0) \quad \text{or} \quad [\theta_1(1 - \theta_0)/\theta_0(1 - \theta_1)]^t > C[(1 - \theta_0)/(1 - \theta_1)]^n,$$

or

$$t \log \frac{(1 - \theta_0)\theta_1}{\theta_0(1 - \theta_1)} > C'_0 = \log C - n \log \frac{1 - \theta_1}{1 - \theta_0}.$$

This is equivalent to

$$t > C_0 \quad \text{for } \theta_1 > \theta_0, \quad \text{and} \quad t < C_0 \quad \text{for } \theta_1 < \theta_0,$$

where $C_0 = C'_0 / \log \frac{(1 - \theta_0)\theta_1}{\theta_0(1 - \theta_1)}$. The requirement in (24) becomes here, respectively,

$$L_1 P_{\theta_0}(X > C_0) = L_2 P_{\theta_1}(X \leq C_0) \quad \text{for } \theta_1 > \theta_0,$$

and

$$L_1 P_{\theta_0}(X < C_0) = L_2 P_{\theta_1}(X \geq C_0) \quad \text{for } \theta_1 < \theta_0,$$

or

$$L_1 P_{\theta_0}(X \leq C_0) + L_2 P_{\theta_1}(X \leq C_0) = L_1 \quad \text{for } \theta_1 > \theta_0,$$

and

(32)

$$L_1 P_{\theta_0}(X \leq C_0 - 1) + L_2 P_{\theta_1}(X \leq C_0 - 1) = L_2 \quad \text{for } \theta_1 < \theta_0,$$

where $X \sim B(n, \theta)$.

Numerical Example Let $n = 20$, and suppose $\theta_0 = 0.50$ and $\theta_1 = 0.75$.

DISCUSSION Here, the first relation in (32) is applicable. Since

$$P_{0.75}(X \leq C_0) = P_{0.25}(X \geq 20 - C_0) = 1 - P_{0.25}(X \leq 19 - C_0), \quad (33)$$

the first relation in (32) becomes

$$L_1 P_{0.50}(X \leq C_0) - L_2 P_{0.25}(X \leq 19 - C_0) = L_1 - L_2,$$

or

$$L_2 = [1 - P_{0.50}(X \leq C_0)]L_1/[1 - P_{0.25}(X \leq 19 - C_0)]. \quad (34)$$

At this point, let us take $L_1 = 1$ and $L_2 = 0.269$. Then the right-hand side of (34) gives, for $C_0 = 13$; $\frac{1 - 0.9423}{1 - 0.7858} = \frac{0.0577}{0.2142} \simeq 0.269 = L_2$; i.e., the first relation in (32) obtains. The minimax decision function $\delta = \delta(\mathbf{x})$ is then given by: $\delta(\mathbf{x}) = 1$ if $x \geq 14$, and $\delta(\mathbf{x}) = 0$ for $x \leq 13$. The level of significance and the power of this test are:

$$P_{0.50}(X \geq 14) = 1 - P_{0.50}(X \leq 13) = 1 - 0.9423 = 0.0577,$$

and, on account of (33),

$$\pi(0.75) = P_{0.75}(X \geq 14) = P_{0.25}(X \leq 6) = 0.7858.$$

Instead of attempting to select $\delta = \delta(\mathbf{x})$ so as to minimize the maximum risk, we may, instead, try to determine δ so that δ minimizes the average risk. This approach calls for choosing the p.d.f.'s f_0 and f_1 according to a probability distribution; choose f_0 with probability p_0 and choose f_1 with probability p_1 ($p_0 + p_1 = 1$), and set $\lambda_0 = \{p_0, p_1\}$. If $R_{\lambda_0}(\delta)$ denotes the corresponding average risk, then, on account of (22), this average is given by:

$$\begin{aligned} R_{\lambda_0}(\delta) &= L_1 P_{f_0}(\mathbf{X} \in R) p_0 + L_2 P_{f_1}(\mathbf{X} \in R^c) p_1 \\ &= p_0 L_1 P_{f_0}(\mathbf{X} \in R) + p_1 L_2 [1 - P_{f_1}(\mathbf{X} \in R)] \\ &= p_1 L_2 + [p_0 L_1 P_{f_0}(\mathbf{X} \in R) - p_1 L_2 P_{f_1}(\mathbf{X} \in R)] \\ &= \begin{cases} p_1 L_2 + \int_R [p_0 L_1 f_0(x_1) \cdots f_0(x_n) - p_1 L_2 f_1(x_1) \cdots f_1(x_n)] dx_1 \cdots dx_n \\ p_1 L_2 + \sum_{\mathbf{x} \in R} [p_0 L_1 f_0(x_1) \cdots f_0(x_n) - p_1 L_2 f_1(x_1) \cdots f_1(x_n)] \end{cases} \end{aligned}$$

for the continuous and the discrete case, respectively. From this last expression, it follows that $R_{\lambda_0}(\delta)$ is minimized, if $p_0 L_1 f_0(x_1) \cdots f_0(x_n) - p_1 L_2 f_1(x_1) \cdots f_1(x_n)$ is < 0 on R . But $\delta(\mathbf{x}) = 1$ on R and $\delta(\mathbf{x}) = 0$ on R^c . Thus, we may restate these equations as follows:

$$\delta(\mathbf{x}) = \begin{cases} 1 & \text{if } f_1(x_1) \cdots f_1(x_n) > \frac{p_0 L_1}{p_1 L_2} f_0(x_1) \cdots f_0(x_n), \\ 0 & \text{otherwise.} \end{cases} \quad (35)$$

Thus, given a probability distribution $\lambda_0 = \{p_0, p_1\}$ on $\{f_0, f_1\}$, there is always a (nonrandomized) decision function δ which minimizes the average risk $R_{\lambda_0}(\delta)$, and this δ is given by (35) and is called a *Bayes decision function*.

THEOREM 3

The Bayes decision function $\delta_{\lambda_0}(\mathbf{x})$ corresponding to the probability distribution $\lambda_0 = \{p_0, p_1\}$ on $\{f_0, f_1\}$ is given by (35). This decision function is, actually, the MP test for testing the hypothesis $H_0: f = f_0$ against the alternative $H_A: f = f_1$ with cutoff point $C = p_0 L_1 / p_1 L_2$ and level of significance α given by:

$$P_{f_0}[f_1(X_1) \cdots f_1(X_n) > C f_0(X_1) \cdots f_0(X_n)] = \alpha. \quad (36)$$

REMARK 4 As mentioned earlier, if the underlying p.d.f. depends on a parameter $\theta \in \Omega$, then the above problem becomes that of testing $H_0: \theta = \theta_0$ against $H_A: \theta = \theta_1$ for some specified θ_0 and θ_1 in Ω .

EXAMPLE 4

DISCUSSION In reference to Example 2 and for the case that $\theta_1 > \theta_0$, $\delta_{\lambda_0}(\mathbf{x}) = 1$ if $\bar{x} > C_0$, $C_0 = \frac{1}{2}(\theta_1 + \theta_0) + \frac{\log C}{n(\theta_1 - \theta_0)}$, $C = p_0 L_1 / p_1 L_2$, as follows from relation (30). For the numerical data of the same example, we obtain $C_0 = 0.50 + 0.04 \log \frac{2p_0}{1-p_0}$. For example, for $p_0 = \frac{1}{2}$, C_0 is $\simeq 0.50 + 0.04 \times 0.693 = 0.52772 \simeq 0.53$, whereas for $p_0 = \frac{1}{4}$, C_0 is $\simeq 0.50 - 0.04 \times 0.405 = 0.4838 \simeq 0.48$. For $C_0 = 0.53$, the level of significance and the power have already

been calculated. For $C_0 = 0.48$, these quantities are, respectively:

$$P_0(\bar{X} > 0.48) = 1 - \Phi(5 \times 0.48) = 1 - \Phi(2.4) = 1 - 0.991802 = 0.008198,$$

$$\pi(1) = P_1(\bar{X} > 0.48) = 1 - \Phi[5(0.48 - 1)] = \Phi(2.6) = 0.995339.$$

In reference to Example 3 and for the case that $\theta_1 > \theta_0$, $\delta_{\lambda_0}(\mathbf{x}) = 1$ if $x > C_0$, $C_0 = (\log C - n \log \frac{1-\theta_1}{1-\theta_0}) / \log \frac{(1-\theta_0)\theta_1}{\theta_0(1-\theta_1)}$, $C = p_0 L_1 / p_1 L_2$. For the numerical data of the same example, we have $C_0 \simeq (15.173 + \log \frac{p_0}{1-p_0}) / 1.099$. For $p = \frac{1}{2}$, C_0 is 13.81, and for $p_0 = \frac{1}{4}$, C_0 is 12.81. In the former case, $\delta_{\lambda_0}(\mathbf{x}) = 1$ for $x \geq 14$, and in the latter case, $\delta_{\lambda_0}(\mathbf{x}) = 0$ for $x \leq 13$. The level of significance and the power have been calculated for the former case. As for the latter case, we have:

$$P_{0.50}(X \geq 13) = 1 - P_{0.50}(X \leq 12) = 1 - 0.8684 = 0.1316,$$

$$\pi(0.75) = P_{0.75}(X \geq 13) = P_{0.25}(X \leq 7) = 0.8982.$$

12.4 Relationship Between Testing Hypotheses and Confidence Regions

In this brief section, we discuss a relationship which connects a testing hypothesis problem and the problem of constructing a confidence region for the underlying parameter. To this effect, suppose X_1, \dots, X_n is a random sample from the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}^r$, $r \geq 1$, and for each θ in Ω , consider the problem of testing the hypothesis, to be denoted by $H_0(\theta)$, that the parameter θ^* , say, in Ω , is actually, equal to the value of θ considered. That is, $H_0(\theta): \theta^* = \theta$ at level of significance α . Denote by $A(\theta)$ the respective acceptance region in \mathfrak{R}^n . As usually, $\mathbf{X} = (X_1, \dots, X_n)$ and $\mathbf{x} = (x_1, \dots, x_n)$ is the observed value of \mathbf{X} . For each $\mathbf{x} \in \mathfrak{R}^n$, define in Ω the region $T(\mathbf{x})$ as follows:

$$T(\mathbf{x}) = \{\theta \in \Omega; \mathbf{x} \in A(\theta)\}. \quad (37)$$

Thus, $T(\mathbf{x})$ consists of all those $\theta \in \Omega$ for which, on the basis of the outcome \mathbf{x} , the hypothesis $H_0(\theta)$ is accepted. On the basis of the definition of $T(\mathbf{x})$ by (37), it is clear that

$$\theta \in T(\mathbf{x}) \text{ if and only if } \mathbf{x} \in A(\theta).$$

Therefore

$$P_\theta[\theta \in T(\mathbf{X})] = P_\theta[\mathbf{X} \in A(\theta)]. \quad (38)$$

But the probability on the right-hand side of (38) is equal to $1 - \alpha$, since the hypothesis $H_0(\theta)$ being tested is of level of significance α . Thus,

$$P_\theta[\theta \in T(\mathbf{X})] = 1 - \alpha,$$

and this means that the region $T(\mathbf{X})$ is a confidence region for θ with confidence coefficient $1 - \alpha$.

Summarizing what has been discussed so far in the form of a theorem, we have the following result.

THEOREM 4

Let X_1, \dots, X_n be a random sample from the p.d.f. $f(\cdot; \theta)$, $\theta \in \Omega \subseteq \mathfrak{R}^r$, $r \geq 1$, and, for each $\theta \in \Omega$, consider the problem of testing the hypothesis $H_0(\theta): \theta^* = \theta$ at level of significance α . Let $A(\theta)$ be the corresponding acceptance region in \mathfrak{R}^n , and for each $\mathbf{x} \in \mathfrak{R}^n$, define the region $T(\mathbf{x})$ in Ω as in (37). Then $T(\mathbf{X})$ is a confidence region for θ with confidence coefficient $1 - \alpha$, where $\mathbf{X} = (X_1, \dots, X_n)$ and $\mathbf{x} = (x_1, \dots, x_n)$ is the observed value of \mathbf{X} .

This result will now be illustrated below by two examples.

EXAMPLE 5

On the basis of a random sample X_1, \dots, X_n from the $N(\theta, \sigma^2)$ distribution with σ known, construct a confidence interval for θ with confidence coefficient $1 - \alpha$, by utilizing Theorem 4.

DISCUSSION For each $\theta \in \Omega = \mathfrak{R}$ and for testing the hypothesis $H_0(\theta)$ that the (unknown) parameter θ^* , say, is, actually, equal to θ , it makes sense to reject $H_0(\theta)$ when \bar{X} is either too far to the left or too far to the right of θ . Equivalently, if $\bar{X} - \theta$ is either $< C_1$ or $\bar{X} - \theta$ is $> C_2$ for some constants C_1, C_2 . If $H_0(\theta)$ is to be of level of significance α , we will have $P_\theta(\bar{X} - \theta < C_1 \text{ or } \bar{X} - \theta > C_2) = \alpha$. But under $H_0(\theta)$, the distribution of \bar{X} is symmetric about θ , so that it is reasonable to take $C_1 = -C_2$, and then $C_2 = z_{\alpha/2}$, $C_1 = -z_{\alpha/2}$. Thus, $H_0(\theta)$ is accepted whenever $-z_{\alpha/2} \leq \bar{x} - \theta \leq z_{\alpha/2}$ or $-z_{\alpha/2} \frac{\sqrt{n}}{\sigma} \leq \frac{\sqrt{n}(\bar{x} - \theta)}{\sigma} \leq z_{\alpha/2} \frac{\sqrt{n}}{\sigma}$, and, of course,

$$P_\theta \left[-z_{\alpha/2} \frac{\sqrt{n}}{\sigma} \leq \frac{\sqrt{n}(\bar{X} - \theta)}{\sigma} \leq z_{\alpha/2} \frac{\sqrt{n}}{\sigma} \right] = 1 - \alpha.$$

Thus,

$$A(\theta) = \left\{ \mathbf{x} \in \mathfrak{R}^n; -z_{\alpha/2} \frac{\sqrt{n}}{\sigma} \leq \frac{\sqrt{n}(\bar{x} - \theta)}{\sigma} \leq z_{\alpha/2} \frac{\sqrt{n}}{\sigma} \right\},$$

and therefore, by (37),

$$\begin{aligned} T(\mathbf{x}) &= \{ \theta \in \mathfrak{R}; \mathbf{x} \in A(\theta) \} \\ &= \left\{ \theta \in \mathfrak{R}; -z_{\alpha/2} \frac{\sqrt{n}}{\sigma} \leq \frac{\sqrt{n}(\bar{x} - \theta)}{\sigma} \leq z_{\alpha/2} \frac{\sqrt{n}}{\sigma} \right\} \\ &= \left\{ \theta \in \mathfrak{R}; \bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \theta \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right\}. \end{aligned}$$

In other words, we ended up with the familiar confidence interval for θ , $\bar{X} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$, we have already constructed in Chapter 10, Example 1(i).

EXAMPLE 6

Let the random sample X_1, \dots, X_n be from the $N(\mu, \sigma^2)$ distribution, where both μ and σ^2 are unknown. Construct a confidence interval for σ^2 with confidence coefficient $1 - \alpha$, again using Theorem 4.

DISCUSSION Here $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ is an estimate of σ^2 , and, therefore, for testing the hypothesis $H_0(\sigma^2)$: variance = σ^2 , it is reasonable to reject $H_0(\sigma^2)$ whenever the ratio of S^2 over the σ^2 specified by the hypothesis is either too small or too large. That is, reject $H_0(\sigma^2)$ when $\frac{S^2}{\sigma^2} < C_1$ or $\frac{S^2}{\sigma^2} > C_2$ for some (>0) constants C_1, C_2 to be specified by the requirement that $P_{\sigma^2}(\frac{S^2}{\sigma^2} < C_1$ or $\frac{S^2}{\sigma^2} > C_2) = \alpha$, or

$$P_{\sigma^2} \left[\frac{(n-1)S^2}{\sigma^2} < C'_1 \quad \text{or} \quad \frac{(n-1)S^2}{\sigma^2} > C'_2 \right] = \alpha,$$

$$C'_1 = \frac{(n-1)C_1}{\sigma^2}, \quad C'_2 = \frac{(n-1)C_2}{\sigma^2}.$$

Since under $H_0(\sigma^2)$, $\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$, we may choose to split the probability α equally between the two tails, in which case $C'_1 = \chi_{n-1; 1-\alpha/2}^2$ and $C'_2 = \chi_{n-1; \alpha/2}^2$, and $H(\sigma^2)$ is accepted whenever

$$\chi_{n-1; 1-\alpha/2}^2 \leq \frac{(n-1)S^2}{\sigma^2} \leq \chi_{n-1; \alpha/2}^2.$$

Of course,

$$P_{\sigma^2} \left[\chi_{n-1; 1-\alpha/2}^2 \leq \frac{(n-1)S^2}{\sigma^2} \leq \chi_{n-1; \alpha/2}^2 \right] = 1 - \alpha.$$

Then, with s^2 denoting the observed value of S^2 ,

$$A(\sigma^2) = \left\{ \mathbf{x} \in \mathfrak{R}^n; \chi_{n-1; 1-\alpha/2}^2 \leq \frac{(n-1)s^2}{\sigma^2} \leq \chi_{n-1; \alpha/2}^2 \right\},$$

and therefore (37) becomes here:

$$\begin{aligned} T(\mathbf{x}) &= \{ \sigma^2 \in (0, \infty); \mathbf{x} \in A(\sigma^2) \} \\ &= \left\{ \sigma^2 \in (0, \infty); \chi_{n-1; 1-\alpha/2}^2 \leq \frac{(n-1)s^2}{\sigma^2} \leq \chi_{n-1; \alpha/2}^2 \right\} \\ &= \left\{ \sigma^2 \in (0, \infty); \frac{(n-1)s^2}{\chi_{n-1; \alpha/2}^2} \leq \sigma^2 \leq \frac{(n-1)s^2}{\chi_{n-1; 1-\alpha/2}^2} \right\}; \end{aligned}$$

that is, we have arrived once again at the familiar confidence interval for σ^2 , $\left[\frac{(n-1)s^2}{\chi_{n-1; \alpha/2}^2}, \frac{(n-1)s^2}{\chi_{n-1; 1-\alpha/2}^2} \right]$, (see Example 3 in Chapter 10).



Chapter 13

A Simple Linear Regression Model

This is a rather extensive chapter on an important subject matter with an abundance of diverse applications. The basic idea involved may be described as follows. There is a stimulus, denoted by x , and a response to it, denoted by y . At different levels of x , one observes the respective responses. How are the resulting (x, y) pairs related, if they are related at all? There are all kind of possibilities, and the one discussed in this chapter is the simplest such possibility, namely, the pairs are linearly related.

In reality, what one, actually, observes at x , due to errors, is a value of a r.v. Y , and then the question arises as to how we would draw a straight line, which would lie “close” to most of the (x, y) pairs. This leads to the Principle of Least Squares. On the basis of this principle, one is able to draw the so-called fitted linear regression line by computing the Least Squares Estimates of parameters involved. Also, some properties of these estimates are established. These things are done in the first two sections of the chapter.

Up to this point, the errors are not required to have any specific distribution, other than having zero mean and finite variance. However, in order to proceed with statistical inference about the parameters involved, such as constructing confidence intervals and testing hypotheses, one has to stipulate a distribution for the errors; this distribution, reasonably enough, is assumed to be Normal. As a consequence of it, one is in a position to specify the distribution of all estimates involved and proceed with the inference problems referred to above. These issues are discussed in Sections 13.3 and 13.4.

In the following section, Section 13.5, the problem of predicting the expected value of the observation Y_0 at a given point x_0 and the problem of predicting a single value of Y_0 are discussed. Suitable predictors are provided, and also confidence intervals for them are constructed.

The chapter is concluded with Section 3.7 indicating extensions of the model discussed in this chapter to more general situations covering a much wider class of applications.

13.1 Setting up the Model – The Principle of Least Squares

As has already been mentioned, Examples 22 and 23 in Chapter 1 provide motivation for the statistical model to be adopted and studied in this chapter. Example 22, in particular, will serve throughout the chapter to illustrate the underlying general results. For convenience, the data related to this example are reproduced here in Table 13.1.

Table 13.1


The Data $x =$
Undergraduate GPA and
 $y =$ Score in the
Graduate Management
Aptitude Test (GMAT);
There Are 34 (x, y)
Pairs Altogether

DATA OF UNDERGRADUATE GPA (x) AND GMAT SCORE (y)					
x	y	x	y	x	y
3.63	447	2.36	399	2.80	444
3.59	588	2.36	482	3.13	416
3.30	563	2.66	420	3.01	471
3.40	553	2.68	414	2.79	490
3.50	572	2.48	533	2.89	431
3.78	591	2.46	509	2.91	446
3.44	692	2.63	504	2.75	546
3.48	528	2.44	336	2.73	467
3.47	552	2.13	408	3.12	463
3.35	520	2.41	469	3.08	440
3.39	543	2.55	538	3.03	419
				3.00	509

The first question which arises is whether the pairs (x, y) are related at all and, if they are, how. An indication that those pairs are, indeed, related is borne out by the scatter plot depicted in Figure 13.1. Indeed, taking into consideration that we are operating in a random environment, one sees a conspicuous, albeit somewhat loose, linear relationship between the pairs (x, y) .

Figure 13.1

Scatter Diagram for
Table 13.1



So, we are not too far off the target by assuming that there is a straight line in the xy -plane which is “close” to most of the pairs (x, y) . The question now is how to quantify the term “close.” The first step toward this end is the adoption of the model described in relation (11) of Chapter 8. Namely, we assume that, for each $i = 1, \dots, 34$, the respective y_i is the observed value of a r.v. Y_i associated with x_i , and if it were not for the random errors involved, the pairs (x_i, y_i) , $i = 1, \dots, 34$ would lie on a straight line $y = \beta_1 + \beta_2 x$; i.e., we would have $y_i = \beta_1 + \beta_2 x_i$, $i = 1, \dots, 34$. Thus, the r.v. Y_i itself, whose y_i are simply observed values, would be equal to $\beta_1 + \beta_2 x_i$ except for fluctuations due to a random error e_i . In other words, $Y_i = \beta_1 + \beta_2 x_i + e_i$. Next, arguing as in Section 8.4, it is reasonable to assume that the e_i 's are independent r.v.'s with $Ee_i = 0$ and $\text{Var}(e_i) = \sigma^2$ for all i 's, so that one arrives at the model described in relation (11) of Chapter 8; namely, Y_1, \dots, Y_{34} are independent r.v.'s having the structure:

$$Y_i = \beta_1 + \beta_2 x_i + e_i, \quad \text{with } Ee_i = 0 \quad \text{and} \quad \text{Var}(e_i) = \sigma^2, \quad i = 1, \dots, 34. \quad (1)$$

Set $EY_i = \eta_i$. Then, because of the errors involved, it is, actually, the pairs (x_i, η_i) , $i = 1, \dots, 34$ which lie on a straight line $y = \beta_1 + \beta_2 x$; i.e., $\eta_i = \beta_1 + \beta_2 x_i$, $i = 1, \dots, 34$. It is in the determination of a particular straight line where the Principle of Least Squares enters the picture. According to this principle, one argues as follows: On the basis of the model described in (1), what we would *expect* to have observed at x_i would be η_i , whereas what is, *actually*, observed is y_i . Thus, there is a deviation measured by $y_i - \eta_i$, $i = 1, \dots, 34$ (see Figure 13.1). Some of these deviations are positive, some are negative, and, perhaps, some are zero. In order to deal with nonnegative numbers, look at $|y_i - \eta_i|$, which is, actually, the distance between the points (x_i, y_i) and (x_i, η_i) . Then, draw the line $y = \beta_1 + \beta_2 x$, so that these distances are simultaneously minimized. More formally, first look at the squares of these distances $(y_i - \eta_i)^2$, as it is much easier to work with squares as opposed to absolute values, and in order to account for the simultaneous minimization mentioned earlier, consider the sum $\sum_{i=1}^{34} (y_i - \eta_i)^2$ and seek its minimization. At this point, replace the observed value y_i by the r.v. Y_i itself and set

$$S(\mathbf{Y}, \boldsymbol{\beta}) = \sum_{i=1}^{34} (Y_i - \eta_i)^2 = \sum_{i=1}^{34} [Y_i - (\beta_1 + \beta_2 x_i)]^2 \left(= \sum_{i=1}^{34} e_i^2 \right), \quad (2)$$

where $\mathbf{Y} = (Y_1, \dots, Y_{34})$ and $\boldsymbol{\beta} = (\beta_1, \beta_2)$.

Then the *Principle of Least Squares* calls for the determination of β_1 and β_2 which *minimize the sum of squares of errors*; i.e., the quantity $S(\mathbf{Y}, \boldsymbol{\beta})$ in (2). The actual minimization is a calculus problem. If there is a unique straight line so determined, then, clearly, this would be the line which lies “close” to most pairs (x_i, Y_i) , $i = 1, \dots, 34$, in the Least Squares sense. It will be seen below that this is, indeed, the case.

In a more general setting, consider the model below:

$Y_i = \beta_1 + \beta_2 x_i + e_i$, where the random errors e_i , $i = 1, \dots, n$ are i.i.d. r.v.'s with

$Ee_i = 0$, and $\text{Var}(e_i) = \sigma^2$, which imply that the r.v.'s Y_i , $i = 1, \dots, n$ are independent, but not identically distributed, with $EY_i = \eta_i = \beta_1 + \beta_2 x_i$ and $\text{Var}(y_i) = \sigma^2$. (3)

Let $\hat{\beta}_1$ and $\hat{\beta}_2$ be the unique values of β_1 and β_2 , respectively, which minimize the sum of squares of errors $S(\mathbf{Y}, \boldsymbol{\beta}) = \sum_{i=1}^n [Y_i - (\beta_1 + \beta_2 x_i)]^2 (= \sum_{i=1}^n e_i^2)$. These values, which are functions of the Y_i 's as well as the x_i 's, are the *Least Squares Estimates* (LSE's) of β_1 and β_2 . Any line $y = \beta_1 + \beta_2 x$ is referred to as a *regression line* and, in particular, the line $\hat{y} = \hat{\beta}_1 + \hat{\beta}_2 x$ is known as the *fitted regression line*. For this line, the \hat{y}_i 's corresponding to the x_i 's are $\hat{y}_i = \hat{\beta}_1 + \hat{\beta}_2 x_i$, $i = 1, \dots, n$.

13.2 The Least Squares Estimates of β_1 and β_2 and Some of their Properties

In this section, the LSE's of β_1 and β_2 are derived and some of their properties are obtained. Also, the (unknown) variance σ^2 is estimated.

THEOREM 1

In reference to the model described in (3), the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ of β_1 and β_2 , respectively, are given by the following expressions (which are also appropriate for computational purposes):

$$\hat{\beta}_1 = \frac{(\sum_{i=1}^n x_i^2)(\sum_{i=1}^n Y_i) - (\sum_{i=1}^n x_i)(\sum_{i=1}^n x_i Y_i)}{n \sum_{i=1}^n x_i^2 - (\sum_{i=1}^n x_i)^2}, \quad (4)$$

and

$$\hat{\beta}_2 = \frac{n \sum_{i=1}^n x_i Y_i - (\sum_{i=1}^n x_i)(\sum_{i=1}^n Y_i)}{n \sum_{i=1}^n x_i^2 - (\sum_{i=1}^n x_i)^2}. \quad (5)$$

PROOF Consider the partial derivatives:

$$\frac{\partial}{\partial \beta_1} S(\mathbf{Y}, \boldsymbol{\beta}) = 2 \sum_{i=1}^n (Y_i - \beta_1 - \beta_2 x_i)(-1) = -2 \left(\sum_{i=1}^n Y_i - n\beta_1 - \beta_2 \sum_{i=1}^n x_i \right), \quad (6)$$

$$\begin{aligned} \frac{\partial}{\partial \beta_2} S(\mathbf{Y}, \boldsymbol{\beta}) &= 2 \sum_{i=1}^n (Y_i - \beta_1 - \beta_2 x_i)(-x_i) \\ &= -2 \left(\sum_{i=1}^n x_i Y_i - \beta_1 \sum_{i=1}^n x_i - \beta_2 \sum_{i=1}^n x_i^2 \right), \end{aligned} \quad (7)$$

and solve the so-called *normal equations*: $\frac{\partial}{\partial \beta_1} S(\mathbf{Y}, \boldsymbol{\beta}) = 0$ and $\frac{\partial}{\partial \beta_2} S(\mathbf{Y}, \boldsymbol{\beta}) = 0$, or $n\beta_1 + (\sum_{i=1}^n x_i)\beta_2 = \sum_{i=1}^n Y_i$ and $(\sum_{i=1}^n x_i)\beta_1 + (\sum_{i=1}^n x_i^2)\beta_2 = \sum_{i=1}^n x_i Y_i$ to find:

$$\hat{\beta}_1 = \frac{\begin{vmatrix} \sum_{i=1}^n Y_i & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i Y_i & \sum_{i=1}^n x_i^2 \end{vmatrix}}{\begin{vmatrix} n & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i^2 \end{vmatrix}} = \frac{(\sum_{i=1}^n x_i^2)(\sum_{i=1}^n Y_i) - (\sum_{i=1}^n x_i)(\sum_{i=1}^n x_i Y_i)}{n \sum_{i=1}^n x_i^2 - (\sum_{i=1}^n x_i)^2},$$

and

$$\hat{\beta}_2 = \frac{\left| \begin{array}{cc} n & \sum_{i=1}^n Y_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i Y_i \end{array} \right|}{n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2} = \frac{n \sum_{i=1}^n x_i Y_i - \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n Y_i \right)}{n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2}.$$

It remains to show that $\hat{\beta}_1$ and $\hat{\beta}_2$, actually, minimize $\mathcal{S}(\mathbf{Y}, \beta)$. From (6) and (7), we get:

$$\begin{aligned} \frac{\partial^2}{\partial \beta_1^2} \mathcal{S}(\mathbf{Y}, \beta) &= 2n, & \frac{\partial^2}{\partial \beta_1 \partial \beta_2} \mathcal{S}(\mathbf{Y}, \beta) &= \frac{\partial^2}{\partial \beta_2 \partial \beta_1} \mathcal{S}(\mathbf{Y}, \beta) = 2 \sum_{i=1}^n x_i, \\ \frac{\partial^2}{\partial \beta_2^2} \mathcal{S}(\mathbf{Y}, \beta) &= 2 \sum_{i=1}^n x_i^2, \end{aligned}$$

and the 2×2 matrix below is positive semidefinite for all β_1, β_2 , since, for all λ_1, λ_2 reals not both 0,

$$\begin{aligned} &(\lambda_1, \lambda_2) \begin{pmatrix} n & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i^2 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} \\ &= \begin{pmatrix} \lambda_1 n + \lambda_2 \sum_{i=1}^n x_i & \lambda_1 \sum_{i=1}^n x_i + \lambda_2 \sum_{i=1}^n x_i^2 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} \\ &= \lambda_1^2 n + 2\lambda_1 \lambda_2 \sum_{i=1}^n x_i + \lambda_2^2 \sum_{i=1}^n x_i^2 \\ &= \lambda_1^2 n + 2n\lambda_1 \lambda_2 \bar{x} + \lambda_2^2 \sum_{i=1}^n x_i^2 \quad \left(\text{where } \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \right) \\ &= \lambda_1^2 n + 2n\lambda_1 \lambda_2 \bar{x} + \lambda_2^2 \left(\sum_{i=1}^n x_i^2 - n\bar{x}^2 \right) + \lambda_2^2 n\bar{x}^2 \\ &= n(\lambda_1^2 + 2\lambda_1 \lambda_2 \bar{x} + \lambda_2^2 \bar{x}^2) + \lambda_2^2 \sum_{i=1}^n (x_i - \bar{x})^2 \\ &= n(\lambda_1 + \lambda_2 \bar{x})^2 + \lambda_2^2 \sum_{i=1}^n (x_i - \bar{x})^2 \geq 0. \end{aligned}$$

This completes the proof of the theorem. \blacktriangle

COROLLARY With $\bar{x} = (x_1 + \cdots + x_n)/n$ and $\bar{Y} = (Y_1 + \cdots + Y_n)/n$, the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ may also be written as follows (useful expressions for noncomputational purposes):

$$\hat{\beta}_1 = \bar{Y} - \hat{\beta}_2 \bar{x}, \quad \hat{\beta}_2 = \frac{\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{1}{\sum_{i=1}^n (x_i - \bar{x})^2} \sum_{i=1}^n (x_i - \bar{x}) Y_i. \quad (8)$$

PROOF First,

$$\begin{aligned} \sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y}) &= \sum_{i=1}^n x_i Y_i - \bar{Y} \sum_{i=1}^n x_i - \bar{x} \sum_{i=1}^n Y_i + n\bar{x}\bar{Y} \\ &= \sum_{i=1}^n x_i Y_i - \frac{1}{n} \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n Y_i \right) - n\bar{x}\bar{Y} + n\bar{x}\bar{Y} \\ &= \frac{1}{n} \left[n \sum_{i=1}^n x_i Y_i - \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n Y_i \right) \right], \end{aligned}$$

and

$$\begin{aligned} n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2 &= n \sum_{i=1}^n x_i^2 - (n\bar{x})^2 = n \left(\sum_{i=1}^n x_i^2 - n\bar{x}^2 \right) \\ &= n \sum_{i=1}^n (x_i - \bar{x})^2. \end{aligned}$$

Therefore

$$\begin{aligned} \frac{\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{\sum_{i=1}^n (x_i - \bar{x})^2} &= \frac{n \sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{n \sum_{i=1}^n (x_i - \bar{x})^2} \\ &= \frac{n \sum_{i=1}^n x_i Y_i - \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n Y_i \right)}{n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2} = \hat{\beta}_2, \text{ on account of (5)}. \end{aligned}$$

The second expression for $\hat{\beta}_2$ as a linear combination of the Y_i 's follows, because

$$\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y}) = \sum_{i=1}^n (x_i - \bar{x})Y_i - \bar{Y} \sum_{i=1}^n (x_i - \bar{x}) = \sum_{i=1}^n (x_i - \bar{x})Y_i,$$

since

$$\sum_{i=1}^n (x_i - \bar{x}) = \sum_{i=1}^n x_i - n\bar{x} = \sum_{i=1}^n x_i - \sum_{i=1}^n x_i = 0.$$

Also,

$$\begin{aligned} \bar{Y} - \hat{\beta}_2 \bar{x} &= \bar{Y} - \frac{n \sum_{i=1}^n x_i Y_i - \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n Y_i \right)}{n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2} \bar{x} \\ &= \bar{Y} - \frac{n \sum_{i=1}^n x_i Y_i - n^2 \bar{x} \bar{Y}}{n \sum_{i=1}^n x_i^2 - n^2 \bar{x}^2} \bar{x} \\ &= \frac{n \bar{Y} \sum_{i=1}^n x_i^2 - n^2 \bar{x}^2 \bar{Y} - n \bar{x} \sum_{i=1}^n x_i Y_i + n^2 \bar{x}^2 \bar{Y}}{n \sum_{i=1}^n (x_i - \bar{x})^2} \\ &= \frac{\left(\sum_{i=1}^n x_i^2 \right) \left(\sum_{i=1}^n Y_i \right) - \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n x_i Y_i \right)}{n \sum_{i=1}^n (x_i - \bar{x})^2} = \hat{\beta}_1 \text{ by (4)}. \quad \blacktriangle \end{aligned}$$

The following notation is suggested (at least in part) by the expressions in the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$, and it will be used extensively and conveniently throughout the rest of this chapter.

Set

$$SS_x = \sum_{i=1}^n (x_i - \bar{x})^2 = \sum_{i=1}^n x_i^2 - n\bar{x}^2 = \sum_{i=1}^n x_i^2 - \frac{1}{n} \left(\sum_{i=1}^n x_i \right)^2,$$

and likewise

$$SS_y = \sum_{i=1}^n (Y_i - \bar{Y})^2 = \sum_{i=1}^n Y_i^2 - n\bar{Y}^2 = \sum_{i=1}^n Y_i^2 - \frac{1}{n} \left(\sum_{i=1}^n Y_i \right)^2, \quad (9)$$

and

$$\begin{aligned} SS_{xy} &= \sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y}) = \sum_{i=1}^n (x_i - \bar{x})Y_i \\ &= \sum_{i=1}^n x_i Y_i - \frac{1}{n} \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n Y_i \right). \end{aligned}$$

Then the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ may be rewritten as follows:

$$\hat{\beta}_1 = \frac{1}{n} \sum_{i=1}^n Y_i - \hat{\beta}_2 \left(\frac{1}{n} \sum_{i=1}^n x_i \right), \quad \hat{\beta}_2 = \frac{SS_{xy}}{SS_x}. \quad (10)$$

Also, recall that the fitted regression line is given by:

$$\hat{y} = \hat{\beta}_1 + \hat{\beta}_2 x \quad \text{and that} \quad \hat{y}_i = \hat{\beta}_1 + \hat{\beta}_2 x_i, \quad i = 1, \dots, n. \quad (11)$$

Before we go any further, let us discuss the example below.

EXAMPLE 1

In reference to Table 13.1, compute the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ and draw the fitted regression line $\hat{y} = \hat{\beta}_1 + \hat{\beta}_2 x$.

DISCUSSION The application of formula (10) calls for the calculation of SS_x and SS_{xy} given in (9). Table 13.2 facilitates the calculations.

$$\sum_i x_i = 100.73, \quad \sum_i y_i = 16,703, \quad \sum_i x_i^2 = 304.7885, \quad \sum_i x_i y_i = 50,066.47,$$

and then

$$\begin{aligned} SS_x &= 304.7885 - \frac{(100.73)^2}{34} \simeq 304.7885 - 298.4274 \simeq 6.361, \\ SS_{xy} &= 50,066.47 - \frac{(100.73) \times (16,703)}{34} \simeq 50,066.47 - 49,485.094 \\ &= 581.376. \end{aligned}$$

Table 13.2

	x	y	x^2	xy	x	y	x^2	xy
	3.63	447	13.1769	1,622.61	3.48	528	12.1104	1,837.44
	3.59	588	12.8881	2,110.92	3.47	552	12.0409	1,915.44
	3.30	563	10.8900	1,857.90	3.35	520	11.2225	1,742.00
	3.40	553	11.5600	1,880.20	3.39	543	11.4921	1,840.77
	3.50	572	12.2500	2,002.00	2.36	399	5.5696	941.64
	3.78	591	14.2884	2,233.98	2.36	482	5.5696	1,137.52
	3.44	692	11.8336	2,380.48	2.66	420	7.0756	1,117.20
	2.68	414	7.1824	1,109.52	3.01	471	9.0601	1,417.71
	2.48	533	6.1504	1,321.84	2.79	490	7.7841	1,367.10
	2.46	509	6.0516	1,252.14	2.89	431	8.3521	1,245.59
	2.63	504	6.9169	1,325.52	2.91	446	8.4681	1,297.86
	2.44	336	5.9536	819.84	2.75	546	7.5625	1,501.50
	2.13	408	4.5369	869.04	2.73	467	7.4529	1,274.91
	2.41	469	5.8081	1,130.29	3.12	463	9.7344	1,444.56
	2.55	538	6.5025	1,371.90	3.08	440	9.4864	1,355.20
	2.80	444	7.8400	1,243.20	3.03	419	9.1809	1,269.57
	3.13	416	9.7969	1,302.08	3.00	509	9.0000	1,527.00
Totals	50.35	8,577	153.6263	25,833.46	50.38	8,126	151.1622	24,233.01

Then


$$\hat{\beta}_2 = \frac{581.376}{6.361} \simeq 91.397 \quad \text{and} \quad \hat{\beta}_1 = \frac{16,703}{34} - (91.397) \times \frac{100.73}{34}$$

$$\simeq 491.265 - 270.809 = 220.456,$$

and the fitted regression line $\hat{y} = 220.456 + 91.397x$ is depicted in the Figure 13.2.

Figure 13.2

The Fitted
Regression Line $\hat{y} =$
 $220.456 + 91.397x$



The LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ have the desirable property of being unbiased, as shown in the following theorem.

THEOREM 2

The LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ are unbiased; i.e., $E\hat{\beta}_1 = \beta_1$ and $E\hat{\beta}_2 = \beta_2$. Furthermore,

$$\text{Var}(\hat{\beta}_1) = \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{SS_x} \right) \quad \text{and} \quad \text{Var}(\hat{\beta}_2) = \frac{\sigma^2}{SS_x},$$

where SS_x is given in (9).

PROOF In this proof and also elsewhere, the range of the summation is not explicitly indicated, since it is always from 1 to n . Consider $\hat{\beta}_2$ as given in (8). Then: $SS_x \hat{\beta}_2 = \sum_i (x_i - \bar{x}) Y_i$, so that, by taking expectations:

$$\begin{aligned} SS_x E\hat{\beta}_2 &= \sum_i (x_i - \bar{x}) EY_i = \sum_i (x_i - \bar{x})(\beta_1 + \beta_2 x_i) \\ &= \beta_1 \sum_i (x_i - \bar{x}) + \beta_2 \sum_i x_i (x_i - \bar{x}) = \beta_2 \sum_i x_i (x_i - \bar{x}) \\ &= \beta_2 \left(\sum_i x_i^2 - n\bar{x}^2 \right) = \beta_2 \sum_i (x_i - \bar{x})^2 = SS_x \beta_2. \end{aligned}$$

Therefore, dividing through by SS_x , we get $E\hat{\beta}_2 = \beta_2$. Next, also from (8),

$$\begin{aligned} E\hat{\beta}_1 &= E(\bar{Y} - \hat{\beta}_2 \bar{x}) = E\bar{Y} - \bar{x} E\hat{\beta}_2 = \frac{1}{n} \sum_i (\beta_1 + \beta_2 x_i) - \bar{x} \beta_2 \\ &= \beta_1 + \beta_2 \bar{x} - \beta_2 \bar{x} = \beta_1. \end{aligned}$$

Regarding the variances, we have from (8): $SS_x \hat{\beta}_2 = \sum_i (x_i - \bar{x}) Y_i$, so that:

$$\begin{aligned} SS_x^2 \text{Var}(\hat{\beta}_2) &= \text{Var} \left(\sum_i (x_i - \bar{x}) Y_i \right) = \sum_i (x_i - \bar{x})^2 \text{Var}(Y_i) \\ &= \sigma^2 \sum_i (x_i - \bar{x})^2 = \sigma^2 SS_x, \end{aligned}$$

so that $\text{Var}(\hat{\beta}_2) = \sigma^2 / SS_x$. Finally, from (8),

$$\hat{\beta}_1 = \bar{Y} - \bar{x} \hat{\beta}_2 = \frac{1}{n} \sum_i Y_i - \frac{\bar{x}}{SS_x} \sum_i (x_i - \bar{x}) Y_i = \sum_i \left[\frac{1}{n} - \frac{\bar{x}(x_i - \bar{x})}{SS_x} \right] Y_i, \quad (12)$$

so that

$$\text{Var}(\hat{\beta}_1) = \sigma^2 \sum_i \left[\frac{1}{n} - \frac{\bar{x}(x_i - \bar{x})}{SS_x} \right]^2 = \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{SS_x^2} SS_x \right) = \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{SS_x} \right). \quad \blacktriangle$$

EXAMPLE 2

DISCUSSION In reference to Example 1, the variances of the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ are given by:

$$\text{Var}(\hat{\beta}_1) \simeq \sigma^2 \left(\frac{1}{34} + \frac{8.777}{6.361} \right) \simeq \sigma^2(0.029 + 1.380) = 1.409\sigma^2,$$

and

$$\text{Var}(\hat{\beta}_2) = \frac{\sigma^2}{6.361} \simeq 0.157\sigma^2.$$

In fitting a regression line, there are various deviations which occur. At this point, these deviations will be suitably attributed to several sources, certain pieces of terminology will be introduced, and also some formal relations will be established. To this end, look at the observable Y_i and split it as follows: $Y_i = \hat{y}_i + (Y_i - \hat{y}_i)$. The component \hat{y}_i represents the point (x_i, \hat{y}_i) which lies on the fitted regression line $\hat{y} = \hat{\beta}_1 + \hat{\beta}_2 x$, and the difference $Y_i - \hat{y}_i$ is the deviation of Y_i from \hat{y}_i . We may refer to the component \hat{y}_i as that part of Y_i which is *due to the linear regression*, or it is *explained by the linear regression*, and the component $Y_i - \hat{y}_i$ of Y_i as the *residual*, or the *deviation from the linear regression*, or *variability unexplained by the linear regression*. We can go through the same arguments with reference to the sample mean \bar{Y} of the Y_i 's. That is, we consider:

$$Y_i - \bar{Y} = (\hat{y}_i - \bar{Y}) + (Y_i - \hat{y}_i).$$

The interpretation of this decomposition is the same as the one given above, but with reference to \bar{Y} . Next, look at the squares of these quantities:

$$(Y_i - \bar{Y})^2, \quad (\hat{y}_i - \bar{Y})^2, \quad (Y_i - \hat{y}_i)^2,$$

and, finally, at their sums:

$$\sum_{i=1}^n (Y_i - \bar{Y})^2, \quad \sum_{i=1}^n (\hat{y}_i - \bar{Y})^2, \quad \sum_{i=1}^n (Y_i - \hat{y}_i)^2.$$

At this point, assume for a moment that:

$$\sum_{i=1}^n (Y_i - \bar{Y})^2 = \sum_{i=1}^n (\hat{y}_i - \bar{Y})^2 + \sum_{i=1}^n (Y_i - \hat{y}_i)^2. \quad (13)$$

Then this relation would state that the *total variability* (of the Y_i 's in reference to their mean \bar{Y}), $\sum_{i=1}^n (Y_i - \bar{Y})^2$, is the sum of the *variability* $\sum_{i=1}^n (\hat{y}_i - \bar{Y})^2$ *due to the linear regression*, or *explained by the linear regression*, and the *residual variability*, $\sum_{i=1}^n (Y_i - \hat{y}_i)^2$, or *variability unexplained by the linear regression*.

We proceed in proving relation (13).

THEOREM 3

Let $SS_T (= SS_y)$, see (9), SS_R , and SS_E , respectively, be the total variability, the variability due to the linear regression (or explained by the linear regression), and the residual variability (or variability not explained by the linear regression); i.e.,

$$SS_T (= SS_y) = \sum_{i=1}^n (Y_i - \bar{Y})^2, \quad SS_R = \sum_{i=1}^n (\hat{y}_i - \bar{Y})^2, \quad SS_E = \sum_{i=1}^n (Y_i - \hat{y}_i)^2, \quad (14)$$

where $\hat{y}_i = \hat{\beta}_1 + \hat{\beta}_2 x_i$, $i = 1, \dots, n$, the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ are given by (10) (or (8)), and \bar{Y} is the mean of the Y_i 's. Then:

$$(i) \quad SS_T = SS_R + SS_E. \quad (15)$$

Furthermore,

$$(ii) \quad SS_T = SS_y, \quad SS_R = \frac{SS_{xy}^2}{SS_x}, \quad \text{and hence} \quad SS_E = SS_y - \frac{SS_{xy}^2}{SS_x}, \quad (16)$$

where SS_x , SS_y , and SS_{xy} are given in (9).

PROOF We have:

$$\begin{aligned} SS_T &= \sum_i (Y_i - \bar{Y})^2 = \sum_i [(\hat{y}_i - \bar{Y}) + (Y_i - \hat{y}_i)]^2 \\ &= \sum_i (\hat{y}_i - \bar{Y})^2 + \sum_i (Y_i - \hat{y}_i)^2 + 2 \sum_i (\hat{y}_i - \bar{Y})(Y_i - \hat{y}_i) \\ &= SS_R + SS_E + 2 \sum_i (\hat{y}_i - \bar{Y})(Y_i - \hat{y}_i). \end{aligned}$$

So, we have to show that the last term on the right-hand side above is equal to 0. To this end, observe that $\hat{y}_i = \hat{\beta}_1 + \hat{\beta}_2 x_i$ and $\hat{\beta}_1 = \bar{Y} - \hat{\beta}_2 \bar{x}$ (by (8)), so that

$$\hat{y}_i - \bar{Y} = \hat{\beta}_1 + \hat{\beta}_2 x_i - \bar{Y} = \bar{Y} - \hat{\beta}_2 \bar{x} + \hat{\beta}_2 x_i - \bar{Y} = \hat{\beta}_2 (x_i - \bar{x}),$$

and

$$Y_i - \hat{y}_i = Y_i - \hat{\beta}_1 - \hat{\beta}_2 x_i = Y_i - \bar{Y} + \hat{\beta}_2 \bar{x} - \hat{\beta}_2 x_i = (Y_i - \bar{Y}) - \hat{\beta}_2 (x_i - \bar{x}),$$

so that

$$\begin{aligned} (\hat{y}_i - \bar{Y})(Y_i - \hat{y}_i) &= \hat{\beta}_2 (x_i - \bar{x}) [(Y_i - \bar{Y}) - \hat{\beta}_2 (x_i - \bar{x})] \\ &= \hat{\beta}_2 (x_i - \bar{x})(Y_i - \bar{Y}) - \hat{\beta}_2^2 (x_i - \bar{x})^2. \end{aligned}$$

Therefore, by (9) and (10):

$$\sum_i (\hat{y}_i - \bar{Y})(Y_i - \hat{y}_i) = \frac{SS_{xy}}{SS_x} \times SS_{xy} - \frac{SS_{xy}^2}{SS_x^2} \times SS_x = \frac{SS_{xy}^2}{SS_x} - \frac{SS_{xy}^2}{SS_x} = 0. \quad (17)$$

Thus, $SS_T = SS_R + SS_E$.

(ii) That $SS_T = SS_y$ is immediate from relations (9) and (14). Next,

$$\hat{y}_i - \bar{Y} = \hat{\beta}_1 + \hat{\beta}_2 x_i - \bar{Y} = \bar{Y} - \hat{\beta}_2 \bar{x} + \hat{\beta}_2 x_i - \bar{Y} = \hat{\beta}_2 (x_i - \bar{x}) \quad (\text{by (8) and (11)}),$$

so that, by (10),

$$SS_R = \sum_i (\hat{y}_i - \bar{Y})^2 = \hat{\beta}_2^2 \sum_i (x_i - \bar{x})^2 = \frac{SS_{xy}^2}{SS_x^2} \times SS_x = \frac{SS_{xy}^2}{SS_x},$$

as was to be seen. Finally,

$$\begin{aligned} SS_E &= \sum_i (Y_i - \hat{y}_i)^2 = \sum_i [(Y_i - \bar{Y}) - (\hat{y}_i - \bar{Y})]^2 \\ &= \sum_i (Y_i - \bar{Y})^2 + \sum_i (\hat{y}_i - \bar{Y})^2 - 2 \sum_i (\hat{y}_i - \bar{Y})(Y_i - \bar{Y}) \\ &= SS_T + SS_R - 2 \sum_i (\hat{y}_i - \bar{Y})(Y_i - \bar{Y}), \end{aligned}$$

and

$$\begin{aligned} \sum_i (\hat{y}_i - \bar{Y})(Y_i - \bar{Y}) &= \sum_i (\hat{y}_i - \bar{Y})[(Y_i - \hat{y}_i) + (\hat{y}_i - \bar{Y})] \\ &= \sum_i (\hat{y}_i - \bar{Y})(Y_i - \hat{y}_i) + \sum_i (\hat{y}_i - \bar{Y})^2 = SS_R \quad (\text{by (17)}). \end{aligned}$$

It follows that $SS_E = SS_T - SS_R = SS_y - \frac{SS_{xy}^2}{SS_x}$, as was to be seen. ▲

This section is closed with some remarks.

REMARK 1

- (i) The quantities SS_T , SS_R , and SS_E , given in (14), are computed by way of SS_x , SS_y , and SS_{xy} given in (9). This is so because of (16).
- (ii) In the next section, an estimate of the (unknown) variance σ^2 will also be given, based on the residual variability SS_E . That this should be the case is intuitively clear by the nature of SS_E , and it will be formally justified in the following section.
- (iii) From the relation $SS_T = SS_R + SS_E$ given in (15) and the definition of the variability due to regression, SS_R , given in (14), it follows that the better the regression fit is, the smaller the value of SS_R is. Then, its ratio to the total variability, SS_T , $r = SS_R/SS_T$, can be used as an index of how good the linear regression fit is.

13.3 Normally Distributed Errors: MLE's of β_1 , β_2 , and σ^2 , Some Distributional Results

It is to be noticed that in the linear regression model as defined in relation (3), no distribution assumption about the errors e_i , and therefore the r.v.'s Y_i , was made. Such an assumption was not necessary, neither for the construction of the LSE's of β_1 , β_2 , nor in proving their unbiasedness and in calculating their variances. However, in order to be able to construct confidence intervals for

β_1 and β_2 and test hypotheses about them, among other things, we have to assume a distribution for the e_i 's. The e_i 's being errors, it is not unreasonable to assume that they are Normally distributed, and we shall do so. Then the model (3) is supplemented as follows:

$$Y_i = \beta_1 + \beta_2 x_i + e_i, \quad \text{where } e_i, i = 1, \dots, n \text{ are independent r.v.'s} \\ \sim N(0, \sigma^2), \quad \text{which implies that } Y_i, i = 1, \dots, n \text{ are} \quad (18) \\ \text{independent r.v.'s and } Y_i \sim N(\beta_1 + \beta_2 x_i, \sigma^2).$$

We now proceed with the following theorem.

THEOREM 4

Under model (18):

- (i) The LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$ of β_1 and β_2 , respectively, are also MLE's.
- (ii) The MLE $\hat{\sigma}^2$ of σ^2 is given by: $\hat{\sigma}^2 = SS_E/n$.
- (iii) The estimates $\hat{\beta}_1$ and $\hat{\beta}_2$ are Normally distributed as follows:

$$\hat{\beta}_1 \sim N\left(\beta_1, \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{SS_x}\right)\right), \quad \hat{\beta}_2 \sim N\left(\beta_2, \frac{\sigma^2}{SS_x}\right),$$

where SS_x is given in (9).

PROOF

- (i) The likelihood function of the Y_i 's is given by:

$$L(y_1, \dots, y_n; \beta_1, \beta_2, \sigma^2) = \left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^n \exp\left[-\frac{1}{2\sigma^2} \sum_i (y_i - \beta_1 - \beta_2 x_i)^2\right].$$

For each fixed σ^2 , maximization of the likelihood function with respect to β_1 and β_2 , is, clearly, equivalent to minimization of $\sum_i (y_i - \beta_1 - \beta_2 x_i)^2$ with respect to β_1 and β_2 , which minimization has produced the LSE's $\hat{\beta}_1$ and $\hat{\beta}_2$.

- (ii) The MLE of σ^2 is to be found by minimizing, with respect to σ^2 , the expression:

$$\log L(y_1, \dots, y_n; \hat{\beta}_1, \hat{\beta}_2, \sigma^2) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log \sigma^2 - \frac{1}{2\sigma^2} SS_E,$$

since, by (14) and (11), $\sum_i (y_i - \hat{\beta}_1 - \hat{\beta}_2 x_i)^2 = \sum_i (y_i - \hat{y}_i)^2 = SS_E$. From this expression, we get:

$$\frac{d}{d\sigma^2} \log L(y_1, \dots, y_n; \hat{\beta}_1, \hat{\beta}_2, \sigma^2) = -\frac{n}{2} \times \frac{1}{\sigma^2} + \frac{SS_E}{2(\sigma^2)^2} = 0,$$

so that $\sigma^2 = SS_E/n$. Since

$$\frac{d^2}{d(\sigma^2)^2} \log L(y_1, \dots, y_n; \hat{\beta}_1, \hat{\beta}_2, \sigma^2) \Big|_{\sigma^2=SS_E/n} = -\frac{n^3}{2SS_E^2} < 0,$$

it follows that $\hat{\sigma}^2 = SS_E/n$ is, indeed, the MLE of σ^2 .

(iii) From (12), we have: $\hat{\beta}_1 = \sum_i [\frac{1}{n} - \frac{\bar{x}(x_i - \bar{x})}{SS_x}] Y_i$, and we have also seen in Theorem 2 that:

$$E\hat{\beta}_1 = \beta_1, \quad \text{Var}(\hat{\beta}_1) = \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{SS_x} \right).$$

Thus, $\hat{\beta}_1$ is Normally distributed as a linear combination of independent Normally distributed r.v.'s, and its mean and variance must be as stated above. Next, from (8), we have that: $\hat{\beta}_2 = \sum_i (\frac{x_i - \bar{x}}{SS_x}) Y_i$, so that, as above, $\hat{\beta}_2$ is Normally distributed. Its mean and variance have been computed in Theorem 2 and they are β_2 and σ^2/SS_x , respectively. ▲

Before proceeding further, we return to Example 1 and compute an estimate for σ^2 . Also, discuss Example 23 in Chapter 1 and, perhaps, an additional example to be introduced here.

EXAMPLE 3

In reference to Example 1, determine the MLE of σ^2 .

DISCUSSION By Theorem 4(ii), this estimate is: $\hat{\sigma}^2 = \frac{SS_E}{n}$. For the computation of SS_E by (16), we have to have the quantity $\sum y_i^2$ from Table 13.2, which is calculated to be:

$$\sum_i y_i^2 = 8,373,295. \quad (19)$$

Then, by (9),

$$SS_y = 8,373,295 - \frac{(16,703)^2}{34} \simeq 8,373,295 - 8,205,594.382 = 167,700.618,$$

and therefore

$$SS_E = 167,700.618 - \frac{(581.376)^2}{6.361} \simeq 167,700.618 - 53,135.993 = 114,564.625;$$

i.e.,

$$SS_E = 114,564.625 \quad \text{and then} \quad \hat{\sigma}^2 = \frac{114,564.625}{34} \simeq 3,369.548.$$

Since $SS_T = SS_y = 167,700.618$ and $SS_R = 53,135.993$, it follows that only $\frac{53,135.993}{167,700.618} \simeq 31.685\%$ of the variability is explained by linear regression and $\frac{114,564.625}{167,700.618} \simeq 68.315\%$ is not explained by linear regression. The obvious outlier (3.44, 692) may be mainly responsible for it.

EXAMPLE 4

In reference to Example 23 in Chapter 1, assume a linear relationship between the dose of a compost fertilizer x and the yield of a crop y . On the basis of the following summary data recorded:

$$n = 15, \quad \bar{x} = 10.8, \quad \bar{y} = 122.7, \quad SS_x = 70.6, \quad SS_y = 98.5, \quad SS_{xy} = 68.3:$$

- (i) Determine the estimates $\hat{\beta}_1$ and $\hat{\beta}_2$, and draw the fitted regression line.
- (ii) Give the MLE $\hat{\sigma}^2$ of σ^2 .
- (iii) Over the range of x values covered in the study, what would your conjecture be regarding the average increase in yield per unit increase in the compost dose?

DISCUSSION

- (i) By (10),


$$\hat{\beta}_2 = \frac{68.3}{70.6} \simeq 0.967 \quad \text{and} \quad \hat{\beta}_1 = 122.7 - 0.967 \times 10.8 \simeq 112.256,$$

$$\text{and} \quad \hat{y} = 112.256 + 0.967x.$$

- (ii) We have: $\hat{\sigma}^2 = \frac{SS_E}{15}$, where $SS_E = 98.5 - \frac{(68.3)^2}{70.6} \simeq 32.425$, so that $\hat{\sigma}^2 = \frac{32.425}{15} = 2.162$.
- (iii) The conjecture would be a number close to the slope of the fitted regression line, which is 0.967 (Figure 13.3).

Figure 13.3

**The Fitted
Regression Line $\hat{y} =$
 $112.256 + 0.967x$**



EXAMPLE 5

In one stage of the development of a new medication for an allergy, an experiment is conducted to study how different dosages of the medication affect the duration of relief from the allergic symptoms. Ten patients are included in the experiment. Each patient receives a specific dosage of the medication and is asked to report back as soon as the protection of the medication seems to wear off. The observations are recorded in Table 13.3, which shows the dosage (x) and respective duration of relief (y) for the 10 patients.

- (i) Draw the scatter diagram of the data in Table 13.3 (which indicate tendency toward linear dependence).
- (ii) Compute the estimates $\hat{\beta}_1$ and $\hat{\beta}_2$, and draw the fitted regression line.
- (iii) What percentage of the total variability is explained by the linear regression and what percentage remains unexplained?
- (iv) Compute the MLE $\hat{\sigma}^2$ of σ^2 .

Table 13.3

Dosage (x) (in milligrams) and the Number of Days of Relief (y) from Allergy for 10 Patients

	x	y	x^2	y^2	xy
	3	9	9	81	27
	3	5	9	25	15
	4	12	16	144	48
	5	9	25	81	45
	6	14	36	196	84
	6	16	36	256	96
	7	22	49	484	154
	8	18	64	324	144
	8	24	64	576	192
	9	22	81	484	198
Totals	59	151	389	2,651	1,003

DISCUSSION

(i),(ii) First, $SS_x = 389 - \frac{59^2}{10} = 40.9$ and $SS_{xy} = 1,003 - \frac{59 \times 151}{10} = 112.1$, and hence:

$$\hat{\beta}_2 = \frac{112.1}{40.9} \simeq 2.741 \quad \text{and} \quad \hat{\beta}_1 = \frac{151}{10} - 2.741 \times \frac{59}{10} \simeq -1.072.$$


Then the fitted regression line is $\hat{y} = -1.072 + 2.741x$ (Figure 13.4).

(iii) Since $SS_T = SS_y = 2,651 - \frac{151^2}{10} = 370.9$ and $SS_R = \frac{(112.1)^2}{40.9} \simeq 307.247$, it follows that $SS_E = 370.9 - 307.247 = 63.653$. Therefore $\frac{307.247}{370.9} \simeq 82.838\%$ of the variability is explained by the linear regression and $\frac{63.653}{370.9} \simeq 17.162\%$ remains unexplained.

(iv) We have: $\hat{\sigma}^2 = \frac{63.653}{10} = 6.3653 \simeq 6.365$.

Figure 13.4

Scatter Diagram and the Fitted Regression Line $\hat{y} = -1.072 + 2.741x$



For the purpose of constructing confidence intervals for the parameters of the model, and also testing hypotheses about them, we have to know the

distribution of SS_E and also establish independence of the statistics $\hat{\beta}_1$ and SS_E , as well as independence of the statistics $\hat{\beta}_2$ and SS_E . The relevant results are stated in the following theorem, whose proof is deferred to Section 13.6.

THEOREM 5

Under model (18):

- (i) The distribution of SS_E/σ^2 is χ_{n-2}^2 .
 (ii) The following statistics are independent:
 (a) SS_E and $\hat{\beta}_2$; (b) \bar{Y} and $\hat{\beta}_2$; (c) SS_E , \bar{Y} and $\hat{\beta}_2$; (d) SS_E and $\hat{\beta}_1$.

PROOF Deferred to Section 13.6.

To this theorem, in conjunction with Theorem 4, there is the following corollary.

COROLLARY Under model (18):

- (i) The MLE $\hat{\sigma}^2$ of σ^2 is a biased estimate of σ^2 , but $\frac{n}{n-2}\hat{\sigma}^2 = \frac{SS_E}{n-2}$, call it S^2 , is an unbiased estimate of σ^2 .

$$(ii) \frac{\hat{\beta}_1 - \beta_1}{S\sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}}} \sim t_{n-2}, \quad (iii) \frac{\hat{\beta}_2 - \beta_2}{S/\sqrt{SS_x}} \sim t_{n-2}, \quad (20)$$

where

$$S^2 = SS_E/(n-2). \quad (21)$$

PROOF

- (i) It has been seen in Theorem 4(ii) that $\hat{\sigma}^2 = \frac{SS_E}{n} = \frac{n-2}{n} \times \frac{SS_E}{n-2}$. Since $\frac{SS_E}{\sigma^2} \sim \chi_{n-2}^2$, it follows that $E(\frac{SS_E}{\sigma^2}) = n-2$, or $E(\frac{SS_E}{n-2}) = \sigma^2$, so that $\frac{SS_E}{n-2}$ is an unbiased estimate of σ^2 . Also, $E\hat{\sigma}^2 = \frac{n-2}{n}E(\frac{SS_E}{n-2}) = \frac{n-2}{n}\sigma^2$, so that $\hat{\sigma}^2$ is biased.

- (ii) By Theorem 4(iii),

$$\frac{\hat{\beta}_1 - \beta_1}{\text{s.d.}(\hat{\beta}_1)} = \frac{\hat{\beta}_1 - \beta_1}{\sigma\sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}}} \sim N(0, 1),$$

and $\frac{SS_E}{\sigma^2} = \frac{(n-2)SS_E}{(n-2)\sigma^2} = \frac{(n-2)S^2}{\sigma^2} \sim \chi_{n-2}^2$. Furthermore, $\frac{\hat{\beta}_1 - \beta_1}{\text{s.d.}(\hat{\beta}_1)}$ and $\frac{SS_E}{\sigma^2}$ are independent, since $\hat{\beta}_1$ and SS_E are so. It follows that:

$$\frac{(\hat{\beta}_1 - \beta_1)/\text{s.d.}(\hat{\beta}_1)}{\sqrt{\frac{SS_E}{\sigma^2}/(n-2)}} \sim t_{n-2}, \quad \text{or} \quad \frac{(\hat{\beta}_1 - \beta_1)/\sigma\sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}}}{\sqrt{S^2/\sigma^2}} \sim t_{n-2},$$

or, finally,

$$\frac{\hat{\beta}_1 - \beta_1}{S\sqrt{\frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2}}} \sim t_{n-2}.$$

(iii) Again by Theorem 4(iii),

$$\frac{\hat{\beta}_2 - \beta_2}{\text{s.d.}(\hat{\beta}_2)} = \frac{\hat{\beta}_2 - \beta_2}{\sigma/\sqrt{SS_x}} \sim N(0, 1),$$

and $\frac{\hat{\beta}_2 - \beta_2}{\text{s.d.}(\hat{\beta}_2)}$ and $\frac{SS_E}{\sigma^2}$ are independent, since $\hat{\beta}_2$ and SS_E are so. Then:

$$\frac{(\hat{\beta}_2 - \beta_2)/\text{s.d.}(\hat{\beta}_2)}{\sqrt{SS_E/\sigma^2}/(n-2)} \sim t_{n-2}, \quad \text{or} \quad \frac{(\hat{\beta}_2 - \beta_2)/\frac{\sigma}{\sqrt{SS_x}}}{\sqrt{SS_E/\sigma^2}} \sim t_{n-2},$$

or, finally, $\frac{\hat{\beta}_2 - \beta_2}{S/\sqrt{SS_x}} \sim t_{n-2}$. ▲

Exercises

3.1 Verify the result $\frac{d}{dt^2} \log(y_1, \dots, y_n; \hat{\beta}_1, \hat{\beta}_2, t) \Big|_{t=SS_E/n} = -\frac{n}{2SS_E^2}$ as claimed in the proof of Theorem 4(ii), where $t = \sigma^2$.

3.2 Consider Table 13.1 and leave out the “outlier” pairs (3.63, 447), (3.44, 692), and (2.44, 336). Then recalculate all quantities below:

$$\sum_i x_i, \quad \sum_i y_i, \quad \sum_i x_i^2, \quad \sum_i x_i y_i, \quad \sum_i y_i^2, \quad SS_x, \quad SS_y, \quad SS_{xy}.$$

3.3 Use the calculations in Exercise 3.2 to compute the estimates $\hat{\beta}_1$, $\hat{\beta}_2$ and the fitted regression line.

3.4 Refer to Exercise 3.2, and compute the variances $\text{Var}(\hat{\beta}_1)$, $\text{Var}(\hat{\beta}_2)$, and the MLE of σ^2 .

3.5 By Theorem 5, the r.v. SS_E/σ^2 is distributed as χ_{n-2}^2 . Therefore, in the usual manner,

$$\left[\frac{SS_E}{\chi_{n-2; \frac{\alpha}{2}}^2}, \quad \frac{SS_E}{\chi_{n-2; 1-\frac{\alpha}{2}}^2} \right]$$

is a confidence interval for σ^2 with confidence coefficient $1 - \alpha$, where SS_E is given in (16) and (9). That is, $SS_E = SS_y - \frac{SS_{xy}^2}{SS_x}$, where SS_x , SS_y , and SS_{xy} are given in (9).

(i) Refer to Example 1 (see also Example 3), and construct a 95% confidence interval for σ^2 .

(ii) Refer to Example 4 and do the same as in part (i).

(iii) Refer to Example 5 and do the same as in part (i).

(iv) Refer to Exercise 3.2 and do the same as in part (i).

3.6 Consider the linear regression model given in relation (18), and let x_0 be an unknown point at which observations Y_{0i} , $i = 1, \dots, m$ are taken. It is assumed that the Y_{0i} 's and the Y_j 's are independent, and set $Y_0 = \frac{1}{m} \sum_{i=1}^m Y_{0i}$. Set $\mathbf{y} = (y_1, \dots, y_n)$, $\mathbf{y}_0 = (y_{01}, \dots, y_{0m})$ for the observed

values of the Y_j 's and Y_{0i} 's, and form their joint log-likelihood function:

$$\begin{aligned}\Lambda &= \Lambda(\beta_1, \beta_2, \sigma^2, x_0) = \log L(\beta_1, \beta_2, \sigma^2, x_0 | \mathbf{y}, \mathbf{y}_0) \\ &= -\frac{m+n}{2} \log(2\pi) - \frac{m+n}{2} \log \sigma^2 \\ &\quad - \frac{1}{\sigma^2} \left[\sum_{j=1}^n (y_j - \beta_1 - \beta_2 x_j)^2 + \sum_{i=1}^m (y_{0i} - \beta_1 - \beta_2 x_0)^2 \right].\end{aligned}$$

- (i) Show that the log-likelihood equations $\frac{\partial \Lambda}{\partial \beta_1} = 0$, $\frac{\partial \Lambda}{\partial \beta_2} = 0$, and $\frac{\partial \Lambda}{\partial x_0} = 0$ produce the equations:

$$(m+n)\beta_1 + (mx_0 + n\bar{x})\beta_2 = my_0 + n\bar{y} \quad (\text{a})$$

$$(mx_0 + n\bar{x})\beta_1 + \left(mx_0^2 + \sum_j x_j^2 \right) \beta_2 = mx_0 y_0 + \sum_j x_j y_j \quad (\text{b})$$

$$\beta_1 + x_0 \beta_2 = y_0. \quad (\text{c})$$

- (ii) In (c), solve for β_1 , $\beta_1 = y_0 - x_0 \beta_2$, replace it in (a) and (b), and solve for β_2 to obtain, by assuming here and in the sequel that all divisions and cancellations are legitimate,

$$\beta_2 = \frac{\bar{y} - y_0}{\bar{x} - x_0}, \quad \beta_2 = \frac{\sum_j x_j y_j - n\bar{x}y_0}{\sum_j x_j^2 - nx_0\bar{x}}. \quad (\text{d})$$

- (iii) Equate the β_2 's in (ii), and solve for x_0 to obtain:

$$\left. \begin{aligned} x_0 &= \left[y_0 \sum_j (x_j - \bar{x})^2 + \bar{x} \sum_j x_j y_j - \bar{y} \sum_j x_j^2 \right] \\ &\quad \left/ \left(\sum_j x_j y_j - n\bar{x}\bar{y} \right) \right. \\ &= \left[ny_0 \sum_j (x_j - \bar{x})^2 + n\bar{x} \sum_j x_j y_j - n\bar{y} \sum_j x_j^2 \right] \\ &\quad \left/ \left[n \sum_j x_j y_j - \left(\sum_j x_j \right) \left(\sum_j y_j \right) \right] \right. \end{aligned} \right\} \quad (\text{e})$$

- (iv) Replace x_0 in the first expression for β_2 in (d) in order to get, after some simplifications:

$$\beta_2 = \frac{n \sum_j x_j y_j - \left(\sum_j x_j \right) \left(\sum_j y_j \right)}{n \sum_j x_j^2 - \left(\sum_j x_j \right)^2}, \quad (\text{f})$$

and observe that this expression is the MLE (LSE) of β_2 calculated on the basis of y_j and x_j , $j = 1, \dots, n$ only (see relation (5) and Theorem 4(i)).

- (v) Replace x_0 and β_2 in the expression $\beta_1 = y_0 - x_0\beta_2$ in order to arrive at the expression

$$\beta_1 = \frac{(\sum_j x_j^2)(\sum_j y_j) - (\sum_j x_j)(\sum_j x_j y_j)}{n \sum_j x_j^2 - (\sum_j x_j)^2}, \quad (g)$$

after some calculations, and observe that this is the MLE (LSE) of β_1 calculated on the basis of y_j and x_j , $j = 1, \dots, n$ only (see relation (4) and Theorem 4(i)).

- (vi) It follows that the MLE's of β_1 , β_2 , and x_0 , to be denoted by $\hat{\beta}_1$, $\hat{\beta}_2$, and \hat{x}_0 , are given by the expressions:

$$\hat{\beta}_1 = \frac{(\sum_j x_j^2)(\sum_j y_j) - (\sum_j x_j)(\sum_j x_j y_j)}{n \sum_j x_j^2 - (\sum_j x_j)^2},$$

$$\hat{\beta}_2 = \frac{n \sum_j x_j y_j - (\sum_j x_j)(\sum_j y_j)}{n \sum_j x_j^2 - (\sum_j x_j)^2},$$

and

$$\hat{x}_0 = \frac{y_0 - \hat{\beta}_1}{\hat{\beta}_2}.$$

- (vii) Differentiate the log-likelihood function with respect to σ^2 , equate the derivative to zero, and replace β_1 , β_2 , and x_0 by their MLE's in order to obtain the MLE $\hat{\sigma}^2$ of σ^2 , which is given by the expression:

$$\hat{\sigma}^2 = \frac{1}{m+n}(SS_E + SS_{0E}),$$

where

$$SS_E = \sum_{j=1}^n (y_j - \hat{y}_j)^2 = \sum_{j=1}^n (y_j - \hat{\beta}_1 - \hat{\beta}_2 x_j)^2,$$

and

$$SS_{0E} = \sum_{i=1}^m (y_{0i} - \hat{\beta}_1 - \hat{\beta}_2 \hat{x}_0)^2 = \sum_{i=1}^m (y_{0i} - y_0)^2.$$

Also, by means of (14) and (16),

$$SS_E = SS_y - \frac{SS_{xy}^2}{SS_x}, \quad \text{where } SS_x = \sum_{j=1}^n x_j^2 - \frac{1}{n} \left(\sum_{j=1}^n x_j \right)^2,$$

$$SS_y = \sum_{j=1}^n y_j^2 - \frac{1}{n} \left(\sum_{j=1}^n y_j \right)^2, \quad SS_{xy} = \sum_{j=1}^n x_j y_j - \frac{1}{n} \left(\sum_{j=1}^n x_j \right) \left(\sum_{j=1}^n y_j \right),$$

and

$$SS_{0E} = \sum_{i=1}^m y_{0i}^2 - \frac{1}{m} \left(\sum_{i=1}^m y_{0i} \right)^2.$$

- (viii) Observe that, by Theorem 5(i), $\frac{SS_E}{\sigma^2} \sim \chi_{n-2}^2$, whereas $\frac{SS_{0E}}{\sigma^2} \sim \chi_{m-1}^2$. Then, by independence of the Y_j 's and the Y_{0i} 's, it follows that $\frac{1}{\sigma^2}(SS_E + SS_{0E}) \sim \chi_{m+n-3}^2$.
- (ix) Observe that, by Theorem 8(i), $\hat{y}_0 = \hat{\beta}_1 + \hat{\beta}_2 x_0 \sim N(\beta_1 + \beta_2 x_0, \sigma^2(\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}))$, whereas $Y_0 \sim N(\beta_1 + \beta_2 x_0, \frac{\sigma^2}{m})$, and \hat{y}_0 and Y_0 are independent, so that the r.v. $V = Y_0 - \hat{y}_0 \sim N(0, \sigma_V^2)$, where $\sigma_V^2 = \sigma^2(\frac{1}{m} + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x})$, and $\frac{V}{\sigma_V} \sim N(0, 1)$.
- (x) Observe that, by Theorem 5, the r.v.'s V/σ_V and $(SS_E + SS_{0E})/\sigma^2$ are independent, so that

$$\begin{aligned} \frac{V/\sigma_V}{\frac{1}{\sigma} \sqrt{\frac{SS_E + SS_{0E}}{m+n-3}}} &= \frac{V/\sigma \sqrt{\frac{1}{m} + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}}{\frac{1}{\sigma} \sqrt{\frac{SS_E + SS_{0E}}{m+n-3}}} \\ &= \frac{\sqrt{m+n-3} V}{\sqrt{\left[\frac{1}{m} + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x} \right] (SS_E + SS_{0E})}} \sim t_{m+n-3}. \end{aligned}$$

13.4 Confidence Intervals and Hypotheses Testing Problems

The results obtained in the corollary to Theorem 5 allow the construction of confidence intervals for the parameters of the model, as well as testing hypotheses about them.

THEOREM 6

Under model (18), $100(1 - \alpha)\%$ confidence intervals for β_1 and β_2 are given, respectively, by:

$$\left[\hat{\beta}_1 - t_{n-2, \alpha/2} S \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}}, \quad \hat{\beta}_1 + t_{n-2, \alpha/2} S \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}} \right], \quad (22)$$

and

$$\left[\hat{\beta}_2 - t_{n-2, \alpha/2} \frac{S}{\sqrt{SS_x}}, \quad \hat{\beta}_2 + t_{n-2, \alpha/2} \frac{S}{\sqrt{SS_x}} \right], \quad (23)$$

where $S = \sqrt{SS_E/(n-2)}$, and SS_E , SS_x , and $\hat{\beta}_1$, $\hat{\beta}_2$ are given by (16), (9), and (10).

PROOF The confidence intervals in (22) and (23) follow immediately from results (ii) and (iii), respectively, in the corollary to Theorem 5, and the familiar procedure of constructing confidence intervals. ▲

REMARK 2 A confidence interval can also be constructed for σ^2 on the basis of the statistic SS_E and the fact that SS_E/σ^2 is distributed as χ_{n-2}^2 .

Procedures for testing some hypotheses are summarized below in the form of a theorem. The tests proposed here have an obvious intuitive interpretation. However, their justification rests in that they are likelihood ratio tests. For the case of simple hypotheses, this fact can be established directly. For composite hypotheses, it follows as a special case of more general results of testing hypotheses regarding the entire mean $\eta = \beta_1 + \beta_2 x$. See, e.g., Chapter 16 and, in particular, Examples 2 and 3 in the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G. G. Roussas.

THEOREM 7

Under model (18):

- (i) For testing the hypothesis $H_0: \beta_1 = \beta_{10}$ against the alternative $H_A: \beta_1 \neq \beta_{10}$ at level of significance α , the null hypothesis H_0 is rejected whenever

$$|t| > t_{n-2, \alpha/2}, \quad \text{where } t = (\hat{\beta}_1 - \beta_{10}) / S \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}}. \quad (24)$$

- (ii) For testing the hypothesis $H_0: \beta_2 = \beta_{20}$ against the alternative $H_A: \beta_2 \neq \beta_{20}$ at level of significance α , the null hypothesis H_0 is rejected whenever

$$|t| > t_{n-2, \alpha/2}, \quad \text{where } t = (\hat{\beta}_2 - \beta_{20}) / \frac{S}{\sqrt{SS_x}}. \quad (25)$$

When the alternative is of the form $H_A: \beta_2 > \beta_{20}$, the null hypothesis is rejected whenever $t > t_{n-2, \alpha}$, and it is rejected whenever $t < -t_{n-2, \alpha}$ if the alternative is of the form $H_A: \beta_2 < \beta_{20}$.

REMARK 3

- (i) In the reference cited above, the test statistic used, actually, has the F -distribution under the null hypothesis. It should be recalled, however, that if t has the t -distribution with r d.f., i.e., $t = Z / \sqrt{\chi_r^2/r}$, where χ_r^2 has the χ^2 -distribution with r d.f. $Z \sim N(0, 1)$ and Z and χ_r^2 are independent then, $t^2 = \frac{Z^2}{\chi_r^2/r}$ has the F -distribution with 1 and r d.f.
- (ii) Hypotheses can also be tested about σ^2 on the basis of the fact that $\frac{SS_E}{\sigma^2} \sim \chi_{n-2}^2$.

EXAMPLE 6

In reference to Example 1:

- (i) Construct 95% confidence intervals for β_1 and β_2 .
- (ii) Test the hypothesis that the GMAT scores increase with increasing GPA scores.

DISCUSSION (i) The required confidence intervals are given by (22) and (23). In the discussion of Example 1, we have found that: $\bar{x} \simeq 2.963$,

$SS_x \simeq 6.361$, $\hat{\beta}_1 \simeq 220.456$, and $\hat{\beta}_2 \simeq 91.397$. Also, in the discussion of Example 3, we saw that $SS_E \simeq 114,564.625$, so that, by (21), $S = (\frac{114,564.625}{32})^{1/2} \simeq 59.834$. Finally, $t_{32;0.025} = 2.0369$. Then

$$\begin{aligned}\hat{\beta}_1 - t_{n-2;\alpha/2} S \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}} &= 220.456 - 2.0369 \times 59.834 \times \sqrt{\frac{1}{34} + \frac{(2.963)^2}{6.361}} \\ &\simeq 220.456 - 121.876 \times 1.187 \simeq 220.456 - 144.667 = 75.789,\end{aligned}$$

and

$$\hat{\beta}_1 + t_{n-2;\alpha/2} S \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}} \simeq 220.456 + 144.667 = 365.123.$$

So the required interval is [75.789, 365.123].

Likewise, $t_{n-2;\alpha/2} \frac{S}{\sqrt{SS_x}} \simeq 1.6939 \times \frac{59.834}{\sqrt{6.361}} = 1.6939 \times 23.725 \simeq 40.188$, and therefore the required interval for β_2 is: [51.209, 131.585].

(ii) Here we are to test $H_0: \beta_2 = 0$ against the alternative $H_A: \beta_2 > 0$. Let us take $\alpha = 0.05$, so that $t_{32;0.05} = 1.6939$. The observed value of the test statistics is:

$$t = \frac{\hat{\beta}_2 - \beta_{20}}{S/\sqrt{SS_x}} \simeq \frac{91.397}{23.725} = 3.852,$$

and the null hypothesis is rejected; the GMAT scores increase along with increasing GPA scores.

EXAMPLE 7

In reference to Example 4:

- (i) Construct 95% confidence intervals for β_1 and β_2 .
- (ii) Test the hypothesis that crop yield increases with increasing compost fertilizer amounts.

DISCUSSION (i) In the discussion of Example 4, we have seen that: $n = 15$, $\bar{x} = 10.8$, $SS_x = 70.6$, $SS_y = 98.5$, $\hat{\beta}_1 \simeq 112.256$, and $\hat{\beta}_2 \simeq 0.967$. It follows that:

$$\begin{aligned}S &= \left(\frac{98.5}{13}\right)^{1/2} \simeq 2.753 \quad \text{and} \quad S \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}} = 2.753 \times \sqrt{\frac{1}{15} + \frac{(10.8)^2}{70.6}} \\ &\simeq 2.753 \times 1.311 \simeq 3.609.\end{aligned}$$

Since $t_{13;0.025} = 1.1604$, it follows that the required observed confidence interval for β_1 is: [112.256 - 1.1604 × 3.609, 112.256 + 1.1604 × 3.609], or [108.068, 116.444].

Next, $\frac{S}{\sqrt{SS_x}} \simeq \frac{2.753}{\sqrt{70.6}} \simeq 0.328$, and $t_{13;0.025} \frac{S}{\sqrt{SS_x}} = 1.1604 \times 0.328 \simeq 0.381$, so that the required observed confidence interval for β_2 is: [0.586, 1.348].

(ii) The hypothesis to be tested is $H_0: \beta_2 = 0$ against the alternative $H_A: \beta_2 > 0$. Take $\alpha = 0.05$, so that $t_{13;0.05} = 1.7709$. The observed value of the test statistic is:

$$t = \frac{\hat{\beta}_2 - \beta_{20}}{S/\sqrt{SS_x}} \simeq \frac{0.967}{0.328} \simeq 2.948,$$

and therefore the null hypothesis is rejected. Consequently, crop yield increases with increasing amounts of compost fertilizer.

EXAMPLE 8

In reference to Example 5:

- (i) Construct 95% confidence intervals for β_1 and β_2 .
- (ii) Test the hypothesis that the duration of relief increases with higher dosages of the medication.

DISCUSSION (i) From the discussion of Example 5, we have: $n=10$, $\bar{x}=5.9$, $SS_x=40.9$, $SS_E=63.653$, $\hat{\beta}_1=-1.072$, and $\hat{\beta}_2 \simeq 2.741$. Then $S = (\frac{63.653}{8})^{1/2} \simeq 2.821$. Also, $t_{8;0.025} = 3.3060$. Therefore:

$$t_{n-2;\alpha/2} S \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}} = 3.306 \times 2.821 \times \sqrt{\frac{1}{10} + \frac{(5.9)^2}{40.9}} \simeq 9.326 \times 0.975 \\ \simeq 9.093.$$

Hence the required observed confidence interval for β_1 is: $[-1.072 - 9.093, -1.072 + 9.093]$, or $[-10.165, 8.021]$. Next,

$$t_{n-2;\alpha/2} \frac{S}{\sqrt{SS_x}} = 3.306 \times \frac{2.821}{\sqrt{40.9}} \simeq 3.306 \times 0.441 \simeq 1.458,$$

and therefore the required observed confidence interval for β_2 is: $[2.741 - 1.458, 2.741 + 1.458]$, or $[1.283, 4.199]$.

(ii) The hypothesis to be tested is $H_0: \beta_2 = 0$ against $H_A: \beta_2 > 0$, and let us take $\alpha = 0.05$, so that $t_{8;0.05} = 1.8595$. The observed value of the test statistic is:

$$t = \frac{\hat{\beta}_2 - \beta_{20}}{S/\sqrt{SS_x}} \simeq \frac{2.741}{0.441} \simeq 6.215,$$

and the null hypothesis is rejected. Thus, increased dosages of medication provide longer duration of relief.

Exercises

- 4.1 Refer to Exercises 3.2 and 3.4, and compute 95% confidence intervals for β_1 and β_2 .
- 4.2 Refer to Exercises 3.3 and 4.1, and test the hypotheses $H_0: \beta_1 = 300$ against $H_A: \beta_1 \neq 300$, and $H_0: \beta_2 = 60$ against $H_A: \beta_2 \neq 60$, each at level of significance $\alpha = 0.05$.

4.3 Refer to Example 5 and:

- (i) Derive 95% confidence intervals for β_1 and β_2 .
- (ii) Test the hypothesis $H_0: \beta_1 = -1$ against the alternative $H_A: \beta_1 \neq -1$ at level of significance $\alpha = 0.05$.
- (iii) Do the same for the hypothesis $H_0: \beta_2 = 3$ against the alternative $H_A: \beta_2 \neq 3$ at the same level $\alpha = 0.05$.

4.4 Suppose the observations Y_1, \dots, Y_n are of the following structure: $Y_i = \beta + \gamma(x_i - \bar{x}) + e_i$, where β and γ are parameters and the e_i 's are independent r.v.'s with mean 0 and unknown variance σ^2 .

- (i) Set $t_i = x_i - \bar{x}$, $i = 1, \dots, n$, and observe that the model $Y_i = \beta + \gamma t_i + e_i$ is of the standard form (1) with $\beta_1 = \beta$, $\beta_2 = \gamma$, and the additional property that $\sum_{i=1}^n t_i = 0$, or $\bar{t} = 0$.
- (ii) Use expressions (5) and (8) to conclude that the LSE's of β and γ are given by:

$$\hat{\beta} = \bar{Y}, \quad \hat{\gamma} = \frac{\sum_{i=1}^n t_i Y_i}{\sum_{i=1}^n t_i^2}.$$

(iii) Employ Theorem 4 in order to conclude that:

$$\hat{\beta} \sim N\left(\beta, \frac{\sigma^2}{n}\right) \quad \text{and} \quad \hat{\gamma} \sim N\left(\gamma, \frac{\sigma^2}{SS_t}\right),$$

where (by (9)) $SS_t = \sum_{i=1}^n t_i^2$.

- (iv) Determine the form of the confidence intervals for β and γ from relations (22) and (23).
- (v) Determine the expression of the test statistics by means of relations (24) and (25).
- (vi) What do the confidence intervals in relation (29) and in Theorem 9 (iii) become here?

4.5 Consider the linear regression models: $Y_i = \beta_1 + \beta_2 x_i + e_i$, $i = 1, \dots, m$ and $Y_j^* = \beta_1^* + \beta_2^* x_j^* + e_j^*$, $j = 1, \dots, n$, where the random errors e_1, \dots, e_m and e_1^*, \dots, e_n^* are i.i.d. r.v.'s distributed as $N(0, \sigma^2)$.

- (i) The independence of e_1, \dots, e_m and e_1^*, \dots, e_n^* implies independence of Y_1, \dots, Y_m and Y_1^*, \dots, Y_n^* . Then write down, the joint likelihood of the Y_i 's and the Y_j^* 's and observe that the MLE's of β_1 , β_2 , β_1^* , β_2^* , and σ^2 , in obvious notation, are given by:

$$\hat{\beta}_1 = \bar{Y} - \hat{\beta}_2 \bar{x}, \quad \hat{\beta}_2 = \frac{m \sum_{i=1}^m x_i Y_i - (\sum_{i=1}^m x_i)(\sum_{i=1}^m Y_i)}{m \sum_{i=1}^m x_i^2 - (\sum_{i=1}^m x_i)^2},$$

$$\hat{\beta}_1^* = \bar{Y}^* - \hat{\beta}_2^* \bar{x}^*, \quad \hat{\beta}_2^* = \frac{n \sum_{j=1}^n x_j^* Y_j^* - (\sum_{j=1}^n x_j^*)(\sum_{j=1}^n Y_j^*)}{n \sum_{j=1}^n x_j^{*2} - (\sum_{j=1}^n x_j^*)^2},$$

$$\hat{\sigma}^2 = (SS_E + SS_E^*) / (m + n),$$

where

$$SS_E = \sum_{i=1}^m (Y_i - \hat{\beta}_1 - \hat{\beta}_2 x_i)^2 = SS_y - \frac{SS_{xy}^2}{SS_x},$$

$$SS_x = \sum_{i=1}^m x_i^2 - \frac{1}{m} \left(\sum_{i=1}^m x_i \right)^2,$$

$$SS_y = \sum_{i=1}^m Y_i^2 - \frac{1}{m} \left(\sum_{i=1}^m Y_i \right)^2,$$

$$SS_{xy} = \sum_{i=1}^m x_i Y_i - \frac{1}{m} \left(\sum_{i=1}^m x_i \right) \left(\sum_{i=1}^m Y_i \right),$$

and

$$SS_E^* = \sum_{j=1}^n (Y_j^* - \hat{\beta}_1^* - \hat{\beta}_2^* x_j^*)^2 = SS_y^* - \frac{SS_{xy}^{*2}}{SS_x^*},$$

$$SS_x^* = \sum_{j=1}^n x_j^{*2} - \frac{1}{n} \left(\sum_{j=1}^n x_j^* \right)^2,$$

$$SS_y^* = \sum_{j=1}^n Y_j^{*2} - \frac{1}{n} \left(\sum_{j=1}^n Y_j^* \right)^2,$$

$$SS_{xy}^* = \sum_{j=1}^n x_j^* Y_j^* - \frac{1}{n} \left(\sum_{j=1}^n x_j^* \right) \left(\sum_{j=1}^n Y_j^* \right).$$

(ii) In accordance with Theorem 4, observe that

$$\hat{\beta}_1 \sim N \left(\beta_1, \sigma^2 \left(\frac{1}{m} + \frac{\bar{x}^2}{SS_x} \right) \right), \quad \hat{\beta}_2 \sim N \left(\beta_2, \frac{\sigma^2}{SS_x} \right),$$

$$\hat{\beta}_1^* \sim N \left(\beta_1^*, \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^{*2}}{SS_x^*} \right) \right), \quad \hat{\beta}_2^* \sim N \left(\beta_2^*, \frac{\sigma^2}{SS_x^*} \right),$$

and

$$\frac{SS_E + SS_E^*}{\sigma^2} \sim \chi_{m+n-4}^2.$$

(iii) From part (ii) and Theorem 5, conclude that

$$\frac{\sqrt{m+n-4} [(\hat{\beta}_1 - \hat{\beta}_1^*) - (\beta_1 - \beta_1^*)]}{\sqrt{(SS_E + SS_E^*) \left(\frac{1}{m} + \frac{1}{n} + \frac{\bar{x}^2}{SS_x} + \frac{\bar{x}^{*2}}{SS_x^*} \right)}} \sim t_{m+n-4},$$

and

$$\frac{\sqrt{m+n-4} [(\hat{\beta}_2 - \hat{\beta}_2^*) - (\beta_2 - \beta_2^*)]}{\sqrt{(SS_E + SS_E^*) \left(\frac{1}{SS_x} + \frac{1}{SS_x^*} \right)}} \sim t_{m+n-4}.$$

- (iv) From part (iii), observe that the two regression lines can be compared through the test of the hypotheses $H_0: \beta_1 = \beta_1^*$ against the alternative $H_A: \beta_1 \neq \beta_1^*$, and $H_0: \beta_2 = \beta_2^*$ against the alternative $H_A: \beta_2 \neq \beta_2^*$ by using the respective test statistics:

$$t = \frac{\sqrt{m+n-4}(\hat{\beta}_1 - \hat{\beta}_1^*)}{\sqrt{(SS_E + SS_E^*)\left(\frac{1}{m} + \frac{1}{n} + \frac{\bar{x}^2}{SS_x} + \frac{\bar{x}^{*2}}{SS_x^*}\right)}},$$

$$t' = \frac{\sqrt{m+n-4}(\hat{\beta}_2 - \hat{\beta}_2^*)}{\sqrt{(SS_E + SS_E^*)\left(\frac{1}{SS_x} + \frac{1}{SS_x^*}\right)}}.$$

At level of significance α , the hypothesis H_0 is rejected when $|t| > t_{m+n-4; \frac{\alpha}{2}}$, and the hypothesis H_0' is rejected when $|t'| > t_{m+n-4; \frac{\alpha}{2}}$.

- (v) Again from part (iii), observe that 95% confidence intervals for $\beta_1 - \beta_1^*$ and $\beta_2 - \beta_2^*$ are given by:

$$(\hat{\beta}_1 - \hat{\beta}_1^*) \pm t_{m+n-4; \frac{\alpha}{2}} \sqrt{\frac{SS_E + SS_E^*}{m+n-4} \left(\frac{1}{m} + \frac{1}{n} + \frac{\bar{x}^2}{SS_x} + \frac{\bar{x}^{*2}}{SS_x^*} \right)},$$

and

$$(\hat{\beta}_2 - \hat{\beta}_2^*) \pm t_{m+n-4; \frac{\alpha}{2}} \sqrt{\frac{SS_E + SS_E^*}{m+n-4} \left(\frac{1}{SS_x} + \frac{1}{SS_x^*} \right)},$$

respectively.

- (vi) Finally, from part (ii) conclude that a 95% confidence interval for σ^2 is given by:

$$\left[\frac{SS_E + SS_E^*}{\chi_{m+n-4; \frac{\alpha}{2}}^2}, \frac{SS_E + SS_E^*}{\chi_{m+n-4; 1-\frac{\alpha}{2}}^2} \right].$$

13.5 Some Prediction Problems

According to model (18), the expectation of the observation Y_i at x_i is $EY_i = \beta_1 + \beta_2 x_i$. Now, suppose x_0 is a point distinct from all x_i 's, but lying in the range that the x_i 's span, and we wish to predict the expected value of the observation Y_0 at x_0 ; i.e., $EY_0 = \beta_1 + \beta_2 x_0$. An obvious predictor for EY_0 is the statistic \hat{y}_0 given by the expression below and modified as indicated:

$$\hat{y}_0 = \hat{\beta}_1 + \hat{\beta}_2 x_0 = (\bar{Y} - \hat{\beta}_2 \bar{x}) + \hat{\beta}_2 x_0 = \bar{Y} + (x_0 - \bar{x}) \hat{\beta}_2. \quad (26)$$

The result below gives the distribution of \hat{y}_0 , which also provides for the construction of a confidence interval for $\beta_1 + \beta_2 x_0$.

THEOREM 8

Under model (18) and with \hat{y}_0 given by (26), we have:

$$(i) \quad \frac{\hat{y}_0 - (\beta_1 + \beta_2 x_0)}{\sigma \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}} \sim N(0, 1). \quad (27)$$

$$(ii) \quad \frac{\hat{y}_0 - (\beta_1 + \beta_2 x_0)}{S \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}} \sim t_{n-2}. \quad (28)$$

(iii) A $100(1 - \alpha)\%$ confidence interval for $\beta_1 + \beta_2 x_0$ is given by:

$$\left[\hat{y}_0 - t_{n-2, \alpha/2} S \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}, \quad \hat{y}_0 + t_{n-2, \alpha/2} S \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}} \right]. \quad (29)$$

It is recalled that $S = \sqrt{SS_E/(n-2)}$, $SS_E = SS_y$, and SS_y and SS_x are given in (9).

PROOF (i) The assumption that $Y_i \sim N(\beta_1 + \beta_2 x_i, \sigma^2)$, $i = 1, \dots, n$ independent implies that $\sum_i Y_i \sim N(n\beta_1 + \beta_2 \sum_i x_i, n\sigma^2)$ and hence

$$\bar{Y} \sim N(\beta_1 + \beta_2 \bar{x}, \sigma^2/n). \quad (30)$$

By Theorem 4(iii), $\hat{\beta}_2 \sim N(\beta_2, \sigma^2/SS_x)$, so that

$$(x_0 - \bar{x})\hat{\beta}_2 \sim N\left((x_0 - \bar{x})\beta_2, \frac{\sigma^2(x_0 - \bar{x})^2}{SS_x}\right). \quad (31)$$

Furthermore, by Theorem 5(ii)(b), \bar{Y} and $\hat{\beta}_2$ are independent. Then, relations (26), (30), and (31) yield:

$$\hat{y}_0 = \bar{Y} + (x_0 - \bar{x})\hat{\beta}_2 \sim N\left(\beta_1 + \beta_2 x_0, \sigma^2\left(\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}\right)\right), \quad (32)$$

and then (27) follows by standardization.

(ii) By Theorem 5(ii)(c), SS_E is independent of \bar{Y} and $\hat{\beta}_2$ and hence independent of \hat{y}_0 because of (26). Furthermore, by Theorem 5(i),

$$\frac{SS_E}{\sigma^2} = \frac{(n-2)S^2}{\sigma^2} \sim \chi_{n-2}^2. \quad (33)$$

Therefore

$$\frac{[\hat{y}_0 - (\beta_1 + \beta_2 x_0)]/\sigma \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}}{\sqrt{\frac{(n-2)S^2}{\sigma^2}/(n-2)}} = \frac{\hat{y}_0 - (\beta_1 + \beta_2 x_0)}{S \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}} \sim t_{n-2},$$

which is relation (28).

(iii) This part follows immediately from part (ii) and the standard procedure of setting up confidence intervals. \blacktriangle

Finally, we would like to consider the problem of predicting a *single response* at a given point x_0 rather than its *expected value*. Call Y_0 the response corresponding to x_0 and, reasonably enough, assume that Y_0 is independent of the Y_i 's. The predictor for Y_0 is \hat{y}_0 , the same as the one given in (26). The objective here is to construct a prediction interval for Y_0 . This is done indirectly in the following result.

THEOREM 9

Under model (18), let Y_0 be the (unobserved) observation at x_0 , and assume that Y_0 is independent of the Y_i 's. Predict Y_0 by $\hat{y}_0 = \hat{\beta}_1 + \hat{\beta}_2 x_0$. Then:

$$(i) \quad \frac{\hat{y}_0 - Y_0}{\sigma \sqrt{1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}} \sim N(0, 1). \quad (34)$$

$$(ii) \quad \frac{\hat{y}_0 - Y_0}{S \sqrt{1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}} \sim t_{n-2}. \quad (35)$$

(iii) A $100(1 - \alpha)\%$ prediction interval for Y_0 is given by:

$$\left[\hat{y}_0 - t_{n-2; \alpha/2} S \sqrt{1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}}, \quad \hat{y}_0 + t_{n-2; \alpha/2} S \sqrt{1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}} \right],$$

where S and SS_x are as in Theorem 8(ii), (iii).

PROOF (i) We have: $Y_0 = \beta_1 + \beta_2 x_0 + e_0$, predicted by $\hat{y}_0 = \hat{\beta}_1 + \hat{\beta}_2 x_0$. Then $EY_0 = \beta_1 + \beta_2 x_0$ and $E\hat{y}_0 = \beta_1 + \beta_2 x_0$, so that $E(\hat{y}_0 - Y_0) = 0$. In deriving the distribution of $\hat{y}_0 - Y_0$, we need its variance. By (26), we have:

$$\text{Var}(\hat{y}_0 - Y_0) = \text{Var}(\bar{Y} + (x_0 - \bar{x})\hat{\beta}_2 - Y_0) = \text{Var}(\bar{Y}) + (x_0 - \bar{x})^2 \text{Var}(\hat{\beta}_2) + \text{Var}(Y_0)$$

(since all three r.v.'s, \bar{Y} , $\hat{\beta}_2$, and Y_0 , are independent)

$$\begin{aligned} &= \frac{\sigma^2}{n} + (x_0 - \bar{x})^2 \times \frac{\sigma^2}{SS_x} + \sigma^2 \quad (\text{by Theorem 2}) \\ &= \sigma^2 \left\{ 1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x} \right\}; \quad \text{i.e.,} \end{aligned}$$

$$E(\hat{y}_0 - Y_0) = 0 \quad \text{and} \quad \text{Var}(\hat{y}_0 - Y_0) = \sigma^2 \left[1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x} \right].$$

Since \hat{y}_0 and Y_0 are independent and $Y_0 \sim N(\beta_1 + \beta_2 x_0, \sigma^2)$, then these facts along with (32) yield:

$$\hat{y}_0 - Y_0 \sim N \left(0, \sigma^2 \left[1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x} \right] \right).$$

Relation (34) follows by standardizing $\hat{y}_0 - Y_0$.

(ii) It has been argued in the proof of Theorem 8(ii) that S and \hat{y}_0 are independent. It follows that S and $\hat{y}_0 - Y_0$ are also independent. Then, dividing the expression on the left-hand side in (34) by $\sqrt{\frac{(n-2)S^2}{\sigma^2}}/(n-2) = \frac{S}{\sigma}$ in (33), we obtain the result in (35), after some simplifications.

(iii) This part follows from part (ii) through the usual procedure of setting up confidence intervals. ▲

Exercises

5.1 Refer to Exercises 3.1, 3.3, 4.1, and:

- (i) Predict EY_0 at $x_0 = 3.25$, and construct a 95% confidence interval of EY_0 .
- (ii) Predict the response Y_0 at $x_0 = 3.25$, and construct a 95% prediction interval for Y_0 .

5.2 In reference to Example 22 in Chapter 1 (see also scatter diagram in Figure 13.1 and Examples 1, 2, 3, and 6 here), do the following:

- (i) Predict the EY_0 , where Y_0 is the response at $x_0 = 3.25$.
- (ii) Construct a 95% confidence interval for $EY_0 = \beta_1 + \beta_2 x_0 = \beta_1 + 3.25\beta_2$.
- (iii) Predict the response Y_0 at $x_0 = 2.5$.
- (iv) Construct a 90% prediction interval for Y_0 .

5.3 In reference to Example 23 in Chapter 1 (see also Examples 4 and 7 here), do the following:

- (i) Predict the EY_0 , where Y_0 is the response at $x_0 = 12$.
- (ii) Construct a 95% confidence interval for $EY_0 = \beta_1 + \beta_2 x_0 = \beta_1 + 12\beta_2$.
- (iii) Predict the response Y_0 at $x_0 = 12$.
- (iv) Construct a 95% prediction interval for Y_0 .

5.4 Refer to Example 5 and:

- (i) Predict the EY_0 at $x_0 = 6$.
- (ii) Construct a 95% confidence interval for $EY_0 = \beta_1 + 6\beta_2$.
- (iii) Predict the response Y_0 at $x_0 = 6$.
- (iv) Construct a 95% prediction interval for Y_0 .

5.5 Suppose that the data given in the table below follow model (18).

x	5	10	15	20	25	30
y	0.10	0.21	0.30	0.35	0.44	0.62

- (i) Determine the MLE's (LSE's) of β_1 , β_2 , and σ^2 .
- (ii) Construct 95% confidence intervals for β_1 , β_2 , and σ^2 .
- (iii) At $x_0 = 17$, predict both EY_0 and Y_0 (the respective observation at x_0), and construct a 95% confidence interval and prediction interval, respectively, for them.

Hint: For a confidence interval for σ^2 , see Exercise 3.5.

5.6 The following table gives the reciprocal temperatures x and the corresponding observed solubilities of a certain chemical substance, and assume that they follow model (18).

x	3.80	3.72	3.67	3.60	3.54
y	1.27	1.20	1.10	0.82	0.65
	1.32	1.26	1.07	0.84	0.57
	1.50			0.80	0.62

- (i) Determine the MLE's (LSE's) of β_1 , β_2 , and σ^2 .
- (ii) Construct 95% confidence intervals for β_1 , β_2 , and σ^2 .
- (iii) At $x_0 = 3.77$, predict both EY_0 and Y_0 (the respective observation at x_0), and construct a 95% confidence interval and prediction interval, respectively, for them.

Note: Here $n = 13$ and $x_1 = x_2 = x_3$, $x_4 = x_5$, $x_6 = x_7$, $x_8 = x_9 = x_{10}$, and $x_{11} = x_{12} = x_{13}$.

13.6 Proof of Theorem 5

This section is solely devoted to justifying Theorem 5. Its proof is presented in considerable detail, and it makes use of some linear algebra results. The sources of those results are cited.

PROOF OF THEOREM 5 For later use, let us set

$$U_i = Y_i - \beta_1 - \beta_2 x_i, \quad \text{so that } \bar{U} = \bar{Y} - \beta_1 - \beta_2 \bar{x}, \quad (36)$$

and

$$U_i - \bar{U} = (Y_i - \bar{Y}) - \beta_2(x_i - \bar{x}) \quad \text{and} \quad Y_i - \bar{Y} = (U_i - \bar{U}) + \beta_2(x_i - \bar{x}). \quad (37)$$

Then, by (10),

$$\hat{\beta}_2 SS_x = SS_{xy} = \sum_i (x_i - \bar{x})(Y_i - \bar{Y}), \quad (38)$$

so, that

$$\begin{aligned} (\hat{\beta}_2 - \beta_2) SS_x &= \sum_i (x_i - \bar{x})(Y_i - \bar{Y}) - \beta_2 SS_x \\ &= \sum_i (x_i - \bar{x})[(U_i - \bar{U}) + \beta_2(x_i - \bar{x})] - \beta_2 SS_x \\ &= \sum_i (x_i - \bar{x})(U_i - \bar{U}) + \beta_2 SS_x - \beta_2 SS_x \\ &= \sum_i (x_i - \bar{x})(U_i - \bar{U}). \end{aligned} \quad (39)$$

Next,

$$\begin{aligned}
SS_E &= \sum_i (Y_i - \hat{Y}_i)^2 = \sum_i (Y_i - \hat{\beta}_1 - \hat{\beta}_2 x_i)^2 \\
&= \sum_i (Y_i - \bar{Y} + \hat{\beta}_2 \bar{x} - \hat{\beta}_2 x_i)^2 \quad (\text{by (8)}) \\
&= \sum_i [(Y_i - \bar{Y}) - \hat{\beta}_2 (x_i - \bar{x})]^2 \\
&= \sum_i [(Y_i - \bar{Y}) - \hat{\beta}_2 (x_i - \bar{x}) + \beta_2 (x_i - \bar{x}) - \beta_2 (x_i - \bar{x})]^2 \\
&= \sum_i \{[(Y_i - \bar{Y}) - \beta_2 (x_i - \bar{x})] - (\hat{\beta}_2 - \beta_2)(x_i - \bar{x})\}^2 \\
&= \sum_i [(U_i - \bar{U}) - (\hat{\beta}_2 - \beta_2)(x_i - \bar{x})]^2 \quad (\text{by (37)}) \\
&= \sum_i (U_i - \bar{U})^2 + (\hat{\beta}_2 - \beta_2)^2 SS_x - 2(\hat{\beta}_2 - \beta_2) \sum_i (x_i - \bar{x})(U_i - \bar{U}) \\
&= \sum_i (U_i - \bar{U})^2 + (\hat{\beta}_2 - \beta_2)^2 SS_x - 2(\hat{\beta}_2 - \beta_2)^2 SS_x \quad (\text{by (39)}) \\
&= \sum_i (U_i - \bar{U})^2 - (\hat{\beta}_2 - \beta_2)^2 SS_x \\
&= \sum_i U_i^2 - n\bar{U}^2 - (\hat{\beta}_2 - \beta_2)^2 SS_x; \text{ i.e.,} \\
SS_E &= \sum_i U_i^2 - n\bar{U}^2 - (\hat{\beta}_2 - \beta_2)^2 SS_x. \tag{40}
\end{aligned}$$

From (18) and (36), we have that the r.v.'s U_1, \dots, U_n are independent and distributed as $N(0, \sigma^2)$. Transform them into the r.v.'s V_1, \dots, V_n by means of an orthogonal transformation \mathbf{C} as described below (see also Remark 4):

$$\mathbf{C} = \begin{pmatrix} \frac{x_1 - \bar{x}}{\sqrt{SS_x}} & \frac{x_2 - \bar{x}}{\sqrt{SS_x}} & \cdots & \frac{x_n - \bar{x}}{\sqrt{SS_x}} \\ \frac{1}{\sqrt{n}} & \frac{1}{\sqrt{n}} & \cdots & \frac{1}{\sqrt{n}} \\ \text{(whatever, subject to the restriction that } \mathbf{C} \text{ is orthogonal)} \end{pmatrix}$$

That is, with “ $'$ ” standing for transpose, we have:

$$(V_1, V_2, \dots, V_n)' = \mathbf{C}(U_1, U_2, \dots, U_n)'. \tag{41}$$

Then, by Theorem 8 in Chapter 8, the r.v.'s V_1, \dots, V_n are independent and distributed as $N(0, \sigma^2)$, whereas by relation (21) in the same chapter

$$\sum_i V_i^2 = \sum_i U_i^2. \tag{42}$$

From (41),

$$V_1 = \frac{1}{\sqrt{SS_x}} \sum_i (x_i - \bar{x})U_i, \quad V_2 = \frac{1}{\sqrt{n}} \sum_i U_i = \sqrt{n} \times \frac{1}{n} \sum_i U_i = \sqrt{n}\bar{U}. \quad (43)$$

But

$$\sum_i (x_i - \bar{x})U_i = \sum_i (x_i - \bar{x})(U_i - \bar{U}) = (\hat{\beta}_2 - \beta_2)SS_x, \quad (\text{by (39)}),$$

so that

$$V_1 = (\hat{\beta}_2 - \beta_2)\sqrt{SS_x}, \quad V_1^2 = (\hat{\beta}_2 - \beta_2)^2 SS_x, \quad \text{and} \quad V_2^2 = n\bar{U}^2. \quad (44)$$

Then, from relations (40), (42), and (44), it follows that

$$SS_E = \sum_{i=1}^n V_i^2 - V_1^2 - V_2^2 = \sum_{i=3}^n V_i^2. \quad (45)$$

We now proceed with the justifications of parts (i) and (ii) of the theorem.

- (i) From (45), $\frac{SS_E}{\sigma^2} = \sum_{i=3}^n \left(\frac{V_i}{\sigma}\right)^2 \sim \chi_{n-2}^2$, since $\frac{V_i}{\sigma}$, $i = 1, \dots, n$ are independent and distributed as $N(0, 1)$.
- (ii) (a) From (44) and (45), $\hat{\beta}_2$ and SS_E are functions of nonoverlapping V_i 's (of V_1 the former, and of V_3, \dots, V_n the latter). Thus, SS_E and $\hat{\beta}_2$ are independent.
- (b) By (36) and (43), $\bar{Y} = \bar{U} + (\beta_1 + \beta_2\bar{x}) = \frac{V_2}{\sqrt{n}} + \beta_1 + \beta_2\bar{x}$, so that \bar{Y} is a function of V_2 and recall that $\hat{\beta}_2$ is a function of V_1 . Then the independence of \bar{Y} and $\hat{\beta}_2$ follows.
- (c) As was seen in (a) and (b), SS_E is a function of V_3, \dots, V_n ; \bar{Y} is a function of V_2 ; and $\hat{\beta}_2$ is a function of V_1 ; i.e., they are functions of nonoverlapping V_i 's, and therefore independent.
- (d) By (8), $\hat{\beta}_1 = \bar{Y} - \hat{\beta}_2\bar{x}$ and the right-hand side is a function of V_1 and V_2 alone, by (44) and part (b). Since SS_E is a function of V_3, \dots, V_n , by (45), the independence of SS_E and $\hat{\beta}_1$ follows. \blacktriangle

REMARK 4 There is always an orthogonal matrix \mathbf{C} with the first two rows as given above. Clearly, the vectors $\mathbf{r}_1 = (x_1 - \bar{x}, \dots, x_n - \bar{x})'$ and $\mathbf{r}_2 = (\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}})'$ are linearly independent. Then supplement them with $n-2$ vectors $\mathbf{r}_3, \dots, \mathbf{r}_n$, so that the vectors $\mathbf{r}_1, \dots, \mathbf{r}_n$ are linearly independent. Finally, use the Gram-Schmidt orthogonalization process (which leaves \mathbf{r}_1 and \mathbf{r}_2 intact) to arrive at an orthogonal matrix \mathbf{C} . (See, e.g., Theorem 1.16 and the discussion following it, in pages 33–34, of the book *Linear Algebra for Undergraduates* (1957), John Wiley & Sons, by D. C. Murdoch.)

13.7 Concluding Remarks

In this chapter, we studied the simplest linear regression model, according to which the response Y at a point x is given by $Y = \beta_1 + \beta_2x + e$. There are extensions of this model to different directions. First, the model may *not* be

linear in the parameters involved; i.e., the expectation $\eta = EY$ is not linear. Here are some such examples.

$$(i) \eta = ae^{bx}; \quad (ii) \eta = ax^b; \quad (iii) \eta = \frac{1}{a + bx}; \quad (iv) \eta = a + b\sqrt{x}.$$

It happens that these particular nonlinear models can be reduced to linear ones by suitable transformations. Thus, in (i), taking the logarithms (always with base e), we have:

$$\log \eta = \log a + bx, \quad \text{or} \quad \eta' = \beta_1 + \beta_2 x',$$

where $\eta' = \log \eta$, $\beta_1 = \log a$, $\beta_2 = b$, and $x' = x$, and the new model is linear. Likewise, in (ii):

$$\log \eta = \log a + b \log x, \quad \text{or} \quad \eta' = \beta_1 + \beta_2 x',$$

where $\eta' = \log \eta$, $\beta_1 = \log a$, $\beta_2 = b$, and $x' = \log x$, and the transformed model is linear. In (iii), simply set $\eta' = \frac{1}{\eta}$ to get $\eta' = a + bx$, or $\eta' = \beta_1 + \beta_2 x'$, where $\beta_1 = a$, $\beta_2 = b$ and $x' = x$. Finally, in (iv), let $x' = \sqrt{x}$ in order to get the linear model $\eta' = \beta_1 + \beta_2 x'$, with $\eta' = \eta$, $\beta_1 = a$, and $\beta_2 = b$.

Another direction of a generalization is the consideration of the so-called *multiple regression* linear models. In such models, there is more than one input variable x and more than two parameters β_1 and β_2 . This simply reflects the fact that the response is influenced by more than one factor each time. For example, the observation may be the systolic blood pressure of the individual in a certain group, and the influencing factors may be weight and age. The general form of a multiple regression linear model is as follows:

$$Y_i = x_{1i}\beta_1 + x_{2i}\beta_2 + \cdots + x_{pi}\beta_p + e_i, \quad i = 1, \dots, n,$$

and the assumptions attached to it are similar to those used in model (18). The analysis of such a model can be done, in principle, along the same lines as those used in analyzing model (18). However, the analysis becomes unwieldy and one has to employ, most efficiently, linear algebra methodology. Such models are referred to as *general linear models* in the statistical literature, and they have proved very useful in a host of applications. The theoretical study of such models can be found, e.g., in Chapter 16 of the book *A Course in Mathematical Statistics*, 2nd edition (1997), Academic Press, by G. G. Roussas.

Two Models of Analysis of Variance

This chapter is about statistical analysis of certain statistical models referred to as Analysis of Variance (ANOVA). There is a great variety of such models, and their detailed study constitutes an interesting branch of statistics. What is done presently is to introduce two of the simplest models of ANOVA, underline the basic concepts involved, and proceed with the analysis of the proposed models.

The first section is devoted to the study of the one-way layout ANOVA with the same number of observations for each combination of the factors involved (cells). The study consists in providing a motivation for the model, in deriving the MLE's of its parameters, and in testing an important hypothesis. In the process of doing so, an explanation is provided for the term ANOVA. Also, several technical results necessary for the analysis are established.

In the second section of the chapter, we construct confidence intervals for all so-called contrasts among the (mean) parameters of the model in Section 14.1.

Section 14.3 is a generalization of the model studied in the first section, in that the outcome of an experiment is due to two factors. Again, a motivation is provided for the model, finally, adopted, and then its statistical analysis is discussed. This analysis consists in deriving the MLE's of the parameters of the model, and also in testing two hypotheses reflecting the actual influence, or lack thereof, of the factors involved in the outcome of the underlying experiment. Again, in the process of the analysis, an explanation is provided for the term ANOVA. Also, a substantial number of technical results are stated that are necessary for the analysis. Their proofs are deferred to a final subsection of this section in order not to disrupt the continuity of arguments.

In all sections, relevant examples are discussed in detail in order to clarify the underlying ideas and apply the results obtained.

14.1 One-Way Layout with the Same Number of Observations per Cell

In this section, we derive the MLE's of the parameters μ_i , $i = 1, \dots, I$ and σ^2 described in relation (13) of Chapter 8. Next, we consider the problem of testing the null hypothesis $H_0: \mu_1 = \dots = \mu_I = \mu$ (unspecified) for which the MLE's of the parameters μ and σ^2 are to be derived under H_0 . Then we set up the likelihood ratio test, which turns out to be an F -test. For the justification of this fact, we have to split sums of squares of variations in a certain way. Actually, it is this splitting from which the name ANOVA derives. Furthermore, the splitting provides insight into what is happening behind the formal analysis.

14.1.1 The MLE's of the Parameters of the Model

First, $Y_{ij} \sim N(\mu_i, \sigma^2)$, $i = 1, \dots, I$, $j = 1, \dots, J$, and all these r.v.'s are independent. Then their likelihood function, to be denoted by $L(\mathbf{y}; \boldsymbol{\mu}, \sigma^2)$, is given by the expression below, where $\mathbf{y} = (y_1, \dots, y_J)$ and $\boldsymbol{\mu} = (\mu_1, \dots, \mu_I)$:

$$\begin{aligned} L(\mathbf{y}; \boldsymbol{\mu}, \sigma^2) &= \prod_{i,j} \left\{ \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{1}{2\sigma^2} (y_{ij} - \mu_i)^2 \right] \right\} \\ &= \prod_i \prod_j \left\{ \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{1}{2\sigma^2} (y_{ij} - \mu_i)^2 \right] \right\} \\ &= \prod_i \left\{ \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^J \exp \left[-\frac{1}{2\sigma^2} \sum_j (y_{ij} - \mu_i)^2 \right] \right\} \\ &= \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^{IJ} \prod_i \left\{ \exp \left[-\frac{1}{2\sigma^2} \sum_j (y_{ij} - \mu_i)^2 \right] \right\} \\ &= \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^{IJ} \exp \left[-\frac{1}{2\sigma^2} \sum_i \sum_j (y_{ij} - \mu_i)^2 \right]; \end{aligned}$$

following common practice, we do not explicitly indicate the range of i and j , since no confusion is possible. Hence

$$\log L(\mathbf{y}; \boldsymbol{\mu}, \sigma^2) = -\frac{IJ}{2} \log(2\pi) - \frac{IJ}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_i \sum_j (y_{ij} - \mu_i)^2. \quad (1)$$

From (1), we see that, for each fixed σ^2 , the log-likelihood is maximized with respect to μ_1, \dots, μ_I , if the exponent

$$S(\mu_1, \dots, \mu_I) = \sum_i \sum_j (y_{ij} - \mu_i)^2$$

is minimized with respect to μ_1, \dots, μ_I . By differentiation, we get

$$\frac{\partial}{\partial \mu_i} S(\mu_1, \dots, \mu_I) = -2 \sum_j y_{ij} + 2J\mu_i = 0, \quad \text{so that } \mu_i = \frac{1}{J} \sum_j y_{ij}, \quad (2)$$

and $\frac{\partial^2}{\partial \mu_i^2} \mathcal{S}(\mu_1, \dots, \mu_I) = 2J$. The resulting $I \times I$ diagonal matrix is positive definite, since, for all $\lambda_1, \dots, \lambda_I$ with $\lambda_1^2 + \dots + \lambda_I^2 > 0$,

$$\begin{aligned} (\lambda_1, \dots, \lambda_I) \begin{pmatrix} 2J & \mathbf{0} \\ \mathbf{0} & 2J \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_I \end{pmatrix} &= (2J\lambda_1, \dots, 2J\lambda_I) \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_I \end{pmatrix} \\ &= 2J(\lambda_1^2 + \dots + \lambda_I^2) > 0. \end{aligned}$$

It follows that the values of μ_i 's given in (2) are, indeed, the MLE's of the μ_i 's. That is,

$$\hat{\mu}_i = y_{i.}, \quad \text{where } y_{i.} = \frac{1}{J} \sum_j y_{ij}, \quad i = 1, \dots, I. \quad (3)$$

Now, in (1), replace the exponent by $\hat{S} = \sum_i \sum_j (y_{ij} - y_{i.})^2$ to obtain in obvious notation

$$\log \hat{L}(\mathbf{y}; \hat{\boldsymbol{\mu}}, \sigma^2) = -\frac{IJ}{2} \log(2\pi) - \frac{IJ}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \hat{S}. \quad (4)$$

Differentiating with respect to σ^2 and equating to 0, we get

$$\frac{d}{d\sigma^2} \log \hat{L}(\mathbf{y}; \hat{\boldsymbol{\mu}}, \sigma^2) = -\frac{IJ}{2\sigma^2} + \frac{\hat{S}}{2\sigma^4} = 0, \quad \text{or } \sigma^2 = \frac{\hat{S}}{IJ}. \quad (5)$$

Since $\frac{d^2}{d(\sigma^2)^2} \log \hat{L}(\mathbf{y}; \hat{\boldsymbol{\mu}}, \sigma^2) = \frac{IJ}{2(\sigma^2)^2} - \frac{2\hat{S}}{2(\sigma^2)^3}$, which evaluated at $\sigma^2 = \hat{S}/IJ$ gives: $-\frac{(IJ)^3}{2\hat{S}^2} < 0$. It follows that the value of σ^2 given in (5) is, indeed, its MLE. That is,

$$\hat{\sigma}^2 = \frac{1}{IJ} SS_e, \quad \text{where } SS_e = \sum_i \sum_j (y_{ij} - y_{i.})^2. \quad (6)$$

The results recorded in (3) and (6) provide the answer to the first objective. That is, we have established the following result.

THEOREM 1

Consider the model described in relation (13) of Chapter 8; that is, $Y_{ij} = \mu_i + e_{ij}$ where the e_{ij} 's are independently $\sim N(0, \sigma^2)$ r.v.'s, $i = 1, \dots, I (\geq 2)$, $j = 1, \dots, J (\geq 2)$. Then the MLE's of the parameters μ_i , $i = 1, \dots, I$ and σ^2 of the model are given by (3) and (6), respectively.

14.1.2 Testing the Hypothesis of Equality of Means

Next, consider the problem of testing the null hypothesis

$$H_0: \mu_1 = \dots = \mu_I = \mu \text{ (unspecified)}. \quad (7)$$

Under H_0 , the expression in (1) becomes:

$$\log L(\mathbf{y}; \mu, \sigma^2) = -\frac{IJ}{2} \log(2\pi) - \frac{IJ}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_i \sum_j (y_{ij} - \mu)^2. \quad (8)$$

Repeating a procedure similar to the one we went through above, we derive the MLE's of μ and σ^2 under H_0 , to be denoted by $\hat{\mu}$ and $\widehat{\sigma}_{H_0}^2$, respectively; i.e.,

$$\hat{\mu} = y_{..}, \quad \text{where } y_{..} = \frac{1}{IJ} \sum_i \sum_j y_{ij}, \quad \widehat{\sigma}_{H_0}^2 = \frac{1}{IJ} SS_T,$$

$$\text{where } SS_T = \sum_i \sum_j (y_{ij} - y_{..})^2. \quad (9)$$

We now proceed with the setting up of the likelihood ratio statistic $\lambda = \lambda(\mathbf{y})$ in order to test the hypothesis H_0 . To this end, first observe that, under H_0 :

$$\exp \left[-\frac{1}{2\widehat{\sigma}_{H_0}^2} \sum_i \sum_j (y_{ij} - y_{..})^2 \right] = \exp \left(-\frac{IJ}{2SS_T} \times SS_T \right) = \exp \left(-\frac{IJ}{2} \right),$$

whereas, under no restrictions imposed,

$$\exp \left[-\frac{1}{2\widehat{\sigma}^2} \sum_i (y_{ij} - y_{i.})^2 \right] = \exp \left(-\frac{IJ}{2SS_e} \times SS_e \right) = \exp \left(-\frac{IJ}{2} \right).$$

Therefore, after cancellations, the likelihood ratio statistic λ is

$$\lambda = \left(\widehat{\sigma}^2 / \widehat{\sigma}_{H_0}^2 \right)^{IJ/2}.$$

Hence $\lambda < C$, if and only if

$$\left(\frac{\widehat{\sigma}^2}{\widehat{\sigma}_{H_0}^2} \right)^{IJ/2} < C, \quad \text{or} \quad \frac{\widehat{\sigma}^2}{\widehat{\sigma}_{H_0}^2} < C^{2/IJ}, \quad \text{or} \quad \frac{\widehat{\sigma}_{H_0}^2}{\widehat{\sigma}^2} > 1/C^{2/IJ} = C_0. \quad (10)$$

At this point, we need the following result.

LEMMA 1 $SS_T = SS_e + SS_H$, where SS_e and SS_T are given by (6) and (9), respectively, and

$$SS_H = \sum_i \sum_j (y_{i.} - y_{..})^2 = J \sum_i (y_{i.} - y_{..})^2. \quad (11)$$

PROOF Deferred to Subsection 14.1.3.

According to this lemma, the last expression in (10) becomes:

$$\frac{SS_T}{SS_e} > C_0, \quad \text{or} \quad \frac{SS_e + SS_H}{SS_e} > C_0, \quad \text{or} \quad \frac{SS_H}{SS_e} > C_1 = C_0 - 1.$$

In other words, the likelihood ratio test rejects H_0 whenever

$$\frac{SS_H}{SS_e} > C_1, \quad \text{where } SS_e \text{ and } SS_H \text{ are given by (6) and (11), respectively.} \quad (12)$$

In order to determine the cutoff point C_1 in (12), we have to have the distribution, under H_0 , of the statistic SS_H/SS_e , where it is tacitly assumed that the observed values have been replaced by the respective r.v.'s. For this purpose, we need the following result.

LEMMA 2 Consider the model described in Theorem 1, and in the expressions SS_e , SS_T , and SS_H , defined by (6), (9), and (11), respectively, replace

the observed values y_{ij} , $y_{i.}$, and $y_{..}$ by the r.v.'s Y_{ij} , $Y_{i.}$, and $Y_{..}$, respectively, but retain the same notation. Then:

- (i) The r.v. SS_e/σ^2 is distributed as $\chi_{I(J-1)}^2$.
- (ii) The statistics SS_e and SS_H are independent.

Furthermore, if the null hypothesis H_0 defined in (7) is true, then:

- (iii) The r.v. SS_H/σ^2 is distributed as χ_{I-1}^2 .
- (iv) The statistic $\frac{SS_H/(I-1)}{SS_e/I(J-1)} \sim F_{I-1, I(J-1)}$.
- (v) The r.v. SS_T/σ^2 is distributed as χ_{IJ-1}^2 .

PROOF Deferred to Subsection 14.1.3.

To this lemma, there is the following corollary, which also encompasses the $\hat{\mu}_i$'s.

COROLLARY

- (i) The MLE's $\hat{\mu}_i = Y_{i.}$ are unbiased estimates of μ_i , $i = 1, \dots, I$.
- (ii) The MLE $\hat{\sigma}^2 = SS_e/IJ$ is biased, but the estimate $MS_e = SS_e/I(J-1)$ is unbiased.

PROOF (i) Immediate; (ii) Follow from Lemma 2(i). ▲

We may conclude that, on the basis of (12) and Lemma 2(iv), in order to test the hypothesis stated in (7), at level of significance α , we reject the null hypothesis H_0 whenever

$$\mathcal{F} = \frac{SS_H/(I-1)}{SS_e/I(J-1)} = \frac{MS_H}{MS_e} > F_{I-1, I(J-1); \alpha}. \quad (13)$$

So, the following result has been established.

THEOREM 2

In reference to the model described in Theorem 1, the null hypothesis H_0 defined in (7) is rejected whenever the inequality in (13) holds; the quantities SS_e and SS_H are given in relations (6) and (11), respectively, and they can be computed by using the formulas in (14) below.

REMARK 1 At this point, it should be recalled that the point $F_{m,n;\alpha}$ is determined, so that $P(X > F_{m,n;\alpha}) = \alpha$, where X is a r.v. distributed as $F_{m,n}$; see Figure 14.1.


REMARK 2 By rewriting analytically the relation in Lemma 1, we have that:

$$\sum_i \sum_j (y_{ij} - y_{..})^2 = \sum_i \sum_j (y_{ij} - y_{i.})^2 + \sum_i \sum_j (y_{i.} - y_{..})^2.$$

That is, the *total variation* of the y_{ij} 's with respect to the grand mean $y_{..}$ is split into two parts: the variation of the y_{ij} 's in each i th group with respect

Figure 14.1

The Graph of the p.d.f. of the $F_{m,n}$ Distribution Along with the Rejection Region of H_0 and the Level of Significance



to their mean y_i in that group (*variation within groups*), and the variation of the I means $y_i, i = 1, \dots, I$ from the grand mean $y_{..}$ (*variation between groups*). By changing the term “variation” to “variance,” we are led to the term ANOVA. The expression SS_e is also referred to as the *error sum of squares* for obvious reasons, and the expression SS_H is also referred to as the *treatment sum of squares*, since it reflects variations due to treatment differences. The subscript H accounts for the fact that this statistic is instrumental in testing H_0 ; see also Lemma 3 below. Finally, the expression SS_T is called the *total sum of squares*, again for obvious reasons.

The various quantities employed in carrying out the test described in (13) are usually gathered together in the form of a table, an ANOVA table, as is done in Table 14.1.

REMARK 3 For computational purposes, we have:

$$SS_H = J \sum_i Y_i^2 - I J Y_{..}^2, \quad SS_e = \sum_i \sum_j Y_{ij}^2 - J \sum_i Y_i^2. \quad (14)$$

Indeed,

$$\begin{aligned} SS_H &= J \sum_i (Y_i - Y_{..})^2 = J \sum_i Y_i^2 + I J Y_{..}^2 - 2 J Y_{..} \sum_i Y_i \\ &= J \sum_i Y_i^2 + I J Y_{..}^2 - 2 J Y_{..} I Y_{..} = J \sum_i Y_i^2 - I J Y_{..}^2. \end{aligned}$$

Table 14.1

Analysis of Variance for One-Way Layout

Source of Variance	Sums of Squares	Degrees of Freedom	Mean Squares
Between groups	$SS_H = J \sum_{i=1}^I (Y_i - Y_{..})^2$	$I - 1$	$MS_H = \frac{SS_H}{I-1}$
Within groups	$SS_e = \sum_{i=1}^I \sum_{j=1}^J (Y_{ij} - Y_i)^2$	$I(J - 1)$	$MS_e = \frac{SS_e}{I(J-1)}$
Total	$SS_T = \sum_{i=1}^I \sum_{j=1}^J (Y_{ij} - Y_{..})^2$	$IJ - 1$	—

Also,

$$SS_e = \sum_i \sum_j (Y_{ij} - Y_i)^2 = \sum_i \sum_j Y_{ij}^2 + J \sum_i Y_i^2 - 2 \sum_i \sum_j Y_i Y_{ij},$$

and

$$\sum_i \sum_j Y_i Y_{ij} = \sum_i Y_i \sum_j Y_{ij} = \left(\sum_i Y_i \right) (JY_i) = J \sum_i Y_i^2,$$

so that the result follows.

EXAMPLE 1

For a numerical example, take $I = 3$, $J = 5$, and let:

$$\begin{array}{lll} y_{11} = 82 & y_{21} = 61 & y_{31} = 78 \\ y_{12} = 83 & y_{22} = 62 & y_{32} = 72 \\ y_{13} = 75 & y_{23} = 67 & y_{33} = 74 \\ y_{14} = 79 & y_{24} = 65 & y_{34} = 75 \\ y_{15} = 78 & y_{25} = 64 & y_{35} = 72. \end{array}$$

- (i) Compute the MLE of μ_i , $i = 1, 2, 3$.
- (ii) Compute the sum of squares SS_H and SS_e , and also the unbiased estimate MS_e of σ^2 .
- (iii) Test the hypothesis $H_0: \mu_1 = \mu_2 = \mu_3 = \mu$ at level of significance $\alpha = 0.05$.
- (iv) Compute the MLE of μ .

DISCUSSION

- (i) $\hat{\mu}_1 = 79.4$, $\hat{\mu}_2 = 63.8$, $\hat{\mu}_3 = 74.2$.
- (ii) Since $y_i = \hat{\mu}_i$, $i = 1, 2, 3$ and $y_{..} \simeq 72.467$, we get, by (14):

$$SS_H \simeq 79,402.2 - 78,771.991 = 630.209,$$

$$SS_e = 79,491 - 79,402.2 = 88.8, \quad \text{and} \quad MS_e = \frac{88.8}{12} = 7.4.$$

Since $MS_H \simeq 315.105$, the test statistic is: $\frac{315.105}{7.4} \simeq 42.582$. On the other hand, $F_{2,12;0.05} = 3.8853$, the hypothesis H_0 is rejected.

- (iii) Finally, $\hat{\mu} = y_{..} = 72.467$.

Here is another example with data from a real experiment.

EXAMPLE 2

In an effort to improve the quality of recording tapes, the effects of four kinds of coatings A, B, C, D on the reproducing quality of sound are compared. Suppose that the measurements of sound distortion given in Table 14.2 are obtained from tapes treated with the four coatings. Look at this problem as a one-way layout ANOVA and carry out the analysis; take as level of significance $\alpha = 0.05$.

DISCUSSION Here $I = 4$, $J = 4$. For the MLE's of the means, we have: $\hat{\mu}_1 = y_{1.} = 11.25$, $\hat{\mu}_2 = y_{2.} = 17.00$, $\hat{\mu}_3 = y_{3.} = 15.50$, $\hat{\mu}_4 = y_{4.} = 14.75$. Also,

$y_{..} = 14.625$. Next, by (14): $SS_H = 4 \times 873.375 - 16 \times 213.890625 = 3,493.50 - 3,422.25 = 71.25$, $SS_e = 3,568 - 4 \times 873.375 = 3,568 - 3,493.5 = 74.50$, so that $MS_H = 23.75$, $MS_e \simeq 6.208$. The observed value of the test statistics is: $\frac{23.75}{6.208} \simeq 3.826$, whereas $F_{3,12;0.05} = 3.4903$. Therefore the null hypothesis about equality of the means is rejected. Finally, $\hat{\mu}_{..} = y_{..} = 14.625$.

Table 14.2

Sound Distortion
Obtained with Four
Types of Coatings

Coating	Observations	Mean	Grand Mean
A	10, 15, 8, 12	$y_{1.} = 11.25$	$y_{..} = 14.625$
B	14, 18, 21, 15	$y_{2.} = 17.00$	
C	17, 16, 14, 15	$y_{3.} = 15.50$	
D	12, 15, 17, 15	$y_{4.} = 14.75$	

The following observations are meant to shed more light on the F test which is used for testing the null hypothesis H_0 . Recall that $\mathcal{F} = \frac{MS_H}{MS_e}$, where, by (11) and (13),

$$MS_H = \frac{J}{I-1} \sum_i (Y_i - Y_{..})^2$$

and that $\frac{SS_H}{\sigma^2} \sim \chi_{I-1}^2$, under H_0 , so that $EMS_H = \sigma^2$. It will be shown below that, regardless whether H_0 is true or not,

$$EMS_H = \sigma^2 + \frac{J}{I-1} \sum_i (\mu_i - \mu)^2, \quad \text{where } \mu = \frac{1}{I} \sum_i \mu_i. \quad (15)$$

Therefore $EMS_H \geq \sigma^2 = E(MS_H | H_0)$ and $EMS_H = \sigma^2$ under H_0 ; also, $EMS_e = \sigma^2$. Thus, on the basis of this average criterion, it makes sense to reject H_0 when MS_H , measured against MS_e , takes large values. For reference purposes, relation (15) is stated below as a lemma.

LEMMA 3 It holds that:

$$\begin{aligned} EMS_H &= \frac{1}{I-1} E \sum_i \sum_j (Y_i - Y_{..})^2 = \frac{J}{I-1} E \sum_i (Y_i - Y_{..})^2 \\ &= \sigma^2 + \frac{J}{I-1} \sum_i (\mu_i - \mu)^2. \end{aligned}$$

PROOF Deferred to Subsection 14.1.3.

14.1.3 Proof of Lemmas in Section 14.1

We now proceed with the justification of Lemmas 1–3 in this section.

PROOF OF LEMMA 1 We have:

$$\begin{aligned} SS_T &= \sum_i \sum_j (y_{ij} - y_{..})^2 = \sum_i \sum_j [(y_{ij} - y_{i.}) + (y_{i.} - y_{..})]^2 \\ &= \sum_i \sum_j (y_{ij} - y_{i.})^2 + \sum_i \sum_j (y_{i.} - y_{..})^2 + 2 \sum_i \sum_j (y_{ij} - y_{i.})(y_{i.} - y_{..}) \end{aligned}$$

$$\begin{aligned}
&= SS_e + SS_H, \quad \text{since} \\
&\quad \sum_i \sum_j (y_{ij} - y_i)(y_i - y_{..}) = \sum_i (y_i - y_{..}) \sum_j (y_{ij} - y_i) \\
&= \sum_i (y_i - y_{..})(Jy_i - Jy_i) = 0. \quad \blacktriangle
\end{aligned}$$

PROOF OF LEMMA 2 At this point, recall that, if X_1, \dots, X_n are independent r.v.'s distributed as $N(\mu, \sigma^2)$, then: (a) \bar{X} and $\sum_i (X_i - \bar{X})^2$ are independent; (b) $\frac{1}{\sigma^2} \sum_i (X_i - \bar{X})^2 \sim \chi_{n-1}^2$. Apply these results as follows:

- (i) For each $i = 1, \dots, I$, $\frac{1}{\sigma^2} \sum_j (Y_{ij} - Y_i)^2 \sim \chi_{J-1}^2$, by (b) above. Furthermore, for $i' \neq i$, $\sum_j (Y_{i'j} - Y_{i'})^2$ and $\sum_j (Y_{ij} - Y_i)^2$ are independent, since they are defined (separately) on sets of independent r.v.'s. It follows that:

$$\frac{SS_e}{\sigma^2} = \frac{1}{\sigma^2} \sum_i \sum_j (Y_{ij} - Y_i)^2 = \sum_i \frac{1}{\sigma^2} \sum_j (Y_{ij} - Y_i)^2 \sim \chi_{I(J-1)}^2.$$

- (ii) For each $i = 1, \dots, I$, the independent r.v.'s $Y_{i1}, \dots, Y_{iJ} \sim N(\mu_i, \sigma^2)$. Hence, by (a) above, $\sum_j (Y_{ij} - Y_i)^2$ and Y_i are independent. Furthermore, $\sum_j (Y_{ij} - Y_i)^2$ and $Y_{i'}$ are also independent for $i' \neq i$, because $Y_{i'}$ is defined on a set of r.v.'s which are independent of $\sum_j (Y_{ij} - Y_i)^2$. Thus, each of the statistics $\sum_j (Y_{ij} - Y_i)^2$, $i = 1, \dots, I$ is independent of the statistics Y_1, \dots, Y_I , and the statistics $\sum_j (Y_{ij} - Y_i)^2$, $i = 1, \dots, I$ are independent, as was seen in part (i). It follows that the sets $\sum_j (Y_{ij} - Y_i)^2$, $i = 1, \dots, I$ and Y_i , $i = 1, \dots, I$ are independent. Then so are functions defined (separately) on them. In particular, the functions $\sum_i \sum_j (Y_{ij} - Y_i)^2$ and $\sum_i \sum_j (Y_i - Y_{..})^2$ are independent, or SS_e and SS_H are independent.
- (iii) Under H_0 , the r.v.'s Y_1, \dots, Y_I are independent and distributed as $N(\mu, \sigma^2/J)$. Therefore $\frac{J}{\sigma^2} \sum_i (Y_i - Y_{..})^2 \sim \chi_{I-1}^2$. Since $\frac{J}{\sigma^2} \sum_i (Y_i - Y_{..})^2 = \frac{1}{\sigma^2} \sum_i \sum_j (Y_i - Y_{..})^2 = \frac{1}{\sigma^2} SS_H$, the result follows.
- (iv) It follows from parts (i)–(iii) and the definition of the F distribution.
- (v) Under H_0 , the r.v.'s Y_{ij} , $i = 1, \dots, I$, $j = 1, \dots, J$ are independent and distributed as $N(\mu, \sigma^2)$. Then, by (b) above, $\frac{1}{\sigma^2} \sum_i \sum_j (Y_{ij} - Y_i)^2$ is distributed as $\chi_{I(J-1)}^2$, or $\frac{SS_e}{\sigma^2} \sim \chi_{I(J-1)}^2$. \blacktriangle

PROOF OF LEMMA 3 Before taking expectations, work with $\sum_i (Y_i - Y_{..})^2$ and rewrite it in a convenient form; namely,

$$\sum_i (Y_i - Y_{..})^2 = \sum_i [(Y_i - \mu) - (Y_{..} - \mu)]^2 = \sum_i (Y_i - \mu)^2 - I(Y_{..} - \mu)^2,$$

because

$$\begin{aligned}
\sum_i (Y_i - \mu) &= \sum_i Y_i - I\mu = \sum_i \frac{1}{J} \sum_j Y_{ij} - I\mu \\
&= \frac{1}{J} \sum_i \sum_j Y_{ij} - I\mu = IY_{..} - I\mu = I(Y_{..} - \mu),
\end{aligned}$$

so that

$$-2 \sum_i (Y_i - \mu)(Y_{..} - \mu) = -2(Y_{..} - \mu) \times I(Y_{..} - \mu) = -2I(Y_{..} - \mu)^2.$$

So,

$$\begin{aligned} E \sum_i (Y_i - Y_{..})^2 &= \sum_i E(Y_i - \mu)^2 - IE(Y_{..} - \mu)^2 = \sum_i E(Y_i - \mu)^2 - I \text{Var}(Y_{..}) \\ &= \sum_i E(Y_i - \mu)^2 - I \frac{\sigma^2}{IJ} = \sum_i E(Y_i - \mu)^2 - \frac{\sigma^2}{J}, \end{aligned}$$

and

$$\begin{aligned} E(Y_i - \mu)^2 &= E[(Y_i - \mu_i) + (\mu_i - \mu)]^2 = E(Y_i - \mu_i)^2 + (\mu_i - \mu)^2 \\ &= \text{Var}(Y_i) + (\mu_i - \mu)^2 = \frac{\sigma^2}{J} + (\mu_i - \mu)^2. \end{aligned}$$

Therefore

$$E \sum_i (Y_i - Y_{..})^2 = \frac{I\sigma^2}{J} + \sum_i (\mu_i - \mu)^2 - \frac{\sigma^2}{J} = \frac{I-1}{J}\sigma^2 + \sum_i (\mu_i - \mu)^2,$$

and, by (11) and (13),

$$EMS_H = \sigma^2 + \frac{J}{I-1} \sum_i (\mu_i - \mu)^2, \quad \text{which is (15).} \quad \blacktriangle$$

REMARK 4 In closing this section, it should be pointed out that an obvious generalization of what was done here is to have different J 's for each $i = 1, \dots, I$; i.e., for each i , we have J_i observations. The analysis conceptually remains the same, only one would have to carry along the J_i 's as opposed to one J .

Exercises

- 1.1** Apply the one-way layout analysis of variance to the data given in the table below. Take $\alpha = 0.05$.

A	B	C
10.0	9.1	9.2
11.5	10.3	8.4
11.7	9.4	9.4

- 1.2** Consider the log-likelihood function (1) as it becomes under the null hypothesis H_0 stated in (7), and show that the MLE's of μ and σ^2 are given by the expression in relation (9).
- 1.3** In reference to the derivation of the likelihood ratio test λ for testing the hypothesis H_0 stated in relation (7), show that λ is, actually, given by any one of the expressions in relation (10).

1.4 In reference to the proof of Lemma 3, show that:

- (i) $E(Y_{..} - \mu_{.})^2 = \frac{\sigma^2}{IJ}$.
- (ii) $E(Y_i - \mu_i)^2 = \frac{\sigma^2}{J}$.
- (iii) $E(Y_i - \mu_{.})^2 = E(Y_i - \mu_i)^2 + (\mu_i - \mu_{.})^2$.

14.2 A Multicomparison Method

Refer to the one-way layout model discussed in the previous section, namely, to the model described in Theorem 1. One of the problems we have studied was that of testing the null hypothesis about equality of the I means; i.e.,

$$H_0: \mu_1 = \cdots = \mu_I = \mu \text{ unspecified.} \quad (16)$$

Suppose now that the hypothesis H_0 is rejected, as, indeed, was the case in Examples 1 and 2. Rejection of H_0 simply means that not *all* of the μ_i 's are equal. Clearly, it would be desirable to know which of the μ_i 's are responsible for the rejection of H_0 . It is true that we do gain some information about it by looking at the estimates $\hat{\mu}_i$. However, we would like to obtain additional information analogous to that provided by a confidence interval for a real-valued parameter. This is the problem to examine in this section.

In relation (16), the hypothesis H_0 compares the parameters involved and, actually, stipulates that they are all equal. This suggests that any attempt to construct a confidence interval should not focus on a single parameter, but rather on two or more parameters simultaneously. For example, we would like to compare all possible pairs (μ_i, μ_j) through the differences $\mu_i - \mu_j$. Or, more generally, to compare one subset of these parameters against the complement of this subset. Thus, in Example 1, where we have three parameters μ_1, μ_2 , and μ_3 , we may wish, e.g., to compare μ_1 against (μ_2, μ_3) , or μ_2 against (μ_1, μ_3) , or μ_3 against (μ_1, μ_2) . One way of doing it is to look at the respective differences: $\mu_1 - \frac{1}{2}(\mu_1 + \mu_2)$, $\mu_2 - \frac{1}{2}(\mu_1 + \mu_3)$, $\mu_3 - \frac{1}{2}(\mu_1 + \mu_2)$.

At this point, it is to be observed that all expressions we looked at above are of the form $c_1\mu_1 + \cdots + c_I\mu_I$ with $c_1 + \cdots + c_I = 0$. This observation leads to the following definition.

DEFINITION 1

In reference to the model described in Theorem 1, any relation among the parameters μ_1, \dots, μ_I of the form $\Psi = \sum_{i=1}^I c_i\mu_i$ with $\sum_{i=1}^I c_i = 0$ is called a *contrast* among the μ_i 's.

It follows from the above discussion that what would be really meaningful here would be the construction of confidence intervals for contrasts among the μ_i 's. In particular, it would be clearly, highly desirable to construct confidence intervals for all possible contrast among the μ_i 's, which would *all* have the same confidence coefficient. This is exactly the content of the theorem stated below.

First, let us introduce some pieces of notation needed. To this end, consider the contrast

$$\Psi = \sum_i c_i \mu_i \left(\sum_i c_i = 0 \right), \quad (17)$$

and let us estimate Ψ by $\hat{\Psi}$, where

$$\hat{\Psi} = \sum_i c_i \hat{\mu}_i = \sum_i c_i Y_i. \quad (18)$$

Clearly,

$$E\hat{\Psi} = \Psi \quad \text{and} \quad \text{Var}(\hat{\Psi}) = \left(\frac{1}{J} \sum_i c_i^2 \right) \sigma^2, \quad (19)$$

and the variance is estimated, in an obvious manner, by

$$\widehat{\text{Var}}(\hat{\Psi}) = \left(\frac{1}{J} \sum_i c_i^2 \right) MS_e, \quad MS_e = SS_e / I(J-1), \quad (20)$$

and SS_e is given in (6) (see also (14)). Finally, define S^2 by:

$$S^2 = (I-1)F_{I-1, I(J-1); \alpha}. \quad (21)$$

Then we have the following important result.

THEOREM 3

With the notation introduced in (17), (18), (20), and (21), the interval

$$\left(\hat{\Psi} - S\sqrt{\widehat{\text{Var}}(\hat{\Psi})}, \quad \hat{\Psi} + S\sqrt{\widehat{\text{Var}}(\hat{\Psi})} \right) \quad (22)$$

is a confidence interval with confidence coefficient $1 - \alpha$ *simultaneously* for all contrasts Ψ .

At this point, it should not come as a surprise that there is an intimate relationship between the null hypothesis H_0 and confidence intervals for contrast. The result stated below as a lemma (but not proved!) articulates this relationship. In its statement, we need a concept defined now.

DEFINITION 2

Let Ψ and $\hat{\Psi}$ be as in (17) and (18), respectively. Then we say that $\hat{\Psi}$ is *significantly different from zero*, if the interval defined in (22) does not contain zero; equivalently, $|\hat{\Psi}| > S\sqrt{\widehat{\text{Var}}(\hat{\Psi})}$.

Then the lemma mentioned above is as follows.

LEMMA 4 The null hypothesis H_0 stated in (16) is rejected, if and only if there is at least one contrast Ψ for which $\hat{\Psi}$ is significantly different from zero.

We do not intend to pursue the proof of Theorem 3 here, which can be found in great detail in Section 17.4 of the book *A Course in Mathematical*

Statistics, 2nd edition, Academic Press (1997), by G. G. Roussas. Suffice it to say that it follows from the maximization, with respect to c_1, \dots, c_I , subject to the contrast constraint $\sum_i c_i = 0$, of the function

$$f(c_1, \dots, c_I) = \frac{1}{\sqrt{\frac{1}{J} \sum_i c_i^2}} \sum_i c_i (Y_i - \mu_i),$$

and that this maximization is obtained by means of the so-called Lagrange multipliers. In the process of doing so, we also need the following two facts.

LEMMA 5

- (i) With $\mu_{.} = \frac{1}{J} \sum_i \mu_i$, the r.v.'s $\sum_j [(Y_{ij} - \mu_i) - (Y_i - \mu_i)]^2$, $i = 1, \dots, I$ are independent.
- (ii) The r.v.'s $\sum_i [(Y_i - Y_{.}) - (\mu_i - \mu_{.})]^2$ and SS_e are independent.
- (iii) Under the null hypothesis H_0 , $\frac{J}{\sigma^2} \sum_i [(Y_i - Y_{.}) - (\mu_i - \mu_{.})]^2 \sim \chi_{I-1}^2$.

PROOF Deferred to the end of the section.

We now consider some examples.

EXAMPLE 3

In reference to Example 1, construct a 95% confidence interval for each of the following contrasts:

$$\begin{aligned} \mu_1 - \mu_2, \quad \mu_1 - \mu_3, \quad \mu_2 - \mu_3, \quad \mu_1 - \frac{1}{2}(\mu_2 + \mu_3), \\ \mu_2 - \frac{1}{2}(\mu_3 + \mu_1), \quad \mu_3 - \frac{1}{2}(\mu_1 + \mu_2). \end{aligned}$$

DISCUSSION Here $I = 3$, $J = 5$, and hence $F_{I-1, I(J-1); \alpha} = F_{2, 12; 0.05} = 3.8853$, $S^2 = (I - 1)F_{I-1, I(J-1); \alpha} = 2 \times 3.8853 = 7.7706$ and $S \simeq 2.788$. Also, $MS_e = 7.4$ from Example 1. From the same example, for $\Psi = \mu_1 - \mu_2$, we have $\hat{\Psi} = Y_1 - Y_2 = 79.4 - 63.8 = 15.6$, $\widehat{Var}(\hat{\Psi}) = \frac{2}{5} \times 7.4 = 2.96$, $\sqrt{\widehat{Var}(\hat{\Psi})} \simeq 1.72$ and $S\sqrt{\widehat{Var}(\hat{\Psi})} = 2.788 \times 1.72 \simeq 4.795$. Therefore the required (observed) confidence interval for $\mu_1 - \mu_2$ is:

$$[15.6 - 4.795, \quad 15.6 + 4.795] = [10.805, \quad 20.395].$$

Likewise, for $\Psi = \mu_1 - \mu_3$, we have $\hat{\Psi} = Y_1 - Y_3 = 79.4 - 74.2 = 5.2$, $\widehat{Var}(\hat{\Psi}) = 2.96$ the same as before, and hence $S\sqrt{\widehat{Var}(\hat{\Psi})} \simeq 4.795$. Then the required (observed) confidence interval for $\mu_1 - \mu_3$ is:

$$[5.2 - 4.795, \quad 5.2 + 4.795] = [0.405, \quad 9.995].$$

Also, for $\Psi = \mu_2 - \mu_3$, we have $\hat{\Psi} = Y_2 - Y_3 = 63.8 - 74.2 = -10.4$. Since the $\widehat{Var}(\hat{\Psi})$ is still 2.96, the required (observed) confidence interval for $\mu_2 - \mu_3$ is:

$$[-10.4 - 4.795, \quad -10.4 + 4.795] = [-15.195, \quad -5.605].$$

Next, let $\Psi = \mu_1 - \frac{1}{2}(\mu_2 + \mu_3)$, so that $\hat{\Psi} = 79.4 - \frac{1}{2}(63.8 + 74.2) = 79.4 - 69 = 10.4$, $\widehat{Var}(\hat{\Psi}) = \frac{3}{10} \times 7.4 = 2.22$, $\sqrt{\widehat{Var}(\hat{\Psi})} \simeq 1.49$ and $S\sqrt{\widehat{Var}(\hat{\Psi})} \simeq 4.154$.

Therefore the required (observed) confidence interval is:

$$[10.4 - 4.154, \quad 10.4 + 4.154] = [6.246, \quad 14.554].$$

For $\Psi = \mu_2 - \frac{1}{2}(\mu_3 + \mu_1)$, we have $\hat{\Psi} = 63.8 - \frac{1}{2}(74.2 + 79.4) = -13$, and therefore the required (observed) confidence interval is:

$$[-13 - 4.154, \quad -13 + 4.154] = [-17.154, \quad -8.846].$$

Finally, for $\Psi = \mu_3 - \frac{1}{2}(\mu_1 + \mu_2)$, we have $\hat{\Psi} = 74.2 - \frac{1}{2}(79.4 + 63.8) = 2.6$, and the required (observed) confidence interval is:

$$[2.6 - 4.154, \quad 2.6 + 4.154] = [-1.554, \quad 6.754].$$

It is noteworthy that of the six contrasts we have entertained in this example, for only one contrast $\Psi = \mu_3 - \frac{1}{2}(\mu_1 + \mu_2)$ the respective quantity $\hat{\Psi} = 2.6$, is *not* significantly different from zero. This is consonant with Lemma 4, since we already know (from Example 1) that H_0 is rejected. For example, for the contrast $\Psi = \mu_1 - \mu_2$, we found the confidence interval (10.805, 20.395), which does not contain 0. This simply says that, at the confidence level considered, μ_1 and μ_2 cannot be equal; thus, H_0 would have to be rejected. Likewise for the contrasts $\mu_1 - \mu_3$ and $\mu_2 - \mu_3$.

EXAMPLE 4

In reference to Example 2, construct a 95% confidence interval for each of the following contrasts:

$$\mu_1 - \mu_2, \quad \mu_1 - \mu_3, \quad \mu_1 - \mu_4, \quad \mu_2 - \mu_3, \quad \mu_2 - \mu_4, \quad \mu_3 - \mu_4.$$

DISCUSSION Here $I = J = 4$, $F_{I-1, I(J-1); \alpha} = F_{3, 12; 0.05} = 3.4903$, $S^2 = (I - 1)F_{I-1, I(J-1); \alpha} = 3 \times 3.4903 = 10.4709$, and $S \simeq 3.236$. For $\Psi = \mu_1 - \mu_2$, we have $\hat{\Psi} = Y_{1.} - Y_{2.} = 11.25 - 17 = -5.75$, and $\widehat{Var}(\hat{\Psi}) = 0.5 \times 6.208 = 3.104$, $\sqrt{\widehat{Var}(\hat{\Psi})} \simeq 1.762$. Thus, $S\sqrt{\widehat{Var}(\hat{\Psi})} = 3.236 \times 1.762 \simeq 5.702$. Then the required (observed) confidence interval for $\mu_1 - \mu_2$ is:

$$[-5.75 - 5.702, \quad -5.75 + 5.702] = [-11.452, \quad -0.048].$$

For $\Psi = \mu_1 - \mu_3$, we have $\hat{\Psi} = 11.25 - 15.50 = -4.25$, and the required (observed) confidence interval for $\mu_1 - \mu_3$ is:

$$[-4.25 - 5.702, \quad -4.25 + 5.702] = [-9.952, \quad 1.452].$$

For $\Psi = \mu_1 - \mu_4$, we have $\hat{\Psi} = 11.25 - 14.75 = -3.5$, and the required (observed) confidence interval is:

$$[-3.5 - 5.702, \quad -3.5 + 5.702] = [-9.202, \quad 2.202].$$

For $\Psi = \mu_2 - \mu_3$, we have $\hat{\Psi} = 17 - 15.5 = 1.5$, and the required (observed) confidence interval is:

$$[1.5 - 5.702, \quad 1.5 + 5.702] = [-4.202, \quad 7.202].$$

For $\Psi = \mu_2 - \mu_4$, we have $\hat{\Psi} = 17 - 14.75 = 2.25$, and the required (observed) confidence interval is:

$$[2.25 - 5.702, \quad 2.25 + 5.702] = [-3.452, \quad 7.952].$$

Finally, for $\Psi = \mu_3 - \mu_4$, we have $\hat{\Psi} = 15.50 - 14.75 = 0.75$, and the required (observed) confidence interval is:

$$[0.75 - 5.702, \quad 0.75 + 5.702] = [-4.952, \quad 6.452].$$

In this example, we have that for *only one* of the six contrasts considered, $\Psi = \mu_1 - \mu_2$, the respective $\hat{\Psi} = -5.75$, is significantly different from zero. This, of course, suffices for the rejection of the hypothesis H_0 (according to Lemma 4), as it actually happened in Lemma 2. So, it appears that the means μ_1 and μ_2 are the culprit here. This fact is also reflected by the estimates of the μ_i 's found in Example 2; namely,

$$\hat{\mu}_1 = 11.25, \quad \hat{\mu}_2 = 17.00, \quad \hat{\mu}_3 = 15.50, \quad \hat{\mu}_4 = 14.75;$$

$\hat{\mu}_1$ and $\hat{\mu}_2$ are the furthest apart.

This section is concluded with the presentation of a justification of Lemma 5.

PROOF OF LEMMA 5

(i) Here

$$\sum_j [(Y_{ij} - \mu_i) - (Y_i - \mu_i)]^2 = \sum_j (Y_{ij} - Y_i)^2,$$

and the statistics $\sum_j (Y_{ij} - Y_i)^2$, $i = 1, \dots, I$ are independent, since they are defined (separately) on independent sets (rows) of r.v.'s.

(ii) The proof of this part is reminiscent of that of Lemma 2(ii). The independent r.v.'s $Y_{i1} - \mu_i, \dots, Y_{iJ} - \mu_i$ are distributed as $N(0, \sigma^2)$. Since $\frac{1}{J} \sum_j (Y_{ij} - \mu_i) = Y_i - \mu_i$, it follows, by an application of (a) in the proof of Lemma 2, that, for each $i = 1, \dots, I$, $\sum_j [(Y_{ij} - \mu_i) - (Y_i - \mu_i)]^2 = \sum_j (Y_{ij} - Y_i)^2$ is independent of $Y_i - \mu_i$. For $i' \neq i$, each of $\sum_j (Y_{ij} - Y_i)^2$ is also independent of $Y_{i'} - \mu_{i'}$. Also, by part (i), $\sum_j (Y_{ij} - Y_i)^2$, $i = 1, \dots, I$ are independent. It follows that the sets of statistics

$$\sum_j (Y_{ij} - Y_i)^2, \quad i = 1, \dots, I \quad \text{and} \quad Y_i - \mu_i, \quad i = 1, \dots, I$$

are independent. Then so are functions defined (separately) on them. In particular, this is true for the functions

$$\sum_i \sum_j (Y_{ij} - Y_i)^2 = SS_e \quad \text{and} \\ \sum_i [(Y_i - \mu_i) - (Y_{..} - \mu_{..})]^2 = \sum_i [(Y_i - Y_{..}) - (\mu_i - \mu_{..})]^2.$$

(iii) For $i = 1, \dots, I$, the r.v.'s $Y_i - \mu_i$ are independent and distributed as $N(0, \sigma^2/J)$, so that the independent r.v.'s $\frac{\sqrt{J}}{\sigma}(Y_i - \mu_i)$, $i = 1, \dots, I$ are distributed as $N(0, 1)$.

Since $\frac{1}{J} \sum_i \frac{\sqrt{J}}{\sigma} (Y_i - \mu_i) = \frac{\sqrt{J}}{\sigma} (Y_{..} - \mu_{..})$, it follows (by (a) in the proof of Lemma 2) that

$$\sum_i \left[\frac{\sqrt{J}}{\sigma} (Y_i - \mu_i) - \frac{\sqrt{J}}{\sigma} (Y_{..} - \mu_{..}) \right]^2 \sim \chi_{I-1}^2,$$

or

$$\frac{J}{\sigma^2} \sum_i [(Y_i - Y_{..}) - (\mu_i - \mu_{..})]^2 \sim \chi_{I-1}^2. \quad \blacktriangle$$

Exercises

- 2.1** Refer to Exercise 1.1, and construct 95% confidence intervals for all contrasts of the μ 's.
- 2.2** Refer to Exercise 1.2, and construct 95% confidence intervals for all contrasts of the μ 's.

14.3 Two-Way Layout with One Observation per Cell

In this section, we pursue the study of the kind of problems considered in Section 14.1, but in a more general framework. Specifically, we consider experiments whose outcomes are influenced by more than one factor. In the model to be analyzed here there will be two such factors, one factor occurring at I levels and the other factor occurring at J levels. The following example will help clarify the underlying ideas and the issues to be resolved.

EXAMPLE 5

Suppose we are interested in acquiring a fairly large number of equipments from among I brands entertained. The available workforce to use the equipments bought consists of J workers. Before a purchase decision is made, an experiment is carried out whereby each one of the J workers uses each one of the I equipments for one day. It is assumed that the one day's production would be a quantity, denoted by μ_{ij} , depending on the i th brand of equipment and the j th worker, except for an error e_{ij} associated with the i th equipment and the j th worker. Thus, the one day's outcome is, actually, an observed value of a r.v. Y_{ij} , which has the following structure: $Y_{ij} = \mu_{ij} + e_{ij}$, $i = 1, \dots, I$, $j = 1, \dots, J$. For the errors e_{ij} the familiar assumptions are made; namely, the r.v.'s e_{ij} , $i = 1, \dots, I$, $j = 1, \dots, J$ are independent and distributed as $N(0, \sigma^2)$. It follows that the r.v.'s Y_{ij} , $i = 1, \dots, I$, $j = 1, \dots, J$ are independent with $Y_{ij} \sim N(\mu_{ij}, \sigma^2)$. At this point, the further reasonable assumption is made that each mean μ_{ij} consists of three *additive* parts: a quantity μ , the *grand mean*, the same for all i and j ; an effect due the i th equipment, denoted by α_i and usually referred to as the *row effect*; and an effect due to the j th worker, denoted by β_j and usually referred to as the *column effect*. So, $\mu_{ij} = \mu + \alpha_i + \beta_j$. Now, it is not unreasonable to assume that some of the α_i effects are positive,

some are negative, and on the whole their sum is zero. Likewise for the β_j effects.

Gathering together the assumptions made so far, we have the following model.

$$\left. \begin{array}{l} Y_{ij} = \mu + \alpha_i + \beta_j + e_{ij}, \sum_{i=1}^I \alpha_i = 0 \text{ and } \sum_{j=1}^J \beta_j = 0, \text{ the r.v.'s} \\ e_{ij}, i = 1, \dots, I (\geq 2), j = 1, \dots, J (\geq 2) \text{ are independent and distributed} \\ \text{as } N(0, \sigma^2). \\ \text{It follows that the r.v.'s } Y_{ij}, i = 1, \dots, I, j = 1, \dots, J \\ \text{are independent with } Y_{ij} \sim N(\mu + \alpha_i + \beta_j, \sigma^2). \end{array} \right\} \quad (23)$$

Of course, once model (23) is arrived at, it can be detached from the specific example which helped motivate the model.

In reference to model (23), the questions which arise naturally are the following: What are the magnitudes of the grand mean μ , of the row effects α_i , of the column effects β_j , and of the error variance σ^2 ? Also, are there, really, any row effects present (does it make a difference, for the output, which equipment is purchased)? Likewise for the column effects. In statistical terminology, the questions posed above translate as follows: Estimate the *parameters* of the model μ , α_i , $i = 1, \dots, I$, β_j , $j = 1, \dots, J$, and σ^2 . The estimates sought will be the MLE's, which for the parameters μ , α_i , and β_j are also LSE's. Test the null hypothesis of no row effects $H_{0,A}: \alpha_1 = \dots = \alpha_I$ (and therefore = 0). Test the null hypothesis of no column effects $H_{0,B}: \beta_1 = \dots = \beta_J$ (and therefore = 0).

14.3.1 The MLE's of the Parameters of the Model

The likelihood function of the Y_{ij} 's, to be denoted by $L(\mathbf{y}; \mu, \alpha, \beta, \sigma^2)$ in obvious notation, is given by the formula below. In this formula and in the sequel, the precise range of i and j will not be indicated explicitly for notational convenience.

$$L(\mathbf{y}; \mu, \alpha, \beta, \sigma^2) = \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^{IJ} \exp \left[-\frac{1}{2\sigma^2} \sum_i \sum_j (y_{ij} - \mu - \alpha_i - \beta_j)^2 \right]. \quad (24)$$

For each fixed σ^2 , maximization of the likelihood function with respect to μ , α_i , and β_j is equivalent to minimization, with respect to these parameters, of the expression:

$$S(\mu, \alpha_1, \dots, \alpha_I, \beta_1, \dots, \beta_J) = S(\mu, \alpha, \beta) = \sum_i \sum_j (y_{ij} - \mu - \alpha_i - \beta_j)^2. \quad (25)$$

Minimization of $S(\mu, \alpha, \beta)$ with respect to μ , α , and β yields the values given in the following result.

LEMMA 6 The unique minimizing values of μ , α_i , and β_j for expression (25) (i.e., the LSE's of μ , α_i , and β_j) are given by:

$$\hat{\mu} = \bar{y}_{..}, \quad \hat{\alpha}_i = \bar{y}_{i.} - \bar{y}_{..}, \quad i = 1, \dots, I, \quad \hat{\beta}_j = \bar{y}_{.j} - \bar{y}_{..}, \quad j = 1, \dots, J, \quad (26)$$

where

$$y_i = \frac{1}{J} \sum_j y_{ij}, \quad y_j = \frac{1}{I} \sum_i y_{ij}, \quad y_{..} = \frac{1}{IJ} \sum_i \sum_j y_{ij}. \quad (27)$$

PROOF Deferred to Subsection 14.3.3.

For the values in (26), the log-likelihood function becomes, with obvious notation:

$$\log L(\mathbf{y}; \hat{\mu}, \hat{\alpha}, \hat{\beta}, \sigma^2) = -\frac{IJ}{2} \log(2\pi) - \frac{IJ}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \hat{S}, \quad (28)$$

where $\hat{S} = \sum_i \sum_j (y_{ij} - y_i - y_j + y_{..})^2$. Relation (28) is of exactly the same type as relation (5), maximization of which produced the value

$$\hat{\sigma}^2 = \frac{1}{IJ} \hat{S} = \frac{1}{IJ} \sum_i \sum_j (y_{ij} - y_i - y_j + y_{..})^2. \quad (29)$$

Combining then the results in (26) and (29), we have the following result. \blacktriangle

THEOREM 4

Under model (23), the MLE's of the parameters of the model are given by relations (26) and (29). Furthermore, the MLE's of μ , α_i , and β_j are also their LSE's.

14.3.2 Testing the Hypothesis of No Row or No Column Effects

First, consider the null hypothesis of no row effects; namely,

$$H_{0,A}: \alpha_1 = \cdots = \alpha_I = 0. \quad (30)$$

Under $H_{0,A}$, the likelihood function in (24), to be denoted for convenience by $L_A(\mathbf{y}; \mu, \beta, \sigma^2)$, becomes:

$$L_A(\mathbf{y}; \mu, \beta, \sigma^2) = \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^{IJ} \exp \left[-\frac{1}{2\sigma^2} \sum_i \sum_j (y_{ij} - \mu - \beta_j)^2 \right]. \quad (31)$$

Maximization of this likelihood with respect to β_j 's and μ , for each fixed σ^2 , is equivalent to minimization, with respect to β_j 's and μ of the expression:

$$S(\mu, \beta_1, \dots, \beta_J) = S(\mu, \beta) = \sum_i \sum_j (y_{ij} - \mu - \beta_j)^2. \quad (32)$$

Working exactly as in (25), we obtain the following MLE's, under $H_{0,A}$, to be denoted by $\hat{\mu}_A$ and $\hat{\beta}_{j,A}$:

$$\hat{\mu}_A = y_{..} = \hat{\mu}, \quad \hat{\beta}_{j,A} = y_{.j} - y_{..} = \hat{\beta}_j, \quad j = 1, \dots, J. \quad (33)$$

Then, repeating the steps in relation (28), we obtain the MLE of σ^2 , under $H_{0,A}$:

$$\hat{\sigma}_A^2 = \frac{1}{IJ} \sum_i \sum_j (y_{ij} - y_{.j})^2. \quad (34)$$

The hypothesis $H_{0,A}$ will be tested by means of the likelihood ratio test. First, observe that:

$$\exp\left[-\frac{1}{2\widehat{\sigma}_A^2} \sum_i \sum_j (y_{ij} - y_j)^2\right] = \exp\left(-\frac{IJ}{2\widehat{\sigma}_A^2} \times \widehat{\sigma}_A^2\right) = \exp\left(-\frac{IJ}{2}\right),$$

and

$$\exp\left[-\frac{1}{2\widehat{\sigma}^2} \sum_i \sum_j (y_{ij} - y_i - y_j + y_{..})^2\right] = \exp\left(-\frac{IJ}{2\widehat{\sigma}^2} \times \widehat{\sigma}^2\right) = \exp\left(-\frac{IJ}{2}\right).$$

Then the likelihood ratio statistic λ is given by:

$$\lambda = (\widehat{\sigma}^2 / \widehat{\sigma}_A^2)^{IJ/2}.$$

$$\text{Hence } \left(\frac{\widehat{\sigma}^2}{\widehat{\sigma}_A^2}\right)^{IJ/2} < C, \quad \text{if and only if } \frac{\widehat{\sigma}_A^2}{\widehat{\sigma}^2} > C_0 = 1/C^{2/IJ}. \quad (35)$$

At this point, use the following notation:

$$SS_e = IJ\widehat{\sigma}^2 = \sum_i \sum_j (y_{ij} - y_i - y_j + y_{..})^2, \quad SS_A = J \sum_i \hat{\alpha}_i^2 = J \sum_i (y_i - y_{..})^2, \quad (36)$$

by means of which it is shown that:

LEMMA 7 With $\widehat{\sigma}_A^2$, SS_e and SS_A defined by (34) and (36), it holds: $IJ\widehat{\sigma}_A^2 = IJ\widehat{\sigma}^2 + SS_A = SS_e + SS_A$.

PROOF Deferred to Subsection 14.3.3.

By means of this lemma, relation (35) becomes:

$$\frac{\widehat{\sigma}_A^2}{\widehat{\sigma}^2} = \frac{IJ\widehat{\sigma}_A^2}{IJ\widehat{\sigma}^2} = \frac{SS_e + SS_A}{SS_e} = 1 + \frac{SS_A}{SS_e} > C_0, \quad \text{or } \frac{SS_A}{SS_e} > C_1 = C_0 - 1.$$

So, the likelihood ratio test rejects $H_{0,A}$ whenever

$$\frac{SS_A}{SS_e} > C_1, \quad \text{where } SS_A \text{ and } SS_e \text{ are given in (36)}. \quad (37)$$

For the determination of the cutoff point C_1 in (37), we need the distribution of the statistic SS_A/SS_e under $H_{0,A}$, where it is tacitly assumed that the observed values have been replaced by the respective r.v.'s. For this purpose, we establish the following result.

LEMMA 8 Consider the expressions SS_e and SS_A defined in (36), and replace the observed values y_{ij} , y_i , y_j , and $y_{..}$ by the respective r.v.'s Y_{ij} , Y_i , Y_j , and $Y_{..}$, but retain the same notation. Then, under model (23):

- (i) The r.v. SS_e/σ^2 is distributed as $\chi_{(I-1)(J-1)}^2$.
- (ii) The statistics SS_e and SS_A are independent.

Furthermore, if the null hypothesis $H_{0,A}$ defined in (30) is true, then:

- (iii) The r.v. SS_A/σ^2 is distributed as χ_{I-1}^2 .
- (iv) The statistic $\frac{SS_A/(I-1)}{SS_e/(I-1)(J-1)} \sim F_{I-1, (I-1)(J-1)}$.

PROOF Deferred to Subsection 14.3.3.

To this lemma, there is the following corollary, which also encompasses the estimates $\hat{\mu}$, $\hat{\alpha}_i$, and $\hat{\beta}_j$.

COROLLARY

- (i) The MLE's $\hat{\mu} = Y_{..}$, $\hat{\alpha}_i = Y_{i.} - Y_{..}$, $i = 1, \dots, I$, and $\hat{\beta}_j = Y_{.j} - Y_{..}$, $j = 1, \dots, J$ are unbiased estimates of the respective parameters μ , α_i , and β_j .
- (ii) The MLE $\hat{\sigma}^2 = SS_e/IJ$ of σ^2 given by (29) and (36) is biased, but the estimate $MS_e = SS_e/(I-1)(J-1)$ is unbiased.

PROOF

- (i) It is immediate from the definition of $Y_{i.}$, $Y_{.j}$, and $Y_{..}$ as (sample) means.
- (ii) From the lemma, $\frac{SS_e}{\sigma^2} \sim \chi_{(I-1)(J-1)}^2$, so that

$$E\left(\frac{SS_e}{\sigma^2}\right) = (I-1)(J-1), \quad \text{or} \quad E\left[\frac{SS_e}{(I-1)(J-1)}\right] = \sigma^2,$$

which proves the unbiasedness asserted. Also,

$$\begin{aligned} E\hat{\sigma}^2 &= E\left(\frac{SS_e}{IJ}\right) = \frac{(I-1)(J-1)}{IJ} E\left[\frac{SS_e}{(I-1)(J-1)}\right] \\ &= \frac{(I-1)(J-1)}{IJ} \sigma^2, \end{aligned}$$

which shows that $\hat{\sigma}^2$ is biased. \blacktriangle

By means then of this lemma and relation (37), we reach the following conclusion: The hypothesis $H_{0,A}$ is rejected at level α whenever

$$\mathcal{F}_A = \frac{SS_A/(I-1)}{SS_e/(I-1)(J-1)} = \frac{MS_A}{MS_e} > F_{I-1, (I-1)(J-1); \alpha}. \quad (38)$$

Next, consider the hypothesis of no column effects; i.e.,

$$H_{0,B}: \beta_1 = \dots = \beta_J = 0. \quad (39)$$

Then, working exactly as in (31) and (32), we obtain:

$$\hat{\mu}_B = y_{..} = \hat{\mu}, \quad \hat{\alpha}_{i,B} = y_{i.} - y_{..} = \hat{\alpha}_i, \quad i = 1, \dots, I, \quad (40)$$

and

$$\hat{\sigma}_B^2 = \frac{1}{IJ} \sum_i \sum_j (y_{ij} - y_{i.})^2. \quad (41)$$

Thus, as in (35), the hypothesis $H_{0,B}$ is rejected whenever

$$\frac{\widehat{\sigma}_B^2}{\widehat{\sigma}^2} > C'_0. \quad (42)$$

Set

$$SS_B = I \sum_j \widehat{\beta}_j^2 = I \sum_j (y_{.j} - y_{..})^2, \quad (43)$$

and consider the following result.

LEMMA 9 With SS_e , $\widehat{\sigma}_B^2$, and SS_B defined by (36), (41), and (43), it holds: $IJ\widehat{\sigma}_B^2 = IJ\widehat{\sigma}^2 + SS_B = SS_e + SS_B$.

PROOF Deferred to Subsection 14.3.3.

By means of this lemma, relation (42) becomes, as in (37): Reject $H_{0,B}$ whenever

$$\frac{SS_B}{SS_e} > C'_1, \quad \text{where } SS_e \text{ and } SS_B \text{ are given in (36) and (43).} \quad (44)$$

Finally, for the determination of the cutoff point C'_1 in (44), a certain distribution is needed. In other words, a lemma analogous to Lemma 7 is needed here.

LEMMA 10 Consider the expressions SS_e and SS_B defined in (36) and (43), and replace the observed values y_{ij} , $y_{i.}$, $y_{.j}$, and $y_{..}$ by the respective r.v.'s Y_{ij} , $Y_{i.}$, $Y_{.j}$, and $Y_{..}$, but retain the same notation. Then, under model (23):

- (i) The r.v. SS_e/σ^2 is distributed as $\chi_{(I-1)(J-1)}^2$.
- (ii) The statistics SS_e and SS_B are independent.

Furthermore, if the null hypothesis $H_{0,B}$ defined in (39) is true, then:

- (iii) The r.v. SS_B/σ^2 is distributed as χ_{J-1}^2 .
- (iv) The statistic $\frac{SS_B/(J-1)}{SS_e/(I-1)(J-1)} \sim F_{J-1, (I-1)(J-1)}$.

PROOF Deferred to Subsection 14.3.3.

By means of this lemma and relation (44), we conclude that: The hypothesis $H_{0,B}$ is rejected at level α whenever

$$\mathcal{F}_B = \frac{SS_B/(J-1)}{SS_e/(I-1)(J-1)} = \frac{MS_B}{MS_e} > F_{J-1, (I-1)(J-1); \alpha}. \quad (45)$$

For computational purposes, we need the following result.

LEMMA 11 Let SS_e , SS_A , and SS_B be given by (36) and (43), and let SS_T be defined by:

$$SS_T = \sum_i \sum_j (y_{ij} - y_{..})^2. \quad (46)$$

Then:

$$(i) \quad SS_A = J \sum_i y_i^2 - IJy_{..}^2, \quad SS_B = I \sum_j y_j^2 - IJy_{..}^2, \quad SS_T = \sum_i \sum_j y_{ij}^2 - IJy_{..}^2. \quad (47)$$

$$(ii) \quad SS_T = SS_e + SS_A + SS_B. \quad (48)$$

PROOF Deferred to Subsection 14.3.3.

Gathering together the hypotheses testing results obtained, we have the following theorem.

THEOREM 5

Under model (23), the hypotheses $H_{0,A}$ and $H_{0,B}$ are rejected at level of significance α whenever inequalities (38) and (45), respectively, hold true. The statistics SS_A , SS_B are computed by means of (47), and the statistic SS_e is computed by means of (47) and (48).

As in Section 14.1, the various quantities employed in testing the hypotheses $H_{0,A}$ and $H_{0,B}$, and also for estimating the error variance σ^2 , are gathered together in a table, an ANOVA table, as in Table 14.3.

Table 14.3 Analysis of Variance for Two-Way Layout with One Observation per Cell

Source of Variance	Sums of Squares	Degrees of Freedom	Mean Squares
Rows	$SS_A = J \sum_{i=1}^I \hat{\alpha}_i^2 = J \sum_{i=1}^I (Y_i - Y_{..})^2$	$I - 1$	$MS_A = \frac{SS_A}{I-1}$
Columns	$SS_B = I \sum_{j=1}^J \hat{\beta}_j^2 = I \sum_{j=1}^J (Y_j - Y_{..})^2$	$J - 1$	$MS_B = \frac{SS_B}{J-1}$
Residual	$SS_e = \sum_{i=1}^I \sum_{j=1}^J (Y_{ij} - Y_i - Y_j + Y_{..})^2$	$(I - 1) \times (J - 1)$	$MS_e = \frac{SS_e}{(I-1)(J-1)}$
Total	$SS_T = \sum_{i=1}^I \sum_{j=1}^J (Y_{ij} - Y_{..})^2$	$IJ - 1$	—

REMARK 5 In the present context, relation (48) is responsible for the term ANOVA. It states that the total variation (variance) $\sum_i \sum_j (Y_{ij} - Y_{..})^2$ (with reference to the grand sample mean $Y_{..}$) is split in three ways: one component $\sum_i \sum_j (Y_i - Y_{..})^2$ associated with the row effects (due to the row effects, or explained by the row effects); one component $\sum_i \sum_j (Y_j - Y_{..})^2$ associated with the column effects (due to the column effects, or explained by the column effects); and the residual component $\sum_i \sum_j (Y_{ij} - Y_i - Y_j + Y_{..})^2 = \sum_i \sum_j [(Y_{ij} - Y_{..}) - (Y_i - Y_{..}) - (Y_j - Y_{..})]^2$ (unexplained by the row and column effects, the sum of squares of errors).

Before embarking on the proof of the lemmas stated earlier in this section, let us illustrate the theory developed by a couple of examples. In the first example, we are presented with a set of numbers, not associated with any specific experiment; in the second example, a real-life experiment is considered.

EXAMPLE 6

Apply the two-way layout ANOVA with one observation per cell for the data given in Table 14.4; take $\alpha = 0.05$.

Table 14.4

**Data for a Two-Way
Layout ANOVA**

	1	2	3	4	$y_{i.}$
1	3	7	5	4	19/4
2	-1	2	0	2	3/4
3	1	2	4	0	7/4
$y_{.j}$	1	11/3	3	2	$y_{..} = \frac{29}{12}$

Here: $\hat{\mu} = y_{..} = \frac{29}{12} \simeq 2.417$, and:

$$\hat{\alpha}_1 = y_{1.} - y_{..} = \frac{19}{4} - \frac{29}{12} = \frac{7}{3} \simeq 2.333 \quad \hat{\beta}_1 = y_{1.} - y_{..} = 1 - \frac{29}{12} = -\frac{17}{12} \simeq -1.417$$

$$\hat{\alpha}_2 = y_{2.} - y_{..} = \frac{3}{4} - \frac{29}{12} = -\frac{5}{3} \simeq -1.667; \quad \hat{\beta}_2 = y_{2.} - y_{..} = \frac{11}{3} - \frac{29}{12} = \frac{5}{4} = 1.25$$

$$\hat{\alpha}_3 = y_{3.} - y_{..} = \frac{7}{4} - \frac{29}{12} = -\frac{2}{3} \simeq -0.667 \quad \hat{\beta}_3 = y_{3.} - y_{..} = 3 - \frac{29}{12} = \frac{7}{12} \simeq 0.583$$

$$\hat{\beta}_4 = y_{4.} - y_{..} = 2 - \frac{29}{12} = -\frac{5}{12} \simeq -0.417.$$

$$SS_A = 4 \times \left[\left(\frac{19}{4} \right)^2 + \left(\frac{3}{4} \right)^2 + \left(\frac{7}{4} \right)^2 \right] - 12 \times \left(\frac{29}{12} \right)^2 = \frac{104}{3} \simeq 34.667,$$

$$SS_B = 3 \times \left[1^2 + \left(\frac{11}{3} \right)^2 + 3^2 + 2^2 \right] - 12 \times \left(\frac{29}{12} \right)^2 = \frac{147}{12} = 12.25,$$

$$\begin{aligned} SS_T &= [3^2 + 7^2 + 5^2 + 4^2 + (-1)^2 + 2^2 + 0^2 + 2^2 + 1^2 + 2^2 + 4^2 + 0^2] - 12 \times \left(\frac{29}{12} \right)^2 \\ &= \frac{707}{12} \simeq 58.917, \end{aligned}$$

so that

$$SS_e = SS_T - SS_A - SS_B = \frac{707}{12} - \frac{104}{3} - \frac{147}{12} = 12.$$

Hence, the unbiased estimate of σ^2 is: $\frac{SS_e}{(I-1)(J-1)} = \frac{12}{6} = 2$. Furthermore,

$$\mathcal{F}_A = \frac{MS_A}{MS_e} = \frac{104/3 \times 2}{12/6} = \frac{26}{3} \simeq 8.667,$$

$$\mathcal{F}_B = \frac{MS_B}{MS_e} = \frac{147/12 \times 3}{12/6} = \frac{147}{72} \simeq 2.042.$$

Since $F_{I-1, (I-1), (J-1); \alpha} = F_{2, 6; 0.05} = 5.1433$, we see that the hypothesis $H_{0,A}$ is rejected, whereas the hypothesis $H_{0,B}$ is not rejected.

EXAMPLE 7

The cutting speeds of four types of tools are being compared by using five materials of varying degrees of hardness. The data pertaining to measurements of cutting time in seconds are given in Table 14.5. Carry out the ANOVA for these data; take $\alpha = 0.05$.

Table 14.5

Data for a Two-Way
Layout ANOVA

	1	2	3	4	5	$y_{i.}$
1	12	2	8	1	7	6
2	20	14	17	12	17	16
3	13	7	13	8	14	11
4	11	5	10	3	6	7
$y_{.j}$	14	7	12	6	11	$y_{..} = 10$

Here $I = 4$, $J = 5$. From the table: $\hat{\mu} = 10$, and:

$$\begin{aligned} \hat{\alpha}_1 &= y_{1.} - y_{..} = 6 - 10 = -4 & \hat{\beta}_1 &= y_{.1} - y_{..} = 14 - 10 = 4 \\ \hat{\alpha}_2 &= y_{2.} - y_{..} = 16 - 10 = 6 & \hat{\beta}_2 &= y_{.2} - y_{..} = 7 - 10 = -3 \\ \hat{\alpha}_3 &= y_{3.} - y_{..} = 11 - 10 = 1 & \hat{\beta}_3 &= y_{.3} - y_{..} = 12 - 10 = 2 \\ \hat{\alpha}_4 &= y_{4.} - y_{..} = 7 - 10 = -3 & \hat{\beta}_4 &= y_{.4} - y_{..} = 6 - 10 = -4 \\ & & \hat{\beta}_5 &= y_{.5} - y_{..} = 11 - 10 = 1. \end{aligned}$$

$$SS_A = 5 \times (6^2 + 16^2 + 11^2 + 7^2) - 20 \times 10^2 = 310,$$

$$SS_B = 4 \times (14^2 + 7^2 + 12^2 + 6^2 + 11^2) - 20 \times 10^2 = 184,$$

$$SS_T = 2,158 - 2,000 = 518,$$

$$SS_E = 518 - 310 - 184 = 24.$$

Hence, the unbiased estimate for σ^2 is: $\frac{SS_E}{(I-1)(J-1)} = \frac{24}{3 \times 4} = 2$. Furthermore,

$$\mathcal{F}_A = \frac{MS_A}{MS_E} = \frac{310/3}{24/12} = \frac{155}{3} \simeq 51.667,$$

$$\mathcal{F}_B = \frac{MS_B}{MS_E} = \frac{184/4}{24/12} = 23.$$

Since $F_{I-1, (I-1)(J-1); \alpha} = F_{3, 12; 0.05} = 3.4903$, it follows that both hypotheses $H_{0,A}$ and $H_{0,B}$ are to be rejected. So, the mean cutting times, either for the tools across the material cut, or for the material cut across the tools used, cannot be assumed to be equal (at the $\alpha = 0.05$ level). Actually, this should not come as a surprise when looking at the margin of the table, which provide estimates of these times.

14.3.3 Proof of Lemmas in Section 14.3

In this subsection, a justification (or an outline thereof) is provided for the lemmas used in this section.

PROOF OF LEMMA 6 Consider the expression $\mathcal{S}(\mu, \alpha, \beta) = \sum_i \sum_j (y_{ij} - \mu - \alpha_i - \beta_j)^2$ and recall that $\sum_i \alpha_i = 0$, $\sum_j \beta_j = 0$. Following the method of Lagrange multipliers, consider the linear combination

$$\mathcal{S}^*(\mu, \alpha, \beta) = \sum_i \sum_j (y_{ij} - \mu - \alpha_i - \beta_j)^2 + \lambda_1 \sum_i \alpha_i + \lambda_2 \sum_j \beta_j,$$

where λ_1, λ_2 are constants, determine the partial derivatives of $\mathcal{S}^*(\mu, \alpha, \beta)$ with respect to μ, α_i , and β_j , equate them to 0, append to them the side constraints $\sum_i \alpha_i = 0$, $\sum_j \beta_j = 0$, and solve the resulting system with respect to μ , the α_i 's, and the β_j 's (and also λ_1, λ_2). By implementing these steps, we get:

$$\left. \begin{aligned} \frac{\partial}{\partial \mu} \mathcal{S}^*(\mu, \alpha, \beta) &= -2 \sum_i \sum_j y_{ij} + 2IJ\mu + J \sum_i \alpha_i + 2I \sum_j \beta_j = 0 \\ \frac{\partial}{\partial \alpha_i} \mathcal{S}^*(\mu, \alpha, \beta) &= -2 \sum_j y_{ij} + 2J\mu + 2J\alpha_i + 2 \sum_j \beta_j + \lambda_1 = 0 \\ \frac{\partial}{\partial \beta_j} \mathcal{S}^*(\mu, \alpha, \beta) &= -2 \sum_i y_{ij} + 2I\mu + 2 \sum_i \alpha_i + 2I\beta_j + \lambda_2 = 0 \\ &\sum_i \alpha_i = 0 \\ &\sum_j \beta_j = 0 \end{aligned} \right\},$$

from which we obtain:

$$\mu = \frac{1}{IJ} \sum_i \sum_j y_{ij} = y_{..}, \quad \alpha_i = y_{i.} - y_{..} - \frac{\lambda_1}{2J}, \quad \beta_j = y_{.j} - y_{..} - \frac{\lambda_2}{2I}.$$

But

$$0 = \sum_i \alpha_i = \sum_i y_{i.} - Iy_{..} - \frac{I\lambda_1}{2J} = Iy_{..} - Iy_{..} - \frac{I\lambda_1}{2J} = -\frac{I\lambda_1}{2J}, \quad \text{so that } \lambda_1 = 0,$$

and likewise for λ_2 by summing up the β_j 's. Thus,

$$\mu = y_{..}, \quad \alpha_i = y_{i.} - y_{..}, \quad i = 1, \dots, I, \quad \beta_j = y_{.j} - y_{..}, \quad j = 1, \dots, J. \quad (49)$$

Now the parameter μ is any real number, the α_i 's span an $(I - 1)$ -dimensional hyperplane, and the β_j 's span a $(J - 1)$ -dimensional hyperplane. It is then clear that the expression $\mathcal{S}(\mu, \alpha, \beta)$ (as a function of μ , the α_i 's, and the β_j 's) does not have a maximum. Then the values in (49) are candidates to produce a minimum of $\mathcal{S}(\mu, \alpha, \beta)$, in which case (26) follows. Again, geometrical considerations suggest that they do produce a minimum, and we will leave it at that presently. ▲

REMARK 6 It should be mentioned at this point that ANOVA models are special cases of the so-called *General Linear Models*, and then the above minimization problem is resolved in a general setting by means of linear algebra methodology. For a glimpse at it, one may consult Chapter 17 in the book *A Course in Mathematical Statistics*, 2nd edition, Academic Press (1997), by G. G. Roussas.

PROOF OF LEMMA 7 Here, we have to establish the relation:

$$\sum_i \sum_j (y_{ij} - y_{.j})^2 = J \sum_i (y_i - y_{..})^2 + \sum_i \sum_j (y_{ij} - y_i - y_{.j} + y_{..})^2.$$

Indeed,

$$\begin{aligned} SS_e &= \sum_i \sum_j (y_{ij} - y_i - y_{.j} + y_{..})^2 = \sum_i \sum_j [(y_{ij} - y_{.j}) - (y_i - y_{..})]^2 \\ &= \sum_i \sum_j (y_{ij} - y_{.j})^2 + J \sum_i (y_i - y_{..})^2 - 2 \sum_i \sum_j (y_i - y_{..})(y_{ij} - y_{.j}) \\ &= I \widehat{J\sigma_A^2} + SS_A - 2SS_A = I \widehat{J\sigma_A^2} - SS_A, \text{ because} \\ &\quad \sum_i \sum_j (y_i - y_{..})(y_{ij} - y_{.j}) = \sum_i (y_i - y_{..}) \sum_j (y_{ij} - y_{.j}) \\ &= \sum_i (y_i - y_{..})(Jy_i - Jy_{..}) = J \sum_i (y_i - y_{..})^2 = SS_A. \quad \blacktriangle \end{aligned}$$

PROOF OF LEMMA 8 There are several ways one may attempt to justify the results in this lemma. One would be to refer to Lemma 2 and suggest that a similar approach be used, but that would do no justice. Another approach would be to utilize the theory of quadratic forms, but that would require an extensive introduction to the subject and the statement and/or proof of a substantial number of related results. Finally, the last approach would be to use a geometric descriptive approach based on fundamental concepts of (finite dimensional) vector spaces. We have chosen to follow this last approach.

All vectors to be used here are column vectors, and the prime notation, “ \prime ”, indicates transpose of a vector. Set $\mathbf{Y} = (Y_{11}, \dots, Y_{1J}; Y_{21}, \dots, Y_{2J}; \dots; Y_{I1}, \dots, Y_{IJ})'$, so that \mathbf{Y} belongs in an $I \times J$ -dimensional vector space to be denoted by $V_{I \times J}$. Also, set

$$\begin{aligned} \boldsymbol{\eta} &= \mathbf{EY} = (\mathbf{EY}_{11}, \dots, \mathbf{EY}_{1J}; \mathbf{EY}_{21}, \dots, \mathbf{EY}_{2J}; \dots; \mathbf{EY}_{I1}, \dots, \mathbf{EY}_{IJ})' \\ &= (\mu + \alpha_1 + \beta_1, \dots, \mu + \alpha_1 + \beta_J; \mu + \alpha_2 + \beta_1, \dots, \mu + \alpha_2 + \beta_J; \dots; \\ &\quad \mu + \alpha_I + \beta_1, \dots, \mu + \alpha_I + \beta_J)'. \end{aligned}$$

Although the vector $\boldsymbol{\eta}$ has $I \times J$ coordinates, due to its form and the fact that $\sum_i \alpha_i = \sum_j \beta_j = 0$, it follows that it lies in an $(I + J - 1)$ -dimensional space, V_{I+J-1} . Finally, if $H_{0,A}: \alpha_1 = \dots = \alpha_I = 0$ holds, then the respective mean vector, to be denoted by $\boldsymbol{\eta}_A$, is

$$\boldsymbol{\eta}_A = (\mu + \beta_1, \dots, \mu + \beta_J; \mu + \beta_1, \dots, \mu + \beta_J; \dots; \mu + \beta_1, \dots, \mu + \beta_J)',$$

and reasoning as above, we conclude that $\boldsymbol{\eta}_A \in V_J$. Thus, we have three vector spaces related as follows: $V_J \subset V_{I+J-1} \subset V_{I \times J}$.

It is clear that, if μ , α_i 's, and β_j 's are replaced by their (least squares) estimates $\hat{\mu}$, $\hat{\alpha}_i$'s, and $\hat{\beta}_j$'s, the resulting random vector $\hat{\boldsymbol{\eta}}$ still lies in V_{I+J-1} , and likewise for the random vector $\hat{\boldsymbol{\eta}}_A$, which we get, if μ and β_j 's are replaced by $\hat{\mu}_A = \hat{\mu}$ and $\hat{\beta}_{j,A} = \hat{\beta}_j$; i.e., $\hat{\boldsymbol{\eta}}_A \in V_J$. We now proceed as follows: Let $\boldsymbol{\alpha}_I, \dots, \boldsymbol{\alpha}_{I+J-1}$

be an orthonormal basis in V_J (i.e., $\alpha'_i \alpha_j = 0$ for $i \neq j$ and $\|\alpha_i\| = 1$), which we extend to an orthonormal basis $\alpha_1, \dots, \alpha_{I-1}, \alpha_I, \dots, \alpha_{I+J-1}$ in V_{I+J+1} , and then to an orthonormal basis

$$\alpha_1, \dots, \alpha_{I-1}, \alpha_I, \dots, \alpha_{I+J-1}, \alpha_{I+J}, \dots, \alpha_{I \times J}$$

in $V_{I \times J}$. This can be done, as has already been mentioned in a similar context in the proof of Lemma 5 in Chapter 13. Also, see Remark 4 in the same chapter. Since $\mathbf{Y} \in V_{I \times J}$, it follows that \mathbf{Y} is a linear combination of the α_i 's with coefficient some r.v.'s Z_i 's. That is, $\mathbf{Y} = \sum_{i=1}^{I \times J} Z_i \alpha_i$. Since $\hat{\boldsymbol{\eta}}$ minimizes the quantity $\|\mathbf{Y} - \boldsymbol{\eta}\|^2 = \sum_i \sum_j (Y_{ij} - \mu - \alpha_i - \beta_j)^2$, it follows that $\hat{\boldsymbol{\eta}}$ is, actually, the projection of \mathbf{Y} into the space V_{I+J-1} . It follows then that $\hat{\boldsymbol{\eta}} = \sum_{i=1}^{I+J-1} Z_i \alpha_i$. Under $H_{0,A}$, the vector $\hat{\boldsymbol{\eta}}_A$ minimizes $\|\mathbf{Y} - \boldsymbol{\eta}_A\|^2 = \sum_i \sum_j (Y_{ij} - \mu - \beta_j)^2$, and therefore is the projection of \mathbf{Y} into the space V_J . Thus, $\hat{\boldsymbol{\eta}}_A = \sum_{i=I}^{I+J-1} Z_i \alpha_i$. Then $\mathbf{Y} - \hat{\boldsymbol{\eta}} = \sum_{i=I+J}^{I \times J} Z_i \alpha_i$, $\mathbf{Y} - \hat{\boldsymbol{\eta}}_A = \sum_{i=1}^{I-1} Z_i \alpha_i + \sum_{i=I+J}^{I \times J} Z_i \alpha_i$, and $\hat{\boldsymbol{\eta}} - \hat{\boldsymbol{\eta}}_A = \sum_{i=1}^{I-1} Z_i \alpha_i$. Because of the orthonormality of the α_i 's, it follows that:

$$\begin{aligned} \|\mathbf{Y} - \hat{\boldsymbol{\eta}}\|^2 &= \left\| \sum_{i=I+J}^{I \times J} Z_i \alpha_i \right\|^2 = \sum_{i=I+J}^{I \times J} Z_i^2, \\ \|\mathbf{Y} - \hat{\boldsymbol{\eta}}_A\|^2 &= \left\| \sum_{i=1}^{I-1} Z_i \alpha_i + \sum_{i=I+J}^{I \times J} Z_i \alpha_i \right\|^2 = \sum_{i=1}^{I-1} Z_i^2 + \sum_{i=I+J}^{I \times J} Z_i^2, \end{aligned}$$

and

$$\|\hat{\boldsymbol{\eta}} - \hat{\boldsymbol{\eta}}_A\|^2 = \left\| \sum_{i=1}^{I-1} Z_i \alpha_i \right\|^2 = \sum_{i=1}^{I-1} Z_i^2.$$

However,

$$\|\mathbf{Y} - \hat{\boldsymbol{\eta}}\|^2 = \sum_i \sum_j (Y_{ij} - \hat{\mu} - \hat{\alpha}_i - \hat{\beta}_j)^2 = \sum_i \sum_j (Y_{ij} - Y_{i.} - Y_{.j} + Y_{..})^2 = SS_e,$$

and

$$\|\hat{\boldsymbol{\eta}} - \hat{\boldsymbol{\eta}}_A\|^2 = \sum_i \sum_j \hat{\alpha}_i^2 = \sum_i \sum_j (Y_{i.} - Y_{..})^2 = J \sum_i (Y_{i.} - Y_{..})^2 = SS_A.$$

Therefore

$$SS_A = \sum_{i=1}^{I-1} Z_i^2, \quad SS_e = \sum_{i=I+J}^{I \times J} Z_i^2. \quad (50)$$

Now, observe that the r.v.'s $Z_1, \dots, Z_{I \times J}$ are the transformation of the r.v.'s $Y_1, \dots, Y_{I \times J}$ under the orthogonal matrix \mathbf{P} whose rows are the vectors α_i 's. This follows immediately from the relation $\mathbf{Y} = \sum_{i=1}^{I \times J} Z_i \alpha_i$, if we multiply (in the inner product sense) by α'_j . We then get $\alpha'_j \mathbf{Y} = \sum_{i=1}^{I \times J} Z_i (\alpha'_j \alpha_i)$, and this is Z_i , if $j = i$, and 0 otherwise. So, $Z_i = \alpha'_i \mathbf{Y}$, $i = 1, \dots, I \times J$. Since the Y_i 's are independent and Normally distributed with (common) variance σ^2 , it follows that the Z_i 's are also independently Normally distributed with specified means and the same variance σ^2 . (See Theorem 8 in Chapter 6.) From the fact that

$\hat{\eta} \in V_{I+J-1}$, it follows that its last $I \times J - (I + J - 1) = (I - 1)(J - 1)$ coordinates are zero. Then so are the respective EZ_i . That is, $Z_i, i = 1, \dots, I \times J$ are independent Normal, $EZ_i = 0$ for the last $(I - 1)(J - 1)$ coordinates, and they all have variance σ^2 . It follows that:

- (i) $\frac{SS_e}{\sigma^2} = \frac{1}{\sigma^2} \sum_{i=I+J}^{I \times J} Z_i^2 \sim \chi_{(I-1)(J-1)}^2$.
- (ii) The statistics SS_e and SS_A are independent, because they are defined in terms of nonoverlapping sets of the independent r.v.'s Z_i 's (see relation (50)).
- (iii) The expectations of the coordinates of $\hat{\eta}$ are $\mu + \alpha_i + \beta_j$, and the expectations of the coordinates of $\hat{\eta}_A$ are $\mu + \beta_j$. It follows that the expectations of the coordinates of $\hat{\eta} - \hat{\eta}_A$ are $(\mu + \alpha_i + \beta_j) - (\mu + \beta_j) = \alpha_i$. Therefore, if $H_{0,A}$ is true, these expectations are 0, and then so are the expectations of $Z_i, i = 1, \dots, I - 1$, since $\hat{\eta} - \hat{\eta}_A = \sum_{i=1}^{I-1} Z_i \alpha_i$. It follows that

$$\frac{SS_A}{\sigma^2} = \frac{1}{\sigma^2} \sum_{i=1}^{I-1} Z_i^2 \sim \chi_{I-1}^2.$$

- (iv) Immediate from parts (i)–(iii) and the definition of the F distribution. ▲

PROOF OF LEMMA 9 We have to show that

$$\sum_i \sum_j (y_{ij} - y_i)^2 = I \sum_j (y_j - y_{..})^2 + \sum_i \sum_j (y_{ij} - y_i - y_j + y_{..})^2.$$

As in the proof of Lemma 7,

$$\begin{aligned} SS_e &= \sum_i \sum_j (y_{ij} - y_i - y_j + y_{..})^2 = \sum_i \sum_j [(y_{ij} - y_i) - (y_j - y_{..})]^2 \\ &= \sum_i \sum_j (y_{ij} - y_i)^2 + \sum_i \sum_j (y_j - y_{..})^2 - 2 \sum_i \sum_j (y_j - y_{..})(y_{ij} - y_i) \\ &= I J \widehat{\sigma_B^2} + SS_B - 2SS_B = I J \widehat{\sigma_B^2} - SS_B, \text{ because} \\ &\quad \sum_i \sum_j (y_j - y_{..})(y_{ij} - y_i) = \sum_j (y_j - y_{..}) \sum_i (y_{ij} - y_i) \\ &= \sum_j (y_j - y_{..})(I y_j - I y_{..}) = I \sum_j (y_j - y_{..})^2 = SS_B. \quad \blacktriangle \end{aligned}$$

PROOF OF LEMMA 10

- (i) It is the same as (i) in Lemma 8.
- (ii) It is done as in Lemma 8(ii), where $H_{0,A}$ is replaced by $H_{0,B}$.
- (iii) Again, it is a repetition of the arguments in Lemma 8(iii).
- (iv) Immediate from parts (i)–(iii). ▲

PROOF OF LEMMA 11

- (i) They are all a direct application of the identity: $\sum_{k=1}^n (X_k - \bar{X})^2 = \sum_k X_k^2 - n\bar{X}^2$.

(ii) Clearly,

$$\begin{aligned}
SS_e &= \sum_i \sum_j (y_{ij} - y_i - y_j + y_{..})^2 \\
&= \sum_i \sum_j [(y_{ij} - y_{..}) - (y_i - y_{..}) - (y_j - y_{..})]^2 \\
&= \sum_i \sum_j (y_{ij} - y_{..})^2 + J \sum_i (y_i - y_{..})^2 + I \sum_j (y_j - y_{..})^2 \\
&\quad - 2 \sum_i \sum_j (y_i - y_{..})(y_{ij} - y_{..}) - 2 \sum_i \sum_j (y_j - y_{..})(y_{ij} - y_{..}) \\
&\quad + 2 \sum_i \sum_j (y_i - y_{..})(y_j - y_{..}) = SS_T - SS_A - SS_B,
\end{aligned}$$

because

$$\begin{aligned}
\sum_i \sum_j (y_i - y_{..})(y_{ij} - y_{..}) &= \sum_i (y_i - y_{..})(Jy_i - Jy_{..}) \\
&= J \sum_i (y_i - y_{..})^2 = SS_A,
\end{aligned}$$

$$\begin{aligned}
\sum_i \sum_j (y_j - y_{..})(y_{ij} - y_{..}) &= \sum_j (y_j - y_{..})(Iy_j - Iy_{..}) \\
&= I \sum_j (y_j - y_{..})^2 = SS_B,
\end{aligned}$$

and

$$\sum_i \sum_j (y_i - y_{..})(y_j - y_{..}) = \sum_i (y_i - y_{..})(Jy_{..} - Jy_{..}) = 0. \quad \blacktriangle$$

REMARK 7 In a two-way layout of ANOVA, we may have $K(\geq 2)$ observations per cell rather than one. The concepts remain the same, but the analysis is somewhat more complicated. The reader may wish to refer to Section 17.3 in Chapter 17 of the book *A Course in Mathematical Statistics*, 2nd edition, Academic Press (1997), by G. G. Roussas. The more general cases, where there is an unequal number of observations per cell or there are more than two factors influencing the outcome, are the subject matter of the ANOVA branch of statistics and are, usually, not discussed in an introductory course.

Exercises

3.1 Apply the two-way layout (with one observation per cell) analysis of variance to the data given in the table below. Take $\alpha = 0.05$.

Levels of Factor A \ Levels of Factor B	Levels of Factor B				
	1	2	3	4	5
1	110	128	48	123	19
2	214	183	115	114	129
3	208	183	130	225	114

- 3.2** Under the null hypothesis $H_{0,A}$ stated in relation (30), show that the MLE's of μ and β_j , $j = 1, \dots, J$ are, indeed, given by the expressions in relation (33).
- 3.3** Under the null hypothesis $H_{0,A}$ stated in relation (30), show that the MLE of σ^2 is, indeed, given by the expression in relation (34).
- 3.4** In reference to the proof of Lemma 8, show that η is an $(I + J - 1)$ -dimensional vector.

Hint: This problem may be approached as follows.

$$\mathbf{X}' = \begin{pmatrix} \overbrace{1 \ 1 \ 0 \ 0 \ \dots \ 0}^I & \overbrace{1 \ 0 \ 0 \ \dots \ 0}^J \\ 1 \ 1 \ 0 \ 0 \ \dots \ 0 & 0 \ 1 \ 0 \ \dots \ 0 \\ \vdots & \vdots \\ 1 \ 1 \ 0 \ 0 \ \dots \ 0 & 0 \ 0 \ 0 \ \dots \ 0 \ 1 \\ 1 \ 0 \ 1 \ 0 \ \dots \ 0 & 1 \ 0 \ 0 \ \dots \ 0 \\ 1 \ 0 \ 1 \ 0 \ \dots \ 0 & 0 \ 1 \ 0 \ \dots \ 0 \\ \vdots & \vdots \\ 1 \ 0 \ 1 \ 0 \ \dots \ 0 & 0 \ 0 \ 0 \ \dots \ 0 \ 1 \\ \vdots & \vdots \\ \vdots & \vdots \\ \vdots & \vdots \\ 1 \ 0 \ 0 \ 0 \ \dots \ 0 & 1 \ 1 \ 0 \ 0 \ \dots \ 0 \\ 1 \ 0 \ 0 \ 0 \ \dots \ 0 & 1 \ 0 \ 1 \ 0 \ \dots \ 0 \\ \vdots & \vdots \\ 1 \ 0 \ 0 \ 0 \ \dots \ 0 & 1 \ 0 \ 0 \ 0 \ \dots \ 0 \ 1 \end{pmatrix} \begin{matrix} J \\ J \\ J \end{matrix}$$

Consider the $IJ \times (I + J + 1)$ matrix \mathbf{X}' given above and let the $1 \times (I + J + 1)$ vector β' be defined by: $\beta' = (\mu, \alpha_1, \dots, \alpha_I, \beta_1, \dots, \beta_J)$. Then do the following:

- (i) Observe that $\eta = \mathbf{X}'\beta$, so that η lies in the vector space generated by the columns (rows) of \mathbf{X}' .
- (ii) For $I \geq 2$ and $J \geq \frac{I+1}{I-1}$, observe that $\text{rank } \mathbf{X}' \leq I + J + 1 = \min\{I + J + 1, IJ\}$.
- (iii) Show that $\text{rank } \mathbf{X}' = I + J - 1$ by showing that: (a) The 1st column of \mathbf{X}' is the sum of the subsequent I columns of \mathbf{X}' . (b) The 2nd column is the (sum of the last J columns) – (sum of the 3rd, 4th, \dots , I th columns). (c) The $I + J - 1$ columns, except for the first two, are linearly independent (by showing that any linear combination of them by scalars is the zero vector if and only if all scalars are zero). It will then follow that the dimension of η is $I + J - 1$.

- 3.5** In reference to the proof of Lemma 8, and under the hypothesis $H_{0,A}$, show that the dimension of the vector η_A is J .

Hint: As in Exercise 3.4, one may use similar steps in showing that η_A belongs in a J -dimensional vector space and thus is of dimension J . To this end, consider the $IJ \times (J + 1)$ matrix \mathbf{X}'_A given below, and let $\beta'_A = (\mu, \beta_1, \dots, \beta_J)$. Then do the following:

$$\mathbf{X}' = \begin{pmatrix} \overbrace{1 & 1 & 0 & 0 & \cdot & \cdot & \cdot & 0}^J \\ \overbrace{1 & 0 & 1 & 0 & \cdot & \cdot & \cdot & 0}^J \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \overbrace{1 & 0 & 0 & 0 & \cdot & \cdot & 0 & 1}^J \\ \overbrace{1 & 1 & 0 & 0 & \cdot & \cdot & \cdot & 0}^J \\ \overbrace{1 & 0 & 1 & 0 & \cdot & \cdot & \cdot & 0}^J \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & 0 & 0 & 0 & \cdot & \cdot & 0 & 1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \overbrace{1 & 1 & 0 & 0 & \cdot & \cdot & \cdot & 0}^J \\ \overbrace{1 & 0 & 1 & 0 & \cdot & \cdot & \cdot & 0}^J \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \overbrace{1 & 0 & 0 & 0 & \cdot & \cdot & 0 & 1}^J \end{pmatrix}$$

- (i) Observe that $\eta_A = \mathbf{X}'_A \beta_A$, so that η_A lies in the vector space generated by the columns (rows) of \mathbf{X}'_A .
- (ii) For $I \geq 2$, it is always true that $J + 1 \leq IJ$, and therefore $\text{rank } \mathbf{X}'_A \leq J + 1 = \min\{J + 1, IJ\}$.
- (iii) Show that $\text{rank } \mathbf{X}'_A = J$ by showing that: (a) The 1st column of \mathbf{X}'_A is the sum of the subsequent J columns. (b) The J columns, except for the 1st one, are linearly independent. It will then follow that the dimension of η_A is J .



Chapter 15

Some Topics in Nonparametric Inference

In Chapters 9, 10, 11, and 12, we concerned ourselves with the question of point estimation, interval estimation, and testing hypotheses about (most of the time) a real-valued parameter θ . This inference was hedged on the basic premise that we were able to stipulate each time a probability model, which was completely known except for a parameter θ (real-valued or of higher dimension).

The natural question which arises is this: What do we do, if there is no sound basis for the stipulation of a probability model from which the observations are drawn? In such a situation, we don't have parametric inference problems to worry about, because, simply, we don't have a parametric model. In certain situations things may not be as bad as this, but they are nearly so. Namely, we are in a position to assume the existence of a parametric model which governs the observations. However, the number of parameters required to render the model meaningful is exceedingly large, and therefore inference about them is practically precluded.

It is in situations like this, where the so-called nonparametric models and nonparametric inference enter the picture. Accordingly, a nonparametric approach starts out with a bare minimum of assumptions, which certainly do not include the existence of a parametric model, and proceeds to derive inference for a multitude of important quantities. This chapter is devoted to discussing a handful of problems of this variety.

Specifically, in the first section confidence intervals are constructed for the mean μ of a distribution, and also the value at x of the d.f. F , $F(x)$. The confidence coefficients are approximately $1 - \alpha$ for large n . Illustrative examples are also provided. In the following section confidence intervals are constructed for the quantiles of a d.f. F . Here the concept of a confidence coefficient is

replaced by that of the coverage probability. In the subsequent two sections, two populations are compared by means of the sign test, when the sample sizes are equal, and the rank sum test and the Wilcoxon–Mann–Whitney test in the general case. Some examples are also discussed. The last section consists of two subsections. One is devoted to estimating (nonparametrically) a p.d.f. and the formulation of a number of desirable properties of the proposed estimate. The other subsection addresses very briefly two very important problems; namely, the problem of regression estimation under a fixed design, and the problem of prediction when the design is stochastic.

15.1 Some Confidence Intervals with Given Approximate Confidence Coefficient

We are in a position to construct a confidence interval for the (unknown) mean μ of n i.i.d. observations X_1, \dots, X_n with very little information as to where these observations are coming from. Specifically, all we have to know is that these r.v.'s have finite mean μ and variance $\sigma^2 \in (0, \infty)$, and nothing else. Then, by the CLT,

$$\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1), \quad \bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i. \quad (1)$$

Suppose first that σ is known. Then, for all sufficiently large n , the normal approximation in (1) yields:

$$P \left[-z_{\frac{\alpha}{2}} \leq \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \leq z_{\frac{\alpha}{2}} \right] \simeq 1 - \alpha,$$

or

$$P \left(\bar{X}_n - z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{X}_n + z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \right) \simeq 1 - \alpha.$$

In other words,

$$\left[\bar{X}_n - z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}, \quad \bar{X}_n + z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \right] \quad (2)$$

is a confidence interval for μ with confidence coefficient approximately $1 - \alpha$ ($0 < \alpha < 1$).

Now, if μ is unknown, it is quite likely that σ is also unknown. What we do then is to estimate σ^2 by

$$S_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}_n^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \frac{(\sum_i X_i)^2}{n^2}, \quad (3)$$

and recall that (by Theorem 7(i) in Chapter 7):

$$S_n^2 \xrightarrow[n \rightarrow \infty]{P} \sigma^2, \quad \text{or} \quad \frac{S_n^2}{\sigma^2} \xrightarrow[n \rightarrow \infty]{P} 1. \quad (4)$$

Then convergences (1) and (4), along with Theorem 6(iii) in Chapter 7, yield:

$$\frac{\sqrt{n}(\bar{X}_n - \mu)/\sigma}{S_n/\sigma} = \frac{\sqrt{n}(\bar{X}_n - \mu)}{S_n} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1).$$

Then, proceeding as before, we obtain that

$$\left[\bar{X}_n - z_{\frac{\alpha}{2}} \frac{S_n}{\sqrt{n}}, \quad \bar{X}_n + z_{\frac{\alpha}{2}} \frac{S_n}{\sqrt{n}} \right] \quad (5)$$

is a confidence interval for μ with confidence coefficient approximately $1 - \alpha$.

Here is an application of formula (5).

EXAMPLE 1

Refer to the GPA's in Example 22 of Chapter 1, where we assume that the given GPA scores are observed values of r.v.'s X_i , $i = 1, \dots, 34$ with (unknown) mean μ and (unknown) variance σ^2 , both finite. Construct a confidence interval for μ with confidence coefficient approximately 95%.

DISCUSSION In the discussion of Example 1 in Chapter 13, we saw that: $\sum_i x_i = 100.73$ and $\sum_i x_i^2 = 304.7885$, so that:

$$\bar{x} = \frac{100.73}{34} \simeq 2.963, \quad s_n^2 = \frac{304.7885}{34} - \frac{(100.73)^2}{34^2} \simeq 0.187, \quad \text{and} \quad s_n \simeq 0.432.$$

Since $z_{0.025} = 1.96$, formula (5) gives:

$$\left[2.9626 - 1.96 \times \frac{0.432}{5.831}, \quad 2.9626 + 1.96 \times \frac{0.432}{5.831} \right] \simeq [2.818, 3.108].$$

Another instance where a nonparametric approach provides a confidence interval is the following. The i.i.d. r.v.'s X_1, \dots, X_n have (unknown) d.f. F , and let $F_n(x)$ be the empirical d.f. based on the X_i 's, as defined in Application 5 to the WLLN in Chapter 7. We saw there that

$$F_n(x) = \sum_{i=1}^n Y_i(x), \quad Y_1(x), \dots, Y_n(x) \text{ independent r.v.'s } \sim B(1, F(x)).$$

Then, by the CLT,

$$\frac{\sqrt{n}[F_n(x) - F(x)]}{\sqrt{F(x)[1 - F(x)]}} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1). \quad (6)$$

Also,

$$F_n(x) \xrightarrow[n \rightarrow \infty]{P} F(x), \quad \text{or} \quad \frac{F_n(x)[1 - F_n(x)]}{F(x)[1 - F(x)]} \xrightarrow[n \rightarrow \infty]{P} 1. \quad (7)$$

From (6) and (7) and Theorem 6(iii) in Chapter 7, it follows that:

$$\frac{\sqrt{n}[F_n(x) - F(x)]/\sqrt{F(x)[1 - F(x)]}}{\sqrt{F_n(x)[1 - F_n(x)]}/\sqrt{F(x)[1 - F(x)]}} = \frac{\sqrt{n}[F_n(x) - F(x)]}{\sqrt{F_n(x)[1 - F_n(x)]}} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1).$$

It follows that, for all sufficiently large n (depending on x), the following interval is a confidence interval for $F(x)$ with confidence coefficient approximately $1 - \alpha$; namely,

$$\left[F_n(x) - z_{\frac{\alpha}{2}} \sqrt{\frac{F_n(x)[1 - F_n(x)]}{n}}, \quad F_n(x) + z_{\frac{\alpha}{2}} \sqrt{\frac{F_n(x)[1 - F_n(x)]}{n}} \right]. \quad (8)$$

As an application of formula (8), consider the following example.

EXAMPLE 2

Refer again to Example 22 in Chapter 1 (see also Example 1 in this chapter), and construct a confidence interval for $F(3)$ with approximately 95% confidence coefficient, where F is the d.f. of the r.v.'s describing the GPA scores.

DISCUSSION In this example, $n = 34$ and the number of the observations which are ≤ 3 are 18 (the following; 2.36, 2.36, 2.66, 2.68, 2.48, 2.46, 2.63, 2.44, 2.13, 2.41, 2.55, 2.80, 2.79, 2.89, 2.91, 2.75, 2.73, and 3.00). Then

$$F_{34}(3) = \frac{18}{34} = \frac{9}{17} \simeq 0.529, \quad \sqrt{\frac{F_{34}(3)[1 - F_{34}(3)]}{34}} \simeq 0.086,$$

and therefore the required (observed) confidence interval is:

$$[0.529 - 1.96 \times 0.086, 0.529 + 1.96 \times 0.086] \simeq [0.360, 0.698].$$

REMARK 1 It should be pointed out that the confidence interval given by (8) is of limited usefulness, because the value of (the large enough) n for which (8) holds depends on x .

15.2 Confidence Intervals for Quantiles of a Distribution Function

In the previous section, we constructed a confidence interval for the mean μ of a distribution, whether its variance is known or not, with confidence coefficient approximately a prescribed number $1 - \alpha$ ($0 < \alpha < 1$). Also, such an interval was constructed for each value $F(x)$ of a d.f. F . Now, we have seen (in Section 3.4 of Chapter 3) that the median, and, more generally, the quantiles of a d.f. F are important quantities through which we gain information about F . It would then be worth investigating the possibility of constructing confidence intervals for quantiles of F . To simplify matters, it will be assumed that F is continuous, and that for each $p \in (0, 1)$, there is a unique p th quantile x_p ; i.e., $F(x_p) = P(X \leq x_p) = p$. The objective is to construct a confidence interval for x_p , and, in particular, for the median $x_{0.50}$. This is done below in a rather neat manner, except that we don't have much control on the confidence coefficient involved. Specifically, the following result is established.

THEOREM 1

Let X_1, \dots, X_n be i.i.d. r.v.'s with d.f. F , and let Y_1, \dots, Y_n be the order statistics of the X_i 's. For $p \in (0, 1)$, let x_p be the unique (by assumption) p th quantile of F . Then, for any $1 \leq i < j \leq n$, the random interval $[Y_i, Y_j]$ is a confidence interval for x_p with confidence coefficient $\sum_{k=i}^{j-1} \binom{n}{k} p^k (1-p)^{n-k}$.

PROOF Define the r.v.'s W_j , $j = 1, \dots, n$ as follows:

$$W_j = \begin{cases} 1 & \text{if } X_j \leq x_p \\ 0 & \text{if } X_j > x_p, \quad j = 1, \dots, n. \end{cases}$$

Then the r.v.'s W_1, \dots, W_n are independent and distributed as $B(n, p)$, since $P(W_j = 1) = P(X_j \leq x_p) = F(x_p) = p$. Therefore

$$P(\text{at least } i \text{ of } X_1, \dots, X_n \text{ are } \leq x_p) = \sum_{k=i}^n \binom{n}{k} p^k (1-p)^{n-k}.$$

However, $P(\text{at least } i \text{ of } X_1, \dots, X_n \text{ are } \leq x_p) = P(Y_i \leq x_p)$. Thus,

$$P(Y_i \leq x_p) = \sum_{k=i}^n \binom{n}{k} p^k (1-p)^{n-k}. \quad (9)$$

Next, for $1 \leq i < j \leq n$, we, clearly, have;

$$\begin{aligned} P(Y_i \leq x_p) &= P(Y_i \leq x_p, Y_j \geq x_p) + P(Y_i \leq x_p, Y_j < x_p) \\ &= P(Y_i \leq x_p \leq Y_j) + P(Y_j < x_p) \\ &= P(Y_i \leq x_p \leq Y_j) + P(Y_j \leq x_p), \end{aligned} \quad (10)$$

since $P(Y_i \leq x_p, Y_j < x_p) = P(Y_i \leq x_p, Y_j \leq x_p) = P(Y_j \leq x_p)$ by the fact that $(Y_j \leq x_p) \subseteq (Y_i \leq x_p)$. Then, relations (9) and (10) yield:

$$\begin{aligned} P(Y_i \leq x_p \leq Y_j) &= \sum_{k=i}^n \binom{n}{k} p^k (1-p)^{n-k} - P(Y_j \leq x_p) \\ &= \sum_{k=i}^n \binom{n}{k} p^k (1-p)^{n-k} - \sum_{k=j}^n \binom{n}{k} p^k (1-p)^{n-k} \\ &= \sum_{k=i}^{j-1} \binom{n}{k} p^k (1-p)^{n-k}. \end{aligned} \quad (11)$$

So, the random interval $[Y_i, Y_j]$ contains the point x_p with probability $\sum_{k=i}^{j-1} \binom{n}{k} p^k (1-p)^{n-k}$, as was to be seen. \blacktriangle

REMARK 2

(i) From relation (11), it is clear that, although p is fixed, we can enlarge the confidence coefficient $\sum_{k=i}^{j-1} \binom{n}{k} p^k (1-p)^{n-k}$ by taking a smaller i and/or a larger j . The price we pay, however, is that of having a larger confidence interval.

(ii) By the fact that the confidence interval $[Y_i, Y_j]$ does not have a prescribed confidence coefficient $1 - \alpha$, as is the case in the usual construction of confidence intervals, we often refer to the probability $\sum_{k=i}^{j-1} \binom{n}{k} p^k (1-p)^{n-k}$ as the *probability of coverage* of x_p by $[Y_i, Y_j]$.

EXAMPLE 3

Consider the i.i.d. r.v.'s X_1, \dots, X_{20} with continuous d.f. F , which has unique $x_{0.50}$, $x_{0.25}$, and $x_{0.75}$, and let Y_1, \dots, Y_{20} be the corresponding order statistics. Then consider several confidence intervals for $x_{0.50}$, $x_{0.25}$, and $x_{0.75}$, and calculate the respective coverage probabilities.

DISCUSSION Using formula (11), we obtain the coverage probabilities listed in Table 15.1 for several confidence intervals for the median $x_{0.50}$ and the first quartile $x_{0.25}$. For the calculation of coverage probabilities for confidence intervals for the third quartile $x_{0.75}$, we employ the following formula, which allows us to use the Binomial tables; namely,

$$\sum_{k=i}^{j-1} \binom{20}{k} (0.75)^k (0.25)^{20-k} = \sum_{r=20-j+1}^{20-i} \binom{20}{r} (0.25)^r (0.75)^{20-r}.$$

Table 15.1

Quantile	Confidence Interval	Coverage Probability
$x_{0.50}$	(Y_9, Y_{12})	0.3364
	(Y_8, Y_{13})	0.6167
	(Y_7, Y_{14})	0.8107
	(Y_6, Y_{15})	0.9216
$x_{0.25}$	(Y_6, Y_6)	0.2024
	(Y_3, Y_7)	0.5606
	(Y_2, Y_8)	0.8069
	(Y_1, Y_9)	0.9348
$x_{0.75}$	(Y_{15}, Y_{17})	0.2024
	(Y_{14}, Y_{18})	0.5606
	(Y_{13}, Y_{19})	0.8069
	(Y_{12}, Y_{20})	0.9348

15.3 The Two-Sample Sign Test

In this brief section, we discuss a technique of comparing two populations by means of the so-called *sign test*. The test requires that the two samples available are of the same size, and makes no direct use of the values observed; instead, what is really used is the relative size of the components in the pairs of r.v.'s. Some cases where such a test would be appropriate include those in which one is interested in comparing the effectiveness of two different drugs used for the treatment of the same disease, the efficiency of two manufacturing processes producing the same item, the response of n customers regarding their preferences toward a certain consumer item, etc.

In more precise terms, let X_1, \dots, X_n be i.i.d. r.v.'s with *continuous* d.f. F , and let Y_1, \dots, Y_n be i.i.d. r.v.'s with *continuous* d.f. G ; the two samples are assumed to be independent. On the basis of the X_i 's and Y_j 's, we wish to test the null hypothesis $H_0: F = G$ against any one of the alternatives $H_A: F > G$, $H'_A: F < G$, $H''_A: F \neq G$. The inequality $F > G$ means that $F(z) \geq G(z)$ for all z , and $F(z) > G(z)$ for at least one z ; likewise for $F < G$. To this end, set

$$Z_i = \begin{cases} 1 & \text{if } X_i < Y_i \\ 0 & \text{if } X_i > Y_i \end{cases} \quad p = P(X_i < Y_i), \quad i = 1, \dots, n, \quad Z = \sum_{i=1}^n Z_i. \quad (12)$$

It is clear that the r.v.'s Z_1, \dots, Z_n are independent and distributed as $B(1, p)$, so that the r.v. Z is distributed as $B(n, p)$. Under the hypothesis H_0 , $p = \frac{1}{2}$, whereas under H_A , H'_A , and H''_A , we have, respectively, $p > \frac{1}{2}$, $p < \frac{1}{2}$, $p \neq \frac{1}{2}$. Thus, the problem of testing H_0 becomes, equivalently, that of testing $\tilde{H}_0: p = \frac{1}{2}$ in the $B(n, p)$ distribution. Formulating the relevant results, and drawing upon Application 1 in Section 11.3 of Chapter 11, we have the following theorem.

THEOREM 2

Consider the independent samples of the i.i.d. r.v.'s X_1, \dots, X_n and Y_1, \dots, Y_n with respective continuous d.f.'s F and G . Then, for testing the null hypothesis $H_0: F = G$ against any one of the alternatives $H_A: F > G$, or $H'_A: F < G$, or $H''_A: F \neq G$, at level of significance α , the hypothesis H_0 is rejected, respectively, whenever

$$Z \geq C, \quad \text{or} \quad Z \leq C', \quad \text{or} \quad Z \leq C_1 \quad \text{or} \quad Z \geq C_2. \quad (13)$$

The cutoff points C, C' , and C_1, C_2 are determined by the relations:

$$\left. \begin{aligned} P(Z > C) + \gamma P(Z = C) &= \alpha, \quad \text{or} \quad P(Z \leq C) - \gamma P(Z = C) = 1 - \alpha, \\ P(Z < C') + \gamma' P(Z = C') &= \alpha, \\ P(Z < C_1) + \gamma_0 P(Z = C_1) &= \frac{\alpha}{2} \quad \text{and} \quad P(Z > C_2) + \gamma_0 P(Z = C_2) = \frac{\alpha}{2}, \\ \text{or} \\ P(Z < C_1) + \gamma_0 P(Z = C_1) &= \frac{\alpha}{2} \quad \text{and} \quad P(Z \leq C_2) - \gamma_0 P(Z = C_2) = 1 - \frac{\alpha}{2}, \end{aligned} \right\} \quad (14)$$

and $Z \sim B(n, 1/2)$ under H_0 .

For large values of n , the CLT applies and the cutoff points are given by the relations:

$$\left. \begin{aligned} C &\simeq \frac{n}{2} + z_\alpha \frac{\sqrt{n}}{2}, & C' &\simeq \frac{n}{2} - z_\alpha \frac{\sqrt{n}}{2} \\ C_1 &\simeq \frac{n}{2} - z_{\frac{\alpha}{2}} \frac{\sqrt{n}}{2}, & C_2 &\simeq \frac{n}{2} + z_{\frac{\alpha}{2}} \frac{\sqrt{n}}{2}. \end{aligned} \right\} \quad (15)$$

EXAMPLE 4

Refer to Example 25 in Chapter 1 regarding the plant height (in 1/8 inches) of cross-fertilized and self-fertilized plants. Denote by X_i 's and Y_i 's, respectively, the heights of cross-fertilized and self-fertilized plants. Then the observed values for the 15 pairs are given in Example 25 of Chapter 1, which are reproduced

in the Table 15.2 for convenience. At the level of significance $\alpha = 0.05$, test the null hypothesis $H_0: F = G$, where F and G are the d.f.'s of the X_i 's and Y_i 's, respectively.

Table 15.2

Pair	Cross-	Self-	Pair	Cross-	Self-
1	188	139	9	146	132
2	96	163	10	173	144
3	168	160	11	186	130
4	176	160	12	168	144
5	153	147	13	177	102
6	172	149	14	184	124
7	177	149	15	96	144
8	163	122			

DISCUSSION From Table 15.2, we have:

$$Z_1 = 0, \quad Z_2 = 1, \quad Z_3 = 0, \quad Z_4 = 0, \quad Z_5 = 0, \quad Z_6 = 0, \quad Z_7 = 0, \quad Z_8 = 0, \\ Z_9 = 0, \quad Z_{10} = 0, \quad Z_{11} = 0, \quad Z_{12} = 0, \quad Z_{13} = 0, \quad Z_{14} = 0, \quad Z_{15} = 1,$$

so that $Z = 2$. Suppose first that the alternative is $H'_A: F \neq G$. Then H_0 is rejected in favor of H'_A whenever $Z \leq C_1$ or $Z \geq C_2$, where:

$$P(Z < C_1) + \gamma_0 P(Z = C_1) = 0.025 \quad \text{and} \quad P(Z \leq C_2) - \gamma_0 P(Z = C_2) = 0.975, \\ \text{and } Z \sim B(15, 1/2).$$

From the Binomial tables, we find $C_1 = 4$, $C_2 = 11$, and $\gamma_0 = \frac{37}{208} \simeq 0.178$. Since $Z = 2 < C_1 (= 4)$, the null hypothesis is rejected. Next, test H_0 against the alternative $H'_A: p < \frac{1}{2}$ again at level $\alpha = 0.05$. Then H_0 is rejected in favor of H'_A whenever $Z \leq C'$, where C' is determined by:

$$P(Z < C') + \gamma' P(Z = C') = 0.05, \quad Z \sim B(15, 1/2).$$

From the Binomial tables, we find $C' = 4$ and $\gamma' = \frac{81}{104} \simeq 0.779$. Since $Z = 2 < C' (= 4)$, H_0 is rejected in favor of H'_A , which is consistent with what the data say.

For the Normal approximation, we get from (15): $z_{0.025} = 1.96$, so that $C_1 = 3.703$, $C_2 \simeq 11.297$, and H_0 is rejected again, since $Z = 2 < C_1 (\simeq 3.703)$. Also, $z_{0.05} = 1.645$, and hence $C' \simeq 4.314$. Again, H_0 is rejected in favor of H'_A , since $Z = 2 < C' (\simeq 4.314)$.

15.4 The Rank Sum and the Wilcoxon–Mann–Whitney Two-Sample Tests

The purpose of this section is the same as that of the previous section; namely, the comparison of the d.f.'s of two independent samples of i.i.d. r.v.'s. However, the technique used in the last section may not apply here, as the two samples may be of different size, and therefore no pairwise comparison is possible (without discarding observations!)

So, what we have here is two independent samples consisting of the i.i.d. r.v.'s X_1, \dots, X_m with *continuous* d.f. F , and the i.i.d. r.v.'s Y_1, \dots, Y_n with *continuous* d.f. G . The problem is that of testing the null hypothesis $H_0: F = G$ against any one of the alternatives $H_A: F > G$, or $H'_A: F < G$, or $H''_A: F \neq G$. The test statistic to be used makes no use of the actual values of the X_i 's and the Y_j 's, but rather of their ranks in the combined sample, which are defined as follows. Consider the combined sample of X_1, \dots, X_m and Y_1, \dots, Y_n , and order them in ascending order. Because of the assumption of continuity of F and G , we are going to have strict inequalities with probability one. Then the *rank* of X_i , to be denoted by $R(X_i)$, is that integer among the integers $1, 2, \dots, m+n$, which corresponds to the position of X_i . The *rank* $R(Y_j)$ of Y_j is defined similarly. Next, consider the *rank sums* R_X and R_Y defined by:

$$R_X = \sum_{i=1}^m R(X_i), \quad R_Y = \sum_{j=1}^n R(Y_j). \quad (16)$$

Then

$$R_X + R_Y = \frac{(m+n)(m+n+1)}{2}, \quad (17)$$

because $R_X + R_Y = \sum_{i=1}^m R(X_i) + \sum_{j=1}^n R(Y_j) = 1 + 2 + \dots + m + n = \frac{(m+n)(m+n+1)}{2}$. Before we go further, let us illustrate the concepts introduced so far by a numerical example.

EXAMPLE 5

Let $m = 5$, $n = 4$, and suppose that:

$$\begin{aligned} X_1 = 78, \quad X_2 = 65, \quad X_3 = 74, \quad X_4 = 45, \quad X_5 = 82, \\ Y_1 = 110, \quad Y_2 = 71, \quad Y_3 = 53, \quad Y_4 = 50. \end{aligned}$$

Combining the X_i 's and the Y_j 's and arranging them in ascending order, we get:

$$\begin{array}{cccccccccc} 45 & 50 & 53 & 65 & 71 & 74 & 78 & 82 & 110 \\ (X) & (Y) & (Y) & (X) & (Y) & (X) & (X) & (X) & (Y). \end{array}$$

$$\begin{aligned} \text{Then: } R(X_1) = 7, \quad R(X_2) = 4, \quad R(X_3) = 6, \quad R(X_4) = 1, \quad R(X_5) = 8, \\ R(Y_1) = 9, \quad R(Y_2) = 5, \quad R(Y_3) = 3, \quad R(Y_4) = 2. \end{aligned}$$

It follows that: $R_X = 26$, $R_Y = 19$, and, of course, $R_X + R_Y = 45 = \frac{9 \times 10}{2}$.

The m ranks $(R(X_1), \dots, R(X_m))$ can be placed in m positions out of $m+n$ possible in $\binom{m+n}{m}$ different ways (the remaining n positions will be taken up by the n ranks $(R(Y_1), \dots, R(Y_n))$), and under the null hypothesis H_0 , each one of them is equally likely to occur. So, each one of the $\binom{m+n}{m}$ positions of $(R(X_1), \dots, R(X_m))$ has probability $1/\binom{m+n}{m}$. The alternative H_A stipulates that $F > G$; i.e., $F(z) \geq G(z)$ for all z , or $P(X \leq z) \geq P(Y \leq z)$ for all z , with the inequalities strict for at least one z , where the r.v.'s X and Y are distributed as the X_i 's and the Y_j 's, respectively. That is, under H_A , the X_i 's tend to be smaller than any z with higher probability than the Y_j 's are smaller than any z . Consequently, since $R_X + R_Y$ is fixed $= (m+n)(m+n+1)/2$, this suggests

that the rank sum R_X would tend to take small values. Therefore, H_0 should be rejected in favor of H_A whenever $R_X < C$. The rejection region is determined as follows: Consider the $\binom{m+n}{m}$ positions of the ranks $(R(X_1), \dots, R(X_m))$, and for each one of them, form the respective sum R_X . We start with the smallest value of R_X and proceed with the next smallest, etc, until we get to the k th smallest, where k is determined by: $k/\binom{m+n}{m} = \alpha$. (In the present setting, the level of significance α is taken to be an integer multiple of $1/\binom{m+n}{m}$, if we wish to have an exact level.) So, the rejection region consists of the k smallest values of R_X , where $k/\binom{m+n}{m}$ equals α .

Likewise, the hypothesis H_0 is rejected in favor of the alternative $H'_A: F < G$ whenever $R_X > C'$, and the rejection region consists of the k largest values of R_X , where $k/\binom{m+n}{m}$ equals α . Also, H_0 is rejected in favor of $H''_A: F \neq G$ whenever $R_X < C_1$ or $R_X > C_2$, and the rejection region consists of the smallest r values of R_X and the largest r values of R_X , where r satisfies the requirement $r/\binom{m+n}{m}$ equals $\alpha/2$.

Summarize these results in the following theorem.

THEOREM 3

Consider the independent samples of the i.i.d. r.v.'s X_1, \dots, X_m and Y_1, \dots, Y_n with respective continuous d.f.'s F and G . Then, for testing the null hypothesis $H_0: F = G$ against any one of the alternatives $H_A: F > G$, or $H'_A: F < G$, or $H''_A: F \neq G$, at level of significance α (so that α or $\frac{\alpha}{2}$ are integer multiples of $1/\binom{m+n}{m}$), the respective rejection regions of the rank sum tests consist of: The k smallest values of the rank sum R_X , where $k/\binom{m+n}{m} = \alpha$; the k largest values of the rank sum R_X , where k is as above; the r smallest and the r largest values of the rank sum R_X , where $r/\binom{m+n}{m} = \frac{\alpha}{2}$.

REMARK 3 In theory, carrying out the test procedures described in Theorem 3 is straightforward and neat. Their practical implementation, however, is another matter. To illustrate the difficulties involved, consider Example 5, where $m = 5$ and $n = 4$, so that $\binom{m+n}{m} = \binom{9}{5} = 126$. Thus, one would have to consider the 126 possible arrangements of the ranks $(R(X_1), \dots, R(X_5))$, form the respective rank sums, and see which ones of its values are to be included in the rejection regions. Clearly, this is not an easy task even for such small sample sizes.

A special interesting case where the rank sum test is appropriate is that where the d.f. G of the Y_i 's is assumed to be of the form:

$$G(x) = F(x - \Delta), \quad x \in \mathfrak{R}, \quad \text{for some unknown } \Delta \in \mathfrak{R}.$$

In such a case, we say that G is a *shift* of F (to the right, if $\Delta > 0$, and to the left, if $\Delta < 0$). Then the hypothesis $H_0: F = G$ is equivalent to testing $\Delta = 0$, and the alternatives $H_A: F > G$, $H'_A: F < G$, $H''_A: F \neq G$ are equivalent to: $\Delta > 0$, $\Delta < 0$, $\Delta \neq 0$.

Because of the difficulties associated with the implementation of the rank sum tests, there is an alternative closely related to it, for which a Normal approximation may be used. This is the *Wilcoxon–Mann–Whitney two-sample test*. For the construction of the relevant test statistic, consider all mn pairs (X_i, Y_j) , and among them, count those for which $X_i > Y_j$. The resulting r.v. is denoted by U and is the statistic to be employed. More formally, let the function u be defined by:

$$u(z) = \begin{cases} 1 & \text{if } z > 0 \\ 0 & \text{if } z < 0. \end{cases} \quad (18)$$

Then, clearly, the statistic U may be written thus:

$$U = \sum_{i=1}^m \sum_{j=1}^n u(X_i - Y_j). \quad (19)$$

The statistics U , R_X , and R_Y are related as follows.

LEMMA 1 Let R_X , R_Y , and U be defined, respectively, by (16) and (19). Then:

$$U = R_X - \frac{m(m+1)}{2} = mn + \frac{n(n+1)}{2} - R_Y. \quad (20)$$

PROOF Deferred to Subsection 15.4.1.

On the basis of (20), Theorem 3 may be rephrased as follows in terms of the U statistics. ▲

THEOREM 4

In the notation of Theorem 3 and for testing the null hypothesis H_0 against any one of the alternatives H_A , or H'_A , or H''_A as described there, at level of significance α , the Wilcoxon–Mann–Whitney test rejects H_0 , respectively:

$$\left. \begin{array}{l} \text{For } U < C, \text{ where } C \text{ is determined by } P(U < C | H_0) = \alpha; \\ \text{or } U > C', \text{ where } C' \text{ is determined by } P(U > C' | H_0) = \alpha; \\ \text{or } U < C_1 \text{ or } U > C_2, \text{ where } C_1 \text{ and } C_2 \text{ are determined} \\ \text{by } P(U < C_1 | H_0) = P(U > C_2 | H_0) = \frac{\alpha}{2}. \end{array} \right\} \quad (21)$$

In determining the cutoff points C , C' , C_1 , and C_2 above, we are faced with the same difficulty as that in the implementation of the rank sum tests. However, presently, there are two ways out of it. First, tables are available for small values of m and n ($n \leq m \leq 10$) (see page 341 in the book *Handbook of Statistical Tables*, Addison-Wesley (1962), by D. B. Owen), and second, for large values of m and n , and under H_0 ,

$$\frac{U - EU}{s.d.(U)} \simeq Z \sim N(0, 1), \quad (22)$$

where, under H_0 ,

$$EU = \frac{mn}{2}, \quad \text{Var}(U) = \frac{mn(m+n+1)}{12}. \quad (23)$$

EXAMPLE 6

Refer to Example 5 and test the hypothesis $H_0: F = G$ against the alternatives $H'_A: F \neq G$ and $H_A: F > G$.

DISCUSSION In Example 5 we saw that $R_X = 26$ (and $R_Y = 19$). Since $m = 5$ and $n = 4$, relation (20) gives: $U = 11$. From the tables cited above, we have: $P(U \leq 2) = P(U \geq 17) = 0.032$. So, $C_1 = 2$, $C_2 = 17$, and H_0 is rejected in favor of H'_A at the level of significance 0.064. From the same tables, we have that $P(U \leq 3) = 0.056$, so that $C = 3$, and H_0 is rejected in favor of H_A at level of significance 0.056. The results stated in (23) are formulated as a lemma below.

LEMMA 2 With U defined by (19), the relations in (23) hold true under H_0 .

PROOF Deferred to Subsection 15.4.1.

For the statistic U , the CLT holds. Namely,

LEMMA 3 With U , EU , and $Var(U)$ defined, respectively, by (19) and (23),

$$\frac{U - EU}{s.d.(U)} \xrightarrow{m, n \rightarrow \infty} Z \sim N(0, 1). \quad (24)$$

PROOF It is omitted. By means of the result in (24), the cutoff points in (21) may be determined approximately, by means of the Normal tables. That is, we have the following corollary.

COROLLARY (to Theorem 4 and Lemma 3) For large m and n , the cutoff points in (21) are given by the following approximate quantities:

$$\left. \begin{aligned} C &\simeq \frac{mn}{2} - z_\alpha \sqrt{\frac{mn(m+n+1)}{12}}, & C' &\simeq \frac{mn}{2} + z_\alpha \sqrt{\frac{mn(m+n+1)}{12}}, \\ C_1 &\simeq \frac{mn}{2} - z_{\frac{\alpha}{2}} \sqrt{\frac{mn(m+n+1)}{12}}, & C_2 &\simeq \frac{mn}{2} + z_{\frac{\alpha}{2}} \sqrt{\frac{mn(m+n+1)}{12}}. \end{aligned} \right\} \quad (25)$$

PROOF Follows immediately, from (21) and (24). \blacktriangle

EXAMPLE 7

Refer to Example 25 in Chapter 1 (see also Example 4 here), and test the null hypothesis $H_0: F = G$ at level of significance $\alpha = 0.05$ by using Theorem 4 and the above corollary.

DISCUSSION Here $m = n = 15$, $z_{0.05} = 1.645$, and $z_{0.025} = 1.96$. Then:

$$C_1 \simeq 112.50 - 1.96 \times 24.105 = 112.50 - 47.2458 \simeq 65.254,$$

$$C_2 \simeq 112.50 + 47.2458 \simeq 159.746,$$

$$C \simeq 112.50 - 1.645 \times 24.105 \simeq 112.50 - 39.653 = 72.847,$$

$$C' \simeq 112.50 + 39.653 = 152.153.$$

Next, comparing all 15×15 pairs in Table 15.3, we get the observed value of $U = 185$.

Table 15.3

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
X	188	96	168	176	153	172	177	163	146	173	186	168	177	184	96
Y	139	163	160	160	147	149	149	122	132	144	130	144	102	124	144

Therefore the hypothesis H_0 is rejected in favor of H'_A , since $U = 185 > 159.746 = C_2$; the null hypothesis is also rejected in favor of H_A , since $U = 185 > 152.153 = C'$; but the null hypothesis is not rejected when the alternative is H_A , because $U = 185 \not\leq 72.847 = C$.

15.4.1 Proofs of Lemmas 1 and 2

PROOF OF LEMMA 1 Let $X_{(1)}, \dots, X_{(m)}$ be the order statistics of the r.v.'s X_1, \dots, X_m , and look at the rank $R(X_{(i)})$ in the combined sample of the X_i 's and the Y_j 's. For each $R(X_{(i)})$, there are $R(X_{(i)}) - 1$ X_i 's and Y_j 's preceding $X_{(i)}$. Of these, $i - 1$ are X_i 's and hence $R(X_{(i)}) - 1 - (i - 1) = R(X_{(i)}) - i$ are Y_j 's. Therefore

$$\begin{aligned} U &= [R(X_{(1)}) - 1] + \dots + [R(X_{(m)}) - m] \\ &= [R(X_{(1)}) + \dots + R(X_{(m)})] - (1 + \dots + m) \\ &= [R(X_1) + \dots + R(X_m)] - \frac{m(m+1)}{2} = R_X - \frac{m(m+1)}{2}, \end{aligned}$$

since $R(X_{(1)}) + \dots + R(X_{(m)})$ is simply a rearrangement of the terms in the rank sum $R(X_1) + \dots + R(X_m) = R_X$. Next, from the result just obtained and (17), we have:

$$\begin{aligned} U &= \frac{(m+n)(m+n+1)}{2} - R_Y - \frac{m(m+1)}{2} \\ &= \frac{(m+n)(m+n+1) - m(m+1)}{2} - R_Y = mn + \frac{n(n+1)}{2} - R_Y. \quad \blacktriangle \end{aligned}$$

PROOF OF LEMMA 2 Recall that all derivations below are carried out under the assumption that $H_0(F = G)$ holds. Next, for any i and j :

$$Eu(X_i - Y_j) = 1 \times P(X_i > Y_j) = \frac{1}{2}, \quad Eu^2(X_i - Y_j) = 1^2 \times P(X_i > Y_j) = \frac{1}{2},$$

so that

$$\text{Var}(u(X_i - Y_j)) = \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$$

Therefore $EU = \sum_{i=1}^m \sum_{j=1}^n \frac{1}{2} = \frac{mn}{2}$, and

$$\begin{aligned} \text{Var}(U) &= \sum_{i=1}^m \sum_{j=1}^n \text{Var}(u(X_i - Y_j)) + \sum_{i=1}^m \sum_{j=1}^n \sum_{k=1}^m \sum_{l=1}^n \text{Cov}(u(X_i - Y_j), u(X_k - Y_l)) \\ &= \frac{mn}{4} + \text{sum of the covariances on the right-hand side above.} \quad (26) \end{aligned}$$

Regarding the covariances, we consider the following cases. First, let $i \neq k$ and $j \neq l$. Then $\text{Cov}(u(X_i - Y_j), u(X_k - Y_l)) = 0$ by independence. Thus, it suffices to restrict ourselves to pairs (X_i, Y_j) and (X_k, Y_l) for which $i = k$ and $j \neq l$, and $i \neq k$ and $j = l$. In order to see how many such pairs there are, consider the following array:

$$\begin{array}{cccc} (X_1, Y_1), & (X_1, Y_2), & \dots, & (X_1, Y_n) \\ (X_2, Y_1), & (X_2, Y_2), & \dots, & (X_2, Y_n) \\ \vdots & \vdots & & \vdots \\ (X_m, Y_1), & (X_m, Y_2), & \dots, & (X_m, Y_n). \end{array}$$

From each one of the m rows, we obtain $\binom{m}{2} \times 2 = n(n-1)$ terms of the form: $\text{Cov}(u(X - Y), u(X - Z))$, where X, Y, Z are independent r.v.'s with d.f. $F = G$. Since $\text{Cov}(u(X - Y), u(X - Z)) = P(X > Y \text{ and } X > Z) - \frac{1}{4}$, we have then $n(n-1)P(X > Y \text{ and } X > Z) - \frac{n(n-1)}{4}$ as a contribution to the sum of the covariances from each row, and therefore from the m rows, the contribution to the sum of the covariances is:

$$mn(n-1)P(X > Y \text{ and } X > Z) - \frac{mn(n-1)}{4}. \quad (27)$$

Next, from each one of the n columns, we obtain $\binom{m}{2} \times 2 = m(m-1)$ terms of the form: $\text{Cov}(u(X - Z), u(Y - Z)) = P(X > Z \text{ and } Y > Z) - \frac{1}{4}$. Therefore the contribution from the n columns to the sum of the covariances is:

$$mn(m-1)P(X > Z \text{ and } Y > Z) - \frac{mn(m-1)}{4}. \quad (28)$$

Now,

$$\begin{aligned} (X > Y \text{ and } X > Z) &= (X > Y, Y > Z, X > Z) \cup (X > Y, Y \leq Z, X > Z) \\ &= (X > Y, Y > Z) \cup (X > Z, Z \geq Y) = (X > Y > Z) \cup (X > Z \geq Y), \end{aligned}$$

since

$$(X > Y, Y > Z) \subseteq (X > Z), \quad \text{and} \quad (X > Z, Z \geq Y) \subseteq (X > Y).$$

Thus,

$$P(X > Y \text{ and } X > Z) = P(X > Y > Z) + P(X > Z > Y). \quad (29)$$

Likewise,

$$\begin{aligned} (X > Z \text{ and } Y > Z) &= (X > Y, Y > Z, X > Z) \cup (X \leq Y, Y > Z, X > Z) \\ &= (X > Y, Y > Z) \cup (Y \geq X, X > Z) = (X > Y > Z) \cup (Y \geq X > Z), \end{aligned}$$

since

$$(X > Y, Y > Z) \subseteq (X > Z), \quad \text{and} \quad (Y \geq X, X > Z) \subseteq (Y > Z).$$

Thus,

$$P(X > Z \text{ and } Y > Z) = P(X > Y > Z) + P(Y > X > Z). \quad (30)$$

The r.v.'s X , Y , and Z satisfy, with probability one, exactly one of the inequalities:

$$\begin{aligned} X > Y > Z, \quad X > Z > Y, \quad Y > X > Z, \\ Y > Z > X, \quad Z > X > Y, \quad Z > Y > X, \end{aligned}$$

and each one of these inequalities has probability $1/6$. Then, the expressions in (27) and (28) become, by means of (29) and (30), respectively:

$$\frac{mn(n-1)}{3} - \frac{mn(n-1)}{4} = \frac{mn(n-1)}{12}, \quad \frac{mn(m-1)}{3} - \frac{mn(m-1)}{4} = \frac{mn(m-1)}{12}$$

Then, formula (26) yields:

$$\text{Var}(U) = \frac{mn}{4} + \frac{mn(n-1)}{12} + \frac{mn(m-1)}{12} = \frac{mn(m+n+1)}{12}. \quad \blacktriangle$$

15.5 Nonparametric Curve Estimation

For quite a few years now, work on nonparametric methodology has switched decisively in what is referred to as nonparametric *curve estimation*. Such estimation includes estimation of d.f.'s, of p.d.f.'s or functions thereof, regression functions, etc. The empirical d.f. is a case of nonparametric estimation of a d.f., although there are others as well. In this section, we are going to describe briefly a way of estimating nonparametrically a p.d.f., and record some of the (asymptotic) desirable properties of the proposed estimate. Also, the problem of estimating, again nonparametrically, a regression function will be discussed very briefly. There is already a huge statistical literature in this area, and research is currently very active.

15.5.1 Nonparametric Estimation of a Probability Density Function

The problem we are faced with here is the following: We are given n i.i.d. r.v.'s X_1, \dots, X_n with p.d.f. f of the *continuous type*, for which very little is known, and we are asked to construct a nonparametric estimate $\hat{f}_n(x)$ of $f(x)$, for each $x \in \mathfrak{R}$, based on the random sample X_1, \dots, X_n . The approach to be used here is the so-called *kernel-estimation* approach. According to this method, we select a (known) p.d.f. to be denoted by K and to be termed a *kernel*, subject to some rather minor requirements. Also, we choose a sequence of positive numbers, denoted by $\{h_n\}$, which has the property that $h_n \rightarrow 0$ as $n \rightarrow \infty$ and also satisfies some additional requirements. The numbers h_n , $n \geq 1$, are referred to as *bandwidth* for a reason to be seen below. Then, on the basis of the random sample X_1, \dots, X_n , the kernel K , and the bandwidths h_n , $n \geq 1$, the proposed estimate of $f(x)$ is $\hat{f}_n(x)$ given by:

$$\hat{f}_n(x) = \frac{1}{nh_n} \sum_{i=1}^n K\left(\frac{x - X_i}{h_n}\right). \quad (31)$$

EXAMPLE 8

Construct the kernel estimate of $f(x)$, for each $x \in \mathfrak{R}$, by using the $U(-1, 1)$ kernel; i.e., by taking

$$K(x) = \frac{1}{2}, \quad \text{for } -1 \leq x \leq 1, \text{ and } 0, \text{ otherwise.}$$

DISCUSSION Here, it is convenient to use the indicator notation, namely, $K(x) = I_{[-1,1]}(x)$ (where, it is recalled, $I_A(x) = 1$ if $x \in A$, and 0 if $x \in A^c$). Then the estimate (31) becomes as follows:

$$\hat{f}_n(x) = \frac{1}{nh_n} \sum_{i=1}^n I_{[-1,1]} \left(\frac{x - X_i}{h_n} \right), \quad x \in R. \quad (32)$$

So, $I_{[-1,1]} \left(\frac{x - X_i}{h_n} \right) = 1$, if and only if $x - h_n \leq X_i \leq x + h_n$; in other words, in forming $\hat{f}_n(x)$, we use only those observations X_i which lie in the window $[x - h_n, x + h_n]$. The breadth of this window is, clearly, determined by h_n , and this is the reason that h_n is referred to as the bandwidth.

Usually, the minimum of assumptions required of the kernel K and the bandwidth h_n , in order for us to be able to establish some desirable properties of the estimate $\hat{f}_n(x)$ given in (31), are the following:

$$\left. \begin{array}{l} K \text{ is bounded; i.e., } \sup \{K(x); x \in \mathfrak{R}\} < \infty. \\ xK(x) \text{ tends to } 0 \text{ as } x \rightarrow \pm\infty; \text{ i.e., } |xK(x)| \xrightarrow{|x| \rightarrow \infty} 0. \\ K \text{ is symmetric about } 0; \text{ i.e., } K(-x) = K(x), \quad x \in R. \end{array} \right\} \quad (33)$$

$$\left. \begin{array}{l} \text{As } n \rightarrow \infty: \text{ (i) } \quad (0 <) h_n \rightarrow 0 \\ \text{(ii) } \quad nh_n \rightarrow \infty \\ \text{(iii) } \quad nh_n^2 \rightarrow \infty. \end{array} \right\} \quad (34)$$

REMARK 4 Observe that requirements (33) are met for the kernel used in (32). Furthermore, the convergences in (34) are satisfied if one takes, e.g., $h_n = n^{-\alpha}$ with $0 < \alpha < 1/2$. Below, we record three (asymptotic) results regarding the estimate $\hat{f}_n(x)$ given in (31).

THEOREM 5

Under assumptions (33) and (34)(i), the estimate $\hat{f}_n(x)$ given in (31) is an *asymptotically unbiased* estimate of $f(x)$ for every $x \in \mathfrak{R}$ at which f is continuous; i.e.,

$$E\hat{f}_n(x) \rightarrow f(x) \text{ as } n \rightarrow \infty.$$

THEOREM 6

Under assumptions (33) and (34)(i), (ii), the estimate $\hat{f}_n(x)$ given in (31) is a *consistent in quadratic mean* estimate of $f(x)$ for every $x \in \mathfrak{R}$ at which f is continuous; i.e.,

$$E[\hat{f}_n(x) - f(x)]^2 \rightarrow 0 \text{ as } n \rightarrow \infty.$$

THEOREM 7

Under assumptions (33) and (34)(i)–(iii), the estimate $\hat{f}_n(x)$ given in (31) is *asymptotically normal*, when properly normalized, for every $x \in \mathfrak{R}$ at which f is continuous; i.e.,

$$\frac{\hat{f}_n(x) - E\hat{f}_n(x)}{\text{s.d.}(\hat{f}_n(x))} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1).$$

We have no intention of even attempting to prove any of the theorems just stated. Their proofs can be found in the second reference given below. In closing this section, it is only fitting to mention that the concept of kernel estimation of a p.d.f. was introduced by Murray Rosenblatt in 1956, and it was popularized by a fundamental paper by E. Parzen in 1962. The relevant references are as follows:

“Remarks on some nonparametric estimates of a density function” by M. Rosenblatt in the *Annals of Mathematical Statistics*, Vol. 27 (1956), pages 823–835. “On estimation of a probability density function and mode” by E. Parzen in the *Annals of Mathematical Statistics*, Vol. 33 (1962), pages 1065–1076.

15.5.2 Nonparametric Regression Estimation

In Chapter 13, a simple linear regression model was studied and its usefulness was demonstrated by means of specific examples. It was also stated that there is a definite need for more general regression models, where the linearity is retained, or it is discarded altogether. This issue is addressed, to a considerable extent, in this section.

Specifically, the model considered here is the following: For each $n = 1, 2, \dots$, consider points x_{n1}, \dots, x_{nm} in \mathfrak{R} , and at each one of them, an observation is taken, to be denoted by Y_{ni} , $i = 1, \dots, n$. It is assumed that Y_{ni} is equal to some unknown function g evaluated at x_{ni} except for an error e_{ni} ; i.e.,

$$Y_{ni} = g(x_{ni}) + e_{ni}, \quad i = 1, \dots, n. \quad (35)$$

On the errors e_{ni} , $i = 1, \dots, n$, we make the usual assumptions that they are i.i.d. r.v.'s with $Ee_{ni} = 0$ and $\text{Var}(e_{ni}) = \sigma^2 < \infty$.

The model in (1) of Chapter 13 is a very special case of the model just described. In the first place, the points where observations are taken are allowed here to depend on n , and second, the regression function in (1) of Chapter 13 is taken from here by setting $g(x) = \beta_1 + \beta_2 x$, so that $\beta_1 + \beta_2 x_i = g(x_i)$, $i = 1, \dots, n$. The function g in (35) is subject only to the requirement that it is defined on a bounded subset S of \mathfrak{R} and that it is continuous.

The objective here is to (nonparametrically) estimate the function $g(x)$, for each $x \in S$, by means of the observations Y_{n1}, \dots, Y_{nm} . The proposed estimate

is the statistic $g_n(x; \mathbf{x}_n)$ defined as follows:

$$g_n(x; \mathbf{x}_n) = \sum_{i=1}^n w_{ni}(x; \mathbf{x}_n) Y_{ni}, \quad (36)$$

where $\mathbf{x}_n = (x_{n1}, \dots, x_{nm})$ and w_{ni} are *weights*, properly chosen, which depend on the particular point x in S and also the points x_{n1}, \dots, x_{nm} , where observations are taken. The weights are required to satisfy certain conditions, and there is considerable flexibility in choosing them. We do not intend to enter here into this kind of detail. Instead, we restrict ourselves to stating three basic properties that the estimate defined in (36) satisfies.

THEOREM 8

Under suitable regularity conditions, the estimate $g_n(x; \mathbf{x}_n)$ defined in (36) is an *asymptotically unbiased estimate* of $g(x)$; i.e.,

$$Eg_n(x; \mathbf{x}_n) \xrightarrow[n \rightarrow \infty]{} g(x), \quad \text{for every } x \in S. \quad (37)$$

THEOREM 9

Under suitable regularity conditions, the estimate $g_n(x; \mathbf{x}_n)$ is a *consistent in quadratic mean estimate* of $g(x)$; i.e.,

$$E[g_n(x; \mathbf{x}_n) - g(x)]^2 \xrightarrow[n \rightarrow \infty]{} 0, \quad \text{for every } x \in S. \quad (38)$$

THEOREM 10

Under suitable regularity conditions, the estimate $g_n(x; \mathbf{x}_n)$, properly normalized, is *asymptotically Normal*; i.e.,

$$\frac{g_n(x; \mathbf{x}_n) - Eg_n(x; \mathbf{x}_n)}{\text{s.d.}(g_n(x; \mathbf{x}_n))} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1), \quad \text{for every } x \in S.$$

Also,

$$\frac{g_n(x; \mathbf{x}_n) - g(x)}{\text{s.d.}(g_n(x; \mathbf{x}_n))} \xrightarrow[n \rightarrow \infty]{d} Z \sim N(0, 1), \quad \text{for every } x \in S. \quad (39)$$

Convergences (37) and (38) provide asymptotic optimal properties for the estimate proposed in (36). If it happens that the error variance σ^2 is known, or an estimate of it is available, then convergence (39) provides a way of constructing confidence interval for $g(x)$ with confidence coefficient approximately equal to $1 - \alpha$ for large n . (See also Exercise 5.2.)

In the regression model considered in Chapter 13, one of the basic tenets was that the point x at which an observation Y is to be made can be chosen, more or less, at will. This, however, need not always be the case. Instead, it may happen that the point x itself is the observed value of a r.v. X . Thus, the setup here is as follows: A r.v. X is observed, and if $X = x$, then an observation is taken at the point x .

In this framework, several questions may be posed. One of the most important is this: Given that $X = x$, construct a *predictor* of Y corresponding to x . The proposed predictor is the conditional expectation of Y , given $X = x$, call it $m(x)$; i.e.,

$$m(x) = E(Y | X = x). \quad (40)$$

The quantity defined in (40) is an *unknown* function of x , since the conditional p.d.f. Y , given $X = x$, is unknown. The problem which then arises is that of estimating $m(x)$. The discussion in the remainder of this section revolves around this question.

Clearly, the estimation of $m(x)$ must be made on the basis of available data. To this effect, we assume that we have at our disposal n pairs of r.v.'s (X_i, Y_i) , $i = 1, \dots, n$ which are independent and distributed as the pair (X, Y) . Then the proposed estimate of $m(x)$, call it $\hat{m}_n(x)$, is the following:

$$\hat{m}_n(x) = \frac{\hat{w}_n(x)}{\hat{f}_n(x)}, \quad \text{where } \hat{w}_n(x) = \frac{1}{nh_n} \sum_{i=1}^n Y_i K\left(\frac{x - X_i}{h_n}\right), \quad (41)$$

and

$$\hat{f}_n(x) \text{ is given in (31); i.e., } \hat{f}_n(x) = \frac{1}{nh_n} \sum_{i=1}^n K\left(\frac{x - X_i}{h_n}\right).$$

The estimated predictor $\hat{m}_n(x)$ has several asymptotic optimal properties of which we single out only one here; namely, asymptotic normality.

THEOREM 11

Let $\hat{m}_n(x)$ be the estimate of the predictor $m(x)$ given by (41) and (40), respectively, and let $\sigma^2(x)$ be defined by:

$$\sigma^2(x) = \frac{\sigma_0^2(x)}{f(x)} \int_{-\infty}^{\infty} K^2(t) dt \quad (\text{for } f(x) > 0), \quad (42)$$

where $\sigma_0^2(x)$ is the conditional variance of Y , given $X = x$. Then the estimated predictor $\hat{m}_n(x)$, properly normalized, is *asymptotically Normal*; i.e.,

$$\sqrt{nh_n}[\hat{m}_n(x) - m(x)] \xrightarrow[n \rightarrow \infty]{d} N(0, \sigma^2(x)). \quad (43)$$

The variance $\sigma^2(x)$ of the limiting normal distribution is unknown, but an estimate of it may be constructed. Then the convergence (43) may be used to set up a confidence interval for $m(x)$ with confidence coefficient approximately equal to $1 - \alpha$ for large n . (See also Exercise 5.3.)

In closing this section, it should be mentioned that its purpose has been not to list detailed assumptions and present proofs (many of which are beyond the assumed level of this book anyway), but rather to point out that there are regression results available in the literature, way beyond the simple linear model studied in Chapter 13. Finally, let us mention a piece a terminology used in the literature, namely, the regression model defined by (35) is referred

to as a *fixed design* regression model, whereas the one defined by (40) is called a *stochastic design* regression model. The reasons for this are obvious. In the former case, the points where observations are taken are *fixed*, whereas in the latter case they are *values of a r.v. X*.

Exercises

5.1 Note: All convergences in this exercise hold for continuity points x of $f(x)$. In Theorem 7.1, it is stated that

$$\frac{\hat{f}_n(x) - E\hat{f}_n(x)}{s.d.(\hat{f}_n(x))} \xrightarrow[n \rightarrow \infty]{d} N(0, 1).$$

By this fact, Theorem 5, and some additional assumptions, it is also shown that

$$\frac{\hat{f}_n(x) - f(x)}{s.d.(\hat{f}_n(x))} \xrightarrow[n \rightarrow \infty]{d} N(0, 1). \quad (44)$$

(i) Use expression (31) in order to show that

$$\begin{aligned} \text{Var}(\hat{f}_n(x)) &= \frac{1}{nh_n^2} \text{Var}\left(K\left(\frac{x - X_1}{h_n}\right)\right), \quad \text{so that} \\ (nh_n)\text{Var}(\hat{f}_n(x)) &= \frac{1}{h_n} \text{Var}\left(K\left(\frac{x - X_1}{h_n}\right)\right). \end{aligned}$$

(ii) Use the formula $\text{Var}(X) = EX^2 - (EX)^2$, and the transformation $\frac{x-y}{h_n} = u$ with $-\infty < u < \infty$, in order to show that

$$\begin{aligned} (nh_n)\text{Var}(\hat{f}_n(x)) &= \int_{-\infty}^{\infty} K^2(u)f(x - h_nu) du \\ &\quad - h_n \left[\int_{-\infty}^{\infty} K(u)f(x - h_nu) du \right]^2. \end{aligned}$$

Now, it can be shown that

$$\int_{-\infty}^{\infty} K(u)f(x - h_nu) du \xrightarrow[n \rightarrow \infty]{} f(x) \int_{-\infty}^{\infty} K(u)du = f(x),$$

and

$$\int_{-\infty}^{\infty} K^2(u)f(x - h_nu)du \xrightarrow[n \rightarrow \infty]{} f(x) \int_{-\infty}^{\infty} K^2(u) du.$$

From these results, assumption (34)(i), and part (ii), it follows then that

$$\sigma_n^2(x) \stackrel{\text{def}}{=} (nh_n)\text{Var}(\hat{f}_n(x)) \xrightarrow[n \rightarrow \infty]{} f(x) \int_{-\infty}^{\infty} K^2(u) du \stackrel{\text{def}}{=} \sigma^2(x).$$

(45)

- (iii) From convergence (45) and Theorem 7, conclude (by means of the Corollary to Theorem 5 in Chapter 7) that

$$\hat{f}_n(x) - E\hat{f}_n(x) \xrightarrow[n \rightarrow \infty]{d} 0, \quad \text{and hence } \hat{f}_n(x) - E\hat{f}_n(x) \xrightarrow[n \rightarrow \infty]{p} 0. \quad (46)$$

- (iv) Use convergence (46) and Theorem 5 in order to conclude that $\hat{f}_n(x)$ is a *consistent* estimate of $f(x)$ (in the probability sense); i.e., $\hat{f}_n(x) \xrightarrow[n \rightarrow \infty]{p} f(x)$.

Set

$$\hat{\sigma}_n^2(x) = \hat{f}_n(x) \int_{-\infty}^{\infty} K^2(u) du. \quad (47)$$

- (v) Use relations (45) and (47) to conclude that

$$\frac{\hat{\sigma}_n^2(x)}{\sigma_n^2(x)} \xrightarrow[n \rightarrow \infty]{p} 1, \quad \text{or} \quad \frac{\hat{\sigma}_n(x)}{\sigma_n(x)} \xrightarrow[n \rightarrow \infty]{p} 1. \quad (48)$$

Since, by (44) and (45),

$$\begin{aligned} \frac{\hat{f}_n(x) - f(x)}{s.d.(\hat{f}_n(x))} &= \frac{\sqrt{nh_n}[\hat{f}_n(x) - f(x)]}{\sqrt{nh_n \text{Var}(\hat{f}_n(x))}} \\ &= \frac{\sqrt{nh_n}[\hat{f}_n(x) - f(x)]}{\sigma_n(x)} \xrightarrow[n \rightarrow \infty]{d} N(0, 1), \end{aligned}$$

it follows from this and (48) (by means of Theorem 6 in Chapter 7) that

$$\frac{\sqrt{nh_n}[\hat{f}_n(x) - f(x)]/\sigma_n(x)}{\hat{\sigma}_n(x)/\sigma_n(x)} = \frac{\sqrt{nh_n}[\hat{f}_n(x) - f(x)]}{\hat{\sigma}_n(x)} \xrightarrow[n \rightarrow \infty]{d} N(0, 1). \quad (49)$$

- (vi) Use convergence (49) in order to conclude that, for all sufficiently large n ,

$$P \left[\hat{f}_n(x) - \frac{\hat{\sigma}_n(x)}{\sqrt{nh_n}} z_{\alpha/2} \leq f(x) \leq \hat{f}_n(x) + \frac{\hat{\sigma}_n(x)}{\sqrt{nh_n}} z_{\alpha/2} \right] \simeq 1 - \alpha;$$

i.e., the interval $[\hat{f}_n(x) - \frac{\hat{\sigma}_n(x)}{\sqrt{nh_n}} z_{\alpha/2}, \hat{f}_n(x) + \frac{\hat{\sigma}_n(x)}{\sqrt{nh_n}} z_{\alpha/2}]$ is a confidence interval for $f(x)$ with confidence coefficient approximately $1 - \alpha$, for all sufficiently large n .

- 5.2** Refer to convergence (39), and set $\sigma_n(x) = s.d.(g_n(x; x_n))$. Use relation (39) in order to conclude that, for all sufficiently large n ,

$$P[g_n(x; x_n) - z_{\alpha/2}\sigma_n(x) \leq g(x) \leq g_n(x; x_n) + z_{\alpha/2}\sigma_n(x)] \simeq 1 - \alpha. \quad (50)$$

Thus, if $\sigma_n(x)$ is known, then expression (50) states that the interval $[g_n(x; x_n) - z_{\alpha/2}\sigma_n(x), g_n(x; x_n) + z_{\alpha/2}\sigma_n(x)]$ is a confidence interval for $g(x)$ with confidence coefficient approximately $1 - \alpha$, for all sufficiently large n . If $\sigma_n(x)$ is not known, but a suitable estimate of it, $\hat{\sigma}_n(x)$, can be constructed, then the interval $[g_n(x; x_n) - z_{\alpha/2}\hat{\sigma}_n(x), g_n(x; x_n) + z_{\alpha/2}\hat{\sigma}_n(x)]$ is

a confidence interval for $g(x)$ with confidence coefficient approximately $1 - \alpha$, for all sufficiently large n . One arrives at this conclusion working as in Exercise 5.1.

5.3 Refer to convergence (43), and go through the usual manipulations to conclude that, for all sufficiently large n ,

$$P \left[\hat{m}_n(x) - \frac{\sigma(x)}{\sqrt{nh_n}} \leq m(x) \leq \hat{m}_n(x) + \frac{\sigma(x)}{\sqrt{nh_n}} \right] \simeq 1 - \alpha. \quad (51)$$

Thus, if $\sigma(x)$ is known, then expression (51) states that the interval $[\hat{m}_n(x) - \frac{\sigma(x)}{\sqrt{nh_n}} z_{\alpha/2}, \hat{m}_n(x) + \frac{\sigma(x)}{\sqrt{nh_n}} z_{\alpha/2}]$ is a confidence interval for $m(x)$ with confidence coefficient approximately $1 - \alpha$, for all sufficiently large n . If $\sigma(x)$ is not known, but a suitable estimate of it, $\hat{\sigma}_n(x)$, can be constructed, then the interval $[\hat{m}_n(x) - \frac{\hat{\sigma}_n(x)}{\sqrt{nh_n}} z_{\alpha/2}, \hat{m}_n(x) + \frac{\hat{\sigma}_n(x)}{\sqrt{nh_n}} z_{\alpha/2}]$ is a confidence interval for $m(x)$ with confidence coefficient approximately $1 - \alpha$, for all sufficiently large n .

Appendix

Tables

Table 1

The Cumulative Binomial Distribution

The tabulated quantity is

$$\sum_{j=0}^k \binom{n}{j} p^j (1-p)^{n-j}.$$

<i>n</i>	<i>k</i>	<i>p</i>							
		1/16	2/16	3/16	4/16	5/16	6/16	7/16	8/16
2	0	0.8789	0.7656	0.6602	0.5625	0.4727	0.3906	0.3164	0.2500
	1	0.9961	0.9844	0.9648	0.9375	0.9023	0.8594	0.8086	0.7500
	2	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
3	0	0.8240	0.6699	0.5364	0.4219	0.3250	0.2441	0.1780	0.1250
	1	0.9888	0.9570	0.9077	0.8437	0.7681	0.6836	0.5933	0.5000
	2	0.9998	0.9980	0.9934	0.9844	0.9695	0.9473	0.9163	0.8750
	3	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
4	0	0.7725	0.5862	0.4358	0.3164	0.2234	0.1526	0.1001	0.0625
	1	0.9785	0.9211	0.8381	0.7383	0.6296	0.5188	0.4116	0.3125
	2	0.9991	0.9929	0.9773	0.9492	0.9065	0.8484	0.7749	0.6875
	3	1.0000	0.9998	0.9988	0.9961	0.9905	0.9802	0.9634	0.9375
	4	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
5	0	0.7242	0.5129	0.3541	0.2373	0.1536	0.0954	0.0563	0.0312
	1	0.9656	0.8793	0.7627	0.6328	0.5027	0.3815	0.2753	0.1875
	2	0.9978	0.9839	0.9512	0.8965	0.8200	0.7248	0.6160	0.5000
	3	0.9999	0.9989	0.9947	0.9844	0.9642	0.9308	0.8809	0.8125
	4	1.0000	1.0000	0.9998	0.9990	0.9970	0.9926	0.9840	0.9687
	5	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
6	0	0.6789	0.4488	0.2877	0.1780	0.1056	0.0596	0.0317	0.0156
	1	0.9505	0.8335	0.6861	0.5339	0.3936	0.2742	0.1795	0.1094
	2	0.9958	0.9709	0.9159	0.8306	0.7208	0.5960	0.4669	0.3437
	3	0.9998	0.9970	0.9866	0.9624	0.9192	0.8535	0.7650	0.6562
	4	1.0000	0.9998	0.9988	0.9954	0.9868	0.9694	0.9389	0.8906
	5	1.0000	1.0000	1.0000	0.9998	0.9991	0.9972	0.9930	0.9844
	6	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
7	0	0.6365	0.3927	0.2338	0.1335	0.0726	0.0373	0.0178	0.0078
	1	0.9335	0.7854	0.6114	0.4449	0.3036	0.1937	0.1148	0.0625
	2	0.9929	0.9537	0.8728	0.7564	0.6186	0.4753	0.3412	0.2266
	3	0.9995	0.9938	0.9733	0.9294	0.8572	0.7570	0.6346	0.5000
	4	1.0000	0.9995	0.9965	0.9871	0.9656	0.9260	0.8628	0.7734
	5	1.0000	1.0000	0.9997	0.9987	0.9952	0.9868	0.9693	0.9375

Table 1 (continued)

n	k	P							
		1/16	2/16	3/16	4/16	5/16	6/16	7/16	8/16
7	6	1.0000	1.0000	1.0000	0.9999	0.9997	0.9990	0.9969	0.9922
	7	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
8	0	0.5967	0.3436	0.1899	0.1001	0.0499	0.0233	0.0100	0.0039
	1	0.9150	0.7363	0.5406	0.3671	0.2314	0.1350	0.0724	0.0352
	2	0.9892	0.9327	0.8238	0.6785	0.5201	0.3697	0.2422	0.1445
	3	0.9991	0.9888	0.9545	0.8862	0.7826	0.6514	0.5062	0.3633
	4	1.0000	0.9988	0.9922	0.9727	0.9318	0.8626	0.7630	0.6367
	5	1.0000	0.9999	0.9991	0.9958	0.9860	0.9640	0.9227	0.8555
	6	1.0000	1.0000	0.9999	0.9996	0.9983	0.9944	0.9849	0.9648
	7	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9987	0.9961
9	0	0.5594	0.3007	0.1543	0.0751	0.0343	0.0146	0.0056	0.0020
	1	0.8951	0.6872	0.4748	0.3003	0.1747	0.0931	0.0451	0.0195
	2	0.9846	0.9081	0.7707	0.6007	0.4299	0.2817	0.1679	0.0898
	3	0.9985	0.9817	0.9300	0.8343	0.7006	0.5458	0.3907	0.2539
	4	0.9999	0.9975	0.9851	0.9511	0.8851	0.7834	0.6506	0.5000
	5	1.0000	0.9998	0.9978	0.9900	0.9690	0.9260	0.8528	0.7461
	6	1.0000	1.0000	0.9998	0.9987	0.9945	0.9830	0.9577	0.9102
	7	1.0000	1.0000	1.0000	0.9999	0.9994	0.9977	0.9926	0.9805
	8	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9994	0.9980
10	0	0.5245	0.2631	0.1254	0.0563	0.0236	0.0091	0.0032	0.0010
	1	0.8741	0.6389	0.4147	0.2440	0.1308	0.0637	0.0278	0.0107
	2	0.9790	0.8805	0.7152	0.5256	0.3501	0.2110	0.1142	0.0547
	3	0.9976	0.9725	0.9001	0.7759	0.6160	0.4467	0.2932	0.1719
	4	0.9998	0.9955	0.9748	0.9219	0.8275	0.6943	0.5369	0.3770
	5	1.0000	0.9995	0.9955	0.9803	0.9428	0.8725	0.7644	0.6230
	6	1.0000	1.0000	0.9994	0.9965	0.9865	0.9616	0.9118	0.8281
	7	1.0000	1.0000	1.0000	0.9996	0.9999	0.9922	0.9773	0.9453
	8	1.0000	1.0000	1.0000	1.0000	0.9998	0.9990	0.9964	0.9893
	9	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9990
11	0	0.4917	0.2302	0.1019	0.0422	0.0162	0.0057	0.0018	0.0005
	1	0.8522	0.5919	0.3605	0.1971	0.0973	0.0432	0.0170	0.0059
	2	0.9724	0.8503	0.6589	0.4552	0.2816	0.1558	0.0764	0.0327
	3	0.9965	0.9610	0.8654	0.7133	0.5329	0.3583	0.2149	0.1133
	4	0.9997	0.9927	0.9608	0.8854	0.7614	0.6014	0.4303	0.2744
	5	1.0000	0.9990	0.9916	0.9657	0.9068	0.8057	0.6649	0.5000
	6	1.0000	0.9999	0.9987	0.9924	0.9729	0.9282	0.8473	0.7256
	7	1.0000	1.0000	0.9999	0.9988	0.9943	0.9807	0.9487	0.8867
	8	1.0000	1.0000	1.0000	0.9999	0.9992	0.9965	0.9881	0.9673
	9	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9983	0.9941
	10	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995
12	0	0.4610	0.2014	0.0828	0.0317	0.0111	0.0036	0.0010	0.0002
	1	0.8297	0.5467	0.3120	0.1584	0.0720	0.0291	0.0104	0.0032
	2	0.9649	0.8180	0.6029	0.3907	0.2240	0.1135	0.0504	0.0193
	3	0.9950	0.9472	0.8267	0.6488	0.4544	0.2824	0.1543	0.0730

(Continued)

Table 1 (continued)

n	k	p								
		1/16	2/16	3/16	4/16	5/16	6/16	7/16	8/16	
12	4	0.9995	0.9887	0.9429	0.8424	0.6900	0.5103	0.3361	0.1938	
	5	1.0000	0.9982	0.9858	0.9456	0.8613	0.7291	0.5622	0.3872	
	6	1.0000	0.9998	0.9973	0.9857	0.9522	0.8822	0.7675	0.6128	
	7	1.0000	1.0000	0.9996	0.9972	0.9876	0.9610	0.9043	0.8062	
	8	1.0000	1.0000	1.0000	0.9996	0.9977	0.9905	0.9708	0.9270	
	9	1.0000	1.0000	1.0000	1.0000	0.9997	0.9984	0.9938	0.9807	
	10	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9992	0.9968	
	11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	
	12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
	13	0	0.4321	0.1762	0.0673	0.0238	0.0077	0.0022	0.0006	0.0001
		1	0.8067	0.5035	0.2690	0.1267	0.0530	0.0195	0.0063	0.0017
		2	0.9565	0.7841	0.5484	0.3326	0.1765	0.0819	0.0329	0.0112
3		0.9931	0.9310	0.7847	0.5843	0.3824	0.2191	0.1089	0.0461	
4		0.9992	0.9835	0.9211	0.7940	0.6164	0.4248	0.2565	0.1334	
5		0.9999	0.9970	0.9778	0.9198	0.8078	0.6470	0.4633	0.2905	
6		1.0000	0.9996	0.9952	0.9757	0.9238	0.8248	0.6777	0.5000	
7		1.0000	1.0000	0.9992	0.9944	0.9765	0.9315	0.8445	0.7095	
8		1.0000	1.0000	0.9999	0.9990	0.9945	0.9795	0.9417	0.8666	
9		1.0000	1.0000	1.0000	0.9999	0.9991	0.9955	0.9838	0.9539	
10		1.0000	1.0000	1.0000	1.0000	0.9999	0.9993	0.9968	0.9888	
11		1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9983	
12		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
14	0	0.4051	0.1542	0.0546	0.0178	0.0053	0.0014	0.0003	0.0001	
	1	0.7833	0.4626	0.2312	0.1010	0.0388	0.0130	0.0038	0.0009	
	2	0.9471	0.7490	0.4960	0.2811	0.1379	0.0585	0.0213	0.0065	
	3	0.9908	0.9127	0.7404	0.5213	0.3181	0.1676	0.0756	0.0287	
	4	0.9988	0.9970	0.8955	0.7415	0.5432	0.3477	0.1919	0.0898	
	5	0.9999	0.9953	0.9671	0.8883	0.7480	0.5637	0.3728	0.2120	
	6	1.0000	0.9993	0.9919	0.9167	0.8876	0.7581	0.5839	0.3953	
	7	1.0000	0.9999	0.9985	0.9897	0.9601	0.8915	0.7715	0.6047	
	8	1.0000	1.0000	0.9998	0.9978	0.9889	0.9615	0.8992	0.7880	
	9	1.0000	1.0000	1.0000	0.9997	0.9976	0.9895	0.9654	0.9102	
	10	1.0000	1.0000	1.0000	1.0000	0.9996	0.9979	0.9911	0.9713	
	11	1.0000	1.0000	1.0000	1.0000	1.0000	0.9997	0.9984	0.9935	
	12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9991	
	13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	0	0.3798	0.1349	0.0444	0.0134	0.0036	0.0009	0.0002	0.0000	
	1	0.7596	0.4241	0.1981	0.0802	0.0283	0.0087	0.0023	0.0005	
	2	0.9369	0.7132	0.4463	0.2361	0.1069	0.0415	0.0136	0.0037	
	3	0.9881	0.8922	0.6946	0.4613	0.2618	0.1267	0.0518	0.0176	
	4	0.9983	0.9689	0.8665	0.6865	0.4729	0.2801	0.1410	0.0592	
	5	0.9998	0.9930	0.9537	0.8516	0.6840	0.4827	0.2937	0.1509	
	6	1.0000	0.9988	0.9873	0.9434	0.8435	0.6852	0.4916	0.3036	
	7	1.0000	0.9998	0.9972	0.9827	0.9374	0.8415	0.6894	0.5000	
	8	1.0000	1.0000	0.9995	0.9958	0.9799	0.9352	0.8433	0.6964	
	9	1.0000	1.0000	0.9999	0.9992	0.9949	0.9790	0.9364	0.8491	
	10	1.0000	1.0000	1.0000	0.9999	0.9990	0.9947	0.9799	0.9408	
	11	1.0000	1.0000	1.0000	1.0000	0.9999	0.9990	0.9952	0.9824	
12	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9992	0.9963		

Table 1 (continued)

n	k	p							
		1/16	2/16	3/16	4/16	5/16	6/16	7/16	8/16
15	13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995
	14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
	15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
16	0	0.3561	0.1181	0.0361	0.0100	0.0025	0.0005	0.0001	0.0000
	1	0.7359	0.3879	0.1693	0.0635	0.0206	0.0057	0.0014	0.0003
	2	0.9258	0.6771	0.3998	0.1971	0.0824	0.0292	0.0086	0.0021
	3	0.9849	0.8698	0.6480	0.4050	0.2134	0.0947	0.0351	0.0106
	4	0.9977	0.9593	0.8342	0.6302	0.4069	0.2226	0.1020	0.0384
	5	0.9997	0.9900	0.9373	0.8103	0.6180	0.4067	0.2269	0.1051
	6	1.0000	0.9981	0.9810	0.9204	0.7940	0.6093	0.4050	0.2272
	7	1.0000	0.9997	0.9954	0.9729	0.9082	0.7829	0.6029	0.4018
	8	1.0000	1.0000	0.9991	0.9925	0.9666	0.9001	0.7760	0.5982
	9	1.0000	1.0000	0.9999	0.9984	0.9902	0.9626	0.8957	0.7728
	10	1.0000	1.0000	1.0000	0.9997	0.9977	0.9888	0.9609	0.8949
	11	1.0000	1.0000	1.0000	1.0000	0.9996	0.9974	0.9885	0.9616
	12	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9975	0.9894
	13	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9979
	14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9997
	15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
16	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
17	0	0.3338	0.1033	0.0293	0.0075	0.0017	0.0003	0.0001	0.0000
	1	0.7121	0.3542	0.1443	0.0501	0.0149	0.0038	0.0008	0.0001
	2	0.9139	0.6409	0.3566	0.1637	0.0631	0.0204	0.0055	0.0012
	3	0.9812	0.8457	0.6015	0.3530	0.1724	0.0701	0.0235	0.0064
	4	0.9969	0.9482	0.7993	0.5739	0.3464	0.1747	0.0727	0.0245
	5	0.9996	0.9862	0.9180	0.7653	0.5520	0.3377	0.1723	0.0717
	6	1.0000	0.9971	0.9728	0.8929	0.7390	0.5333	0.3271	0.1662
	7	1.0000	0.9995	0.9927	0.9598	0.8725	0.7178	0.5163	0.3145
	8	1.0000	0.9999	0.9984	0.9876	0.9484	0.8561	0.7002	0.5000
	9	1.0000	1.0000	0.9997	0.9969	0.9828	0.9391	0.8433	0.6855
	10	1.0000	1.0000	1.0000	0.9994	0.9954	0.9790	0.9323	0.8338
	11	1.0000	1.0000	1.0000	0.9999	0.9990	0.9942	0.9764	0.9283
	12	1.0000	1.0000	1.0000	1.0000	0.9998	0.9987	0.9935	0.9755
	13	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9987	0.9936
	14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9988
	15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999
16	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
18	0	0.3130	0.0904	0.0238	0.0056	0.0012	0.0002	0.0000	0.0000
	1	0.6885	0.3228	0.1227	0.0395	0.0108	0.0025	0.0005	0.0001
	2	0.9013	0.6051	0.3168	0.1353	0.0480	0.0142	0.0034	0.0007
	3	0.9770	0.8201	0.5556	0.3057	0.1383	0.0515	0.0156	0.0038
	4	0.9959	0.9354	0.7622	0.5187	0.2920	0.1355	0.0512	0.0154
	5	0.9994	0.9814	0.8958	0.7175	0.4878	0.2765	0.1287	0.0481
	6	0.9999	0.9957	0.9625	0.8610	0.6806	0.4600	0.2593	0.1189
	7	1.0000	0.9992	0.9889	0.9431	0.8308	0.6486	0.4335	0.2403
	8	1.0000	0.9999	0.9973	0.9807	0.9247	0.8042	0.6198	0.4073
	9	1.0000	1.0000	0.9995	0.9946	0.9721	0.9080	0.7807	0.5927
	10	1.0000	1.0000	0.9999	0.9988	0.9915	0.9640	0.8934	0.7597
11	1.0000	1.0000	1.0000	0.9998	0.9979	0.9885	0.9571	0.8811	

(Continued)

Table 1 (continued)

n	k	p							
		1/16	2/16	3/16	4/16	5/16	6/16	7/16	8/16
18	12	1.0000	1.0000	1.0000	1.0000	0.9996	0.9970	0.9860	0.9519
	13	1.0000	1.0000	1.0000	1.0000	0.9999	0.9994	0.9964	0.9846
	14	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9993	0.9962
	15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9993
	16	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999
	17	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
19	0	0.2934	0.0791	0.0193	0.0042	0.0008	0.0001	0.0000	0.0000
	1	0.6650	0.2938	0.1042	0.0310	0.0078	0.0016	0.0003	0.0000
	2	0.8880	0.5698	0.2804	0.1113	0.0364	0.0098	0.0021	0.0004
	3	0.9722	0.7933	0.5108	0.2631	0.1101	0.0375	0.0103	0.0022
	4	0.9947	0.9209	0.7235	0.4654	0.2440	0.1040	0.0356	0.0096
	5	0.9992	0.9757	0.8707	0.6678	0.4266	0.2236	0.0948	0.0318
	6	0.9999	0.9939	0.9500	0.8251	0.6203	0.3912	0.2022	0.0835
	7	1.0000	0.9988	0.9840	0.9225	0.7838	0.5779	0.3573	0.1796
	8	1.0000	0.9998	0.9957	0.9713	0.8953	0.7459	0.5383	0.3238
	9	1.0000	1.0000	0.9991	0.9911	0.9573	0.8691	0.7103	0.5000
	10	1.0000	1.0000	0.9998	0.9977	0.9854	0.9430	0.8441	0.0672
	11	1.0000	1.0000	1.0000	0.9995	0.9959	0.9793	0.9292	0.8204
	12	1.0000	1.0000	1.0000	0.9999	0.9990	0.9938	0.9734	0.9165
	13	1.0000	1.0000	1.0000	1.0000	0.9998	0.9985	0.9919	0.9682
14	1.0000	1.0000	1.0000	1.0000	1.0000	0.9997	0.9980	0.9904	
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9996	0.9978	
16	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9996	
17	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
18	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
20	0	0.2751	0.0692	0.0157	0.0032	0.0006	0.0001	0.0000	0.0000
	1	0.6148	0.2669	0.0883	0.0243	0.0056	0.0011	0.0002	0.0000
	2	0.8741	0.5353	0.2473	0.0913	0.0275	0.0067	0.0013	0.0002
	3	0.9670	0.7653	0.4676	0.2252	0.0870	0.0271	0.0067	0.0013
	4	0.9933	0.9050	0.6836	0.4148	0.2021	0.0790	0.0245	0.0059
	5	0.9989	0.9688	0.8431	0.6172	0.3695	0.1788	0.0689	0.0207
	6	0.9999	0.9916	0.9351	0.7858	0.5598	0.3284	0.1552	0.0577
	7	1.0000	0.9981	0.9776	0.8982	0.7327	0.5079	0.2894	0.1316
	8	1.0000	0.9997	0.9935	0.9591	0.8605	0.6829	0.4591	0.2517
	9	1.0000	0.9999	0.9984	0.9861	0.9379	0.8229	0.6350	0.4119
	10	1.0000	1.0000	0.9997	0.9961	0.9766	0.9153	0.7856	0.5881
	11	1.0000	1.0000	0.9999	0.9991	0.9926	0.9657	0.8920	0.7483
	12	1.0000	1.0000	1.0000	0.9998	0.9981	0.9884	0.9541	0.8684
	13	1.0000	1.0000	1.0000	1.0000	0.9996	0.9968	0.9838	0.9423
14	1.0000	1.0000	1.0000	1.0000	0.9999	0.9993	0.9953	0.9793	
15	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9989	0.9941	
16	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9987	
17	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	
18	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
21	0	0.2579	0.0606	0.0128	0.0024	0.0004	0.0001	0.0000	0.0000
	1	0.6189	0.2422	0.0747	0.0190	0.0040	0.0007	0.0001	0.0000
	2	0.8596	0.5018	0.2175	0.0745	0.0206	0.0046	0.0008	0.0001
	3	0.9612	0.7366	0.4263	0.1917	0.0684	0.0195	0.0044	0.0007
	4	0.9917	0.8875	0.6431	0.3674	0.1662	0.0596	0.0167	0.0036

Table 1 (continued)

<i>n</i>	<i>k</i>	<i>P</i>								
		1/16	2/16	3/16	4/16	5/16	6/16	7/16	8/16	
21	5	0.9986	0.9609	0.8132	0.5666	0.3172	0.1414	0.0495	0.0133	
	6	0.9998	0.9888	0.9179	0.7436	0.5003	0.2723	0.1175	0.0392	
	7	1.0000	0.9973	0.9696	0.8701	0.6787	0.4405	0.2307	0.0946	
	8	1.0000	0.9995	0.9906	0.9439	0.8206	0.6172	0.3849	0.1917	
	9	1.0000	0.9999	0.9975	0.9794	0.9137	0.7704	0.5581	0.3318	
	10	1.0000	1.0000	0.9995	0.9936	0.9645	0.8806	0.7197	0.5000	
	11	1.0000	1.0000	0.9999	0.9983	0.9876	0.9468	0.8454	0.6682	
	12	1.0000	1.0000	1.0000	0.9996	0.9964	0.9799	0.9269	0.8083	
	13	1.0000	1.0000	1.0000	0.9999	0.9991	0.9936	0.9708	0.9054	
	14	1.0000	1.0000	1.0000	1.0000	0.9998	0.9983	0.9903	0.9605	
	15	1.0000	1.0000	1.0000	1.0000	1.0000	0.9996	0.9974	0.9867	
	16	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9994	0.9964	
	17	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9993	
	18	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	
	19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
	20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
	22	0	0.2418	0.0530	0.0104	0.0018	0.0003	0.0000	0.0000	0.0000
		1	0.5963	0.2195	0.0631	0.0149	0.0029	0.0005	0.0001	0.0000
		2	0.8445	0.4693	0.1907	0.0606	0.0154	0.0031	0.0005	0.0001
		3	0.9548	0.7072	0.3871	0.1624	0.0535	0.0139	0.0028	0.0004
4		0.9898	0.8687	0.6024	0.3235	0.1356	0.0445	0.0133	0.0022	
5		0.9981	0.9517	0.7813	0.5168	0.2700	0.1107	0.0352	0.0085	
6		0.9997	0.9853	0.8983	0.6994	0.4431	0.2232	0.0877	0.0267	
7		1.0000	0.9963	0.9599	0.8385	0.6230	0.3774	0.1812	0.0669	
8		1.0000	0.9992	0.9866	0.9254	0.7762	0.5510	0.3174	0.1431	
9		1.0000	0.9999	0.9962	0.9705	0.8846	0.7130	0.4823	0.2617	
10		1.0000	1.0000	0.9991	0.9900	0.9486	0.8393	0.6490	0.4159	
11		1.0000	1.0000	0.9998	0.9971	0.9804	0.9220	0.7904	0.5841	
12		1.0000	1.0000	1.0000	0.9993	0.9936	0.9675	0.8913	0.7383	
13		1.0000	1.0000	1.0000	0.9999	0.9982	0.9885	0.9516	0.8569	
14		1.0000	1.0000	1.0000	1.0000	0.9996	0.9966	0.9818	0.9331	
15		1.0000	1.0000	1.0000	1.0000	0.9999	0.9991	0.9943	0.9739	
16		1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9985	0.9915	
17		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9997	0.9978	
18		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9995	
19		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	
20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
23	0	0.2266	0.0464	0.0084	0.0013	0.0002	0.0000	0.0000	0.0000	
	1	0.5742	0.1987	0.0532	0.0116	0.0021	0.0003	0.0000	0.0000	
	2	0.8290	0.4381	0.1668	0.0492	0.0115	0.0021	0.0003	0.0000	
	3	0.9479	0.6775	0.3503	0.1370	0.0416	0.0099	0.0018	0.0002	
	4	0.9876	0.8485	0.5621	0.2832	0.1100	0.0330	0.0076	0.0013	
	5	0.9976	0.9413	0.7478	0.4685	0.2280	0.0859	0.0247	0.0053	
	6	0.9996	0.9811	0.8763	0.6537	0.3890	0.1810	0.0647	0.0173	
	7	1.0000	0.9949	0.9484	0.8037	0.5668	0.3196	0.1403	0.0466	
	8	1.0000	0.9988	0.9816	0.9037	0.7283	0.4859	0.2578	0.1050	
	9	1.0000	0.9998	0.9944	0.9592	0.8507	0.6522	0.4102	0.2024	
	10	1.0000	1.0000	0.9986	0.9851	0.9286	0.7919	0.5761	0.3388	
11	1.0000	1.0000	0.9997	0.9954	0.9705	0.8910	0.7285	0.5000		

(Continued)

Table 1 (continued)

n	k	p							
		1/16	2/16	3/16	4/16	5/16	6/16	7/16	8/16
23	12	1.0000	1.0000	0.9999	0.9988	0.9895	0.9504	0.8471	0.6612
	13	1.0000	1.0000	1.0000	0.9997	0.9968	0.9806	0.9252	0.7976
	14	1.0000	1.0000	1.0000	0.9999	0.9992	0.9935	0.9686	0.8950
	15	1.0000	1.0000	1.0000	1.0000	0.9998	0.9982	0.9888	0.9534
	16	1.0000	1.0000	1.0000	1.0000	1.0000	0.9996	0.9967	0.9827
	17	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9992	0.9947
	18	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9987
	19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998
	20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
	21	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
	24	0	0.2125	0.0406	0.0069	0.0010	0.0001	0.0000	0.0000
1		0.5524	0.1797	0.0448	0.0090	0.0015	0.0002	0.0000	0.0000
2		0.8131	0.4082	0.1455	0.0398	0.0086	0.0014	0.0002	0.0000
3		0.9405	0.6476	0.3159	0.1150	0.0322	0.0070	0.0011	0.0001
4		0.9851	0.8271	0.5224	0.2466	0.0886	0.0243	0.0051	0.0008
5		0.9970	0.9297	0.7130	0.4222	0.1911	0.0661	0.0172	0.0033
6		0.9995	0.9761	0.8522	0.6074	0.3387	0.1453	0.0472	0.0113
7		0.9999	0.9932	0.9349	0.7662	0.5112	0.2676	0.1072	0.0320
8		1.0000	0.9983	0.9754	0.8787	0.6778	0.4235	0.2064	0.0758
9		1.0000	0.9997	0.9920	0.9453	0.8125	0.5898	0.3435	0.1537
10		1.0000	0.9999	0.9978	0.9787	0.9043	0.7395	0.5035	0.2706
11		1.0000	1.0000	0.9995	0.9928	0.9574	0.8538	0.6618	0.4194
12		1.0000	1.0000	0.9999	0.9979	0.9835	0.9281	0.7953	0.5806
13		1.0000	1.0000	1.0000	0.9995	0.9945	0.9693	0.8911	0.7294
14	1.0000	1.0000	1.0000	0.9999	0.9984	0.9887	0.9496	0.8463	
15	1.0000	1.0000	1.0000	1.0000	0.9996	0.9964	0.9799	0.9242	
16	1.0000	1.0000	1.0000	1.0000	0.9999	0.9990	0.9932	0.9680	
17	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9981	0.9887	
18	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9996	0.9967	
19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9992	
20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	
21	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
22	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
25	0	0.1992	0.0355	0.0056	0.0008	0.0001	0.0000	0.0000	0.0000
	1	0.5132	0.1623	0.0377	0.0070	0.0011	0.0001	0.0000	0.0000
	2	0.7968	0.3796	0.1266	0.0321	0.0064	0.0010	0.0001	0.0000
	3	0.9325	0.6176	0.2840	0.0962	0.0248	0.0049	0.0007	0.0001
	4	0.9823	0.8047	0.4837	0.2137	0.0710	0.0178	0.0033	0.0005
	5	0.9962	0.9169	0.6772	0.3783	0.1591	0.0504	0.0119	0.0028
	6	0.9993	0.9703	0.8261	0.5611	0.2926	0.1156	0.0341	0.0073
	7	0.9999	0.9910	0.9194	0.7265	0.4573	0.2218	0.0810	0.0216
	8	1.0000	0.9977	0.9678	0.8506	0.6258	0.3651	0.1630	0.0539
	9	1.0000	0.9995	0.9889	0.9287	0.7704	0.5275	0.2835	0.1148
	10	1.0000	0.9999	0.9967	0.9703	0.8756	0.6834	0.4335	0.2122
	11	1.0000	1.0000	0.9992	0.9893	0.9408	0.8110	0.5926	0.3450
	12	1.0000	1.0000	0.9998	0.9966	0.9754	0.9003	0.7369	0.5000
	13	1.0000	1.0000	1.0000	0.9991	0.9911	0.9538	0.8491	0.6550
	14	1.0000	1.0000	1.0000	0.9998	0.9972	0.9814	0.9240	0.7878
	15	1.0000	1.0000	1.0000	1.0000	0.9992	0.9935	0.9667	0.8852
	16	1.0000	1.0000	1.0000	1.0000	0.9998	0.9981	0.9874	0.9462
17	1.0000	1.0000	1.0000	1.0000	1.0000	0.9995	0.9960	0.9784	

Table 2

The Cumulative Poisson Distribution

The tabulated quantity is

$$\sum_{j=0}^k e^{-\lambda} \frac{\lambda^j}{j!}.$$

		λ					
k	0.001	0.005	0.010	0.015	0.020	0.025	
0	0.9990 0050	0.9950 1248	0.9900 4983	0.9851 1194	0.9801 9867	0.9753 099	
1	0.9999 9950	0.9999 8754	0.9999 5033	0.9998 8862	0.9998 0264	0.9996 927	
2	1.0000 0000	0.9999 9998	0.9999 9983	0.9999 9945	0.9999 9868	0.9999 974	
3		1.0000 0000	1.0000 0000	1.0000 0000	0.9999 9999	1.0000 000	
4					1.0000 0000	1.0000 000	
		λ					
k	0.030	0.035	0.040	0.045	0.050	0.055	
0	0.970 446	0.965 605	0.960 789	0.955 997	0.951 229	0.946 485	
1	0.999 559	0.999 402	0.999 221	0.999 017	0.998 791	0.998 542	
2	0.999 996	0.999 993	0.999 990	0.999 985	0.999 980	0.999 973	
3	1.000 000	1.000 000	1.000 000	1.000 000	1.000 000	1.000 000	
		λ					
k	0.060	0.065	0.070	0.075	0.080	0.085	
0	0.941 765	0.937 067	0.932 394	0.927 743	0.923 116	0.918 512	
1	0.998 270	0.997 977	0.997 661	0.997 324	0.996 966	0.996 586	
2	0.999 966	0.999 956	0.999 946	0.999 934	0.999 920	0.999 904	
3	0.999 999	0.999 999	0.999 999	0.999 999	0.999 998	0.999 998	
4	1.000 000	1.000 000	1.000 000	1.000 000	1.000 000	1.000 000	
		λ					
k	0.090	0.095	0.100	0.200	0.300	0.400	
0	0.913 931	0.909 373	0.904 837	0.818 731	0.740 818	0.670 320	
1	0.996 185	0.995 763	0.995 321	0.982 477	0.963 064	0.938 448	
2	0.999 886	0.999 867	0.999 845	0.998 852	0.996 401	0.992 074	
3	0.999 997	0.999 997	0.999 996	0.999 943	0.999 734	0.999 224	
4	1.000 000	1.000 000	1.000 000	0.999 998	0.999 984	0.999 939	
5				1.000 000	0.999 999	0.999 996	
6					1.000 000	1.000 000	
		λ					
k	0.500	0.600	0.700	0.800	0.900	1.000	
0	0.606 531	0.548 812	0.496 585	0.449 329	0.406 329	0.367 879	
1	0.909 796	0.878 099	0.844 195	0.808 792	0.772 482	0.735 759	
2	0.985 612	0.976 885	0.965 858	0.952 577	0.937 143	0.919 699	
3	0.998 248	0.996 642	0.994 247	0.990 920	0.986 541	0.981 012	
4	0.999 828	0.999 606	0.999 214	0.998 589	0.997 656	0.996 340	
5	0.999 986	0.999 961	0.999 910	0.999 816	0.999 657	0.999 406	
6	0.999 999	0.999 997	0.999 991	0.999 979	0.999 957	0.999 917	
7	1.000 000	1.000 000	0.999 999	0.999 998	0.999 995	0.999 990	
8			1.000 000	1.000 000	1.000 000	0.999 999	
9						1.000 000	

Table 3

The Normal Distribution

The tabulated quantity is

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt.$$

$$[\Phi(-x) = 1 - \Phi(x)].$$

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0.00	0.500000	0.45	0.673645	0.90	0.815940	1.35	0.911492
0.01	0.503989	0.46	0.677242	0.91	0.818589	1.36	0.913085
0.02	0.507978	0.47	0.680822	0.92	0.821214	1.37	0.914657
0.03	0.511966	0.48	0.684386	0.93	0.823814	1.38	0.916207
0.04	0.515953	0.49	0.687933	0.94	0.826391	1.39	0.917736
0.05	0.519939	0.50	0.691462	0.95	0.828944	1.40	0.919243
0.06	0.523922	0.51	0.694974	0.96	0.831472	1.41	0.920730
0.07	0.527903	0.52	0.698468	0.97	0.833977	1.42	0.922196
0.08	0.531881	0.53	0.701944	0.98	0.836457	1.43	0.923641
0.09	0.535856	0.54	0.705401	0.99	0.838913	1.44	0.925066
0.10	0.539828	0.55	0.708840	1.00	0.841345	1.45	0.926471
0.11	0.543795	0.56	0.712260	1.01	0.843752	1.46	0.927855
0.12	0.547758	0.57	0.715661	1.02	0.846136	1.47	0.929219
0.13	0.551717	0.58	0.719043	1.03	0.848495	1.48	0.930563
0.14	0.555670	0.59	0.722405	1.04	0.850830	1.49	0.931888
0.15	0.559618	0.60	0.725747	1.05	0.853141	1.50	0.933193
0.16	0.563559	0.61	0.729069	1.06	0.855428	1.51	0.934478
0.17	0.567495	0.62	0.732371	1.07	0.857690	1.52	0.935745
0.18	0.571424	0.63	0.735653	1.08	0.859929	1.53	0.936992
0.19	0.575345	0.64	0.738914	1.09	0.862143	1.54	0.938220
0.20	0.579260	0.65	0.742154	1.10	0.864334	1.55	0.939429
0.21	0.583166	0.66	0.745373	1.11	0.866500	1.56	0.940620
0.22	0.587064	0.67	0.748571	1.12	0.868643	1.57	0.941792
0.23	0.590954	0.68	0.751748	1.13	0.870762	1.58	0.942947
0.24	0.594835	0.69	0.754903	1.14	0.872857	1.59	0.944083
0.25	0.598706	0.70	0.758036	1.15	0.874928	1.60	0.945201
0.26	0.602568	0.71	0.761148	1.16	0.876976	1.61	0.946301
0.27	0.606420	0.72	0.764238	1.17	0.879000	1.62	0.947384
0.28	0.610261	0.73	0.767305	1.18	0.881000	1.63	0.948449
0.29	0.614092	0.74	0.770350	1.19	0.882977	1.64	0.949497
0.30	0.617911	0.75	0.773373	1.20	0.884930	1.65	0.950529
0.31	0.621720	0.76	0.776373	1.21	0.886861	1.66	0.951543
0.32	0.625516	0.77	0.779350	1.22	0.888768	1.67	0.952540
0.33	0.629300	0.78	0.782305	1.23	0.890651	1.68	0.953521
0.34	0.633072	0.79	0.785236	1.24	0.892512	1.69	0.954486
0.35	0.636831	0.80	0.788145	1.25	0.894350	1.70	0.955435
0.36	0.640576	0.81	0.791030	1.26	0.896165	1.71	0.956367
0.37	0.644309	0.82	0.793892	1.27	0.897958	1.72	0.957284
0.38	0.648027	0.83	0.796731	1.28	0.899727	1.73	0.958185
0.39	0.651732	0.84	0.799546	1.29	0.901475	1.74	0.959070
0.40	0.655422	0.85	0.802337	1.30	0.903200	1.75	0.959941
0.41	0.659097	0.86	0.805105	1.31	0.904902	1.76	0.960796
0.42	0.662757	0.87	0.807850	1.32	0.906582	1.77	0.961636
0.43	0.666402	0.88	0.810570	1.33	0.908241	1.78	0.962462
0.44	0.670031	0.89	0.813267	1.34	0.909877	1.79	0.963273

Table 3 (continued)

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
1.80	0.964070	2.30	0.989276	2.80	0.997445	3.30	0.999517
1.81	0.964852	2.31	0.989556	2.81	0.997523	3.31	0.999534
1.82	0.965620	2.32	0.989830	2.82	0.997599	3.32	0.999550
1.83	0.966375	2.33	0.990097	2.83	0.997673	3.33	0.999566
1.84	0.967116	2.34	0.990358	2.84	0.997744	3.34	0.999581
1.85	0.967843	2.35	0.990613	2.85	0.997814	3.35	0.999596
1.86	0.968557	2.36	0.990863	2.86	0.997882	3.36	0.999610
1.87	0.969258	2.37	0.991106	2.87	0.997948	3.37	0.999624
1.88	0.969946	2.38	0.991344	2.88	0.998012	3.38	0.999638
1.89	0.970621	2.39	0.991576	2.89	0.998074	3.39	0.999651
1.90	0.971283	2.40	0.991802	2.90	0.998134	3.40	0.999663
1.91	0.971933	2.41	0.992024	2.91	0.998193	3.41	0.999675
1.92	0.972571	2.42	0.992240	2.92	0.998250	3.42	0.999687
1.93	0.973197	2.43	0.992451	2.93	0.998305	3.43	0.999698
1.94	0.973810	2.44	0.992656	2.94	0.998359	3.44	0.999709
1.95	0.974412	2.45	0.992857	2.95	0.998411	3.45	0.999720
1.96	0.975002	2.46	0.993053	2.96	0.998462	3.46	0.999730
1.97	0.975581	2.47	0.993244	2.97	0.998511	3.47	0.999740
1.98	0.976148	2.48	0.993431	2.98	0.998559	3.48	0.999749
1.99	0.976705	2.49	0.993613	2.99	0.998605	3.49	0.999758
2.00	0.977250	2.50	0.993790	3.00	0.998650	3.50	0.999767
2.01	0.977784	2.51	0.993963	3.01	0.998694	3.51	0.999776
2.02	0.978308	2.52	0.994132	3.02	0.998736	3.52	0.999784
2.03	0.978822	2.53	0.994297	3.03	0.998777	3.53	0.999792
2.04	0.979325	2.54	0.994457	3.04	0.998817	3.54	0.999800
2.05	0.979818	2.55	0.994614	3.05	0.998856	3.55	0.999807
2.06	0.980301	2.56	0.994766	3.06	0.998893	3.56	0.999815
2.07	0.980774	2.57	0.994915	3.07	0.998930	3.57	0.999822
2.08	0.981237	2.58	0.995060	3.08	0.998965	3.58	0.999828
2.09	0.981691	2.59	0.995201	3.09	0.998999	3.59	0.999835
2.10	0.982136	2.60	0.995339	3.10	0.999032	3.60	0.999841
2.11	0.982571	2.61	0.995473	3.11	0.999065	3.61	0.999847
2.12	0.982997	2.62	0.995604	3.12	0.999096	3.62	0.999853
2.13	0.983414	2.63	0.995731	3.13	0.999126	3.63	0.999858
2.14	0.983823	2.64	0.995855	3.14	0.999155	3.64	0.999864
2.15	0.984222	2.65	0.995975	3.15	0.999184	3.65	0.999869
2.16	0.984614	2.66	0.996093	3.16	0.999211	3.66	0.999874
2.17	0.984997	2.67	0.996207	3.17	0.999238	3.67	0.999879
2.18	0.985371	2.68	0.996319	3.18	0.999264	3.68	0.999883
2.19	0.985738	2.69	0.996427	3.19	0.999289	3.69	0.999888
2.20	0.986097	2.70	0.996533	3.20	0.999313	3.70	0.999892
2.21	0.986447	2.71	0.996636	3.21	0.999336	3.71	0.999896
2.22	0.986791	2.72	0.996736	3.22	0.999359	3.72	0.999900
2.23	0.987126	2.73	0.996833	3.23	0.999381	3.73	0.999904
2.24	0.987455	2.74	0.996928	3.24	0.999402	3.74	0.999908
2.25	0.987776	2.75	0.997020	3.25	0.999423	3.75	0.999912
2.26	0.988089	2.76	0.997110	3.26	0.999443	3.76	0.999915
2.27	0.988396	2.77	0.997197	3.27	0.999462	3.77	0.999918
2.28	0.988696	2.78	0.997282	3.28	0.999481	3.78	0.999922
2.29	0.988989	2.79	0.997365	3.29	0.999499	3.79	0.999925

(Continued)

Table 3 (*continued*)

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
3.80	0.999928	3.85	0.999941	3.90	0.999952	3.95	0.999961
3.81	0.999931	3.86	0.999943	3.91	0.999954	3.96	0.999963
3.82	0.999933	3.87	0.999946	3.92	0.999956	3.97	0.999964
3.83	0.999936	3.88	0.999948	3.93	0.999958	3.98	0.999966
3.84	0.999938	3.89	0.999950	3.94	0.999959	3.99	0.999967

Table 4

Let t_r be a random variable having the Student's t -distribution with r degrees of freedom. Then the tabulated quantities are the numbers x for which

$$P(t_r \leq x) = \gamma.$$

**Critical Values for
Student's t -Distribution**

r	0.75	0.90	0.95	γ	0.975	0.99	0.995
1	1.0000	3.0777	6.3138		12.7062	31.8207	63.6574
2	0.8165	1.8856	2.9200		4.3027	6.9646	9.9248
3	0.7649	1.6377	2.3534		3.1824	4.5407	5.8409
4	0.7407	1.5332	2.1318		2.7764	3.7649	4.6041
5	0.7267	1.4759	2.0150		2.5706	3.3649	4.0322
6	0.7176	1.4398	1.9432		2.4469	3.1427	3.7074
7	0.7111	1.4149	1.8946		2.3646	2.9980	3.4995
8	0.7064	1.3968	1.8595		2.3060	2.8965	3.3554
9	0.7027	1.3830	1.8331		2.2622	2.8214	3.2498
10	0.6998	1.3722	1.8125		2.2281	2.7638	3.1693
11	0.6974	1.3634	1.7959		2.2010	2.7181	3.1058
12	0.6955	1.3562	1.7823		2.1788	2.6810	3.0545
13	0.6938	1.3502	1.7709		2.1604	2.6503	3.0123
14	0.6924	1.3450	1.7613		2.1448	2.6245	2.9768
15	0.6912	1.3406	1.7531		2.1315	2.6025	2.9467
16	0.6901	1.3368	1.7459		2.1199	2.5835	2.9208
17	0.6892	1.3334	1.7396		2.1098	2.5669	2.8982
18	0.6884	1.3304	1.7341		2.1009	2.5524	2.8784
19	0.6876	1.3277	1.7291		2.0930	2.5395	2.8609
20	0.6870	1.3253	1.7247		2.0860	2.5280	2.8453
21	0.6864	1.3232	1.7207		2.0796	2.5177	2.8314
22	0.6858	1.3212	1.7171		2.0739	2.5083	2.8188
23	0.6853	1.3195	1.7139		2.0687	2.4999	2.8073
24	0.6848	1.3178	1.7109		2.0639	2.4922	2.7969
25	0.6844	1.3163	1.7081		2.0595	2.4851	2.7874
26	0.6840	1.3150	1.7056		2.0555	2.4786	2.7787
27	0.6837	1.3137	1.7033		2.0518	2.4727	2.7707
28	0.6834	1.3125	1.7011		2.0484	2.4671	2.7633
29	0.6830	1.3114	1.6991		2.0452	2.4620	2.7564
30	0.6828	1.3104	1.6973		2.0423	2.4573	2.7500
31	0.6825	1.3095	1.6955		2.0395	2.4528	2.7440
32	0.6822	1.3086	1.6939		2.0369	2.4487	2.7385
33	0.6820	1.3077	1.6924		2.0345	2.4448	2.7333
34	0.6818	1.3070	1.6909		2.0322	2.4411	2.7284
35	0.6816	1.3062	1.6896		2.0301	2.4377	2.7238
36	0.6814	1.3055	1.6883		2.0281	2.4345	2.7195
37	0.6812	1.3049	1.6871		2.0262	2.4314	2.7154
38	0.6810	1.3042	1.6860		2.0244	2.4286	2.7116
39	0.6808	1.3036	1.6849		2.0227	2.4258	2.7079
40	0.6807	1.3031	1.6839		2.0211	2.4233	2.7045
41	0.6805	1.3025	1.6829		2.0195	2.4208	2.7012
42	0.6804	1.3020	1.6820		2.0181	2.4185	2.6981
43	0.6802	1.3016	1.6811		2.0167	2.4163	2.6951
44	0.6801	1.3011	1.6802		2.0154	2.4141	2.6923
45	0.6800	1.3006	1.6794		2.0141	2.4121	2.6896

(Continued)

Table 4 (continued)

r	0.75	0.90	0.95 γ	0.975	0.99	0.995
46	0.6799	1.3002	1.6787	2.0129	2.4102	2.6870
47	0.6797	1.2998	1.6779	2.0117	2.4083	2.6846
48	0.6796	1.2994	1.6772	2.0106	2.4066	2.6822
49	0.6795	1.2991	1.6766	2.0096	2.4069	2.6800
50	0.6794	1.2987	1.6759	2.0086	2.4033	2.6778
51	0.6793	1.2984	1.6753	2.0076	2.4017	2.6757
52	0.6792	1.2980	1.6747	2.0066	2.4002	2.6737
53	0.6791	1.2977	1.6741	2.0057	2.3988	2.6718
54	0.6791	1.2974	1.6736	2.0049	2.3974	2.6700
55	0.6790	1.2971	1.6730	2.0040	2.3961	2.6682
56	0.6789	1.2969	1.6725	2.0032	2.3948	2.6665
57	0.6788	1.2966	1.6720	2.0025	2.3936	2.6649
58	0.6787	1.2963	1.6716	2.0017	2.3924	2.6633
59	0.6787	1.2961	1.6711	2.0010	2.3912	2.6618
60	0.6786	1.2958	1.6706	2.0003	2.3901	2.6603
61	0.6785	1.2956	1.6702	1.9996	2.3890	2.6589
62	0.6785	1.2954	1.6698	1.9990	2.3880	2.6575
63	0.6784	1.2951	1.6694	1.9983	2.3870	2.6561
64	0.6783	1.2949	1.6690	1.9977	2.3860	2.6549
65	0.6783	1.2947	1.6686	1.9971	2.3851	2.6536
66	0.6782	1.2945	1.6683	1.9966	2.3842	2.6524
67	0.6782	1.2943	1.6679	1.9960	2.3833	2.6512
68	0.6781	1.2941	1.6676	1.9955	2.3824	2.6501
69	0.6781	1.2939	1.6672	1.9949	2.3816	2.6490
70	0.6780	1.2938	1.6669	1.9944	2.3808	2.6479
71	0.6780	1.2936	1.6666	1.9939	2.3800	2.6469
72	0.6779	1.2934	1.6663	1.9935	2.3793	2.6459
73	0.6779	1.2933	1.6660	1.9930	2.3785	2.6449
74	0.6778	1.2931	1.6657	1.9925	2.3778	2.6439
75	0.6778	1.2929	1.6654	1.9921	2.3771	2.6430
76	0.6777	1.2928	1.6652	1.9917	2.3764	2.6421
77	0.6777	1.2926	1.6649	1.9913	2.3758	2.6412
78	0.6776	1.2925	1.6646	1.9908	2.3751	2.6403
79	0.6776	1.2924	1.6644	1.9905	2.3745	2.6395
80	0.6776	1.2922	1.6641	1.9901	2.3739	2.6387
81	0.6775	1.2921	1.6639	1.9897	2.3733	2.6379
82	0.6775	1.2920	1.6636	1.9893	2.3727	2.6371
83	0.6775	1.2918	1.6634	1.9890	2.3721	2.6364
84	0.6774	1.2917	1.6632	1.9886	2.3716	2.6356
85	0.6774	1.2916	1.6630	1.9883	2.3710	2.6349
86	0.6774	1.2915	1.6628	1.9879	2.3705	2.6342
87	0.6773	1.2914	1.6626	1.9876	2.3700	2.6335
88	0.6773	1.2912	1.6624	1.9873	2.3695	2.6329
89	0.6773	1.2911	1.6622	1.9870	2.3690	2.6322
90	0.6772	1.2910	1.6620	1.9867	2.3685	2.6316

Table 5

**Critical Values for the
Chi-Square Distribution**

Let χ_r^2 be a random variable having the chi-square distribution with r degrees of freedom. Then the tabulated quantities are the numbers x for which

$$P(\chi_r^2 \leq x) = \gamma.$$

r	γ					
	0.005	0.01	0.025	0.05	0.10	0.25
1	—	—	0.001	0.004	0.016	0.102
2	0.010	0.020	0.051	0.103	0.211	0.575
3	0.072	0.115	0.216	0.352	0.584	1.213
4	0.207	0.297	0.484	0.711	1.064	1.923
5	0.412	0.554	0.831	1.145	1.610	2.675
6	0.676	0.872	1.237	1.635	2.204	3.455
7	0.989	1.239	1.690	2.167	2.833	4.255
8	1.344	1.646	2.180	2.733	3.490	5.071
9	1.735	2.088	2.700	3.325	4.168	5.899
10	2.156	2.558	3.247	3.940	4.865	6.737
11	2.603	3.053	3.816	4.575	5.578	7.584
12	3.074	3.571	4.404	5.226	6.304	9.438
13	3.565	4.107	5.009	5.892	7.042	9.299
14	4.075	4.660	5.629	6.571	7.790	10.165
15	4.601	5.229	6.262	7.261	8.547	11.037
16	5.142	5.812	6.908	7.962	9.312	11.912
17	5.697	6.408	7.564	8.672	10.085	12.792
18	6.265	7.015	8.231	9.390	10.865	13.675
19	6.844	7.633	8.907	10.117	11.651	14.562
20	7.434	8.260	9.591	10.851	12.443	15.452
21	8.034	8.897	10.283	11.591	13.240	16.344
22	8.643	9.542	10.982	12.338	14.042	17.240
23	9.260	10.196	11.689	13.091	14.848	18.137
24	9.886	10.856	12.401	13.848	15.659	19.037
25	10.520	11.524	13.120	14.611	16.473	19.939
26	11.160	12.198	13.844	15.379	17.292	20.843
27	11.808	12.879	14.573	16.151	18.114	21.749
28	12.461	13.565	15.308	16.928	18.939	22.657
29	13.121	14.257	16.047	17.708	19.768	23.567
30	13.787	14.954	16.791	18.493	20.599	24.478
31	14.458	15.655	17.539	19.281	21.434	25.390
32	15.134	16.362	18.291	20.072	22.271	26.304
33	15.815	17.074	19.047	20.867	23.110	27.219
34	16.501	17.789	19.806	21.664	23.952	28.136
35	17.192	18.509	20.569	22.465	24.797	29.054
36	17.887	19.233	21.336	23.269	25.643	29.973
37	18.586	19.960	22.106	24.075	26.492	30.893
38	19.289	20.691	22.878	24.884	27.343	31.815
39	19.996	21.426	23.654	25.695	28.196	32.737
40	20.707	22.164	24.433	26.509	29.051	33.660
41	21.421	22.906	25.215	27.326	29.907	34.585
42	22.138	23.650	25.999	28.144	30.765	35.510
43	22.859	24.398	26.785	28.965	31.625	36.436
44	23.584	25.148	27.575	29.787	32.487	37.363
45	24.311	25.901	28.366	30.612	33.350	38.291

(Continued)

Table 5 (continued)

r	0.75	0.90	0.95 γ	0.975	0.99	0.995
1	1.323	2.706	3.841	5.024	6.635	7.879
2	2.773	4.605	5.991	7.378	9.210	10.597
3	4.108	6.251	7.815	9.348	11.345	12.838
4	5.385	7.779	9.488	11.143	13.277	14.860
5	6.626	9.236	11.071	12.833	15.086	16.750
6	7.841	10.645	12.592	14.449	16.812	18.548
7	9.037	12.017	14.067	16.013	18.475	20.278
8	10.219	13.362	15.507	17.535	20.090	21.955
9	11.389	14.684	16.919	19.023	21.666	23.589
10	12.549	15.987	18.307	20.483	23.209	25.188
11	13.701	17.275	19.675	21.920	24.725	26.757
12	14.845	18.549	21.026	23.337	26.217	28.299
13	15.984	19.812	23.362	24.736	27.688	29.819
14	17.117	21.064	23.685	26.119	29.141	31.319
15	18.245	22.307	24.996	27.488	30.578	32.801
16	19.369	23.542	26.296	28.845	32.000	34.267
17	20.489	24.769	27.587	30.191	33.409	35.718
18	21.605	25.989	28.869	31.526	34.805	37.156
19	22.718	27.204	30.144	32.852	36.191	38.582
20	23.828	28.412	31.410	34.170	37.566	39.997
21	24.935	29.615	32.671	35.479	38.932	41.401
22	26.039	30.813	33.924	36.781	40.289	42.796
23	27.141	32.007	35.172	38.076	41.638	44.181
24	28.241	33.196	36.415	39.364	42.980	45.559
25	29.339	34.382	37.652	40.646	44.314	46.928
26	30.435	35.563	38.885	41.923	45.642	48.290
27	31.528	36.741	40.113	43.194	46.963	49.645
28	32.620	37.916	41.337	44.461	48.278	50.993
29	33.711	39.087	42.557	45.722	49.588	52.336
30	34.800	40.256	43.773	46.979	50.892	53.672
31	35.887	41.422	44.985	48.232	51.191	55.003
32	36.973	42.585	46.194	49.480	53.486	56.328
33	38.058	43.745	47.400	50.725	54.776	57.648
34	39.141	44.903	48.602	51.966	56.061	58.964
35	40.223	46.059	49.802	53.203	57.342	60.275
36	41.304	47.212	50.998	54.437	58.619	61.581
37	42.383	48.363	52.192	55.668	59.892	62.883
38	43.462	49.513	53.384	56.896	61.162	64.181
39	44.539	50.660	54.572	58.120	62.428	65.476
40	45.616	51.805	55.758	59.342	63.691	66.766
41	46.692	52.949	56.942	60.561	64.950	68.053
42	47.766	54.090	58.124	61.777	66.206	69.336
43	48.840	55.230	59.304	62.990	67.459	70.616
44	49.913	56.369	60.481	64.201	68.710	71.893
45	50.985	57.505	61.656	65.410	69.957	73.166

Table 6

Let F_{r_1, r_2} be a random variable having the F -distribution with r_1, r_2 degrees of freedom. Then the tabulated quantities are the numbers x for which

$$P(F_{r_1, r_2} \leq x) = \gamma.$$

Critical Values for the F -Distribution

		r_1						
	γ	1	2	3	4	5	6	γ
1	0.500	1.0000	1.5000	1.7092	1.8227	1.8937	1.9422	0.500
	0.750	5.8285	7.5000	8.1999	8.5810	8.8198	8.9833	0.750
	0.900	39.864	49.500	53.593	55.833	57.241	58.204	0.900
	0.950	161.45	199.50	215.71	224.58	230.16	233.99	0.950
	0.975	647.79	799.50	864.16	899.58	921.85	937.11	0.975
	0.990	4052.2	4999.5	5403.3	5624.6	5763.7	5859.0	0.990
	0.995	16211	20000	21615	22500	23056	23437	0.995
2	0.500	0.66667	1.0000	1.1349	1.2071	1.2519	1.2824	0.500
	0.750	2.5714	3.0000	3.1534	3.2320	3.2799	3.3121	0.750
	0.900	8.5623	9.0000	9.1618	9.2434	9.2926	9.3255	0.900
	0.950	18.513	19.000	19.164	19.247	19.296	19.330	0.950
	0.975	38.506	39.000	39.165	39.248	39.298	39.331	0.975
	0.990	98.503	99.000	99.166	99.249	99.299	99.332	0.990
	0.995	198.50	199.00	199.17	199.25	199.30	199.33	0.995
3	0.500	0.58506	0.88110	1.0000	1.0632	1.1024	1.1289	0.500
	0.750	2.0239	2.2798	2.3555	2.3901	2.4095	2.4218	0.750
	0.900	5.5383	5.4624	5.3908	5.3427	5.3092	5.2847	0.900
	0.950	10.128	9.5521	9.2766	9.1172	9.0135	8.9406	0.950
	0.975	17.443	16.044	15.439	15.101	14.885	14.735	0.975
	0.990	34.116	30.817	29.457	28.710	28.237	27.911	0.990
	0.995	55.552	49.799	47.467	46.195	45.392	44.838	0.995
4	0.500	0.54863	0.82843	0.94054	1.0000	1.0367	1.0617	0.500
	0.750	1.8074	2.0000	2.0467	2.0642	2.0723	2.0766	0.750
	0.900	4.5448	4.3246	4.1908	4.1073	4.0506	4.0098	0.900
	0.950	7.7086	6.9443	6.5914	6.3883	6.2560	6.1631	0.950
	0.975	12.218	10.649	9.9792	9.6045	9.3645	9.1973	0.975
	0.990	21.198	18.000	16.694	15.977	15.522	15.207	0.990
	0.995	31.333	26.284	24.259	23.155	22.456	21.975	0.995
5	0.500	0.52807	0.79877	0.90715	0.96456	1.0000	1.0240	0.500
	0.750	1.6925	1.8528	1.8843	1.8927	1.8947	1.8945	0.750
	0.900	4.0604	3.7797	3.6195	3.5202	3.4530	3.4045	0.900
	0.950	6.6079	5.7861	5.4095	5.1922	5.0503	4.9503	0.950
	0.975	10.007	8.4336	7.7636	7.3879	7.1464	6.9777	0.975
	0.990	16.258	13.274	12.060	11.392	10.967	10.672	0.990
	0.995	22.785	18.314	16.530	15.556	14.940	14.513	0.995
6	0.500	0.51489	0.77976	0.88578	0.94191	0.97654	1.0000	0.500
	0.750	1.6214	1.7622	1.7844	1.7872	1.7852	1.7821	0.750
	0.900	3.7760	3.4633	3.2888	3.1808	3.1075	3.0546	0.900
	0.950	5.9874	5.1433	4.7571	4.5337	4.3874	4.2839	0.950
	0.975	8.8131	7.2598	6.5988	6.2272	5.9876	5.8197	0.975
	0.990	13.745	10.925	9.7795	9.1483	8.7459	8.4661	0.990
	0.995	18.635	14.544	12.917	12.028	11.464	11.073	0.995

(Continued)

Table 6 (continued)

		r_1									
		γ	7	8	9	10	11	12	γ		
1		0.500	1.9774	2.0041	2.0250	2.0419	2.0558	2.0674	0.500	1	
		0.750	9.1021	9.1922	9.2631	9.3202	9.3672	9.4064	0.750		
		0.900	58.906	59.439	59.858	60.195	60.473	60.705	0.900		
		0.950	236.77	238.88	240.54	241.88	242.99	243.91	0.950		
		0.975	948.22	956.66	963.28	968.63	973.04	976.71	0.975		
		0.990	5928.3	5981.1	6022.5	6055.8	6083.3	6106.3	0.990		
		0.995	23715	23925	24091	24224	24334	24426	0.995		
2		0.500	1.3045	1.3213	1.3344	1.3450	1.3537	1.3610	0.500	2	
		0.750	3.3352	3.3526	3.3661	3.3770	3.3859	3.3934	0.750		
		0.900	9.3491	9.3668	9.3805	9.3916	9.4006	9.4081	0.900		
		0.950	19.353	19.371	19.385	19.396	19.405	19.413	0.950		
		0.975	39.355	39.373	39.387	39.398	39.407	39.415	0.975		
		0.990	99.356	99.374	99.388	99.399	99.408	99.416	0.990		
		0.995	199.36	199.37	199.39	199.40	199.41	199.42	0.995		
3		0.500	1.1482	1.1627	1.1741	1.1833	1.1909	1.1972	0.500	3	
		0.750	2.4302	2.4364	2.4410	2.4447	2.4476	2.4500	0.750		
		0.900	5.2662	5.2517	5.2400	5.2304	5.2223	5.2156	0.900		
		0.950	8.8868	8.8452	8.8123	8.7855	8.7632	8.7446	0.950		
		0.975	14.624	14.540	14.473	14.419	14.374	14.337	0.975		
		0.990	27.672	27.489	27.345	27.229	27.132	27.052	0.990		
		0.995	44.434	44.126	43.882	43.686	43.523	43.387	0.995		
4		0.500	1.0797	1.0933	1.1040	1.1126	1.1196	1.1255	0.500	4	
		0.750	2.0790	2.0805	2.0814	2.0820	2.0823	2.0826	0.750		
		0.900	3.9790	3.9549	3.9357	3.9199	3.9066	3.8955	0.900		
		0.950	6.0942	6.0410	5.9988	5.9644	5.9357	5.9117	0.950		
		0.975	9.0741	8.9796	8.9047	8.8439	8.7933	8.7512	0.975		
		0.990	14.976	14.799	14.659	14.546	14.452	14.374	0.990		
		0.995	21.622	21.352	21.139	20.967	20.824	20.705	0.995		
5		0.500	1.0414	1.0545	1.0648	1.0730	1.0798	1.0855	0.500	5	
		0.750	1.8935	1.8923	1.8911	1.8899	1.8887	1.8877	0.750		
		0.900	3.3679	3.3393	3.3163	3.2974	3.2815	3.2682	0.900		
		0.950	4.8759	4.8183	4.7725	4.7351	4.7038	4.6777	0.950		
		0.975	6.8531	6.7572	6.6810	6.6192	6.5676	6.5246	0.975		
		0.990	10.456	10.289	10.158	10.051	9.9623	9.8883	0.990		
		0.995	14.200	13.961	13.772	13.618	13.490	13.384	0.995		
6		0.500	1.0169	1.0298	1.0398	1.0478	1.0545	1.0600	0.500	6	
		0.750	1.7789	1.7760	1.7733	1.7708	1.7686	1.7668	0.750		
		0.900	3.0145	2.9830	2.9577	2.9369	2.9193	2.9047	0.900		
		0.950	4.2066	4.1468	4.0990	4.0600	4.0272	3.9999	0.950		
		0.975	5.6955	5.5996	5.5234	5.4613	5.4094	5.3662	0.975		
		0.990	8.2600	8.1016	7.9761	7.8741	7.7891	7.7183	0.990		
		0.995	10.786	10.566	10.391	10.250	10.132	10.034	0.995		

Table 6 (continued)

		r_1									
		γ	13	14	15	18	20	24	γ		
1		0.500	2.0773	2.0858	2.0931	2.1104	2.1190	2.1321	0.500	1	
		0.750	9.4399	9.4685	9.4934	9.5520	9.5813	9.6255	0.750		
		0.900	60.903	61.073	61.220	61.567	61.740	62.002	0.900		
		0.950	244.69	245.37	245.95	247.32	248.01	249.05	0.950		
		0.975	979.85	982.54	984.87	990.36	993.10	997.25	0.975		
		0.990	6125.9	6142.7	6157.3	6191.6	6208.7	6234.6	0.990		
		0.995	24504	24572	24630	24767	24836	24940	0.995		
2		0.500	1.3672	1.3725	1.3771	1.3879	1.3933	1.4014	0.500	2	
		0.750	3.3997	3.4051	3.4098	3.4208	3.4263	3.4345	0.750		
		0.900	9.4145	9.4200	9.4247	9.4358	9.4413	9.4496	0.900		
		0.950	19.419	19.424	19.429	19.440	19.446	19.454	0.950		
		0.975	39.421	39.426	39.431	39.442	39.448	39.456	0.975		
		0.990	99.422	99.427	99.432	99.443	99.449	99.458	0.990		
		0.995	199.42	199.43	199.43	199.44	199.45	199.46	0.995		
3		0.500	1.2025	1.2071	1.2111	1.2205	1.2252	1.2322	0.500	3	
		0.750	2.4520	2.4537	2.4552	2.4585	2.4602	2.4626	0.750		
		0.900	5.2097	5.2047	5.2003	5.1898	5.1845	5.1764	0.900		
		0.950	8.7286	8.7148	8.7029	8.6744	8.6602	8.6385	0.950		
		0.975	14.305	14.277	14.253	14.196	14.167	14.124	0.975		
		0.990	26.983	26.923	26.872	26.751	26.690	26.598	0.990		
		0.995	43.271	43.171	43.085	42.880	42.778	42.622	0.955		
4		0.500	1.1305	1.1349	1.1386	1.1473	1.1517	1.1583	0.500	4	
		0.750	2.0827	2.0828	2.0829	2.0828	2.0828	2.0827	0.750		
		0.900	3.8853	3.8765	3.8689	3.8525	3.8443	3.8310	0.900		
		0.950	5.8910	5.8732	5.8578	5.8209	5.8025	5.7744	0.950		
		0.975	8.7148	8.6836	8.6565	8.5921	8.5599	8.5109	0.975		
		0.990	14.306	14.248	14.198	14.079	14.020	13.929	0.990		
		0.995	20.602	20.514	20.438	20.257	20.167	20.030	0.995		
5		0.500	1.0903	1.0944	1.0980	1.1064	1.1106	1.1170	0.500	5	
		0.750	1.8867	1.8858	1.8851	1.8830	1.8820	1.8802	0.750		
		0.900	3.2566	3.2466	3.2380	3.2171	3.2067	3.1905	0.900		
		0.950	4.6550	4.6356	4.6188	4.5783	4.5581	4.5272	0.950		
		0.975	6.4873	6.4554	6.4277	6.3616	6.3285	6.2780	0.975		
		0.990	9.8244	9.7697	9.7222	9.6092	9.5527	9.4665	0.990		
		0.995	13.292	13.214	13.146	12.984	12.903	12.780	0.995		
6		0.500	1.0647	1.0687	1.0722	1.0804	1.0845	1.0907	0.500	6	
		0.750	1.7650	1.7634	1.7621	1.7586	1.7569	1.7540	0.750		
		0.900	2.8918	2.8808	2.8712	2.8479	2.8363	2.8183	0.900		
		0.950	3.9761	3.9558	3.9381	3.8955	3.8742	3.8415	0.950		
		0.975	5.3287	5.2966	5.2687	5.2018	5.1684	5.1172	0.975		
		0.990	7.6570	7.6045	7.5590	7.4502	7.3958	7.3127	0.990		
		0.995	9.9494	9.8769	9.8140	9.6639	9.5888	9.4741	0.995		

(Continued)

Table 6 (continued)

		γ	30	40	48	r_1		∞	γ		
						60	120				
1		0.500	2.1452	2.1584	2.1650	2.1716	2.1848	2.1981	0.500	1	
		0.750	9.6698	9.7144	9.7368	9.7591	9.8041	9.8492	0.750		
		0.900	62.265	62.529	62.662	62.794	63.061	63.328	0.900		
		0.950	250.09	251.14	251.67	252.20	253.25	254.32	0.950		
		0.975	1001.4	1005.6	1007.7	1009.8	1014.0	1018.3	0.975		
		0.990	6260.7	6286.8	6299.9	6313.0	6339.4	6366.0	0.990		
		0.995	25044	25148	25201	25253	25359	25465	0.995		
2		0.500	1.4096	1.4178	1.4220	1.4261	1.4344	1.4427	0.500	2	
		0.750	3.4428	3.4511	3.4553	3.4594	3.4677	3.4761	0.750		
		0.900	9.4579	9.4663	9.4705	9.4746	9.4829	9.4913	0.900		
		0.950	19.462	19.471	19.475	19.479	19.487	19.496	0.950		
		0.975	39.465	39.473	39.477	39.481	39.490	39.498	0.975		
		0.990	99.466	99.474	99.478	99.483	99.491	99.499	0.990		
		0.995	199.47	199.47	199.47	199.48	199.49	199.51	0.995		
3		0.500	1.2393	1.2464	1.2500	1.2536	1.2608	1.2680	0.500	3	
		0.750	2.4650	2.4674	2.4686	2.4697	2.4720	2.4742	0.750		
		0.900	5.1681	5.1597	5.1555	5.1512	5.1425	5.1337	0.900		
		0.950	8.6166	8.5944	8.5832	8.5720	8.5494	8.5265	0.950		
		0.975	14.081	14.037	14.015	13.992	13.947	13.902	0.975		
		0.990	26.505	26.411	26.364	26.316	26.221	26.125	0.990		
		0.995	42.466	42.308	42.229	42.149	41.989	41.829	0.995		
4		0.500	1.1649	1.1716	1.1749	1.1782	1.1849	1.1916	0.500	4	
		0.750	2.0825	2.0821	2.0819	2.0817	2.0812	2.0806	0.750		
		0.900	3.8174	3.8036	3.7966	3.7896	3.7753	3.7607	0.900		
		0.950	5.7459	5.7170	5.7024	5.6878	5.6581	5.6281	0.950		
		0.975	8.4613	8.4111	8.3858	8.3604	8.3092	8.2573	0.975		
		0.990	13.838	13.745	13.699	13.652	13.558	13.463	0.990		
		0.995	19.892	19.752	19.682	19.611	19.468	19.325	0.995		
5		0.500	1.1234	1.1297	1.1329	1.1361	1.1426	1.1490	0.500	5	
		0.750	1.8784	1.8763	1.8753	1.8742	1.8719	1.8694	0.750		
		0.900	3.1741	3.1573	3.1488	3.1402	3.1228	3.1050	0.900		
		0.950	4.4957	4.4638	4.4476	4.4314	4.3984	4.3650	0.950		
		0.975	6.2269	6.1751	6.1488	6.1225	6.0693	6.0153	0.975		
		0.990	9.3793	9.2912	9.2466	9.2020	9.1118	0.0204	0.990		
		0.995	12.656	12.530	12.466	12.402	12.274	12.144	0.995		
6		0.500	1.0969	1.1031	1.1062	1.1093	1.1156	1.1219	0.500	6	
		0.750	1.7510	1.7477	1.7460	1.7443	1.7407	1.7368	0.750		
		0.900	2.8000	2.7812	2.7716	2.7620	2.7423	2.7222	0.900		
		0.950	3.8082	3.7743	3.7571	3.7398	3.7047	3.6688	0.950		
		0.975	5.0652	5.0125	4.9857	4.9589	4.9045	4.9491	0.975		
		0.990	7.2285	7.1432	7.1000	7.0568	6.9690	6.8801	0.990		
		0.995	9.3583	9.2408	9.1814	9.1219	9.0015	8.8793	0.995		

Table 6 (continued)

		r_1								
	γ	1	2	3	4	5	6		γ	
7	0.500	0.50572	0.76655	0.87095	0.92619	0.96026	0.98334	0.500	7	
	0.750	1.5732	1.7010	1.7169	1.7157	1.7111	1.7059	0.750		
	0.900	3.5894	3.2574	3.0741	2.9605	2.8833	2.8274	0.900		
	0.950	5.5914	4.7374	4.3468	4.1203	3.9715	3.8660	0.950		
	0.975	8.0727	6.5415	5.8898	5.5226	5.2852	5.1186	0.975		
	0.990	12.246	9.5466	8.4513	7.8467	7.4604	7.1914	0.990		
	0.995	16.236	12.404	10.882	10.050	9.5221	9.1554	0.995		
8	0.500	0.49898	0.75683	0.86004	0.91464	0.94831	0.97111	0.500	8	
	0.750	1.5384	1.6569	1.6683	1.6642	1.6575	1.6508	0.750		
	0.900	3.4579	3.1131	2.9238	2.8064	2.7265	2.6683	0.900		
	0.950	5.3177	4.4590	4.0662	3.8378	3.6875	3.5806	0.950		
	0.975	7.5709	6.0595	5.4160	5.0526	4.8173	4.6517	0.975		
	0.990	11.259	8.6491	7.5910	7.0060	6.6318	6.3707	0.990		
	0.995	14.688	11.042	9.5965	8.8051	8.3018	7.9520	0.995		
9	0.500	0.49382	0.74938	0.85168	0.90580	0.93916	0.96175	0.500	9	
	0.750	1.5121	1.6236	1.6315	1.6253	1.6170	1.6091	0.750		
	0.900	3.3603	3.0065	2.8129	2.6927	2.6106	2.5509	0.900		
	0.950	5.1174	4.2565	3.8626	3.6331	3.4817	3.3738	0.950		
	0.975	7.2093	5.7147	5.0781	4.7181	4.4844	4.3197	0.975		
	0.990	10.561	8.0215	6.9919	6.4221	6.0569	5.8018	0.990		
	0.995	13.614	10.107	8.7171	7.9559	7.4711	7.1338	0.995		
10	0.500	0.48973	0.74349	0.84508	0.89882	0.93193	0.95436	0.500	10	
	0.750	1.4915	1.5975	1.6028	1.5949	1.5853	1.5765	0.750		
	0.900	3.2850	2.9245	2.7277	2.6053	2.5216	2.4606	0.900		
	0.950	4.9646	4.1028	3.7083	3.4780	3.3258	3.2172	0.950		
	0.975	6.9367	5.4564	4.8256	4.4683	4.2361	4.0721	0.975		
	0.990	10.044	7.5594	6.5523	5.9943	5.6363	5.3858	0.990		
	0.995	12.826	9.4270	8.0807	7.3428	6.8723	6.5446	0.995		
11	0.500	0.48644	0.73872	0.83973	0.89316	0.92608	0.94837	0.500	11	
	0.750	1.4749	1.5767	1.5798	1.5704	1.5598	1.5502	0.750		
	0.900	3.2252	2.8595	2.6602	2.5362	2.4512	2.3891	0.900		
	0.950	4.8443	3.9823	3.5874	3.3567	3.2039	3.0946	0.950		
	0.975	6.7241	5.2559	4.6300	4.2751	4.0440	3.8807	0.975		
	0.990	9.6460	7.2057	6.2167	5.6683	5.3160	5.0692	0.990		
	0.995	12.226	8.9122	7.6004	6.8809	6.4217	6.1015	0.995		
12	0.500	0.48369	0.73477	0.83530	0.88848	0.92124	0.94342	0.500	12	
	0.750	1.4613	1.5595	1.5609	1.5503	1.5389	1.5286	0.750		
	0.900	3.1765	2.8068	2.6055	2.4801	2.3940	2.3310	0.900		
	0.950	4.7472	3.8853	3.4903	3.2592	3.1059	2.9961	0.950		
	0.975	6.5538	5.0959	4.4742	4.1212	3.8911	3.7283	0.975		
	0.990	9.3302	6.9266	5.9526	5.4119	5.0643	4.8206	0.990		
	0.995	11.754	8.5096	7.2258	6.5211	6.0711	5.7570	0.995		

(Continued)

Table 6 (continued)

		r_1									
		γ	7	8	9	10	11	12	γ		
7		0.500	1.0000	1.0216	1.0224	1.0304	1.0369	1.0423	0.500	7	
		0.750	1.7011	1.6969	1.6931	1.6898	1.6868	1.6843	0.750		
		0.900	2.7849	2.7516	2.7247	2.7025	2.6837	2.6681	0.900		
		0.950	3.7870	3.7257	3.6767	3.6365	3.6028	3.5747	0.950		
		0.975	4.9949	4.8994	4.8232	4.7611	4.7091	4.6658	0.975		
		0.990	6.9928	6.8401	6.7188	6.6201	6.5377	6.4691	0.990		
		0.995	8.8854	8.6781	8.5138	8.3803	8.2691	8.1764	0.995		
8		0.500	0.98757	1.0000	1.0097	1.0175	1.0239	1.0293	0.500	8	
		0.750	1.6448	1.6396	1.6350	1.6310	1.6274	1.6244	0.750		
		0.900	2.6241	2.5893	2.5612	2.5380	2.5184	2.5020	0.900		
		0.950	3.5005	3.4381	3.3881	3.3472	3.3127	3.2840	0.950		
		0.975	4.5286	4.4332	4.3572	4.2951	4.2431	4.1997	0.975		
		0.990	6.1776	6.0289	5.9106	5.8143	5.7338	5.6668	0.990		
		0.995	7.6942	7.4960	7.3386	7.2107	7.1039	7.0149	0.995		
9		0.500	0.97805	0.99037	1.0000	1.0077	1.0141	1.0194	0.500	9	
		0.750	1.6022	1.5961	1.5909	1.5863	1.5822	1.5788	0.750		
		0.900	2.5053	2.4694	2.4403	2.4163	2.3959	2.3789	0.900		
		0.950	3.2927	3.2296	3.1789	3.1373	3.1022	3.0729	0.950		
		0.975	4.1971	4.1020	4.0260	3.9639	3.9117	3.8682	0.975		
		0.990	5.6129	5.4671	5.3511	5.2565	5.1774	5.1114	0.990		
		0.995	6.8849	6.6933	6.5411	6.4171	6.3136	6.2274	0.995		
10		0.500	0.97054	0.98276	0.99232	1.0000	1.0063	1.0166	0.500	10	
		0.750	1.5688	1.5621	1.5563	1.5513	1.5468	1.5430	0.750		
		0.900	2.4140	2.3772	2.3473	2.3226	2.3016	2.2841	0.900		
		0.950	3.1355	3.0717	3.0204	2.9782	2.9426	2.9130	0.950		
		0.975	3.9498	3.8549	3.7790	3.7168	3.6645	3.6209	0.975		
		0.990	5.2001	5.0567	4.9424	4.8492	4.7710	4.7059	0.990		
		0.995	6.3025	6.1159	5.9676	5.8467	5.7456	5.6613	0.995		
11		0.500	0.96445	0.97661	0.98610	0.99373	0.99999	1.0052	0.500	11	
		0.750	1.5418	1.5346	1.5284	1.5230	1.5181	1.5140	0.750		
		0.900	2.3416	2.3040	2.2735	2.2482	2.2267	2.2087	0.900		
		0.950	3.0123	2.9480	2.8962	2.8536	2.8176	2.7876	0.950		
		0.975	3.7586	3.6638	3.5879	3.5257	3.4733	3.4296	0.975		
		0.990	4.8861	4.7445	4.6315	4.5393	4.4619	4.3974	0.990		
		0.995	5.8648	5.6821	5.5368	5.4182	5.3190	5.2363	0.995		
12		0.500	0.95943	0.97152	0.98097	0.98856	0.99480	1.0000	0.500	12	
		0.750	1.5197	1.5120	1.5054	1.4996	1.4945	1.4902	0.750		
		0.900	2.2828	2.2446	2.2135	2.1878	2.1658	2.1474	0.900		
		0.950	2.9134	2.8486	2.7964	2.7534	2.7170	2.6866	0.950		
		0.975	3.6065	3.5118	3.4358	3.3736	3.3211	3.2773	0.975		
		0.990	4.6395	4.4994	4.3875	4.2961	4.2193	4.1553	0.990		
		0.995	5.5245	5.3451	5.2021	5.0855	4.9878	4.9063	0.995		

Table 6 (continued)

		r_1							
γ	13	14	15	18	20	24	γ		
7	0.500	1.0469	1.0509	1.0543	1.0624	1.0664	1.0724	0.500	7
	0.750	1.6819	1.6799	1.6781	1.6735	1.6712	1.6675	0.750	
	0.900	2.6543	2.6425	2.6322	2.6072	2.5947	2.5753	0.900	
	0.950	3.5501	3.5291	3.5108	3.4666	3.4445	3.4105	0.950	
	0.975	4.6281	4.5958	4.5678	4.5004	4.4667	4.4150	0.975	
	0.990	6.4096	6.3585	6.3143	6.2084	6.1554	6.0743	0.990	
	0.995	8.0962	8.0274	7.9678	7.8253	7.7540	7.6450	0.995	
8	0.500	1.0339	1.0378	1.0412	1.0491	1.0531	1.0591	0.500	8
	0.750	1.6216	1.6191	1.6170	1.6115	1.6088	1.6043	0.750	
	0.900	2.4875	2.4750	2.4642	2.4378	2.4246	2.4041	0.900	
	0.950	3.2588	3.2371	3.2184	3.1730	3.1503	3.1152	0.950	
	0.975	4.1618	4.1293	4.1012	4.0334	3.9995	3.9472	0.975	
	0.990	5.6085	5.5584	5.5151	5.4111	5.3591	5.2793	0.990	
	0.995	6.9377	6.8716	6.8143	6.6769	6.6082	6.5029	0.995	
9	0.500	1.0239	1.0278	1.0311	1.0390	1.0429	1.0489	0.500	9
	0.750	1.5756	1.5729	1.5705	1.5642	1.5611	1.5560	0.750	
	0.900	2.3638	2.3508	2.3396	2.3121	2.2989	2.2768	0.900	
	0.950	3.0472	3.0252	3.0061	2.9597	2.9365	2.9005	0.950	
	0.975	3.8302	3.7976	3.7694	3.7011	3.6669	3.6142	0.975	
	0.990	5.0540	5.0048	4.9621	4.8594	4.8080	4.7290	0.990	
	0.995	6.1524	6.0882	6.0325	5.8987	5.8318	5.7292	0.995	
10	0.500	1.0161	1.0199	1.0232	1.0310	1.0349	1.0408	0.500	10
	0.750	1.5395	1.5364	1.5338	1.5269	1.5235	1.5179	0.750	
	0.900	2.2685	2.2551	2.2435	2.2150	2.2007	2.1784	0.900	
	0.950	2.8868	2.8644	2.8450	2.7977	2.7740	2.7372	0.950	
	0.975	3.5827	3.5500	3.5217	3.4530	3.4186	3.3654	0.975	
	0.990	4.6491	4.6004	4.5582	4.4563	4.4054	4.3269	0.990	
	0.995	5.5880	5.5252	5.4707	5.3396	5.2740	5.1732	0.995	
11	0.500	1.0097	1.0135	1.0168	1.0245	1.0284	1.0343	0.500	11
	0.750	1.5102	1.5069	1.5041	1.4967	1.4930	1.4869	0.750	
	0.900	2.1927	2.1790	2.1671	2.1377	2.1230	2.1000	0.900	
	0.950	2.7611	2.7383	2.7186	2.6705	2.6464	2.6090	0.950	
	0.975	3.3913	3.3584	3.3299	3.2607	3.2261	3.1725	0.975	
	0.990	4.3411	4.2928	4.2509	4.1496	4.0990	4.0209	0.990	
	0.995	5.1642	5.1024	5.0489	4.9198	4.8552	4.7557	0.995	
12	0.500	1.0044	1.0082	1.0115	1.0192	1.0231	1.0289	0.500	12
	0.750	1.4861	1.4826	1.4796	1.4717	1.4678	1.4613	0.750	
	0.900	2.1311	2.1170	2.1049	2.0748	2.0597	2.0360	0.900	
	0.950	2.6598	2.6368	2.6169	2.5680	2.5436	2.5055	0.950	
	0.975	3.2388	3.2058	3.1772	3.1076	3.0728	3.0187	0.975	
	0.990	4.0993	4.0512	4.0096	3.9088	3.8584	3.7805	0.990	
	0.995	4.8352	4.7742	4.7214	4.5937	4.5299	4.4315	0.995	

(Continued)

Table 6 (continued)

		r_1									
		γ	30	40	48	60	120	∞	γ		
7		0.500	1.0785	1.0846	1.0877	1.0908	1.0969	1.1031	0.500	7	
		0.750	1.6635	1.6593	1.6571	1.6548	1.6502	1.6452	0.750		
		0.900	2.5555	2.5351	2.5427	2.5142	2.4928	2.4708	0.900		
		0.950	3.3758	3.3404	3.3224	3.3043	3.2674	3.2298	0.950		
		0.975	4.3624	4.3089	4.2817	4.2544	4.1989	4.1423	0.975		
		0.990	5.9921	5.9084	5.8660	5.8236	5.7372	5.6495	0.990		
		0.995	7.5345	7.4225	7.3657	7.3088	7.1933	7.0760	0.995		
8		0.500	1.0651	1.0711	1.0741	1.0771	1.0832	1.0893	0.500	8	
		0.750	1.5996	1.5945	1.5919	1.5892	1.5836	1.5777	0.750		
		0.900	2.3830	2.3614	2.3503	2.3391	2.3162	2.2926	0.900		
		0.950	3.0794	3.0428	3.0241	3.0053	2.9669	2.9276	0.950		
		0.975	3.8940	3.8398	3.8121	3.7844	3.7279	3.6702	0.975		
		0.990	5.1981	5.1156	5.0736	5.0316	4.9460	4.8588	0.990		
		0.995	6.3961	6.2875	6.2324	6.1772	6.0649	5.9505	0.995		
9		0.500	1.0548	1.0608	1.0638	1.0667	1.0727	1.0788	0.500	9	
		0.750	1.5506	1.5450	1.5420	1.5389	1.5325	1.5257	0.750		
		0.900	2.2547	2.2320	2.2203	2.2085	2.1843	2.1592	0.900		
		0.950	2.8637	2.8259	2.8066	2.7872	2.7475	2.7067	0.950		
		0.975	3.5604	3.5055	3.4774	3.4493	3.3918	3.3329	0.975		
		0.990	4.6486	4.5667	4.5249	4.4831	4.3978	4.3105	0.990		
		0.995	5.6248	5.5186	5.4645	5.4104	5.3001	5.1875	0.995		
10		0.500	1.0467	1.0526	1.0556	1.0585	1.0645	1.0705	0.500	10	
		0.750	1.5119	1.5056	1.5023	1.4990	1.4919	1.4843	0.750		
		0.900	2.1554	1.1317	2.1195	2.1072	2.0818	2.0554	0.900		
		0.950	2.6996	2.6609	2.6410	2.6211	2.5801	2.5379	0.950		
		0.975	3.3110	3.2554	3.2269	3.1984	3.1399	3.0798	0.975		
		0.990	4.2469	4.1653	4.1236	4.0819	3.9965	3.9090	0.990		
		0.995	5.0705	4.9659	4.9126	4.8592	4.7501	4.6385	0.995		
11		0.500	1.0401	1.0460	1.0490	1.0519	1.0578	1.0637	0.500	11	
		0.750	1.4805	1.4737	1.4701	1.4664	1.4587	1.4504	0.750		
		0.900	2.0762	2.0516	2.0389	2.0261	1.9997	1.9721	0.900		
		0.950	2.5705	2.5309	2.5105	2.4901	2.4480	2.4045	0.950		
		0.975	3.1176	3.0613	3.0324	3.0035	2.9441	2.8828	0.975		
		0.990	3.9411	3.8596	3.8179	3.7761	3.6904	3.6025	0.990		
		0.995	4.6543	4.5508	4.4979	4.4450	4.3367	4.2256	0.995		
12		0.500	1.0347	1.0405	1.0435	1.0464	1.0523	1.0582	0.500	12	
		0.750	1.4544	1.4471	1.4432	1.4393	1.4310	1.4221	0.750		
		0.900	2.0115	1.9861	1.9729	1.9597	1.9323	1.9036	0.900		
		0.950	2.4663	2.4259	2.4051	2.3842	2.3410	2.2962	0.950		
		0.975	2.9633	2.9063	2.8771	2.8478	2.7874	2.7249	0.975		
		0.990	3.7008	3.6192	3.5774	3.5355	3.4494	3.3608	0.990		
		0.995	4.3309	4.2282	4.1756	4.1229	4.0149	3.9039	0.995		

Table 6 (continued)

		r_1								
γ	1	2	3	4	5	6	γ			
13	0.500	0.48141	0.73145	0.83159	0.88454	0.91718	0.93926	0.500	13	
	0.750	1.4500	1.5452	1.5451	1.5336	1.5214	1.5105	0.750		
	0.900	3.1362	2.7632	2.5603	2.4337	2.3467	2.2830	0.900		
	0.950	4.6672	3.8056	3.4105	3.1791	3.0254	2.9153	0.950		
	0.975	6.4143	4.9653	4.3472	3.9959	3.7667	3.6043	0.975		
	0.990	9.0738	6.7010	5.7394	5.2053	4.8616	4.6204	0.990		
	0.995	11.374	8.1865	6.9257	6.2335	5.7910	5.4819	0.995		
14	0.500	0.47944	0.72862	0.82842	0.88119	0.91371	0.93573	0.500	14	
	0.750	1.4403	1.5331	1.5317	1.5194	1.5066	1.4952	0.750		
	0.900	3.1022	2.7265	2.5222	2.3947	2.3069	2.2426	0.900		
	0.950	4.6001	3.7389	3.3439	3.1122	2.9582	2.8477	0.950		
	0.975	6.2979	4.8567	4.2417	3.8919	3.6634	3.5014	0.975		
	0.990	8.8616	6.5149	5.5639	5.0354	4.6950	4.4558	0.990		
	0.995	11.060	7.9216	6.6803	5.9984	5.5623	5.2574	0.995		
15	0.500	0.47775	0.72619	0.82569	0.87830	0.91073	0.93267	0.500	15	
	0.750	1.4321	1.5227	1.5202	1.5071	1.4938	1.4820	0.750		
	0.900	3.0732	2.6952	2.4898	2.3614	2.2730	2.2081	0.900		
	0.950	4.5431	3.6823	3.2874	3.0556	2.9013	2.7905	0.950		
	0.975	6.1995	4.7650	4.1528	3.8043	3.5764	3.4147	0.975		
	0.990	8.6831	6.3589	5.4170	4.8932	4.5556	4.3183	0.990		
	0.995	10.798	7.7008	6.4760	5.8029	5.3721	5.0708	0.995		
16	0.500	0.47628	0.72406	0.82330	0.87578	0.90812	0.93001	0.500	16	
	0.750	1.4249	1.5137	1.5103	1.4965	1.4827	1.4705	0.750		
	0.900	3.0481	2.6682	2.4618	2.3327	2.2438	2.1783	0.900		
	0.950	4.4940	3.6337	3.2389	3.0069	2.8524	2.7413	0.950		
	0.975	6.1151	4.6867	4.0768	3.7294	3.5021	3.3406	0.975		
	0.990	8.5310	6.2262	5.2922	4.7726	4.4374	4.2016	0.990		
	0.995	10.575	7.5138	6.3034	5.6378	5.2117	4.9134	0.995		
17	0.500	0.47499	0.72219	0.82121	0.87357	0.90584	0.92767	0.500	17	
	0.750	1.4186	1.5057	1.5015	1.4873	1.4730	1.4605	0.750		
	0.900	3.0262	2.6446	2.4374	2.3077	2.2183	2.1524	0.900		
	0.950	4.4513	3.5915	3.1968	2.9647	2.8100	2.6987	0.950		
	0.975	6.0420	4.6189	4.0112	3.6648	3.4379	3.2767	0.975		
	0.990	8.3997	6.1121	5.1850	4.6690	4.3359	4.1015	0.990		
	0.995	10.384	7.3536	6.1556	5.4967	5.0746	5.7789	0.995		
18	0.500	0.47385	0.72053	0.81936	0.87161	0.90381	0.92560	0.500	18	
	0.750	1.4130	1.4988	1.4938	1.4790	1.4644	1.4516	0.750		
	0.900	3.0070	2.6239	2.4160	2.2858	2.1958	1.1296	0.900		
	0.950	4.4139	3.5546	3.1599	2.9277	2.7729	2.6613	0.950		
	0.975	5.9781	4.5597	3.9539	3.6083	3.3820	3.2209	0.975		
	0.990	8.2854	6.0129	5.0919	4.5790	4.2479	4.0146	0.990		
	0.995	10.218	7.2148	6.0277	5.3746	4.9560	4.6627	0.995		

(Continued)

Table 6 (continued)

		r_1									
		γ	7	8	9	10	11	12	γ		
13	γ	0.500	0.95520	0.96724	0.97665	0.98421	0.99042	0.99560	0.500	13	γ
		0.750	1.5011	1.4931	1.4861	1.4801	1.4746	1.4701	0.750		
		0.900	2.2341	2.1953	2.1638	1.1376	1.1152	2.0966	0.900		
		0.950	2.8321	2.7669	2.7144	2.6710	2.6343	2.6037	0.950		
		0.975	3.4827	3.3880	3.3120	3.2497	3.1971	3.1532	0.975		
		0.990	4.4410	4.3021	4.1911	4.1003	4.0239	3.9603	0.990		
		0.995	5.2529	5.0761	4.9351	4.8199	4.7234	4.6429	0.995		
14	γ	0.500	0.95161	0.96360	0.97298	0.98051	0.98670	0.99186	0.500	14	γ
		0.750	1.4854	1.4770	1.4697	1.4634	1.4577	1.4530	0.750		
		0.900	2.1931	2.1539	2.1220	2.0954	2.0727	2.0537	0.900		
		0.950	2.7642	2.6987	2.6548	2.6021	2.5651	2.5342	0.950		
		0.975	3.3799	2.2853	3.2093	3.1469	3.0941	3.0501	0.975		
		0.990	4.2779	4.1399	4.0297	3.9394	3.8634	3.8001	0.990		
		0.995	5.0313	4.8566	4.7173	4.6034	4.5078	4.4281	0.995		
15	γ	0.500	0.94850	0.96046	0.96981	0.97732	0.98349	0.98863	0.500	15	γ
		0.750	1.4718	1.4631	1.4556	1.4491	1.4432	1.4383	0.750		
		0.900	2.1582	2.1185	2.0862	2.0593	2.0363	2.0171	0.900		
		0.950	2.7066	2.6408	2.5876	2.5437	2.5064	2.4753	0.950		
		0.975	3.2934	3.1987	3.1227	3.0602	3.0073	2.9633	0.975		
		0.990	4.1415	4.0045	3.8948	3.8049	3.7292	3.6662	0.990		
		0.995	4.8473	4.6743	4.5364	4.4236	4.3288	4.2498	0.995		
16	γ	0.500	0.94580	0.95773	0.96705	0.97454	0.98069	0.98582	0.500	16	γ
		0.750	1.4601	1.4511	1.4433	1.4366	1.4305	1.4255	0.750		
		0.900	2.1280	2.0880	2.0553	2.0281	2.0048	1.9854	0.900		
		0.950	2.6572	2.5911	2.5377	2.4935	2.4560	2.4247	0.950		
		0.975	3.2194	3.1248	3.0488	2.9862	2.9332	2.8890	0.975		
		0.990	4.0259	3.8896	3.7804	3.6909	3.6155	3.5527	0.990		
		0.995	4.6920	4.5207	4.3838	4.2719	4.1778	4.0994	0.995		
17	γ	0.500	0.94342	0.95532	0.96462	0.97209	0.97823	0.98334	0.500	17	γ
		0.750	1.4497	1.4405	1.4325	1.4256	1.4194	1.4142	0.750		
		0.900	2.1017	2.0613	2.0284	2.0009	1.9773	1.9577	0.900		
		0.950	2.6143	2.5480	2.4943	2.4499	2.4122	2.3807	0.950		
		0.975	3.1556	3.0610	2.9849	2.9222	2.8691	2.8249	0.975		
		0.990	3.9267	3.7910	3.6822	3.5931	3.5179	3.4552	0.990		
		0.995	4.5594	4.3893	4.2535	4.1423	4.0488	3.9709	0.995		
18	γ	0.500	0.94132	0.95319	0.96247	0.96993	0.97606	0.98116	0.500	18	γ
		0.750	1.4406	1.4312	1.4320	1.4159	1.4095	1.4042	0.750		
		0.900	2.0785	2.0379	2.0047	1.9770	1.9532	1.9333	0.900		
		0.950	2.5767	2.5102	2.4563	2.4117	2.3737	2.3421	0.950		
		0.975	3.0999	3.0053	2.9291	2.8664	2.8132	2.7689	0.975		
		0.990	3.8406	3.7054	3.5971	3.5082	3.4331	3.3706	0.990		
		0.995	4.4448	4.2759	4.1410	4.0305	3.9374	3.8599	0.995		

These tables have been adapted from Donald B. Owen's *Handbook of Statistical Tables*, published by Addison-Wesley, by permission of the publishers.

Table 7 Table of Selected Discrete and Continuous Distributions and Some of Their Characteristics

Distribution	PROBABILITY DENSITY FUNCTIONS IN ONE VARIABLE		
	Probability Density Function	Mean	Variance
Binomial, $B(n, p)$	$f(x) = \binom{n}{x} p^x q^{n-x}, x = 0, 1, \dots, n;$ $0 < p < 1, q = 1 - p$	np	npq
(Bernoulli, $B(1, p)$)	$f(x) = p^x q^{1-x}, x = 0, 1$	p	pq
Geometric	$f(x) = pq^{x-1}, x = 1, 2, \dots;$ $0 < p < 1, q = 1 - p$	$\frac{1}{p}$	$\frac{q}{p^2}$
Poisson, $P(\lambda)$	$f(x) = e^{-\lambda} \frac{\lambda^x}{x!}, x = 0, 1, \dots; \lambda > 0$	λ	λ
Hypergeometric	$f(x) = \frac{\binom{m}{x} \binom{n}{r-x}}{\binom{m+n}{r}}, \text{ where}$ $x = 0, 1, \dots, r \left(\binom{m}{r} = 0, r > m \right)$	$\frac{mr}{m+n}$	$\frac{mnr(m+n-r)}{(m+n)^2(m+n-1)}$
Gamma	$f(x) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} \exp\left(-\frac{x}{\beta}\right), x > 0;$ $\alpha, \beta > 0$	$\alpha\beta$	$\alpha\beta^2$
Negative Exponential	$f(x) = \lambda \exp(-\lambda x), x > 0; \lambda > 0; \text{ or}$ $f(x) = \frac{1}{\mu} e^{-x/\mu}, x > 0; \mu > 0$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$
Chi-Square	$f(x) = \frac{1}{\Gamma\left(\frac{r}{2}\right) 2^{r/2}} x^{r/2-1} \exp\left(-\frac{x}{2}\right), x > 0;$ $r > 0 \text{ integer}$	r	$2r$
Normal, $N(\mu, \sigma^2)$	$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2}\right],$ $x \in \Re; \mu \in \Re, \sigma > 0$	μ	σ^2
(Standard Normal, $N(0, 1)$)	$f(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right), x \in \Re$	0	1)
Uniform, $U(\alpha, \beta)$	$f(x) = \frac{1}{\beta - \alpha}, \alpha \leq x \leq \beta;$ $-\infty < \alpha < \beta < \infty$	$\frac{\alpha + \beta}{2}$	$\frac{(\alpha - \beta)^2}{12}$


(Continued)

Table 7 (continued)

Distribution	PROBABILITY DENSITY FUNCTIONS IN MANY VARIABLES		
	Probability Density Function	Means	Variations
Multinomial	$f(x_1, \dots, x_k) = \frac{n!}{x_1!x_2! \cdots x_k!} \times$ $p_1^{x_1} p_2^{x_2} \cdots p_k^{x_k}, x_i \geq 0 \text{ integers,}$ $x_1 + x_2 + \cdots + x_k = n; p_j > 0, j = 1,$ $2, \dots, k, p_1 + p_2 + \cdots + p_k = 1$	np_1, \dots, np_k	$np_1q_1, \dots, np_kq_k.$ $q_i = 1 - p_i, j = 1, \dots, k$
Bivariate Normal	$f(x_1, x_2) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \exp\left(-\frac{q}{2}\right),$ $q = \frac{1}{1-\rho^2} \left[\left(\frac{x_1-\mu_1}{\sigma_1}\right)^2 - 2\rho\left(\frac{x_1-\mu_1}{\sigma_1}\right) \right.$ $\left. \times \left(\frac{x_2-\mu_2}{\sigma_2}\right) + \left(\frac{x_2-\mu_2}{\sigma_2}\right)^2 \right],$ $x_1, x_2, \in \mathfrak{R}; \mu_1, \mu_2 \in \mathfrak{R}, \sigma_1, \sigma_2 > 0, -1 \leq \rho \leq 1, \rho = \text{correlation coefficient}$	μ_1, μ_2	σ_1^2, σ_2^2
k-Variate Normal, $N(\mu, \Sigma)$	$f(\mathbf{x}) = (2\pi)^{-k/2} \Sigma ^{-1/2} \times$ $\exp\left[-\frac{1}{2}(\mathbf{x} - \mu)' \Sigma^{-1}(\mathbf{x} - \mu),\right.$ $\mathbf{x} \in \mathfrak{R}^k; \mu \in \mathfrak{R}^k, \Sigma : k \times k$ <p>nonsingular symmetric matrix</p>	μ_1, \dots, μ_k	Covariance matrix: Σ

Table 7 (continued)

Distribution	Moment Generating Function
Binomial, $B(n, p)$ (Bernoulli, $B(1, p)$)	$M(t) = (pe^t + q)^n, t \in \mathfrak{R}$ $M(t) = pe^t + q, t \in \mathfrak{R}$
Geometric	$M(t) = \frac{pe^t}{1 - qe^t}, t < -\log q$
Poisson, $P(\lambda)$	$M(t) = \exp(\lambda e^t - \lambda), t \in \mathfrak{R}$
Hypergeometric	—
Gamma	$M(t) = \frac{1}{(1 - \beta t)^\alpha}, t < \frac{1}{\beta}$
Negative Exponential	$M(t) = \frac{\lambda}{\lambda - t}, t < \lambda; \text{ or } M(t) = \frac{1}{1 - \mu t}, t < \frac{1}{\mu}$
Chi-Square	$M(t) = \frac{1}{(1 - 2t)^{r/2}}, t < \frac{1}{2}$
Normal, $N(\mu, \sigma^2)$	$M(t) = \exp\left(\mu t + \frac{\sigma^2 t^2}{2}\right), t \in \mathfrak{R}$
(Standard Normal, $N(0, 1)$)	$M(t) = \exp\left(\frac{t^2}{2}\right), t \in \mathfrak{R}$
Uniform, $U(\alpha, \beta)$	$M(t) = \frac{e^{t\beta} - e^{t\alpha}}{t(\beta - \alpha)}, t \in \mathfrak{R}$
Multinomial	$M(t_1, \dots, t_k) = (p_1 e^{t_1} + \dots + p_k e^{t_k})^n,$ $t_1, \dots, t_k \in \mathfrak{R}$
Bivariate Normal	$M(t_1, t_2) = \exp\left[\mu_1 t_1 + \mu_2 t_2 + \frac{1}{2}\left(\sigma_1^2 t_1^2 + 2\rho\sigma_1\sigma_2 t_1 t_2 + \sigma_2^2 t_2^2\right)\right],$ $t_1, t_2 \in \mathfrak{R}$
k -Variate Normal, $N(\mu, \Sigma)$	$M(\mathbf{t}) = \exp\left(\mathbf{t}'\mu + \frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}\right),$ $\mathbf{t} \in \mathfrak{R}^k$




Some Notation and Abbreviations

\mathfrak{R}	real line
$\mathfrak{R}^k, k \geq 1$	k -dimensional Euclidean space
\uparrow, \downarrow	increasing (nondecreasing) and decreasing (nonincreasing), respectively
\mathcal{S}	sample space; also, sure (or certain) event
\emptyset	empty set; also, impossible event
$A \subseteq B$	event A is contained in event B (event A implies event B)
A^c	complement of event A
$A \cup B$	union of events A and B
$A \cap B$	intersection of events A and B
$A - B$	difference of events A and B (in this order)
r.v.	random variable
I_A	indicator of the set A : $I_A(x) = 1$ if $x \in A$, $I_A(x) = 0$ if $x \notin A$
$(X \in B) = X^{-1}(B)$	inverse image of the set B under X : $X^{-1}(B) = \{s \in \mathcal{S}; X(s) \in B\}$
$X(\mathcal{S})$	range of X
P	probability function (measure)
$P(A)$	probability of the event A
P_X	probability distribution of X (or just distribution of X)
F_X	distribution function (d.f.) of X
f_X	probability density function (p.d.f.) of X
$P(A B)$	conditional probability of A , given B
$\binom{n}{k}$	combinations of n objects taken k at a time
$P_{n,k}$	permutations of n objects taken k at a time
$n!$	n factorial
EX or $\mu(X)$ or μ_X or just μ	expectation (mean value, mean) of X
$\text{Var}(X)$ or $\sigma^2(X)$ or σ_X^2 or just σ^2	variance of X

$\sqrt{\text{Var}(X)}$ or $\sigma(X)$ or σ_X or just σ	standard deviation (s.d.) of X
M_X or just M	moment generating function (m.g.f.) of X
$B(n, p)$	Binomial distribution with parameters n and p
$P(\lambda)$	Poisson distribution with parameter λ
χ_r^2	Chi-Square distribution with r degrees of freedom (d.f.)
$N(\mu, \sigma^2)$	Normal distribution with parameters μ and σ^2
Φ	distribution function (d.f.) of the standard $N(0, 1)$ distribution
$U(\alpha, \beta)$ or $R(\alpha, \beta)$	Uniform (or Rectangular) distribution with parameters α and β
$X \sim B(n, p)$ etc.	the r.v. X has the distribution indicated
$\chi_{r;\alpha}^2$	the point for which $P(X > \chi_{r;\alpha}^2) = \alpha$, $X \sim \chi_r^2$
z_α	the point for which $P(Z > z_\alpha) = \alpha$, where $Z \sim N(0, 1)$
P_{X_1, \dots, X_n} or P_X	joint probability distribution of the r.v.'s X_1, \dots, X_n or probability distribution of the random vector \mathbf{X}
F_{X_1, \dots, X_n} or F_X	joint d.f. of the r.v.'s X_1, \dots, X_n or d.f. of the random vector \mathbf{X}
f_{X_1, \dots, X_n} or f_X	joint p.d.f. of the r.v.'s X_1, \dots, X_n or p.d.f. of the random vector \mathbf{X}
M_{X_1, \dots, X_n} or M_X	joint m.g.f. of the r.v.'s X_1, \dots, X_n or m.g.f. of the random vector \mathbf{X}
i.i.d. (r.v.'s)	independent identically distributed (r.v.'s)
$f_{X Y}(\cdot y)$ or $f_{X Y}(\cdot y)$	conditional p.d.f. of X , given $Y = y$
$E(X Y = y)$	conditional expectation of X , given $Y = y$
$\text{Var}(X Y = y)$ or $\sigma^2(X Y = y)$	conditional variance of X , given $Y = y$
$\text{Cov}(X, Y)$	covariance of X and Y
$\rho(X, Y)$ or $\rho_{X,Y}$	correlation coefficient of X and Y
t_r	(Student's) t distribution with r degrees of freedom (d.f.)
$t_{r;\alpha}$	the point for which $P(X > t_{r;\alpha}) = \alpha$, $X \sim t_r$
F_{r_1, r_2}	F distribution with r_1 and r_2 degrees of freedom (d.f.)
$F_{r_1, r_2; \alpha}$	the point for which $P(X > F_{r_1, r_2; \alpha}) = \alpha$, $X \sim F_{r_1, r_2}$
$X_{(j)}$ or Y_j	j th order statistic of X_1, \dots, X_n
$\xrightarrow{P}, \xrightarrow{d}, \xrightarrow{\text{q.m.}}$	convergence in probability, distribution, quadratic mean, respectively
WLLN	Weak Law of Large Numbers
CLT	Central Limit Theorem
θ	letter used for a one-dimensional parameter
$\boldsymbol{\theta}$	symbol used for a multidimensional parameter
Ω	letter used for a parameter space
ML	maximum likelihood
MLE	maximum likelihood estimate
UMV	uniformly minimum variance
UMVU	uniformly minimum variance unbiased
LS	least squares
LSE	least squares estimate
H_0	null hypothesis
H_A	alternative hypothesis

φ	letter used for a test function
α	letter used for level of significance
$\beta(\theta)$ or $\beta(\boldsymbol{\theta})$	probability of type II error at $\theta(\boldsymbol{\theta})$
$\pi(\theta)$ or $\pi(\boldsymbol{\theta})$	power of a test at $\theta(\boldsymbol{\theta})$
MP	most powerful (test)
UMP	uniformly most powerful (test)
LR	likelihood ratio
$\lambda = \lambda(x_1, \dots, x_n)$	likelihood ratio test function
$\log x$	the logarithm of $x(>0)$ with base always e whether it is so explicitly stated or not



Answers to Even-Numbered Exercises

Chapter 1

Section 1.2

- 2.2 (i) $S = \{(r, r, r), (r, r, b), (r, r, g), (r, b, r), (r, b, b), (r, b, g), (r, g, r), (r, g, b), (r, g, g), (b, r, r), (b, r, b), (b, r, g), (b, b, r), (b, b, b), (b, b, g), (b, g, r), (b, g, b), (b, g, g), (g, r, r), (g, r, b), (g, r, g), (g, b, r), (g, b, b), (g, b, g), (g, g, r), (g, g, b), (g, g, g)\}$.
- (ii) $A = \{(r, b, g), (r, g, b), (b, r, g), (b, g, r), (g, r, b), (g, b, r)\}$,
 $B = \{(r, r, b), (r, r, g), (r, b, r), (r, b, b), (r, g, r), (r, g, g), (b, r, r), (b, r, b), (b, b, r), (b, b, g), (b, g, b), (b, g, g), (g, r, r), (g, r, g), (g, b, b), (g, b, g), (g, g, r), (g, g, b)\}$,
 $C = A \cup B = S - \{(r, r, r), (b, b, b), (g, g, g)\}$.
- 2.4 (i) Denoting by (x_1, x_2) the cars sold in the first and the second sale, we have:
- $$S = \{(a_1, a_1), (a_1, a_2), (a_1, a_3), (a_2, a_1), (a_2, a_2), (a_2, a_3), (a_3, a_1), (a_3, a_2), (a_3, a_3), (a_1, b_1), (a_1, b_2), (a_2, b_1), (a_2, b_2), (a_3, b_1), (a_3, b_2), (a_1, c), (a_2, c), (a_3, c), (b_1, a_1), (b_1, a_2), (b_1, a_3), (b_2, a_1), (b_2, a_2), (b_2, a_3), (b_1, b_1), (b_1, b_2), (b_2, b_1), (b_2, b_2), (b_1, c), (b_2, c), (c, a_1), (c, a_2), (c, a_3), (c, b_1), (c, b_2), (c, c)\}.$$
- (ii) $A = \{(a_1, a_1), (a_1, a_2), (a_1, a_3), (a_2, a_1), (a_2, a_2), (a_2, a_3), (a_3, a_1), (a_3, a_2), (a_3, a_3)\}$,
 $B = \{(a_1, b_1), (a_1, b_2), (a_2, b_1), (a_2, b_2), (a_3, b_1), (a_3, b_2)\}$,

$$C = B \cup \{(b_1, a_1), (b_1, a_2), (b_1, a_3), (b_2, a_1), (b_2, a_2), (b_2, a_3)\},$$

$$D = \{(c, b_1), (c, b_2), (b_1, c), (b_2, c)\}.$$

$$2.6 \quad E = A^c, F = C - D = C \cap D^c, G = B - C = B \cap C^c,$$

$$H = A^c - B = A^c \cap B^c = (A \cup B)^c, I = B^c.$$

$$2.8 \quad \text{(i)} \quad B_0 = A_1^c \cap A_2^c \cap A_3^c.$$

$$\text{(ii)} \quad B_1 = (A_1 \cap A_2^c \cap A_3^c) \cup (A_1^c \cap A_2 \cap A_3^c) \cup (A_1^c \cap A_2^c \cap A_3).$$

$$\text{(iii)} \quad B_2 = (A_1 \cap A_2 \cap A_3^c) \cup (A_1 \cap A_2^c \cap A_3) \cup (A_1^c \cap A_2 \cap A_3).$$

$$\text{(iv)} \quad B_3 = A_1 \cap A_2 \cap A_3.$$

$$\text{(v)} \quad C = B_0 \cup B_1 \cup B_2.$$

$$\text{(vi)} \quad D = B_1 \cup B_2 \cup B_3 = A_1 \cup A_2 \cup A_3.$$

2.10 If $A = \emptyset$, then $A \cap B^c = \emptyset$, $A^c \cap B = S \cap B = B$, so that $(A \cap B^c) \cup (A^c \cap B) = B$ for every B . Next, let $(A \cap B^c) \cup (A^c \cap B) = B$ and take $B = \emptyset$ to obtain $A \cap B^c = A$, $A^c \cap B = \emptyset$, so that $A = \emptyset$.

2.12 $A \subseteq B$ implies that, for every $s \in A$, we have $s \in B$, whereas $B \subseteq C$ implies that, for every $s \in B$, we have $s \in C$. Thus, for every $s \in A$, we have $s \in C$, so that $A \subseteq C$.

2.14 For $s \in \cup_j A_j$, let $j_0 \geq 1$ be the first j for which $s \in A_{j_0}$. Then, if $j_0 = 1$, it follows that $s \in A_1$ and therefore s belongs in the right-hand side of the relation. If $j_0 > 1$, then $s \notin A_j$, $j = 1, \dots, j_0 - 1$, but $s \in A_{j_0}$, so that $s \in A_1^c \cap \dots \cap A_{j_0-1}^c \cap A_{j_0}$ and hence s belongs to the right-hand side of the relation. Next, let s belong to the right-hand side event. Then, if $s \in A_1$, it follows that $s \in \cup_j A_j$. If $s \notin A_j$ for $j = 1, \dots, j_0 - 1$ but $s \in A_{j_0}$, it follows that $s \in \cup_j A_j$. The identity is established.

2.16 (i) Since $-5 + \frac{1}{n+1} < -5 + \frac{1}{n}$ and $20 - \frac{1}{n} < 20 - \frac{1}{n+1}$, it follows that $(-5 + \frac{1}{n}, 20 - \frac{1}{n}) \subset (-5 + \frac{1}{n+1}, 20 - \frac{1}{n+1})$, or $A_n \subset A_{n+1}$, so that $\{A_n\}$ is increasing. Likewise, $7 + \frac{3}{n+1} < 7 + \frac{3}{n}$, so that $(0, 7 + \frac{3}{n+1}) \subset (0, 7 + \frac{3}{n})$, or $B_{n+1} \subset B_n$; thus, $\{B_n\}$ is decreasing.

$$\text{(ii)} \quad \cup_{n=1}^{\infty} A_n = \cup_{n=1}^{\infty} (-5 + \frac{1}{n}, 20 - \frac{1}{n}) = (-5, 20), \text{ and } \cap_{n=1}^{\infty} B_n = \cap_{n=1}^{\infty} (0, 7 + \frac{3}{n}) = (0, 7].$$

Section 1.3

3.2 Each one of the r.v.'s X_i , $i = 1, 2, 3$ takes on the values: 0, 1, 2, 3 and $X_1 + X_2 + X_3 = 3$.

3.4 X takes on the values: -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7,

$$(X \leq 2) = \{(-3, 0), (-3, 1), (-3, 2), (-3, 3), (-3, 4), (-2, 0), (-2, 1),$$

$$(-2, 2), (-2, 3), (-2, 4), (-1, 0), (-1, 1), (-1, 2), (-1, 3),$$

$$(0, 0), (0, 1), (0, 2), (1, 0), (1, 1), (2, 0)\},$$

$$(3 < X \leq 5) = (4 \leq X \leq 5) = (X = 4 \text{ or } X = 5)$$

$$= \{(0, 4), (1, 3), (1, 4), (2, 2), (2, 3), (3, 1), (3, 2)\},$$

$$(X > 6) = (X \geq 7) = \{(3, 4)\}.$$

- 3.6 (i) $S = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (2, 4), (3, 1), (3, 2), (3, 3), (3, 4), (4, 1), (4, 2), (4, 3), (4, 4)\}$.
- (ii) The values of X are: 2, 3, 4, 5, 6, 7, 8.
- (iii) $(X \leq 3) = (X = 2 \text{ or } X = 3) = \{(1, 1), (1, 2), (2, 1)\}$,
 $(2 \leq X < 5) = (2 \leq X \leq 4) = (X = 2 \text{ or } X = 3 \text{ or } X = 4) =$
 $\{(1, 1), (1, 2), (2, 1), (1, 3), (2, 2), (3, 1)\}$, $(X > 8) = \emptyset$.
- 3.8 (i) $S = [8:00, 8:15]$.
- (ii) The values of X consist of the interval $[8:00, 8:15]$.
- (iii) The event described is the interval $[8:10, 8:15]$.

Chapter 2

Section 2.1

- 1.2 Since $A \cup B \supseteq A$, we have $P(A \cup B) \geq P(A) = \frac{3}{4}$. Also, $A \cap B \subseteq B$ implies $P(A \cap B) \leq P(B) = \frac{3}{8}$. Finally, $P(A \cap B) = P(A) + P(B) - P(A \cup B) = \frac{3}{4} + \frac{3}{8} - P(A \cup B) = \frac{9}{8} - P(A \cup B) \geq \frac{9}{8} - 1 = \frac{1}{8}$.
- 1.4 We have: $A^c \cap B = B \cap A^c = B - A$ and $A \subset B$. Therefore $P(A^c \cap B) = P(B - A) = P(B) - P(A) = \frac{5}{12} - \frac{1}{4} = \frac{1}{6} \approx 0.167$. Likewise, $A^c \cap C = C - A$ with $A \subset C$, so that $P(A^c \cap C) = P(C - A) = P(C) - P(A) = \frac{7}{12} - \frac{1}{4} = \frac{1}{3} \approx 0.333$, $B^c \cap C = C - B$ with $B \subset C$, so that $P(B^c \cap C) = P(C - B) = P(C) - P(B) = \frac{7}{12} - \frac{5}{12} = \frac{1}{6} \approx 0.167$. Next, $A \cap B^c \cap C^c = A \cap (B^c \cap C^c) = A \cap (B \cup C)^c = A \cap C^c = A - C = \emptyset$, so that $P(A \cap B^c \cap C^c) = 0$, and $A^c \cap B^c \cap C^c = (A \cup B \cup C)^c = C^c$, so that $P(A^c \cap B^c \cap C^c) = P(C^c) = 1 - P(C) = 1 - \frac{7}{12} = \frac{5}{12} \approx 0.417$.
- 1.6 The event A is defined as follows: $A = "x = 7n, n = 1, \dots, 28,"$ so that $P(A) = \frac{28}{200} = \frac{7}{50} = 0.14$. Likewise, $B = "x = 3n + 10, n = 1, \dots, 63,"$ so that $P(B) = \frac{63}{200} = 0.315$, and $C = "x^2 + 1 \leq 375" = "x^2 \leq 374" = "x \leq \sqrt{374}" = "x \leq 19,"$ and then $P(C) = \frac{19}{200} = 0.095$.
- 1.8 Denote by $A, B,$ and C the events that a student reads news magazines $A, B,$ and $C,$ respectively. Then the required probability is $P(A^c \cap B^c \cap C^c)$. However,

$$\begin{aligned}
 P(A^c \cap B^c \cap C^c) &= P((A \cup B \cup C)^c) = 1 - P(A \cup B \cup C) \\
 &= 1 - [P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) \\
 &\quad - P(B \cap C) + P(A \cap B \cap C)] \\
 &= 1 - (0.20 + 0.15 + 0.10 - 0.05 - 0.04 - 0.03 + 0.02) \\
 &= 1 - 0.35 = 0.65.
 \end{aligned}$$

- 1.10 From the definition of $A, B,$ and $C,$ we have:

$$A = \{(0, 4), (0, 6), (1, 3), (1, 5), (1, 9), (2, 2), (2, 4), (2, 8), (3, 1), (3, 3), (3, 7), (4, 0), (4, 2), (4, 6), (5, 1), (5, 5), (6, 0), (6, 4)\},$$

$$B = \{(0, 0), (1, 2), (2, 4), (3, 6), (4, 8)\},$$

$$C = \{(0, 1), (0, 2), (0, 3), (0, 4), (0, 5), (0, 6), (0, 7), (0, 8), (0, 9), (1, 0), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (1, 7), (1, 8), (1, 9), (2, 0), (2, 1), (2, 3), (2, 4), (2, 5), (2, 6), (2, 7), (2, 8), (2, 9), (3, 0), (3, 1), (3, 2), (3, 4), (3, 5), (3, 6), (3, 7), (3, 8), (3, 9), (4, 0), (4, 1), (4, 2), (4, 3), (4, 5), (4, 6), (4, 7), (4, 8), (4, 9), (5, 0), (5, 1), (5, 2), (5, 3), (5, 4), (5, 6), (5, 7), (5, 8), (5, 9), (6, 0), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 7), (6, 8), (6, 9)\}$$

or

$$C^c = \{(0, 0), (1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6)\}.$$

Therefore, since the number of points in \mathcal{S} is $7 \times 10 = 70$, we have:

$$P(A) = \frac{18}{70} = \frac{9}{35} \simeq 0.257, \quad P(B) = \frac{5}{70} = \frac{1}{14} \simeq 0.071,$$


$$P(C) = \frac{63}{70} = \frac{9}{10} = 0.9, \quad \text{or} \quad P(C) = 1 - P(C^c) = 1 - \frac{7}{70} = \frac{63}{70} = 0.9.$$

Section 2.2

2.2 (i) For $0 < x \leq 2$, $f(x) = \frac{d}{dx}(2c(x^2 - \frac{1}{3}x^3)) = 2c(2x - x^2)$. Thus, $f(x) = 2c(2x - x^2)$, $0 < x \leq 2$ (and 0 elsewhere).

(ii) From $\int_0^2 2c(2x - x^2) dx = 1$, we get $\frac{8c}{3} = 1$, so that $c = 3/8$.

2.4 (i)



(ii) $P(X \leq 6.5) = 0.7$, $P(X > 8.1) = 1 - P(X \leq 8.1) = 1 - 0.9 = 0.1$,
 $P(5 < X < 8) = P(X < 8) - P(X \leq 5) = 0.7 - 0.4 = 0.3$.

2.6 (i) We need two relations which are provided by: $\int_0^1 (cx + d) dx = 1$ and $\int_{1/2}^1 (cx + d) dx = 1/3$, or: $c + 2d = 2$ and $9c + 12d = 8$, and hence $c = -\frac{4}{3}$, $d = \frac{5}{3}$.

(ii) For $0 \leq x \leq 1$, $F(x) = \int_0^x (-\frac{4}{3}t + \frac{5}{3}) dt = -\frac{2x^2}{3} + \frac{5x}{3}$. Thus,

$$F(x) = \begin{cases} 0, & x < 0 \\ -\frac{2x^2}{3} + \frac{5x}{3}, & 0 \leq x \leq 1 \\ 1, & x > 1. \end{cases}$$

2.8 From $\sum_{x=0}^{\infty} c\alpha^x = c \sum_{x=0}^{\infty} \alpha^x = c \times \frac{1}{1-\alpha} = 1$, we get $c = 1 - \alpha$.

2.10 (i) $\sum_{x=0}^{\infty} c(\frac{1}{3})^x = c[1 + \frac{1}{3} + (\frac{1}{3})^2 + \dots] = \frac{c}{1-\frac{1}{3}} = \frac{3c}{2} = 1$ and $c = \frac{2}{3}$.

(ii) $P(X \geq 3) = \frac{2}{3} \sum_{x=3}^{\infty} (\frac{1}{3})^x = \frac{2}{3} \times \frac{1/3^3}{1-\frac{1}{3}} = \frac{1}{27} \simeq 0.037$.

2.12 (i) $\int_0^{\infty} ce^{-cx} dx = -\int_0^{\infty} de^{-cx} = -e^{-cx}|_0^{\infty} = -(0-1) = 1$ for every $c > 0$.

(ii) $P(X \geq 10) = \int_{10}^{\infty} ce^{-cx} dx = -e^{-cx}|_{10}^{\infty} = -(0 - e^{-10c}) = e^{-10c}$.

(iii) $P(X \geq 10) = 0.5$ implies $e^{-10c} = \frac{1}{2}$, so that $-10c = -\log 2$ and $c = \frac{1}{10} \log 2 \simeq \frac{0.693}{10} \simeq 0.069$.

2.14 (i) From $\sum_{j=0}^{\infty} \frac{c}{3^j} = c \sum_{j=0}^{\infty} \frac{1}{3^j} = c \times \frac{1}{1-\frac{1}{3}} = \frac{3c}{2} = 1$, we get $c = \frac{2}{3}$.

(ii) $P(X \geq 3) = c \sum_{j \geq 3} \frac{1}{3^j} = c \times \frac{1/3^3}{1-\frac{1}{3}} = c \times \frac{1}{2 \times 3^2} = \frac{2}{3} \times \frac{1}{2 \times 3^2} = \frac{1}{3^3} = \frac{1}{27} \simeq 0.037$.

(iii) $P(X = 2k + 1, k = 0, 1, \dots) = c \sum_{k=0}^{\infty} \frac{1}{3^{2k+1}} = c(\frac{1}{3} + \frac{1}{3^3} + \frac{1}{3^5} + \dots) = c \times \frac{1/3}{1-\frac{1}{9}} = c \times \frac{3}{8} = \frac{2}{3} \times \frac{3}{8} = 0.25$.

(iv) $P(X = 3k + 1, k = 0, 1, \dots) = c \sum_{k=0}^{\infty} \frac{1}{3^{3k+1}} = c(\frac{1}{3} + \frac{1}{3^4} + \frac{1}{3^7} + \dots) = c \times \frac{1/3}{1-\frac{1}{27}} = c \times \frac{9}{26} = \frac{2}{3} \times \frac{9}{26} = \frac{3}{13} \simeq 0.231$.

2.16 (i) $P(\text{no items are sold}) = f(0) = \frac{1}{2} = 0.5$.

(ii) $P(\text{more than 3 items are sold}) = \sum_{x=4}^{\infty} (\frac{1}{2})^{x+1} = (\frac{1}{2})^5 \times \frac{1}{1-\frac{1}{2}} = \frac{1}{16} = 0.0625$.

(iii) $P(\text{an odd number of items are sold}) = (\frac{1}{2})^2 + (\frac{1}{2})^4 + (\frac{1}{2})^6 + \dots = (\frac{1}{2})^2 \times \frac{1}{1-\frac{1}{4}} = \frac{1}{3} \simeq 0.333$.

2.18 (i) Since $\int_0^{\infty} c^2 x e^{-cx} dx = -c x e^{-cx}|_0^{\infty} - e^{-cx}|_0^{\infty} = 1$ for all $c > 0$, the given function is a p.d.f. for all $c > 0$.

(ii) From part (i),

$$P(X \geq t) = -c x e^{-cx}|_t^{\infty} - e^{-cx}|_t^{\infty} = c(te^{-ct} + e^{-ct}) = \frac{c(t+1)}{e^{ct}}.$$

(iii) Here $c(t+1) = 0.2 \times 11 = 2.2$, $ct = 0.2 \times 10 = 2$, so that $\frac{c(t+1)}{e^{ct}} = \frac{2.2}{e^2} \simeq 0.297$.

2.20 We have:

$$\begin{aligned} P(X > x_0) &= \int_{x_0}^1 n(1-x)^{n-1} dx = -\int_{x_0}^1 d(1-x)^n \\ &= -(1-x)^n|_{x_0}^1 = (1-x_0)^n, \text{ and it is given that this probability} \end{aligned}$$

is $1/10^{2n}$. Thus,

$$(1-x_0)^n = \frac{1}{10^{2n}}, \text{ or } 1-x_0 = \frac{1}{100} \text{ and } x_0 = 0.99.$$

Section 2.3

3.2 We have: $P(A|A \cup B) = \frac{P(A \cap (A \cup B))}{P(A \cup B)} = \frac{P(A)}{P(A \cup B)} = \frac{P(A)}{P(A) + P(B)}$ (since $A \cap B = \emptyset$), and likewise, $P(B|A \cup B) = \frac{P(B \cap (A \cup B))}{P(A \cup B)} = \frac{P(B)}{P(A) + P(B)}$.

3.4 (i) $P(b_2|b_1) = 15/26 \approx 0.577$; (ii) $P(g_2|g_1) = 13/24 \approx 0.542$;

(iii) $P(b_2) = 0.52$; (iv) $P(b_1 \cap g_2) = 0.22$.

3.6 Parts (i) and (ii) follow without any calculations by using the fact that $P(\cdot|B)$ and $P(\cdot|C)$ are probability functions, or directly as follows:

$$(i) P(A^c|B) = \frac{P(A^c \cap B)}{P(B)} = \frac{P(B - A \cap B)}{P(B)} = \frac{P(B) - P(A \cap B)}{P(B)} = 1 - \frac{P(A \cap B)}{P(B)} = 1 - P(A|B).$$

$$(ii) P(A \cup B|C) = \frac{P((A \cup B) \cap C)}{P(C)} = \frac{P((A \cap C) \cup (B \cap C))}{P(C)} \\ = \frac{P(A \cap C) + P(B \cap C) - P(A \cap B \cap C)}{P(C)} = \frac{P(A \cap C)}{P(C)} + \frac{P(B \cap C)}{P(C)} - \frac{P(A \cap B \cap C)}{P(C)} \\ = P(A|C) + P(B|C) - P(A \cap B|C).$$

(iii) In the sample space $\mathcal{S} = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$ with all outcomes being equally likely, define the events:

$$A = \text{“the \# of } H\text{'s is } \leq 2\text{”} = \{TTT, TTH, THT, HTT, THH, HTH, HHT\},$$

$$B = \text{“the \# of } H\text{'s is } > 1\text{”} = \{HHT, HTH, THH, HHH\}.$$

Then $B^c = \{HTT, THT, TTH, TTT\}$, $A \cap B^c = B^c$, $A \cap B = \{HHT, HTH, THH\}$, so that:

$$P(A|B^c) = \frac{P(A \cap B^c)}{P(B^c)} = \frac{P(B^c)}{P(B^c)} = 1 \text{ and } 1 - P(A|B) = 1 - \frac{P(A \cap B)}{P(B)} = 1 - \frac{3/8}{4/8} = 1 - \frac{3}{4} = \frac{1}{4}. \text{ Thus, } P(A|B^c) \neq 1 - P(A|B).$$

(iv) In the sample space $\mathcal{S} = \{1, 2, 3, 4, 5\}$ with all outcomes being equally likely, consider the events $A = \{1, 2\}$, $B = \{3, 4\}$, and $C = \{2, 3\}$, so that $A \cap B = \emptyset$ and $A \cup B = \{1, 2, 3, 4\}$. Then:

$$P(C|A \cup B) = \frac{P(C \cap (A \cup B))}{P(A \cup B)} = \frac{2/5}{4/5} = \frac{2}{4} = \frac{1}{2}, \text{ whereas}$$

$$P(C|A) = \frac{P(A \cap C)}{P(A)} = \frac{1/5}{2/5} = \frac{1}{2}, P(C|B) = \frac{P(B \cap C)}{P(B)} = \frac{1/5}{2/5} = \frac{1}{2}, \text{ so that}$$

$$P(C|A \cup B) \neq P(C|A) + P(C|B).$$

3.8 For $n = 2$, the theorem is true since $P(A_2|A_1) = \frac{P(A_1 \cap A_2)}{P(A_1)}$ yields $P(A_1 \cap A_2) = P(A_2|A_1)P(A_1)$. Next, assume $P(A_1 \cap \dots \cap A_k) = P(A_k|A_1 \cap \dots \cap A_{k-1}) \dots P(A_2|A_1)P(A_1)$ and show that $P(A_1 \cap \dots \cap A_{k+1}) = P(A_{k+1}|A_1 \cap \dots \cap A_k)P(A_k|A_1 \cap \dots \cap A_{k-1}) \dots P(A_2|A_1)P(A_1)$. Indeed, $P(A_1 \cap \dots \cap A_{k+1}) = P((A_1 \cap \dots \cap A_k) \cap A_{k+1}) = P(A_{k+1}|A_1 \cap \dots \cap A_k)P(A_1 \cap \dots \cap A_k)$ (by applying the theorem for two events $A_1 \cap \dots \cap A_k$ and A_{k+1}) $= P(A_{k+1}|A_1 \cap \dots \cap A_k)P(A_k|A_1 \cap \dots \cap A_{k-1}) \dots P(A_2|A_1)P(A_1)$ (by the induction hypothesis).

3.10 With obvious notation, we have: $P(\text{1st white and 4th white}) = P(W_1 \cap W_2 \cap W_3 \cap W_4) + P(W_1 \cap W_2 \cap B_3 \cap W_4) + P(W_1 \cap B_2 \cap W_3 \cap W_4) + P(W_1 \cap B_1 \cap B_2 \cap W_4) = P(W_4|W_1 \cap W_2 \cap W_3)P(W_3|W_1 \cap W_2)P(W_2|W_1)P(W_1) + P(W_4|$

$$\begin{aligned}
& W_1 \cap W_2 \cap B_3)P(B_3|W_1 \cap W_2)P(W_2|W_1)P(W_1) + P(W_4|W_1 \cap B_2 \cap W_3)P(W_3| \\
& W_1 \cap B_2)P(B_2|W_1)P(W_1) + P(W_4|W_1 \cap B_1 \cap B_2)P(B_2|W_1 \cap B_1)P(B_1|W_1) \times \\
P(W_1) &= \frac{7}{12} \times \frac{8}{13} \times \frac{9}{14} \times \frac{10}{15} + \frac{8}{12} \times \frac{5}{13} \times \frac{9}{14} \times \frac{10}{15} + \frac{8}{12} \times \frac{9}{13} \times \frac{5}{14} \times \frac{10}{15} + \frac{9}{12} \times \frac{4}{13} \times \\
\frac{5}{14} \times \frac{10}{15} &= \frac{1}{12 \times 13 \times 14 \times 15} (7 \times 8 \times 9 \times 10 + 5 \times 8 \times 9 \times 10 \times 2 + 4 \times 5 \times 9 \times 10) \\
&= \frac{9 \times 10 \times 156}{12 \times 13 \times 14 \times 15} = \frac{3}{7} \approx 0.429.
\end{aligned}$$

3.12 (i) $P(+)=0.01188$; (ii) $P(D|+)=\frac{190}{1188} \approx 0.16$.

3.14 Let I = “switch I is open,” Π = “switch II is open,” S = “signal goes through.” Then: (i) $P(S)=0.48$; (ii) $P(I|S^c)=\frac{5}{13} \approx 0.385$; (iii) $P(\Pi|S^c)=\frac{10}{13} \approx 0.769$.

3.16 With F = “an individual is female,” M = “an individual is male,” C = “an individual is color-blind,” we have:
 $P(F)=0.52$, $P(M)=0.48$, $P(C|F)=0.25$, $P(C|M)=0.05$, and therefore
 $P(C)=0.154$, $P(M|C)=\frac{12}{77} \approx 0.156$.

3.18 With obvious notation, we have:

(i) $P(D)=0.029$; (ii) $P(I|D)=\frac{12}{29} \approx 0.414$; (iii) $P(\Pi|D)=\frac{9}{29} \approx 0.310$,
and $P(\text{III}|D)=\frac{8}{29} \approx 0.276$.

3.20 (i) $P(X > t) = \int_t^\infty \lambda e^{-\lambda x} dx = -\int_t^\infty de^{-\lambda x} = -e^{-\lambda x}|_t^\infty = e^{-\lambda t}$.

(ii) $P(X > s + t | X > s) = \frac{P(X > s + t, X > s)}{P(X > s)} = \frac{P(X > s + t)}{P(X > s)}$
 $= \frac{e^{-\lambda(s+t)}}{e^{-\lambda s}}$ (by part (i))
 $= e^{-\lambda t}$.

(iii) The conditional probability that X is greater than t units beyond s , given that it has been greater than s , does not depend on s and is the same as the (unconditional) probability that X is greater than t . That is, this distribution has some sort of “memoryless” property.

Section 2.4

4.2 Here $P(A) = P(A \cap A) = P(A)P(A) = [P(A)]^2$, and this happens if $P(A) = 0$, whereas, if $P(A) \neq 0$, it happens only if $P(A) = 1$.

4.4 Since $P(A_1 \cap A_2) = P(A_1)P(A_2)$, we have to show that:

$$\begin{aligned}
P(A_1 \cap (B_1 \cup B_2)) &= P(A_1)P(B_1 \cup B_2), P(A_2 \cap (B_1 \cup B_2)) \\
&= P(A_2)P(B_1 \cup B_2), P(A_1 \cap A_2 \cap (B_1 \cup B_2)) \\
&= P(A_1)P(A_2)P(B_1 \cup B_2).
\end{aligned}$$


Indeed, $P(A_1 \cap (B_1 \cup B_2)) = P((A_1 \cap B_1) \cup (A_1 \cap B_2))$

$$\begin{aligned}
&= P(A_1 \cap B_1) + P(A_1 \cap B_2) = P(A_1)P(B_1) + P(A_1)P(B_2) \\
&= P(A_1)P(B_1 \cup B_2), \text{ and similarly for } P(A_2 \cap (B_1 \cup B_2)).
\end{aligned}$$

Finally,

$$\begin{aligned} P(A_1 \cap A_2 \cap (B_1 \cup B_2)) &= P((A_1 \cap A_2 \cap B_1) \cup (A_1 \cap A_2 \cap B_2)) \\ &= P(A_1 \cap A_2 \cap B_1) + P(A_1 \cap A_2 \cap B_2) \\ &= P(A_1)P(A_2)P(B_1) + P(A_1)P(A_2)P(B_2) \\ &= P(A_1)P(A_2)P(B_1 \cup B_2). \end{aligned}$$

- 4.6 (i) Clearly, $A = (A \cap B \cap C) \cup (A \cap B^c \cap C) \cup (A \cap B \cap C^c) \cup (A \cap B^c \cap C^c)$ and hence $P(A) = 0.6875$. Likewise, $P(B) = 0.4375$, $P(C) = 0.5625$.
(ii) A , B , and C are not independent.
(iii) $P(A \cap B) = \frac{4}{16}$, and then $P(A|B) = \frac{4}{7} \simeq 0.571$.
(iv) A and B are not independent.
- 4.8 (i) $S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$, $A = \{HHH, TTT\}$ with $P(A) = p^3 + q^3$ ($q = 1 - p$).
(ii) $P(A) = 0.28$.
- 4.10 (i) $c = 1/25$.
(ii) See figure.



- (iii) $P(A) = P(X > 5) = 0.50$, $P(B) = P(52 \times 27.5) = 0.375$.
(iv) $P(B|A) = 0.75$; (v) A and B are not independent.
- 4.12 (i) $P((W_1 \cap R_1^c \cap \dots \cap R_{n-2}^c \cap R_{n-1}) \cup (W_1^c \cap W_2 \cap R_1^c \cap \dots \cap R_{n-3}^c \cap R_{n-2}) \cup \dots \cup (W_1^c \cap \dots \cap W_{n-2}^c \cap W_{n-1} \cap R_n)) = 0.54 \sum_{i=1}^{n-1} (0.1)^{i-1} (0.4)^{n-i-1}$.
(ii) For $n = 5$, the probability in part (i) is 0.0459.
- 4.14 (i) $P(\text{no circuit is closed}) = (1 - p_1) \cdots (1 - p_n)$.
(ii) $P(\text{at least 1 circuit is closed}) = 1 - (1 - p_1) \cdots (1 - p_n)$.
(iii) $P(\text{exactly 1 circuit is closed}) = p_1(1 - p_2) \cdots (1 - p_n) + (1 - p_1)p_2 \times (1 - p_3) \cdots (1 - p_n) + \dots + (1 - p_1) \cdots (1 - p_{n-1})p_n$.
(iv) The answers above are: $(1 - p)^n$, $1 - (1 - p)^n$, $np(1 - p)^{n-1}$.
(v) The numerical values are: 0.01024, 0.98976, 0.0768.

Section 2.5

5.2 (i) $3 \times 4 \times 5 = 60$; (ii) $1 \times 2 \times 5 = 10$; (iii) $3 \times 4 \times 1 = 12$.

5.4 (i) $3 \times 2 \times 3 \times 2 \times 3 = 108$; (ii) $3 \times 2 \times 2 \times 1 \times 1 = 12$.

5.6 2^n ; $2^5 = 32$, $2^{10} = 1,024$, $2^{15} = 32,768$, $2^{20} = 1,048,576$,
 $2^{25} = 33,554,432$.

5.8 The required probability is: $\frac{1}{360} \simeq 0.003$.

5.10 Start with $\frac{\binom{n+1}{m+1}}{\binom{n}{m}}$, expand in terms of factorial, do the cancellations, and you end up with $(n+1)/(m+1)$.5.12 Selecting r out of $m+n$ in $\binom{m+n}{r}$ ways is equivalent to selecting x out of m in $\binom{m}{x}$ ways and $r-x$ out of n in $\binom{n}{r-x}$ ways where $x = 0, 1, \dots, r$. Then $\binom{m+n}{r} = \sum_{x=0}^r \binom{m}{x} \binom{n}{r-x}$.5.14 The required number is $\binom{n}{3}$, which for $n = 10$ becomes $\binom{10}{3} = 120$.

5.16 The required probability is:

$$\frac{\binom{10}{2} \times \binom{15}{3} \times \binom{30}{4} \times \binom{5}{1}}{\binom{60}{10}} = \frac{2,480,625}{66,661,386} \simeq 0.037.$$

5.18 The required probability is: $\frac{\binom{n-1}{m}}{\binom{n}{m}} = 1 - \frac{m}{n}$.5.20 The required probability is: $(0.5)^{2n} \sum_{m=0}^n \binom{n}{m}^2$, which for $n = 5$, becomes:
 $252 \times (0.5)^{10} \simeq 0.246$.5.22 The required probability is: $\sum_{x=5}^{10} \binom{10}{x} (0.2)^x (0.8)^{10-x} = 0.03279$.5.24 (a) (i) $\left(\frac{n_B}{n}\right)^3$; (ii) $1 - \left(\frac{n_B+n_W}{n}\right)^3$; (iii) $\frac{6n_B n_B n_W}{n^3}$.(b) (i) $\frac{\binom{n_B}{3}}{\binom{n}{3}}$; (ii) $1 - \frac{\binom{n_B+n_W}{3}}{\binom{n}{3}}$; (iii) $\frac{\binom{n_B}{1} \binom{n_B}{1} \binom{n_W}{1}}{\binom{n}{3}}$.

Chapter 3

Section 3.1

1.2 (i) $EX = 0$, $EX^2 = c^2$, and $\text{Var}(X) = c^2$.

(ii) $P(|X - EX| \leq c) = P(-c \leq X \leq c) = P(X = -c, X = c) = 1 = \frac{c^2}{c^2} = \frac{\text{Var}(X)}{c^2}$.

1.4 If Y is the net loss to the company, then $EY = \$600$, and if P is the premium to be charged, then $P = \$700$.

- 1.6 $\text{Var}(X) = EX^2 - (EX)^2$. by expanding and taking expectations. Also, $E[X(X-1)] = \text{Var}(X) + (EX)^2 - EX$ by expanding, taking expectations, and using the first result. That $\text{Var}(X) = E[X(X-1)] + EX - (EX)^2$ follows from the first two results.
- 1.8 (i) $EX = 2$, $E[X(X-1)] = 4$; (ii) $\text{Var}(X) = 2$.
- 1.10 $EX = \frac{4}{3}$, $EX^2 = 2$, so that $\text{Var}(X) = \frac{2}{9}$ and s.d. of $X = \frac{\sqrt{2}}{3} \simeq 0.471$.
- 1.12 $c_1 = -1/12$, $c_2 = 5/3$.
- 1.14 (i) by adding and subtracting μ , we get: $E(X-c)^2 = \text{Var}(X) + (\mu-c)^2$;
(ii) Immediate from part (i).
- 1.16 (i) $\int_{-\infty}^{\infty} \frac{dx}{1+x^2} = \arctan x|_{-\infty}^{\infty} = \arctan(\infty) - \arctan(-\infty) = \pi$, so that $\int_{-\infty}^{\infty} \frac{1}{\pi} \times \frac{dx}{1+x^2} = 1$.
(ii) $\frac{1}{\pi} \int_{-\infty}^{\infty} x \times \frac{dx}{1+x^2} = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{d(1+x^2)}{1+x^2} = \frac{1}{2\pi} \log(1+x^2)|_{-\infty}^{\infty} = \frac{1}{2\pi}(\infty - \infty)$.
- 1.18 For the discrete case, $X \geq c$ means $x_i \geq c$ for all values x_i of X . Then $x_i f_X(x_i) \geq c f_X(x_i)$ and hence $\sum_{x_i} x_i f_X(x_i) \geq \sum_{x_i} c f_X(x_i)$. But $\sum_{x_i} x_i f_X(x_i) = EX$ and $\sum_{x_i} c f_X(x_i) = c \sum_{x_i} f_X(x_i) = c$. Thus, $EX \geq c$. The particular case follows, of course, by taking $c = 0$. In the continuous case, summation signs are replaced by integrals.

Section 3.2

- 2.2 (i) $c = \sigma/\sqrt{1-\alpha}$; (ii) $c = \frac{1}{\sqrt{0.05}} \simeq 4.464$.
- 2.4 (i) By the Tchebichev inequality, $P(|X - \mu| \geq c) = 0$ for all $c > 0$.
(ii) Consider a sequence $0 < c_n \downarrow 0$ as $n \rightarrow \infty$. Then $P(|X - \mu| \geq c_n) = 0$ for all n , or equivalently, $P(|X - \mu| < c_n) = 1$ for all n , whereas, clearly, $\{|X - \mu| < c_n\}$ is a nonincreasing sequence of events and its limits is $\cap_{n=1}^{\infty} \{|X - \mu| < c_n\}$. Then, by Theorem 2 in Chapter 2, $1 = \lim_{n \rightarrow \infty} P(|X - \mu| < c_n) = P(\cap_{n=1}^{\infty} \{|X - \mu| < c_n\})$. However, it is clear that $\cap_{n=1}^{\infty} \{|X - \mu| < c_n\} = \{|X - \mu| \leq 0\} = \{X = \mu\}$. Thus, $P(X = \mu) = 1$, as was to be seen.

Section 3.3

- 3.2 (i) It follows by using the identity $\binom{n+1}{x} = \binom{n}{x} + \binom{n}{x-1}$.
(ii) $B(26, 0.25; 10) = 0.050725$.
- 3.4 If X is the number of those favoring the proposal, then $X \sim B(15, 0.4375)$.
Therefore: (i) $P(X \geq 5) = 0.859$; (ii) $P(X \geq 8) = 0.3106$.
- 3.6 If X is the number of times the bull's eye is hit, then $X \sim B(100, p)$.
Therefore:
(i) $P(X \geq 40) = \sum_{x=40}^{100} \binom{100}{x} p^x q^{100-x}$ ($q = 1 - p$).
(ii) $P(X \geq 40) = \sum_{x=40}^{100} \binom{100}{x} (0.25)^x (0.75)^{100-x}$.

- (iii) $EX = np = 100p$, $Var(X) = npq = 100pq$, and for $p = 0.25$,
 $EX = 25$, $Var(X) = 18.75$, s.d. of $X = \sqrt{18.75} \simeq 4.33$.

3.8 From the Tchebichev inequality $n = 8,000$.

- 3.10 (i) Writing $\binom{n}{x}$ in terms of factorials, and after cancellations, we get:

$$EX = np \sum_{y=0}^{n-1} \binom{n-1}{y} p^y q^{(n-1)-y} = np \times 1 = np. \text{ Likewise,}$$

$$E[X(X-1)] = n(n-1)p^2 \sum_{y=0}^{n-2} \binom{n-2}{y} p^y q^{(n-2)-y} = n(n-1)p^2 \times 1$$

$$= n(n-1)p^2.$$

- (ii) From Exercise 1.6, $Var(X) = n(n-1)p^2 + np - (np)^2 = npq$.

- 3.12 (i) $P(X \leq 10) = 1 - q^{10}$; (ii) $1 - (0.8)^{10} \simeq 0.893$.

3.14 If X is the number of tosses to the first success, then X has the Geometric distribution with $p = 1/6$. Then:

(i) $P(X = 3) = \frac{25}{216} \simeq 0.116$;

(ii) $P(X \geq 5) = \left(\frac{5}{6}\right)^4 \simeq 0.482$.

- 3.16 (i) $EX = \frac{1}{p}$, $E[X(X-1)] = \frac{2q}{p^2}$; (ii) $Var(X) = \frac{q}{p^2}$.

3.18 $\lambda = 2$.

3.20 $f(x+1) = e^{-\lambda} \frac{\lambda^{x+1}}{(x+1)!} = \frac{\lambda}{x+1} \times e^{-\lambda} \frac{\lambda^x}{x!} = \frac{\lambda}{x+1} f(x)$.

- 3.22 (i) $M_X(t) = e^{\lambda(e^t-1)}$, $t \in \mathfrak{R}$, by applying the definition of M_X .

(ii) $EX = \frac{d}{dt} M_X(t)|_{t=0} = \lambda$, $EX^2 = \frac{d^2}{dt^2} M_X(t)|_{t=0} = \lambda(\lambda+1)$, so that
 $Var(X) = \lambda$.

- 3.24 (i) $\frac{\binom{70}{5} \binom{10}{0}}{\binom{80}{5}} \simeq 0.503$; (ii) $\frac{1}{\binom{80}{5}} [\binom{70}{3} \binom{10}{2} + \binom{70}{4} \binom{10}{1} + \binom{70}{5} \binom{10}{0}] \simeq 0.987$.

3.26 Writing the combinations in terms of factorials, and recombining the terms, we get the result.

- 3.28 (i) By integration, using the definition of $\Gamma(\alpha)$ and the recursive relation for $\Gamma(\alpha+1)$, we get $EX = \frac{\beta}{\Gamma(\alpha)} \Gamma(\alpha+1) = \alpha\beta$. Likewise, $EX^2 = \frac{\beta^2}{\Gamma(\alpha)} \Gamma(\alpha+2) = \alpha(\alpha+1)\beta^2$, so that $Var(X) = \alpha\beta^2$.

(ii) $EX = 1/\lambda$, $Var(X) = 1/\lambda^2$ from part (i).

(iii) $EX = r$, $Var(X) = 2r$ from part (i).

- 3.30 (i) (a) With $g(X) = cX$, we have $Eg(X) = c/\lambda$.

(b) With $g(X) = c(1 - 0.5e^{-\alpha X})$, we have $Eg(X) = \frac{(\alpha+0.5\lambda)c}{\alpha+\lambda}$.

(ii) (a) 10; (b) 1.5.

3.32 Indeed, $P(T > t) = P(0 \text{ events occurred in the time interval } (0, t)) = e^{-\lambda t} \frac{(\lambda t)^0}{0!} = e^{-\lambda t}$. So, $1 - F_T(t) = e^{-\lambda t}$, $t > 0$, and hence $f_T(t) = \lambda e^{-\lambda t}$, $t > 0$, and T is as described.

3.34 (i) $\int_0^\infty \alpha \beta x^{\beta-1} e^{-\alpha x^\beta} dx = -\int_0^\infty de^{-\alpha x^\beta} = -e^{-\alpha x^\beta} \Big|_0^\infty = 1$.

(ii) $\beta = 1$ and any $\alpha > 0$.

(iii) For $n = 1, 2, \dots$, $EX^n = \Gamma(\frac{n}{\beta} + 1)/\alpha^{n/\beta}$. Then $EX = \Gamma(\frac{1}{\beta} + 1)/\alpha^{1/\beta}$, $EX^2 = \Gamma(\frac{2}{\beta} + 1)/\alpha^{2/\beta}$, and hence $\text{Var}(X) = \{\Gamma(\frac{2}{\beta} + 1) - [\Gamma(\frac{1}{\beta} + 1)]^2\}/\alpha^{2/\beta}$.

3.36 All parts (i)–(iv) are immediate.

3.38 (i) $P(X \leq c) = 0.875$, and $c = \mu + 1.15\sigma$; (ii) $c = 7.30$.

3.40 (i) 0.997020; (ii) 0.10565; (iii) 0.532807.

3.42 Let X be the diameter of a ball bearing, and let p be the probability that a ball bearing is defective. Then $p = P(X > 0.5 + 0.0006 \text{ or } X < 0.5 - 0.0006)$, and by splitting the probabilities and normalizing, we have that it is 0.583921.

3.44 (i) By the hint,

$$\begin{aligned} I^2 &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-\frac{x^2+y^2}{2}} dx dy = \frac{1}{2\pi} \left(\int_0^{2\pi} d\theta \right) \left(\int_0^{\infty} r e^{-r^2/2} dr \right) \\ &= \frac{1}{2\pi} \times 2\pi (-e^{-r^2/2} \Big|_0^{\infty}) = 1. \end{aligned}$$

$$(ii) \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} e^{-y^2/2} \sigma dy = I = 1.$$

3.46 (i) $M_X(t) = e^{t^2/2} \times \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(x-t)^2}{2}} dt = e^{t^2/2} \times 1 = e^{t^2/2}$, $t \in \mathfrak{R}$.

(ii) With $Z = \frac{X-\mu}{\sigma}$, $e^{t^2/2} = M_Z(t) = M_{\frac{1}{\sigma}X + \frac{\mu}{\sigma}}(t) = e^{-\frac{t\mu}{\sigma}} M_X\left(\frac{t}{\sigma}\right)$, so that $M_X\left(\frac{t}{\sigma}\right) = e^{\frac{t\mu}{\sigma} + \frac{t^2}{2}}$, and $M_X(t) = e^{\mu t + \frac{\sigma^2 t^2}{2}}$ by replacing $\frac{t}{\sigma}$ by $t \in \mathfrak{R}$.

(iii) By differentiation of $M_X(t)$ and evaluating at 0, we get: $EX = \mu$, $EX^2 = \mu^2 + \sigma^2$, so that $\text{Var}(X) = \sigma^2$.

3.48 (i) $EX^{2n+1} = 0$, and by the hint $EX^{2n} = (2n-1)(2n-3)\cdots 1 = \frac{1 \times 2 \times \cdots \times (2n-1) \times (2n)}{(2 \times 1) \times \cdots \times [2 \times (n-1)] \times (2 \times n)} = \frac{(2n)!}{2^n [1 \cdots (n-1)n]} = \frac{(2n)!}{2^n (n!)}$.

(ii) $EX = 0$, $EX^2 = 1$, so that $\text{Var}(X) = 1$.

(iii) With $Z = \frac{X-\mu}{\sigma}$, $0 = EZ = \frac{1}{\sigma}(EX - \mu)$, so that $EX = \mu$, and $1 = \text{Var}(Z) = \frac{1}{\sigma^2} \text{Var}(X)$, so that $\text{Var}(X) = \sigma^2$.

3.50 (i) $P(-1 < X < 2) = \frac{3}{2\alpha} = 0.75$ and $\alpha = 2$.

(ii) $P(|X| < 1) = P(|X| > 2)$ is equivalent to $\frac{1}{\alpha} = 1 - \frac{2}{\alpha}$ from which $\alpha = 3$.

3.52 $EX = \frac{1}{3}$, so that: (i) -0.5 and (ii) $2(e-1) \simeq 3.44$.

Section 3.4

4.2 (i) $x_p = [(n+1)p]^{1/(n+1)}$; (ii) For $p = 0.5$ and $n = 3$: $x_{0.5} = 2^{1/4} \simeq 1.189$.

4.4 (i) $c_1 = c_2 = 1$; (ii) $x_{1/3} = 0$.

4.6 (i)

x	2	3	4	5	6	7	8	9	10	11	12
$f(x)$	1/36	2/36	3/36	4/36	5/36	6/36	5/36	4/36	3/36	2/36	1/36

(ii) $EX = 7$; (iii) median = mode = mean = 7.4.8 Mode = 25 and $f(25) = \binom{100}{25}(\frac{1}{4})^{25}(\frac{3}{4})^{75}$; one would bet on $X = 25$.4.10 By the hint, $P(X \leq c) = \int_{-\infty}^c f(x) dx = \int_0^{\infty} f(c-y) dy$, and $P(X \geq c) = \int_c^{\infty} f(x) dx = \int_0^{\infty} f(c+y) dy$. Since $f(c-y) = f(c+y)$, it follows that $P(X \leq c) = P(X \geq c)$, and hence c is the median.4.12 (i) $p = P(Y \leq y_p) = P[g(X) \leq y_p] = P[X \leq g^{-1}(y_p)]$, so that $g^{-1}(y_p) = x_p$ and $y_p = g(x_p)$.(ii) $x_p = -\log(1-p)$.(iii) $y_p = 1/(1-p)$.(iv) $x_{0.5} = -\log(0.5) \simeq 0.693$, and $y_{0.5} = 2$.

Chapter 4

Section 4.1

1.2 $P(X=0, Y=1) = P(X=0, Y=2) = P(X=1, Y=2) = 0$,
 $P(X=0, Y=0) = 0.3$, $P(X=1, Y=0) = 0.2$, $P(X=1, Y=1) = 0.2$,
 $P(X=2, Y=0) = 0.075$, $P(X=2, Y=1) = 0.15$, $P(X=2, Y=2) = 0.075$.1.4 (i) $\int_0^2 \int_0^1 (x^2 + \frac{xy}{2}) dx dy = \frac{6}{7} \times \frac{7}{6} = 1$; $P(X > Y) = \frac{15}{56} \simeq 0.268$.1.6 (i) $P(X \leq x) = 1 - e^{-x}$, $x > 0$; (ii) $P(Y \leq y) = 1 - e^{-y}$, $y > 0$;
(iii) $P(X < Y) = 0.5$; (iv) $P(X+Y < 3) = 1 - 4e^{-3} \simeq 0.801$.1.8 $c = 1/\sqrt{2\pi}$.1.10 $c = 6/7$.

Section 4.2

2.2 $f_X(0) = 0.3$, $f_X(1) = 0.4$, $f_X(2) = 0.3$; $f_Y(0) = 0.575$, $f_Y(1) = 0.35$, $f_Y(2) = 0.075$.2.4 (i) $f_X(1) = 7/36$, $f_X(2) = 17/36$, $f_X(3) = 12/36$; $f_Y(1) = 7/36$, $f_Y(2) = 14/36$, $f_Y(3) = 15/36$.(ii) $f_{X|Y}(1|1) = 2/7$, $f_{X|Y}(2|1) = 2/7$, $f_{X|Y}(3|1) = 3/7$; $f_{X|Y}(1|2) = 1/14$, $f_{X|Y}(2|2) = 10/14$, $f_{X|Y}(3|2) = 3/14$;

$$f_{X|Y}(1|3) = 4/15, f_{X|Y}(2|3) = 5/15, f_{X|Y}(3|3) = 6/15;$$

$$f_{Y|X}(1|1) = 2/7, f_{Y|X}(2|1) = 2/7, f_{Y|X}(3|1) = 3/7;$$

$$f_{Y|X}(1|2) = 2/17, f_{Y|X}(2|2) = 10/17, f_{Y|X}(3|2) = 5/17;$$

$$f_{Y|X}(1|3) = 3/12, f_{Y|X}(2|3) = 3/12, f_{Y|X}(3|3) = 6/12.$$

$$2.6 \quad (i) \quad f_X(x) = \frac{2x}{n(n+1)}, \quad x = 1, \dots, n; \quad f_Y(y) = \frac{2(n-y+1)}{n(n+1)}, \quad y = 1, \dots, n;$$

$$(ii) \quad f_{X|Y}(x|y) = \frac{1}{n-y+1}, \quad x = 1, \dots, n; \quad f_{Y|X}(y|x) = \frac{1}{x}, \quad y = 1, \dots, x;$$

$$y = 1, \dots, x \qquad \qquad \qquad x = 1, \dots, n.$$

$$(iii) \quad E(X|Y = y) = \frac{n(n+1) - (y-1)y}{2(n-y+1)}, \quad y = 1, \dots, n;$$

$$E(Y|X = x) = \frac{x+1}{2}, \quad x = 1, \dots, n.$$

$$2.8 \quad f_X(x) = \frac{6}{5}x + \frac{2}{5}, \quad 0 \leq x \leq 1; \quad f_Y(y) = \frac{6}{5}y^2 + \frac{3}{5}, \quad 0 \leq y \leq 1.$$

$$2.10 \quad (i) \quad f_X(x) = xe^{-x}, \quad x > 0; \quad f_Y(y) = e^{-y}, \quad y > 0.$$

$$(ii) \quad f_{Y|X}(y|x) = e^{-y}, \quad x > 0, \quad y > 0;$$

$$(iii) \quad P(X > \log 4) = \frac{1 + \log 4}{4} \simeq 0.597.$$

$$2.12 \quad (i) \quad f_X(x) = \frac{6x}{7}(2x+1), \quad 0 < x \leq 1; \quad f_Y(y) = \frac{3y}{14} + \frac{2}{7}, \quad 0 \leq y \leq 2;$$

$$f_{Y|X}(y|x) = \frac{2x+y}{4x+2}, \quad 0 < x \leq 1, \quad 0 \leq y \leq 2.$$

$$(ii) \quad EY = \frac{8}{7}; \quad E(Y|X = x) = \frac{2}{3} \times \frac{3x+2}{2x+1}, \quad 0 < x \leq 1.$$

(iii) It follows by a direct integration.

$$(iv) \quad P(Y > \frac{1}{2}|X < \frac{1}{2}) = \frac{207}{280} \simeq 0.739.$$

$$2.14 \quad f_{X|Y}(x|y) = \frac{1}{2}ye^{\frac{y}{2}}e^{-\frac{y}{2}x}, \quad 0 < y < x.$$

2.16 (i)

$$f_X(x) = \begin{cases} 6x/7, & 0 < x \leq 1 \\ 6x(2-x)/7, & 1 < x < 2 \\ 0, & \text{elsewhere.} \end{cases}$$

(ii) $f_Y(y|x)$ is 1 for $0 < x < 1$, and is $1/(2-x)$ for $1 < x < 2$ (and 0 otherwise), whereas $1 \leq x + y < 2$.

2.18 (i) $f_{X|Y}(\cdot|y)$ is the Poisson p.d.f. with parameter y .

$$(ii) \quad f_{X,Y}(x, y) = e^{-2y} \frac{y^x}{x!}, \quad x = 0, 1, \dots;$$

$$(iii) \quad f_X(x) = \frac{1}{2^{x+1}}, \quad x = 0, 1, \dots$$

2.20 (i), (ii) follow by applying the definitions.

Section 4.3

3.2 It follows by an application of the definition and properties of a m.g.f.

3.4 Apply the exercise cited in the hint with $Z = X - Y$ and $Z = X + Y$.

- 3.6 (i) $EX = 1, EY = 0.5, EX^2 = 1.6, EY^2 = 0.65$, so that $\text{Var}(X) = 0.6$ and $\text{Var}(Y) = 0.4$.
- (ii) $E(XY) = 0.8$, so that $\text{Cov}(X, Y) = 0.3$ and $\rho(X, Y) = 1.25\sqrt{0.24} \simeq 0.613$.
- (iii) The r.v.'s X and Y are positively correlated.
- 3.8 (i) $EX = \frac{77}{36}, EY = \frac{20}{9}, EX^2 = \frac{183}{36}, EY^2 = \frac{99}{18}$, so that $\text{Var}(X) = 659/36^2$ and $\text{Var}(Y) = 728/36^2$.
- (ii) $E(XY) = \frac{171}{36}$, so that $\text{Cov}(X, Y) = -\frac{4}{36^2}$, and $\rho(X, Y) = -\frac{2}{\sqrt{182 \times 659}} \simeq -0.006$.
- 3.10 $EX = 0, \text{Var}(X) = 10/4, EY = 5/2, EY^2 = 34/4, \text{Var}(Y) = 9/4, E(XY) = 0$, so that $\text{Cov}(X, Y) = 0$ and $\rho(X, Y) = 0$.
- 3.12 (i) $EX = EY = 7/12$;
- (ii) $EX^2 = EY^2 = 5/12$, so that $\text{Var}(X) = \text{Var}(Y) = 11/144$.
- (iii) $E(XY) = \frac{1}{3}$, so that $\text{Cov}(X, Y) = -\frac{1}{144}$ and $\rho(X, Y) = -\frac{1}{11}$.
- (iv) X and Y are negatively correlated.
- 3.14 With $\text{Var}(X) = \sigma^2$, we get $\text{Cov}(X, Y) = a\sigma^2$ and $\rho(X, Y) = \frac{a}{|a|}$. Thus, $|\rho(X, Y)| = 1$, and $\rho(X, Y) = 1$ if and only if $a > 0$, and $\rho(X, Y) = -1$ if and only if $a < 0$.
- 3.16 By differentiation, with respect to α and β , of the function $g(\alpha, \beta) = E[Y - (\alpha X + \beta)]^2$, and by equating the derivatives to 0, we find: $\hat{\alpha} = \frac{\sigma_Y}{\sigma_X} \rho(X, Y)$, $\hat{\beta} = EY - \hat{\alpha}EX$. The 2×2 matrix M of the second-order derivatives is given by: $M = 4 \begin{pmatrix} EX^2 & EX \\ EX & 1 \end{pmatrix}$, which is positive definite. Then $\hat{\alpha}$ and $\hat{\beta}$ are minimizing values.

Section 4.4

4.2 $M_{X_1, X_2, X_3}(t_1, t_2, t_3) = c^3 / (c - t_1)(c - t_2)(c - t_3)$, provided t_1, t_2, t_3 are $< c$.

4.4 Follows by applying properties of expectations.

Section 4.5

5.2 If X_1, X_2 , and X_3 are the numbers of customers buying brand A , brand B , or just browsing, then X_1, X_2, X_3 have the Multinomial distribution with parameters $n = 10, p_1 = 0.25, p_2 = 0.40$, and $p_3 = 0.35$. Therefore:

(i) $P(X_1 = 2, X_2 = 3, X_3 = 5) = \frac{10!}{2!3!5!} (0.25)^2 \times (0.40)^3 \times (0.35)^5 \simeq 0.053$.

(ii) $P(X_1 = 1, X_2 = 3 | X_3 = 6) = \frac{4!}{1!3!} \left(\frac{5}{13}\right)^1 \left(\frac{8}{13}\right)^3 \simeq 0.358$.

5.4 They follow by taking the appropriate derivatives and evaluating them at 0.

5.6 The second line in (51) follows from the first line by adding and subtracting the quantity $\rho^2 \left(\frac{x_1 - \mu_1}{\sigma_1}\right)^2$. The expression in the following line follows by the fact that the first three terms on the previous line form a perfect

square. What follows is obvious and results from a suitable regrouping of the entities involved.

- 5.8 In Exercise 5.7, it was found that $E(XY) = \mu_1\mu_2 + \rho\sigma_1\sigma_2$, where $\mu_1 = EX$, $\mu_2 = EY$, $\sigma_1 = \text{s.d. of } X$ and $\sigma_2 = \text{s.d. of } Y$. Then

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_1\sigma_2} = \frac{(\mu_1\mu_2 + \rho\sigma_1\sigma_2) - \mu_1\mu_2}{\sigma_1\sigma_2} = \rho.$$

- 5.10 They follow by differentiating the m.g.f. and evaluating the derivatives at 0.

- 5.12 (i) It is not a Bivariate Normal p.d.f., because, for x and y outside the interval $[-1, 1]$, the given p.d.f. becomes $f(x, y) = \frac{1}{2\pi} \exp[-(x^2 + y^2)/2]$, which is the Bivariate Normal with $\mu_1 = \mu_2 = 0$, $\sigma_1 = \sigma_2 = 1$, and $\rho = 0$, whereas $f(-1, -1) = \frac{1}{\pi e} \neq \frac{1}{2\pi e}$, the value of the Bivariate Normal just mentioned evaluated at $x = y = -1$.
- (ii) $f_2(y) = \frac{1}{\sqrt{2\pi}} e^{-y^2/2}$, which is the p.d.f. of the $N(0, 1)$ distribution. Similarly, $f_1(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$.

Chapter 5

Section 5.1

- 1.2 The relation $f_{X,Y}(x, y) = f_X(x)f_Y(y)$ holds true for all values of x and y , and therefore X and Y are independent.
- 1.4 The r.v.'s X and Y are not independent, since, e.g., $f_{X,Y}(0.1, 0.1) = 0.132 \neq 0.31824 = 0.52 \times 0.612 = f_X(0.1)f_Y(0.1)$.
- 1.6 (i) $f_X(x) = \frac{6}{5}(x^2 + \frac{1}{2})$, $0 \leq x \leq 1$; $f_Y(y) = \frac{6}{5}(y + \frac{1}{3})$, $0 \leq y \leq 1$.
- (ii) The r.v.'s are not independent, since, e.g., $f_{X,Y}(\frac{1}{2}, \frac{1}{4}) = \frac{3}{5} \neq \frac{9}{10} \times \frac{7}{10} = f_X(\frac{1}{2})f_Y(\frac{1}{4})$.
- 1.8 (i) $f_X(x) = 2x$, $0 < x < 1$; $f_Y(y) = 2y$, $0 < y < 1$; $f_Z(z) = 2z$, $0 < z < 1$.
- (ii) The r.v.'s are independent because, clearly,

$$f_{X,Y,Z}(x, y, z) = f_X(x)f_Y(y)f_Z(z),$$

- (iii) $P(X < Y < Z) = 1/6$.
- 1.10 (i) c can be any positive constant.
- (ii) $f_{X,Y}(x, y) = c^2 e^{-cx-cy}$, $x > 0$, $y > 0$, and likewise for $f_{X,Z}$ and $f_{Y,Z}$.
- (iii) $f_X(x) = ce^{-cx}$, $x > 0$, and likewise for f_Y and f_Z .
- (iv) The r.v.'s X and Y are independent, and likewise for the r.v.'s X, Z and Y, Z . Finally, from part (iii), it follows that the r.v.'s X, Y , and Z are also independent.

- 1.12 (i) $EX = 200$ days; (ii) $M_{X+Y}(t) = 1/(1 - 200t)^2$, $t < 0.005$, and $f_{X+Y}(t) = (0.005)^2 t e^{-0.005t}$, $t > 0$.
 (iii) $P(X + Y > 500) = 2.5e^{-2.5} + e^{-2.5} \simeq 0.287$.
- 1.14 (i) $M_U(t) = \exp[(a\mu_1 + b)t + \frac{(a\sigma_1)^2 t^2}{2}]$ which is the m.g.f. of the $N(a\mu_1 + b, (a\sigma_1)^2)$ distribution. Likewise for V .
 (ii) $M_{U,V}(t_1, t_2) = \exp[(a\mu_1 + b)t_1 + \frac{(a\sigma_1)^2 t_1^2}{2} + (c\mu_2 + d)t_2 + \frac{(c\sigma_2)^2 t_2^2}{2}]$.
 (iii) Follows from parts (i) and (ii), since $M_U(t_1)M_V(t_2) = M_{U,V}(t_1, t_2)$ for all t_1, t_2 .
- 1.16 $M_{\bar{X}}(t) = [M(\frac{t}{n})]^n$,
- 1.18 (i) $E\bar{X} = p$ and $\text{Var}(\bar{X}) = pq/n$; (ii) $n = 10,000$.
- 1.20 (i) $f_X(-1) = 2\alpha + \beta$, $f_X(0) = 2\beta$, $f_X(1) = 2\alpha + \beta$;
 $f_Y(-1) = 2\alpha + \beta$, $f_Y(0) = 2\beta$, $f_Y(1) = 2\alpha + \beta$.
 (ii) $EX = EY = 0$, and $E(XY) = 0$; (iii) $\text{Cov}(X, Y) = 0$.
 (iv) The r.v.'s are not independent, since, e.g., $f(0, 0) = 0 \neq (2\beta) \times (2\beta) = f_X(0)f_Y(0)$.
- 1.22 (i) $E\bar{X} = \mu$ and $\text{Var}(\bar{X}) = \sigma^2/n$.
 (ii) For $k = 1$, $n = 100$; for $k = 2$, $n = 25$; and for $k = 3$, $n = 12$.
- 1.24 (i) $E\bar{X} = \mu$ and $\text{Var}(\bar{X}) = \sigma^2/n$.
 (ii) The smallest n which is $\geq 1/(1 - \alpha)c^2$.
 (iii) For $c = 0.1$, the required n is its smallest value $\geq 100/(1 - \alpha)$. For $\alpha = 0.90$, $n = 1,000$; for $\alpha = 0.95$, $n = 2,000$; for $\alpha = 0.99$, $n = 10,000$.

Section 5.2

- 2.2 $X + Y \sim B(30, 1/6)$ and $P(X + Y \leq 10) = \sum_{t=0}^{10} \binom{30}{t} (\frac{1}{6})^t (\frac{5}{6})^{30-t}$.
- 2.4 (i) $S_n \sim B(n, p)$;
 (ii) $EX_i = p$, $\text{Var}(X_i) = pq$ ($q = 1 - p$).
 (iii) $ES_n = np$, $\text{Var}(S_n) = npq$.
- 2.6 If X be the r.v. denoting the breakdown voltage, then $X \sim N(40, 1.5^2)$, and therefore:
 (i) $P(39 < X < 42) = 0.656812$; (ii) $= 0.382$.
- 2.8 (i) $X_1 + \dots + X_n \sim P(\lambda_1 + \dots + \lambda_n)$.
 (ii) $E\bar{X} = (\lambda_1 + \dots + \lambda_n)/n$, $\text{Var}(\bar{X}) = (\lambda_1 + \dots + \lambda_n)/n^2$.
 (iii) $E\bar{X} = \lambda$, $\text{Var}(\bar{X}) = \lambda/n$.
- 2.10 (i) $P(X_1 = x_1 | T = t) = \binom{t}{x_1} (\frac{\lambda_1}{\lambda})^{x_1} (1 - \frac{\lambda_1}{\lambda})^{t-x_1}$, so the $X_1 | T = t \sim B(t, \frac{\lambda_1}{\lambda})$, and likewise for the other r.v.'s.
 (ii) Here $\lambda = nc$, and therefore $X_1 | T = t \sim B(t, \frac{1}{n})$.

$$2.12 \quad (\text{i}) P(\bar{X} > \bar{Y}) = 1 - \Phi\left(-\frac{\mu_1 - \mu_2}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}}\right); \quad (\text{ii}) P(\bar{X} > \bar{Y}) = 0.5.$$

$$2.14 \quad (\text{i}) \text{ For } r > 0, P(R \leq r) = P(U \leq r^2/\sigma^2), U \sim \chi_2^2.$$

(ii) For $\sigma = 1$ and the given values of r , the respective probabilities are: 0.75, 0.90, 0.95, 0.975, 0.99, and 0.995.

Chapter 6

Section 6.1

$$1.2 \quad (\text{i}) X \sim N\left(\frac{\mu-160}{9}, \frac{25\sigma^2}{81}\right); \quad (\text{ii}) a \simeq 32.222, b = 35.$$

$$(\text{iii}) a_k = \frac{5\mu-160}{9} - k\frac{5\sigma}{9}, b_k = \frac{5\mu-160}{9} + k\frac{5\sigma}{9}.$$

$$1.4 \quad f_Y(y) = \lambda y^{-(\lambda+1)}, y > 1; f_Z(z) = \lambda e^{z-\lambda e^z}, z \in \mathfrak{R}.$$

$$1.6 \quad (\text{i}) f_Y(y) = \frac{1}{2}e^{-y/2}, y > 0, \text{ which is the p.d.f. of a } \chi_2^2.$$

(ii) $\sum_{i=1}^n Y_i \sim \chi_{2n}^2$, since $Y_i \sim \chi_2^2, i = 1, \dots, n$ independent.

$$1.8 \quad f_Y(y) = \frac{1}{\Gamma(\frac{3}{2})m^{3/2}} y^{\frac{3}{2}-1} e^{-y/m}, y > 0.$$

1.10 The results follow by implementing the suggestions in the hint.

1.12 The results follow by implementing the suggestions in the hint.

Section 6.2

$$2.2 \quad (\text{i}) f_{U,V}(u, v) = \frac{u}{(1+v)^2} e^{-u}, u > 0, v > 0.$$

$$(\text{ii}) f_U(u) = u e^{-u}, u > 0; f_V(v) = 1/(1+v)^2, v > 0.$$

(iii) U and V are independent.

$$2.4 \quad (\text{i}) f_{U,V}(u, v) = \frac{1}{|ac|} f_X\left(\frac{u-b}{a}\right) f_Y\left(\frac{v-d}{c}\right), (u, v) \in T.$$

(ii) $f_{U,V}(u, v) = \frac{1}{\sqrt{2\pi}|a|\sigma_1} \exp\left\{-\frac{[u-(a\mu_1+b)]^2}{2(a\sigma_1)^2}\right\} \times \frac{1}{\sqrt{2\pi}|c|\sigma_2} \exp\left\{-\frac{[v-(c\mu_2+d)]^2}{2(c\sigma_2)^2}\right\}$, and therefore U and V are independently distributed as $N(a\mu_1 + b, (a\sigma_1)^2)$ and $N(c\mu_2 + d, (c\sigma_2)^2)$, respectively.

$$2.6 \quad (\text{i}) f_{U,V}(u, v) = \frac{1}{\sqrt{2\pi}} e^{-u^2/2} \times \frac{1}{\sqrt{2\pi}} e^{-v^2/2}, u, v \in \mathfrak{R}.$$

(ii) $U \sim N(0, 1), V \sim N(0, 1)$.

(iii) U and V are independent.

(iv) By parts (ii) and (iii), $X + Y \sim N(0, 2)$ and $X - Y \sim N(0, 2)$.

$$2.8 \quad f_U(u) = 1, 0 \leq u \leq 1.$$

Section 6.3

- 3.2 It follows by forming the inner products of the row vectors.
- 3.4 It follows from the joint p.d.f. $f_{X,Y}$, the transformation $u = \frac{x-\mu_1}{\sigma_1}$, $v = \frac{y-\mu_2}{\sigma_2}$, and the fact that the Jacobian $J = \sigma_1\sigma_2$.
- 3.6 (i) It follows from the joint p.d.f. $f_{X,Y}$, the transformation $u = x + y$, $v = x - y$, and the fact that the Jacobian $J = -1/2$.
 (ii) U and V are independent by the fact that they have the Bivariate Normal distribution and their correlation coefficient is 0.
 (iii) It follows from part (i) as marginals of the Bivariate Normal.
- 3.8 (i) $P(a\mu < \bar{X} < b\mu, 0 < S^2 < c\sigma^2) = [\Phi(k(b-1)\sqrt{n}) - \Phi(k(a-1)\sqrt{n})] \times P(\chi_{n-1}^2 < c(n-1))$; (ii) The probability is 0.89757.

Section 6.5

- 5.2 $EY_1 = \frac{1}{n+1}$, $EY_n = \frac{n}{n+1}$, and $EY_n \rightarrow 1$ as $n \rightarrow \infty$.
- 5.4 $E(Y_1Y_n) = \frac{1}{n+2}$. Therefore, by Exercise 5.2, $\text{Cov}(Y_1, Y_n) = \frac{1}{(n+1)^2(n+2)}$.
- 5.6 $f_Z(z) = \lambda e^{-\lambda z}$, $z > 0$.
- 5.8 (i) $g_n(y_n) = n\lambda e^{-\lambda y_n} (1 - e^{-\lambda y_n})^{n-1}$, $y_n > 0$.
 (ii) For $n = 2$, $EY_2 = 3/2\lambda$, and for $n = 3$, $EY_3 = 11/6\lambda$.
- 5.10 $g_{1n}(y_1, y_n) = n(n-1)[F(y_n) - F(y_1)]^{n-2} f(y_1)f(y_n)$, $a < y_1 < y_n < b$.

Chapter 7

Section 7.1

- 1.2 For every $\varepsilon > 0$, $P(|X_n| > \varepsilon) = P(X_n = 1) = p_n$, and therefore $X_n \xrightarrow{P} 0$ if and only if $p_n \rightarrow 0$ as $n \rightarrow \infty$.
- 1.4 (i) $P(|Y_{1,n}| > \varepsilon) = (1 - \varepsilon)^n \rightarrow 0$, as $n \rightarrow \infty$.
 (ii) $P(|Y_{n,n} - 1| > \varepsilon) = 1 - P(|Y_{n,n} - 1| \leq \varepsilon)$ and $P(|Y_{n,n} - 1| \leq \varepsilon) = 1 - (1 - \varepsilon)^n \rightarrow 1$, so that $P(|Y_{n,n} - 1| > \varepsilon) \rightarrow 0$, as $n \rightarrow \infty$.
- 1.6 $E\bar{X}_n = \mu$ and $E(\bar{X}_n - \mu)^2 = \text{Var}(\bar{X}_n) = \frac{\sigma^2}{n} \rightarrow 0$, as $n \rightarrow \infty$.
- 1.8 $E(Y_n - X)^2 = E(Y_n - X_n)^2 + E(X_n - X)^2 + 2E[(Y_n - X_n)(X_n - X)] \rightarrow 0$, as $n \rightarrow \infty$, by the assumptions made and the fact that $|E[(Y_n - X_n)(X_n - X)]| \leq E^{1/2}|X_n - Y_n|^2 \times E^{1/2}|X_n - X|^2$.


Section 7.2

2.2 (i) $M_X(t) = (1 - \alpha)/(1 - \alpha e^t)$, $t < -\log \alpha$;

(ii) $EX = \alpha/(1 - \alpha)$.

(iii) $M_{\bar{X}_n}(t) = \left(\frac{1-\alpha}{1-\alpha e^{t/n}}\right)^n = \left\{1 - \frac{\alpha t/(1-\alpha) + [\alpha/(1-\alpha)]nR(\frac{t}{n})}{n}\right\}^{-n} \xrightarrow{n \rightarrow \infty} e^{\alpha t/(1-\alpha)}$,
 since $\frac{n}{t}R(\frac{t}{n}) \xrightarrow{n \rightarrow \infty} 0$ for fixed t , and $e^{\alpha t/(1-\alpha)}$ is the m.g.f. of $\frac{\alpha}{1-\alpha}$.

2.4 Since $X \sim B(1,000, p)$, we have;

(i) $P(1,000p - 50 \leq X \leq 1,000p + 50) = \sum_{x=1,000p-50}^{1,000p+50} \binom{1,000}{x} p^x q^{1,000-x}$,
 $q = 1 - p$. For $p = \frac{1}{2}$ and $p = \frac{1}{4}$:

$$P(450 \leq X \leq 550) = \sum_{x=450}^{550} \binom{1,000}{x} (0.5)^{1,000},$$

$$P(200 \leq X \leq 300) = \sum_{x=200}^{300} \binom{1,000}{x} (0.25)^x \times (0.75)^{1,000-x}.$$

(ii) For $p = \frac{1}{2}$ and $p = \frac{1}{4}$, the approximate probabilities are $2\Phi(3.16) - 1 = 0.998422$ and $2\Phi(3.65) - 1 = 0.999738$.

2.6 $EX_i = \frac{7}{2}$, $EX_i^2 = \frac{91}{6}$, so that $\text{Var}(X_i) = \frac{35}{12}$. Therefore $P(150 \leq X \leq 200) \simeq 2\Phi(2.07) - 1 = 0.961548$.

2.8 Since $X \sim B(1,000, 0.03)$, the required approximate probability is:
 $P(X \leq 50) \simeq \Phi(3.71) = 0.999896$.

2.10 $P(|\frac{X}{n} - 0.53| \leq 0.02) \simeq 2\Phi(\frac{0.02\sqrt{n}}{\sqrt{0.2491}}) - 1 = 0.99$, so that $n = 4,146$.

2.12 With $S_n = \sum_{i=1}^n X_i$, we have:

(i) $P(S_n \leq \lambda n) \simeq \Phi(\sqrt{\lambda n}) - 0.50$.

(ii) $P(S_n \geq \lambda n) \simeq \Phi(\sqrt{\lambda n}) - \Phi(1/\sqrt{\lambda n})$.

(iii) $P(\frac{\lambda n}{2} \leq S_n \leq \frac{3\lambda n}{2}) \simeq \Phi(\sqrt{\lambda n}/2) - \Phi(\sqrt{\lambda n}/4)$.

(iv) $P(S_n \leq 100) \simeq 0.50$, $P(S_n \geq 100) \simeq 0.460172$, $P(50 \leq S_n \leq 75) \simeq 0.00621$.

2.14 The total life time is $X = \sum_{i=1}^{50} X_i$, where X_i 's are independently distributed as Negative Exponential with $\lambda = 1/1,500$. Then $P(X \geq 80,000) \simeq 1 - \Phi(0.47) = 0.319178$.

2.16 (i) $P(a \leq \bar{X} \leq b) \simeq \Phi((2b - 1)\sqrt{3n}) - \Phi((2a - 1)\sqrt{3n})$.

(ii) Here $(2b - 1)\sqrt{3n} = 0.75$, $(2a - 1)\sqrt{3n} = -0.75$, and the above probability is: $2\Phi(0.75) - 1 = 0.546746$.

2.18 $P(|\bar{X} - \mu| \leq 0.0001) \simeq 2\Phi(0.2\sqrt{n}) - 1 = 0.99$, and then $n = 167$.

2.20 (i) $P(|\bar{X}_n - \mu| < k\sigma) \simeq 2\Phi(k\sqrt{n}) - 1 = p$, so that $n = [\frac{1}{k}\Phi^{-1}(\frac{1+p}{2})]^2$.

(ii) Here n is the smallest integer $\geq 1/(1 - p)k^2$.

(iii) For $p = 0.90$, $p = 0.95$, and $p = 0.99$, and the respective values of k , we determine the values of n by means of the CLT and the Tchebichev inequality.

Then, for the various values of k , the respective values of n are given in the following table for part (i).

$k \backslash p$	0.90	0.95	0.99
0.50	11	16	27
0.25	44	62	107
0.10	271	385	664

For the Tchebichev inequality, the values of n are given by the entries of the table below.

$k \backslash p$	0.90	0.95	0.99
0.50	40	80	400
0.25	160	320	1,600
0.10	1,000	2,000	10,000

- 2.22 (i) $P(|\bar{X} - \bar{Y}| \leq 0.25\sigma) = P(|\bar{Z}| \leq 0.25\sigma) \simeq 2\Phi\left(\frac{0.25\sqrt{n}}{\sqrt{2}}\right) - 1 = 0.95$ and then $n = 123$.
- (ii) From $1 - \frac{2}{0.0625n} \geq 0.95$, we find $n = 640$.
- 2.24 (i) $P(|X - 20| \leq 2) \simeq \Phi\left(\frac{22-np}{\sqrt{npq}}\right) - \Phi\left(\frac{18-np}{\sqrt{npq}}\right)$, and taking $\frac{18-np}{\sqrt{npq}} = -\frac{22-np}{\sqrt{npq}}$, we find $n = 50$.
- (ii) For $n = 50$, the approximate probability in part (i) is 0.438086.
- (iii) As in the hint.

Chapter 9

Section 9.1

- 1.2 The matrix is negative definite, because for λ_1, λ_2 with $\lambda_1^2 + \lambda_2^2 \neq 0$, we have:

$$(\lambda_1, \lambda_2) \begin{pmatrix} -n/s^2 & 0 \\ 0 & -n/2s^4 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} = -\lambda_1^2 \frac{n}{s^2} - \lambda_2^2 \frac{n}{2s^4} < 0.$$

- 1.4 With $Y = \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma}\right)^2 \sim \chi_{n-1}^2$ and $S^2 = \sigma^2 Y / (n-1)$, we have $\text{Var}_{\sigma^2}(S^2) = 2\sigma^4 / (n-1)$.

- 1.6 With $L(\theta | \mathbf{x}) = \theta^n (1-\theta)^{n(\bar{x}-1)}$, $\mathbf{x} = (x_1, \dots, x_n)$, we have $\frac{\partial}{\partial \theta} \log L(\theta | \mathbf{x}) = 0$ produces $\theta = 1/\bar{x}$, and $\frac{\partial^2}{\partial \theta^2} \log L(\theta | \mathbf{x}) = -\frac{n}{\theta^2} - \frac{n(\bar{x}-1)}{(1-\theta)^2} < 0$.

- 1.8 The MLE of θ is $\hat{\theta} = -n / \sum_{i=1}^n \log x_i$.

- 1.10 (i) It follows from what it is given;
 (ii) $\hat{\theta} = 2/\bar{x}$.

1.12 (i) The first two expressions are immediate. The third follows from $\frac{d}{d\sigma_1^2} \left(\frac{1}{\sigma_1^2}\right) = -\frac{1}{2\sigma_1^3}$, and similarly for the fourth. The fifth follows from $\frac{d}{d\rho} \left(\frac{1}{1-\rho^2}\right) = \frac{2\rho}{(1-\rho^2)^2}$ and $\frac{d}{d\rho} \left(\frac{\rho}{1-\rho^2}\right) = \frac{1+\rho^2}{(1-\rho^2)^3}$.

(ii) Immediate from part (i).

(iii) $\tilde{\mu}_1 = \bar{x}$ and $\tilde{\mu}_2 = \bar{y}$ follow by solving the equations:

$$\sigma_2\mu_1 - \sigma_1\rho\mu_2 = \sigma_2\bar{x} - \sigma_1\rho\bar{y}, \quad \sigma_2\rho\mu_1 - \sigma_1\mu_2 = \sigma_2\rho\bar{x} - \sigma_1\bar{y}$$

following from the first two likelihood equations.

(iv) They follow from part (ii).

(v) They follow from part (iv) by solving for σ_1^2 , σ_2^2 , and ρ .

1.14 (i) Immediate by the fact that $\tilde{d}_{13}(=\tilde{d}_{31}) = \tilde{d}_{14}(=\tilde{d}_{41}) = \tilde{d}_{15}(=\tilde{d}_{51}) = \tilde{d}_{23}(=\tilde{d}_{32}) = \tilde{d}_{24}(=\tilde{d}_{42}) = \tilde{d}_{25}(=\tilde{d}_{52}) = 0$.

(ii) Immediate by the fact that $\tilde{d}_{12} = \tilde{d}_{21}$ and $\tilde{d}_{34} = \tilde{d}_{43}$.

(iii) Immediate by the fact that $\tilde{d}_{21} = \tilde{d}_{12}$, $\tilde{d}_{43} = \tilde{d}_{34}$, $\tilde{d}_{54} = \tilde{d}_{45}$, and $\tilde{d}_{53} = \tilde{d}_{35}$.

(iv) $\tilde{D}_1 = -\frac{n\beta}{\delta}$, $\tilde{D}_2 = \frac{n^2}{\delta}$, $\tilde{D}_3 = -\frac{n^3(\alpha\beta + \delta)}{4\alpha^2\delta^2}$, $\tilde{D}_4 = \frac{n^4}{4\alpha\beta\delta}$.

(v) $A = \frac{\alpha^3\beta n^2}{2\delta^3}$, $B = -\frac{\alpha\beta\gamma^2 n^2}{2\delta^3}$, $C = \frac{\alpha^{1/2}\gamma n^2}{4\beta^{1/2}\delta^2}$.

(vi) $\tilde{D}_5 = -\frac{\alpha\beta n^5}{4\delta^4}$.

(vii) $\tilde{D}_0 = 1 > 0$, $\tilde{D}_1 = -\frac{n\beta}{\delta} < 0$, $\tilde{D}_2 = \frac{n^2}{\delta} > 0$, $\tilde{D}_3 = -\frac{n^3(\alpha\beta + \delta)}{4\alpha^2\delta^2} < 0$, $\tilde{D}_4 = \frac{n^4}{4\alpha\beta\delta} > 0$, and $\tilde{D}_5 = -\frac{\alpha\beta n^5}{4\delta^4} < 0$.

Section 9.2

2.2 (i) $R(x; \theta) = e^{-x/\theta}$; (ii) The MLE of $R(x; \theta)$ is $e^{-x/\bar{x}}$.

2.4 (i) $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n x_i^\gamma$; (ii) $\hat{\theta} = \bar{x}$.

2.6 (i) It follows by integration.

(ii) $\prod_{i=1}^n X_i$ is a sufficient statistic for θ , and so is $\sum_{i=1}^n \log X_i$.

2.8 $\sum_{i=1}^n |X_i|$ is a sufficient statistic for θ .

2.10 Here (X_1, \dots, X_r) is a set of statistics sufficient for (p_1, \dots, p_r) , or (X_1, \dots, X_{r-1}) is a set of statistics sufficient for (p_1, \dots, p_{r-1}) . Furthermore, $(X_{(1)}, X_{(n)})$ is a set of statistics sufficient for (α, β) .

2.12 They follow by integration.

Section 9.3

3.2 (i) $E_\theta \bar{X} = \frac{1}{\theta}$, $E_\theta(nY_1) = \frac{1}{\theta}$.

(ii) $\text{Var}_\theta(\bar{X}) = \frac{1}{n\theta^2} \leq \frac{1}{\theta^2} = \text{Var}_\theta(nY_1)$.

3.4 (i) $g_1(y) = \frac{n}{\theta_2 - \theta_1} \left(\frac{\theta_2 - y}{\theta_2 - \theta_1}\right)^{n-1}$, $\theta_1 \leq y \leq \theta_2$,

$g_n(y) = \frac{n}{\theta_2 - \theta_1} \left(\frac{y - \theta_1}{\theta_2 - \theta_1}\right)^{n-1}$, $\theta_1 \leq y \leq \theta_2$.

Then, with $\theta = (\theta_1, \theta_2)$,

$$E_{\theta} Y_1 = \frac{n\theta_1 + n\theta_2}{n+1}, \quad E_{\theta} Y_n = \frac{\theta_1 + n\theta_2}{n+1}, \text{ and}$$

$$E_{\theta} \left(\frac{Y_1 + Y_2}{2} \right) = \frac{\theta_1 + \theta_2}{2}, \quad E_{\theta} \left[\frac{n+1}{n-1} (Y_n - Y_1) \right] = \theta_2 - \theta_1.$$

- (ii) The pair of statistics (Y_1, Y_n) is sufficient for the pair of the parameters (θ_1, θ_2) .

3.6 They follow by integration and by using part (ii) of Exercise 3.5.

3.8 (i) The $\text{Var}_{\theta}(U_1)$ and $\text{Var}_{\theta}(U_2)$ follow from the $\text{Var}_{\theta}(Y_1)$ and $\text{Var}_{\theta}(Y_n)$ in Exercise 3.6, and the $\text{Cov}_{\theta}(Y_1, Y_n)$ from Exercise 3.7(ii).

(ii) Immediate by comparing variances.

3.10 (i) The required condition is $\sum_{i=1}^n c_i = 1$.

(ii) $c_i = \frac{1}{n}$, $i = 1, \dots, n$.

3.12 (i) $E_{\theta} \bar{X} = \theta$ and $\text{Var}_{\theta}(\bar{X}) = \frac{1}{n \times \frac{1}{\sigma^2}} = \frac{1}{nI(\theta)}$.

(ii) $E_{\theta} S^2 = \theta$ and $\text{Var}_{\theta}(S^2) = \frac{1}{n \times \frac{1}{2\theta^2}} = \frac{1}{nI(\theta)}$.

3.14 It follows by the hint and Exercise 3.13.

3.16 (i) See application to Theorem 3.

(ii) $E_{\theta} h(X) = 0$ is equivalent to $\sum_{x=0}^n h(x) \binom{n}{x} t^x = 0$ ($t = \frac{\theta}{1-\theta}$) from which it follows $h(x) = 0$, $x = 0, 1, \dots, n$.

(iii) It follows by the fact that $\bar{X} = \frac{T}{n}$ and $T \sim B(n, \theta)$ is sufficient and complete.

3.18 (i) X is sufficient by the Factorization Theorem, and also complete, because $E_{\theta} h(X) = \frac{1-t}{t} \sum_{x=1}^{\infty} h(x) t^x = 0$ ($t = 1 - \theta$) implies $h(x) = 0$, $x = 1, 2, \dots$

(ii) U is unbiased, because $E_{\theta} U = 1 \times P_{\theta}(U = 1) = P_{\theta}(U = 1) = P_{\theta}(X = 1) = \theta$.

(iii) U is *UMVU*, because it is unbiased and depends only on the sufficient statistic X .

(iv) $\text{Var}_{\theta}(U) = \theta(1 - \theta) > \theta^2(1 - \theta) = \frac{1}{I(\theta)}$ ($0 < \theta < 1$).

3.20 (i) Sufficiency of Y_n follows from Exercise 2.9(iii); completeness cannot be established here.

(ii) It follows from part (i) and Exercise 3.3(ii).

(iii) Because the function $L(\theta | x) (= f(x; \theta))$ is not differentiable at $\theta = x$.

Section 9.4

4.2 It follows by the fact that $\int_0^1 \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1} dx = 1$ and the recursive relation of the Gamma function.

4.4 Immediate from the hint.

4.6 $R(\theta; d) = \frac{1}{n}$, independent of θ , and then Theorem 9 applies.

Section 9.5

- 5.2 (i) $\tilde{\theta} = (1 - 2\bar{X})/(\bar{X} - 1)$; (ii) $\tilde{\theta} = 3$ and $\hat{\theta} \simeq 3.116$.
- 5.4 Follows from Exercise 3.15(i).
- 5.6 (i) The EX and EX^2 follow by integration, and $\text{Var}(X) = \beta^2$.
(ii) $\tilde{\alpha} = \bar{X} - S$, $\tilde{\beta} = S$, where $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$.
- 5.8 By equating the second-order sample moment to $\frac{(2\theta)^2}{12}$, we get $\tilde{\theta} = (\frac{3}{n} \sum_{i=1}^n X_i^2)^{1/2}$.
- 5.10 (i) It follows by integration;
(ii) They also follow by integration.
(iii) $\tilde{\theta} = 3\bar{X}$ and hence $E_{\tilde{\theta}}\tilde{\theta} = \theta$ and $\text{Var}_{\tilde{\theta}}(\tilde{\theta}) = \theta^2/2n$.
- 5.12 (i) and (ii) follow by straightforward calculations.
(ii) It follows by the expression of $\hat{\rho}_n(X, Y)$ and the WLLN applied to $n^{-1} \sum_{i=1}^n (X_i Y_i)$, \bar{X} , \bar{Y} , $n^{-1} \sum_{i=1}^n (X_i - \bar{X})^2$, and $n^{-1} \sum_{i=1}^n (Y_i - \bar{Y})^2$.
- 5.14 $\bar{x} = \frac{2,423}{15} \simeq 161.533$, $s_x = \frac{\sqrt{175,826}}{15} \simeq 27.954$, $\bar{y} = 140.6$, and $s_y = \frac{\sqrt{56,574}}{15} \simeq 15.857$.

Chapter 10

Section 10.1

- 1.2 $m = 4n$.
- 1.4 (i) $n = (2z_{\alpha/2}\sigma/l)^2$; (ii) $n = 1,537$.
- 1.6 (i) $M_U(t) = 1/(1 - \theta t)^n$, $t < \frac{1}{\theta}$, which is the m.g.f. of the Gamma distribution with $\alpha = n$ and $\beta = \theta$.
(ii) $M_V(t) = 1/(1 - 2t)^{2n/2}$, $t < \frac{1}{2}$, which is the m.g.f. of the χ_{2n}^2 distribution, so that $V \sim \chi_{2n}^2$.
(iii) $[2b^{-1} \sum_{i=1}^n X_i, 2a^{-1} \sum_{i=1}^n X_i]$.
- 1.8 (i) For $x > \theta$, $F(x; \theta) = 1 - e^{-(x-\theta)}$ and hence $g(y; \theta) = ne^{-n(y-\theta)}$, $y > \theta$.
(ii) By setting $t = 2n(y - \theta)$, we get $f_T(t; \theta) = \frac{1}{2}e^{-t/2}$, $t > 0$, which is the p.d.f. of the χ_2^2 distribution.
(iii) It follows by the usual arguments.
- 1.10 (i) The transformation $y = |x|$ which yields $x = y$ for $x > 0$, $x = -y$ for $x < 0$, and $|\frac{dx}{dy}| = 1$. Then, by Theorem 6, $g(y; \theta) = \frac{1}{\theta}e^{-y/\theta}$, $y > 0$.
(ii) and (iii) are as in Exercise 1.9(ii) and (iii), respectively.
- 1.12 (i) $e^{-a} - e^{-b} = 1 - \alpha$.

- (ii) Immediate by part (i) and the fact that T has the p.d.f. e^{-t} , $t > 0$.
 (iii) Follows from the hint.

1.14 From the transformations $r = y_n - y_1$, and $s = y_1$, we get: $y_1 = s$, $y_n = r + s$, and $|J| = 1$. Then $f_{R,S}(r, s; \theta) = \frac{n(n-1)}{\theta^n} r^{n-2}$, $0 < r < \theta$, $0 < s < \theta - r$, and then $f_R(r; \theta) = \frac{n(n-1)}{\theta^n} r^{n-2}(\theta - r)$, $0 < r < \theta$.

- 1.16 (i) It follows by using the transformation $t = r/\theta$.
 (ii) The required confidence interval is $[R, \frac{R}{C}]$. The relation $c^{n-1}[n - (n-1)c] = \alpha$ follows from:

$$1 - \alpha = P_\theta(c \leq T \leq 1) = \int_c^1 n(n-1)t^{n-2}(1-t) dt \\ = n - nc^{n-1} - (n-1) + (n-1)c^n.$$

1.18 Follows by the usual procedure and the fact that $\frac{mS_X^2}{\sigma_1^2} \sim \chi_m^2$, $\frac{nS_Y^2}{\sigma_2^2} \sim \chi_n^2$ independent, so that $\frac{S_Y^2/\sigma_2^2}{S_X^2/\sigma_1^2} \sim F_{n,m}$.

Section 10.2

2.2 The required interval is $[a \frac{S_X^2}{S_Y^2}, b \frac{S_X^2}{S_Y^2}]$, where $0 < a < b$ with $P(a \leq X \leq b) = 1 - \alpha$, $X \sim F_{n-1, m-1}$. In particular, $a = F_{n-1, m-1; 1-\frac{\alpha}{2}}$, $b = F_{n-1, m-1; \frac{\alpha}{2}}$.

Section 10.4

- 4.2 (i) The required confidence interval is $\bar{X}_n \pm z_{\frac{\alpha}{2}} \frac{S_n}{\sqrt{n}}$.
 (ii) Here $\bar{X}_{100} \pm 0.196 S_{100}$.
 (iii) The length is $2z_{\frac{\alpha}{2}} \frac{S_n}{\sqrt{n}}$, which converges in probability to 0 since $S_n \xrightarrow{P} \sigma$.
- 4.4 (i) $P(Y_i \leq x_p) = P(\text{at least } i \text{ of } X_1, \dots, X_n \leq x_p) = \sum_{k=i}^n \binom{n}{k} p^k q^{n-k}$, and also $P(Y_i \leq x_p) = P(Y_i \leq x_p \leq Y_j) + P(Y_j \leq x_p)$, so that $P(Y_i \leq x_p \leq Y_j) = \sum_{k=i}^n \binom{n}{k} p^k q^{n-k} - \sum_{k=j}^n \binom{n}{k} p^k q^{n-k} - \sum_{k=i}^{j-1} \binom{n}{k} p^k q^{n-k}$.
 (ii) and (iii) follow from part (i) and the Binomial tables.
 (iv) It follows from the hint and the Binomial tables.

Chapter 11

Section 11.1

1.2 H_{0i} and H_{Ai} , $i = 1, \dots, 5$ are all composite; H_{06} and H_{A6} are both simple.

Section 11.2


- 2.2 (i) The required n is determined by solving for n (and C) the two equations: $\frac{C\sqrt{n}}{\sigma} = \Phi^{-1}(1 - \alpha)$ and $\frac{\sqrt{n(C-1)}}{\sigma} = \Phi^{-1}(1 - \pi(1))$.
 (ii) $n = 9$ (and $C \simeq 0.562$).

- 2.4 (i) The MP test rejects H_0 when $\bar{x} > C_n$, where $C_n = \mu_0 + \frac{\sigma}{\sqrt{n}} \Phi^{-1}(1 - \alpha_n)$.
(ii) The required C_n and n are given by:

$$C_n = \mu_0 + \frac{(\mu_1 - \mu_0)\Phi^{-1}(1 - \alpha_n)}{\Phi^{-1}(1 - \alpha_n) - \Phi^{-1}(1 - \pi_n)},$$

$$n = \left\{ \frac{\sigma}{\mu_1 - \mu_0} [\Phi^{-1}(1 - \alpha_n) - \Phi^{-1}(1 - \pi_n)] \right\}^2.$$

- (iii) That $\alpha_n \xrightarrow{n \rightarrow \infty} 0$ follows from $1 - \Phi\left[\frac{\sqrt{n}(C_n - \mu_0)}{\sigma}\right] = \alpha_n$ and the fact that $\mu_0 < C_n$; that $\pi_n \xrightarrow{n \rightarrow \infty} 1$ follows from $1 - \Phi^{-1}\left[\frac{\sqrt{n}(C_n - \mu_1)}{\sigma}\right] = \pi_n$, and the fact that $C_n < \mu_1$.
(iv) $n = 33$ and $C_{33} \simeq 0.546$.
- 2.6 (i) $C(\theta) = \frac{1}{\theta}$, $Q(\theta) = -\frac{1}{\theta}$ strictly increasing, $T(x) = x$, and $h(x) = I_{(0, \infty)}(x)$.
(ii) The UMP test rejects H_0 when $\sum_{i=1}^n x_i < C$, where C is determined by $P_{\theta_0}(\sum_{i=1}^n X_i < C) = \alpha$.
(iii) $M_{\sum_{i=1}^n X_i}(t) = \frac{1}{(1 - \theta t)^n}$ ($t < \frac{1}{\theta}$), and $M_{2(\sum_{i=1}^n X_i)/\theta}(t) = M_{\sum_{i=1}^n X_i}\left(\frac{2t}{\theta}\right) = \frac{1}{(1 - \theta \cdot \frac{2t}{\theta})^n} = \frac{1}{(1 - 2t)^{2n/2}}$, which is the m.g.f. of the χ_{2n}^2 distribution.
(iv) $C = \frac{\theta_0}{2} \chi_{2n; 1-\alpha}^2$, $\pi(\theta_1) = P(X < \frac{2C}{\theta_1})$, $X \sim \chi_{2n}^2$.
(v) The closest value we can get by means of the χ^2 -tables is $n = 23$.
- 2.8 (i) The MP test rejects H_0 when $\frac{f_1(x)}{f_0(x)} > C$, $C \simeq 0.842$.
(ii) $\pi \simeq 0.823$.
(iii) By means of geometric considerations, we find $\pi = 0.822$.



- 2.10 (i) The MP test rejects H_0 when $x \in \{x \geq 0 \text{ integer}; 1.36 \times \frac{x!}{2^x} \geq C\}$, where C is determined by $P_{H_0}(\{1.36 \times \frac{x!}{2^x} \geq C\}) = \alpha$.
- (ii) For $C = 2$, it follows that $\alpha \simeq 0.02$.

Section 11.3

- 3.2 (ii) The UMP test is given by:

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } \sum_{i=1}^n x_i < C \\ \gamma & \text{if } \sum_{i=1}^n x_i = C \\ 0 & \text{if } \sum_{i=1}^n x_i > C, \end{cases}$$

where C and γ are determined by:

$$P_{p_0}(X < C) + \gamma P_{p_0}(X = C) = \alpha, \quad X \sim B(n, p_0).$$

- (iii) Here $C \simeq 3.139$, and H_0 is rejected when $X \leq 3$.

- 3.4 (i) With each student associate a $B(1, p)$ r.v. X_i , $i = 1, \dots, 400$, so that $X = \sum_{i=1}^{400} X_i \sim B(400, p)$. Then the UMP test is given by:

$$\varphi(x_1, \dots, x_{400}) = \begin{cases} 1 & \text{if } \sum_{i=1}^{400} x_i < C \\ \gamma & \text{if } \sum_{i=1}^{400} x_i = C \\ 0 & \text{if } \sum_{i=1}^{400} x_i > C, \end{cases}$$

where C and γ are determined by:

$$P_{0.25}(X < C) + \gamma P_{0.25}(X = C) = 0.05, \quad X \sim B(400, 0.25).$$

- (ii) Here $C \simeq 85.7543$, and H_0 is rejected when $\sum_{i=1}^{400} x_i \leq 85$.
- 3.6 (i) The UMP test is given by relation (19) with C and γ defined by relation (20).
- (ii) Here $C = 5$ and $\gamma \simeq 0.519$.
- (iii) $\pi(0.375) \simeq 0.22$ and $\pi(0.500) \simeq 0.505$.
- (iv) For $\theta > 0.5$, $\pi(\theta) = P_{1-\theta}(X \leq n - C - 1) + \gamma P_{1-\theta}(X = n - C)$.
- (v) $\pi(0.625) \simeq 0.787$, and $\pi(0.875) \simeq 0.998$.
- (vi) $n = 62$.

- 3.8 (i) $H_0 : \lambda = 10$, $H_A : \lambda < 10$.

- (ii) The UMP test is given by:

$$\varphi(x) = \begin{cases} 1 & \text{if } x < C \\ \gamma & \text{if } x = C, \\ 0 & \text{if } x > C \end{cases} \quad \begin{aligned} P_{10}(X \leq C - 1) + \gamma P_{10}(X = C) \\ = 0.01, X \sim P(10). \end{aligned}$$

From the Poisson tables, $C = 3$ and $\gamma = 0.96$, and since $x = 4$, H_0 is not rejected.

- 3.10 (i) H_0 is rejected when $\sum_{i=1}^n x_i < C$, $C = n\mu_0 - z_\alpha \sigma \sqrt{n}$.

$$(ii) \pi(\mu) = \Phi\left(\frac{C - n\mu}{\sigma\sqrt{n}}\right).$$

$$(iii) C = 43,252.5, H_0 \text{ is rejected, and } \pi(1,700) = 0.841345.$$

3.12 H_0 is rejected when $C_1 < \sum_{i=1}^4 z_i < C_2$, where C_1 and C_2 are determined by $\Phi(0.4 + x) - \Phi(0.4 - x) = 0.05$, $x = \frac{C}{10}$. From the Normal tables, $x \simeq 0.07$, so that $C=0.7$, and H_0 is rejected when $-0.7 < z_1 + z_2 + z_3 + z_4 < 0.7$.

3.14 Here $H_0 : \sigma > 0.04$, $H_A : \sigma \leq 0.04$, and the UMP test rejects H_0 when $\sum_{i=1}^n (x_i - \mu)^2 < C$, $C = \sigma_0^2 \chi_{n-1, 1-\alpha}^2$. For the numerical data, $C = 0.0127392 \simeq 0.013$, and since $\sum_{i=1}^{16} (x_i - \mu)^2 = 0.04$, the hypothesis H_0 is not rejected.

Section 11.4

4.2 (i) $\lambda = (0.25)^t (0.75)^{3-t} / (\frac{t}{3})^t (1 - \frac{t}{3})^{3-t}$, $t = 0, 1, 2, 3$, and H_0 is rejected when $\lambda < C$ where C is determined by $P_{0.25}(\lambda < C) = 0.02$.

(ii) At level $\alpha = 0.02$, H_0 is outright rejected when $\lambda = 0.015625$ (which is equivalent to $t = 3$) and is rejected with probability $0.02 - 0.0156 = 0.0044$ when $\lambda = 0.31640625$ (which is equivalent to $t = 2$).

4.4 (ii) With $t(\mathbf{z}) = \sqrt{n}\bar{z} / \sqrt{\frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2}$, $\mathbf{z} = (z_1, \dots, z_n)$, H_0 is rejected when $t(\mathbf{z}) < -t_{n-1; \frac{\alpha}{2}}$ or $t(\mathbf{z}) > t_{n-1; \frac{\alpha}{2}}$.

(iii) Here $t_{89; 0.025} = 1.9870$, and therefore H_0 is rejected when

$$|3\sqrt{10}\bar{z}| / \sqrt{\frac{1}{89} \sum_{i=1}^{90} (z_i - \bar{z})^2} > 1.9870.$$

4.6 Here $H_0 : \mu = 2.5$ and $H_A : \mu \neq 2.5$, and H_0 is rejected when $\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} < -z_{\alpha/2}$ or $\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} > z_{\alpha/2}$. Since $z_{0.025} = 1.96$ and $\frac{\sqrt{n}(\bar{x} - \mu_0)}{\sigma} = -0.8$, H_0 is not rejected.

4.8 Here $H_0 : \mu_1 = \mu_2$, $H_A : \mu_1 \neq \mu_2$, and H_0 is rejected when $t(\mathbf{x}, \mathbf{y}) < -t_{m+n-2; \alpha/2}$ or $t(\mathbf{x}, \mathbf{y}) > t_{m+n-2; \alpha/2}$, where

$$t(\mathbf{x}, \mathbf{y}) = \frac{\sqrt{\frac{mn}{mn}}(\bar{x} - \bar{y})}{\sqrt{\frac{1}{m+n-2} [\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2]}}$$
, $\mathbf{x} = (x_1, \dots, x_m)$, $\mathbf{y} = (y_1, \dots, y_n)$. Since $t_{48; 0.025} = 2.0106$ and $t(\mathbf{x}, \mathbf{y}) = -2.712$, the hypothesis H_0 is rejected.

4.10 H_0 is rejected when $u(\mathbf{x}, \mathbf{y}) \leq F_{m-1, n-1; 1-\frac{\alpha}{2}}$ or $u(\mathbf{x}, \mathbf{y}) \geq F_{m-1, n-1; \frac{\alpha}{2}}$, where $u(\mathbf{x}, \mathbf{y}) = \frac{\frac{1}{m-1} \sum_{i=1}^m (x_i - \bar{x})^2 / \frac{1}{n-1} \sum_{j=1}^n (y_j - \bar{y})^2}{\sum_{i=1}^m (x_i - \bar{x})^2 + \sum_{j=1}^n (y_j - \bar{y})^2}$, $\mathbf{x} = (x_1, \dots, x_m)$, $\mathbf{y} = (y_1, \dots, y_n)$. Here $F_{3, 3; 0.975} \simeq 0.065$, $F_{3, 3; 0.025} = 15.439$, and $u(\mathbf{x}, \mathbf{y}) \simeq 2.168$. Therefore H_0 is not rejected.

4.12 The LR test is the same as that given in Exercise 4.8. For the given numerical data, $t_{8; 0.025} = 3.3060$, $t(\mathbf{x}, \mathbf{y}) \simeq 2.014$, and therefore H_0 is not rejected.

4.14 They follow by integrating by parts.

4.16 With $\lambda(u) = e^{n/2} u^{n/2} e^{-mu/2}$, $u \geq 0$, we have $\lambda'(u) = \frac{d\lambda(u)}{du} = \frac{n}{2} e^{n/2} u^{\frac{n}{2}-1} e^{-mu/2} \times (1-u)$, so that $\lambda'(u) \geq 0$ if $(0 \leq) u \leq 1$ and $\lambda'(u) \leq 0$ if $u > 1$. It follows that $\lambda(u)$ is strictly increasing for $u \leq 1$ and strictly decreasing for $u > 1$.

Since $\lambda'(u) = 0$ gives $u = 1$ and $\frac{d^2}{du^2}\lambda(u)|_{u=1} = -\frac{n}{2} < 0$, it follows that $\lambda(u)$ is maximized for $u = 1$. That $\lambda(0) = 0$ is immediate, and that $\lambda(u) \rightarrow 0$ as $u \rightarrow \infty$ follows by taking the limit of the ratio of derivatives of sufficiently high order.

- 4.18 Maximization of (58) is equivalent to maximization of $g = g(\mu, \tau) = -\frac{m+n}{2} \log \tau - \frac{1}{2\tau} [\sum_{i=1}^m (x_i - \mu)^2 + \sum_{j=1}^n (y_j - \mu)^2]$, where $\tau = \sigma^2$. From $\frac{\partial g}{\partial \mu} = 0$, $\frac{\partial g}{\partial \tau} = 0$, we find $\hat{\mu}_\omega = \frac{n\bar{x} + n\bar{y}}{m+n}$, $\hat{\tau} = \frac{1}{m+n} [\sum_{i=1}^m (x_i - \hat{\mu}_\omega)^2 + \sum_{j=1}^n (y_j - \hat{\mu}_\omega)^2]$. Next, $\frac{\partial^2 g}{\partial \mu^2}$, $\frac{\partial^2 g}{\partial \mu \partial \tau} = \frac{\partial^2 g}{\partial \tau \partial \mu}$, and $\frac{\partial^2 g}{\partial \tau^2}$ evaluated at $\mu = \hat{\mu}_\omega$ and $\tau = \hat{\tau}$, yield, respectively; $-\frac{m+n}{\hat{\tau}}$, 0, and $-\frac{m+n}{2\hat{\tau}^2}$. Setting \mathbf{C} for the 2×2 matrix of the second-order derivatives of g , we have, for λ_1, λ_2 with $\lambda_1^2 + \lambda_2^2 \neq 0$:

$$(\lambda_1, \lambda_2)\mathbf{C} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} = -\frac{m+n}{\hat{\tau}} \left(\lambda_1^2 + \frac{\lambda_2^2}{2} \right) < 0,$$

so that \mathbf{C} is negative definite, and hence $\hat{\mu}_\omega$ and $\hat{\tau}$ are the MLE of μ and τ , respectively.

- 4.20 From the assumptions made, it follows that:

$$\frac{\bar{X} - \bar{Y}}{\sigma \sqrt{\frac{1}{m} + \frac{1}{n}}} \sim N(0, 1); \quad \sum_{i=1}^m \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 \sim \chi_{m-1}^2, \quad \sum_{j=1}^n \left(\frac{Y_j - \bar{Y}}{\sigma} \right)^2 \sim \chi_{n-1}^2$$

independent, so that their sum is $\sim \chi_{m+n-2}^2$. This sum is also independent of $\bar{X} - \bar{Y}$. It follows that $\frac{\bar{X} - \bar{Y}}{\sigma \sqrt{\frac{1}{m} + \frac{1}{n}}}$ divided by $[\sum_{i=1}^m (\frac{X_i - \bar{X}}{\sigma})^2 + \sum_{j=1}^n (\frac{Y_j - \bar{Y}}{\sigma})^2] / (m+n-2)$ is distributed as t_{m+n-2} . The cancellation of σ leads to the assertion made.

- 4.22 Set $c = (m+n)^{\frac{m+n}{2}} / m^{\frac{m}{2}} n^{\frac{n}{2}}$ and $d = \frac{m-1}{n-1}$, so that $\lambda = \lambda(u) = c(du)^{m/2} / (1+du)^{(m+n)/2}$. That $\lambda(0) = 0$ is immediate. Next, $\lambda \rightarrow 0$ as $u \rightarrow \infty$ is also clear. Furthermore, $\frac{d\lambda(u)}{du} = \frac{cd}{2} \times \frac{(du)^{\frac{m}{2}-1}}{(1+du)^{(m+n+2)/2}} \times (m-n)du = 0$ yields $u = \frac{m}{nd} = \frac{m(n-1)}{n(m-1)}$, call it u_0 . Also, $\frac{d\lambda(u)}{du} > 0$ for $u < u_0$, and $\frac{d\lambda(u)}{du} < 0$ for $u > u_0$, so that $\lambda(u)$ increases for $u < u_0$ and decreases for $u > u_0$. It follows that $\lambda(u)$ attains its maximum for $u = u_0$. This maximum is 1.
- 4.24 (i) Since $\frac{df(r)}{dr} = -\frac{n}{2}(1-r)^{\frac{n}{2}-1} < 0$, $f(r)$ is decreasing in r .
 (iv) Since $\frac{dw(r)}{dr} = \frac{\sqrt{n-2}}{(1-r^2)^{3/2}} > 0$, $w(r)$ is increasing in r .

Chapter 12

Section 12.1

- 1.2 (ii), (iii) $-2 \log \lambda \simeq 2(\sum_{i=1}^{12} x_i \log x_i - 335,490.304)$, where x_i is the number of births falling into the i th month. Finally, $-2 \log \lambda = 78.776$ and $\chi_{11;0.01}^2 = 24.725$. The hypothesis H_0 is rejected.

- 1.4 (iii) Here $-2 \log \lambda \simeq 27.952$, and $\chi_{4;0.05}^2 = 9.488$. The hypothesis H_0 is rejected. Also, $\chi_{4;0.01}^2 = 13.277$ and the hypothesis H_0 is still rejected.

Section 12.2

- 2.2 Here $\chi_{\omega}^2 \simeq 72.455 > 24.725 = \chi_{11;0.01}^2$, and H_0 is rejected.
- 2.4 Here $\chi_{\omega}^2 \simeq 28.161 > 9.488 = \chi_{4;0.05}^2$, and H_0 is rejected. Also, $\chi_{\omega}^2 \simeq 28.161 > 13.277 = \chi_{4;0.01}^2$, and H_0 is still rejected.
- 2.6 (i) Here $\chi_{\omega}^2 = 4 > 2.706 = \chi_{1;0.1}^2$, and H_0 is rejected.
(ii) The P -value is approximately 0.047.
- 2.8 Here $\chi_{\omega}^2 = 1.2 < 5.991 = \chi_{2;0.05}^2$, and H_0 is not rejected.
- 2.10 (ii) $p_{10} = 0.251429$, $p_{20} = 0.248571$, $p_{30} = 0.248571$,
 $p_{40} = 0.15967$, $p_{50} = 0.069009$, $p_{60} = 0.02275$.
(iii) $\chi_{\omega}^2 \simeq 51.161 > 11.071 = \chi_{5;0.05}^2$, and H_0 is rejected.
- 2.12 Here $\chi_{\omega}^2 \simeq 1.668 < 11.071 = \chi_{5;0.05}^2$, and H_0 is not rejected.

Chapter 13

Section 13.3

- 3.2 $\sum_i x_i = 91.22$, $\sum_i y_i = 15,228$, $\sum_i x_i^2 = 273.8244$, $\sum_i x_i y_i = 45,243.54$.
Also, $SS_x \simeq 5.402$, $SS_y \simeq 101,339.419$, and $SS_{xy} \simeq 433.922$.
- 3.4 $\text{Var}(\hat{\beta}_1) \simeq 0.577\sigma^2$, $\text{Var}(\hat{\beta}_2) \simeq 0.185\sigma^2$, and $\hat{\sigma}^2 \simeq 2,144.701$.
- 3.6 (i) It follows by differentiating and equating the derivatives to 0.
(ii) $-(x)$ As indicated.

Section 13.4

- 4.2 From relations (24) and (25), we find $t \simeq -0.737$ and $t \simeq 0.987$. Since $t_{29;0.025} = 2.0452$, none of the hypotheses is rejected.
- 4.4 (ii) Replacing the x_i 's by the t_i 's and taking into consideration that $\bar{t} = 0$, we get from (5) and (8) the values specified for $\hat{\gamma}$ and $\hat{\beta}$.
(iii) Immediate by the fact that $\bar{t} = 0$.
(iv) $\hat{\beta} \pm t_{n-2; \frac{\alpha}{2}} \frac{S}{\sqrt{n}}$ and $\hat{\gamma} \pm t_{n-2; \frac{\alpha}{2}} \frac{S}{\sqrt{SS_t}}$, where $S = \sqrt{SS_E/(n-2)}$, $SS_E = SS_y - \frac{SS_{xy}^2}{SS_x}$, $SS_y = \sum_i Y_i^2 - (\frac{1}{n} \sum_i Y_i)^2$, $SS_{xy} = \sum_i t_i Y_i$, $SS_x = SS_t$.
(v) $t = \frac{\hat{\beta} - \beta_0}{S/\sqrt{n}}$, $t = \frac{\hat{\gamma} - \gamma_0}{S/\sqrt{SS_t}}$.
(vi) $\hat{\gamma}_0 \pm t_{n-2; \frac{\alpha}{2}} S \sqrt{\frac{1}{n} + \frac{t_0^2}{SS_t}}$, $\hat{\gamma}_0 \pm t_{n-2; \frac{\alpha}{2}} S \sqrt{1 + \frac{1}{n} + \frac{t_0^2}{SS_t}}$.

Section 13.5

- 5.2 (i) $\hat{y}_0 = 517.474$.
 (ii) $S \simeq 59.834$, $\sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{SS_x}} \simeq 0.205$, and $t_{32;0.025} = 2.0369$. The observed confidence interval is [492.489, 542.459].
 (iii) $\hat{y}_0 = 448.937$.
 (iv) The observed confidence interval is [344.442, 553.432].
- 5.4 (i) $\hat{y}_0 = 15.374$.
 (ii) The observed confidence interval is [14.329, 16.419].
 (iii) $\hat{y}_0 = 15.374$.
 (iv) The observed confidence interval is [14.278, 16.470].
- 5.6 (i) $\hat{\beta}_1 \simeq -9.768$, $\hat{\beta}_2 \simeq 2.941$, $\hat{\sigma}^2 \simeq 0.004$.
 (ii) $t_{11;0.025} = 2.201$, $S \simeq 0.071$, $\sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SS_x}} \simeq 10.575$, so that the observed confidence intervals for β_1 and β_2 are: [-11.421, -8.115] and [2.490, 3.392], respectively. Since $\chi_{11;0.025}^2 = 21.92$ and $\chi_{11;0.975}^2 = 3.816$, the observed confidence interval for σ^2 is [0.003, 0.015].
 (iii) Both EY_0 and Y_0 are predicted by 1.32. The respective observed confidence intervals are [1.254, 1.386] and [1.15, 1.49].

Chapter 14

Section 14.1

1.2 In (1), set $\mu_1 = \cdots = \mu_I = \mu$ to obtain:

$$\log L(\mathbf{y}; \mu, \sigma^2) = -\frac{IJ}{2} \log(2\pi) - \frac{IJ}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_i \sum_j (y_{ij} - \mu)^2.$$

Set $S(\mu) = \sum_i \sum_j (y_{ij} - \mu)^2$, and observe that $\frac{d}{d\mu} S(\mu) = -2 \sum_i \sum_j (y_{ij} - \mu) = 0$ gives $\mu = \frac{1}{IJ} \sum_i \sum_j y_{ij} = y_{..}$, and $\frac{d^2}{d\mu^2} S(\mu) = 2IJ > 0$ for all values of μ , so that $\hat{\mu} = y_{..}$ minimizes $S(\mu)$. Replacing μ by $\hat{\mu}$ in the above expressions, and setting

$$\hat{S} = S(\hat{\mu}), \quad \log \hat{L}(\mathbf{y}; \hat{\mu}, \sigma^2) = -\frac{IJ}{2} \log(2\pi) - \frac{IJ}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \hat{S},$$

we obtain $\sigma_{H_0}^2 = \frac{\hat{S}}{IJ}$ from $\frac{d}{d\sigma^2} \log \hat{L}(\mathbf{y}; \hat{\mu}, \sigma^2) = 0$. Also,

$$\frac{d^2}{d(\sigma^2)^2} \log \hat{L}(\mathbf{y}; \hat{\mu}, \sigma^2) \Big|_{\sigma^2 = \sigma_{H_0}^2} = -\frac{IJ}{(\sigma_{H_0}^2)^2} < 0,$$

so that $\widehat{\sigma_{H_0}^2} = \frac{\hat{S}}{IJ} = \frac{1}{IJ} \sum_i \sum_j (y_{ij} - y_{..})^2 = \frac{SS_T}{IJ}$ is the MLE of σ^2 under H_0 .

1.4 Recall that for $i = 1, \dots, I$ and $j = 1, \dots, J$, the r.v.'s Y_{ij} are independent with $EY_{ij} = \mu_i$ and $\text{Var}(Y_{ij}) = \sigma^2$.

- (i) From $Y_{..} = \frac{1}{IJ} \sum_j \sum_i Y_{ij}$, we have then $EY_{..} = \frac{1}{IJ} \sum_i \sum_j EY_{ij} = \frac{1}{J} \times j \times \frac{1}{I} \sum_i \mu_i = \mu_{.}$, and therefore $E(Y_{..} - \mu_{.})^2 = \text{Var}(Y_{..}) = \text{Var}(\frac{1}{IJ} \sum_i \sum_j Y_{ij}) = \frac{1}{(IJ)^2} IJ\sigma^2 = \frac{\sigma^2}{IJ}$.
- (ii) From $Y_{i.} = \frac{1}{J} \sum_j Y_{ij}$, we have $EY_{i.} = \frac{1}{J} \sum_j \mu_i = \frac{1}{J} \times J\mu_i = \mu_i$, so that $E(Y_{i.} - \mu_i)^2 = \text{Var}(Y_{i.}) = \text{Var}(\frac{1}{J} \sum_j Y_{ij}) = \frac{1}{J^2} \sum_j \sigma^2 = \frac{\sigma^2}{J}$.
- (iii) $E(Y_{i.} - \mu_{.})^2 = E[(Y_{i.} - \mu_i) + (\mu_i - \mu_{.})]^2 = E(Y_{i.} - \mu_i)^2 + (\mu_i - \mu_{.})^2$, because $E[(Y_{i.} - \mu_i)(\mu_i - \mu_{.})] = (\mu_i - \mu_{.})E(Y_{i.} - \mu_i) = (\mu_i - \mu_{.}) \times 0 = 0$.

Section 14.2

2.2 We have: $\hat{\mu}_1 = 11.25$, $\hat{\mu}_2 = 17.00$, $\hat{\mu}_3 = 15.50$, $\hat{\mu}_4 = 14.75$, so that $\hat{\psi} = 11.25c_1 + 17.00c_2 + 15.50c_3 + 14.75c_4$. Also, $\widehat{\text{Var}}(\hat{\psi}) \simeq 1.552(\sum_{i=1}^4 c_i^2)$, and $S^2 = 10.4709$, so that $S\sqrt{\widehat{\text{Var}}(\hat{\psi})} \simeq 4.031\sqrt{\sum_{i=1}^4 c_i^2}$. Therefore the required observed confidence interval is:

$$\left[11.25c_1 + 17.00c_2 + 15.50c_3 + 14.75c_4 \pm 4.031 \sqrt{\sum_{i=1}^4 c_i^2} \right].$$

Section 14.3

3.2 By (31) and (32), $L_A(\mathbf{y}; \mu, \beta, \sigma^2) = (\frac{1}{\sqrt{2\pi\sigma^2}})^{IJ} \exp[-S(\mu, \beta)]$, where $S(\mu, \beta) = \sum_i \sum_j (y_{ij} - \mu - \beta_j)^2$. For each fixed σ^2 , minimize $S(\mu, \beta)$ with respect to μ and the β_j 's subject to the restriction $\sum_j \beta_j = 0$. Doing this minimization by using Lagrange multipliers, we find the required MLE's; namely, $\hat{\mu}_A = y_{.}$, $\hat{\beta}_{j,A} = y_{.j} - y_{.}$, $j = 1, \dots, J$.

- 3.4 (i) That $\boldsymbol{\eta} = \mathbf{X}'\boldsymbol{\beta}$ is immediate, and from this it follows that $\boldsymbol{\eta}$ lies in the vector space generated by the columns (rows) of \mathbf{X}' .
- (ii) Here $I + J + 1 \leq IJ$, or $J \geq \frac{I+1}{I-1}$, provided $I \geq 2$. Thus, for $I \geq 2$ and $J \geq (I+1)/(I-1)$, it follows that $\min\{I + J + 1, IJ\} = I + J + 1$, and hence $\text{rank } \mathbf{X}' \leq I + J + 1$.
- (iii) Parts (a) and (b) are immediate. It then follows that $\text{rank } \mathbf{X}' \leq I + J - 1$. To see part (c), multiply the columns specified by the respective scalars $a_1, a_2, \dots, a_{I-1}, b_1, \dots, b_J$ and add them up to obtain

$$(b_1, b_2, \dots, b_J, a_1 + b_1, a_1 + b_2, \dots, a_1 + b_J, \dots, a_{I-1} + b_1, a_{I-1} + b_2, \dots, a_{I-1} + b_J),$$

and this vector is zero if and only if $b_1 = \dots = b_J = 0 = a_1 = \dots = a_{I-1}$. The conclusion of independence follows.

So, $\boldsymbol{\eta}$, although it has IJ coordinates, belongs in an $(I + J - 1)$ -dimensional space ($I + J - 1 \leq IJ$), and therefore the dimension of $\boldsymbol{\eta}$ is $I + J - 1$.

INDEX

2-dimensional normal distribution, 143, 147
2-dimensional random vector, 110, 138

A

a priori probabilities, 45, 271
Acceptance region, 233, 361
Analysis of variance, 235, 397
 basics, 236–238
 column effects, 418
 contrasts, 237, 407, 408
 examples, 403–404, 409–413, 419–420
 general linear models, 421
 multicomparison method, 407–412
 one-way (classification) layout model.
 See One-way (classification) layout
 model
 row effects, 418
 tables for, 402, 418
 two models of, 397–427
 two-way (classification) layout model.
 See Two-way (classification) layout
 model
 uses, 236–237
ANOVA. *See* Analysis of variance
Asymptotically
 normal, 444, 445, 446
 unbiased, 443, 445
Axiomatic definition of probability, 25–26

B

Bayes approach, 272
 decision function, 359
 estimates, 271, 274, 278
 formula, 46–48
Best linear predictor, 137
Beta
 p.d.f., 273
 expectation, 275
Binomial distribution, 79–81
 application, 315–316
 cumulative table, 450–457

 expectation, 81, 477
 graphs, 80
 m.g.f., 81, 479
 point estimation and, 228
 Poisson distribution relationship to, 84–85,
 98
 p.d.f., 79, 477
 variance, 81, 477
Binomial experiment, 79, 140
Bivariate normal distribution, 143–146
 correlation coefficient, 148
 example, 145–146
 expectations, 148, 478
 graph, 143
 m.g.f., 146, 479
 p.d.f., 143, 478
 variances, 148, 478

C

Cauchy distribution, 76, 325
Cauchy-Schwarz inequality, 131
Center of gravity, 69
Central limit theorem (CLT), 90, 208, 210–213,
 225
 applications, 213–215
 binomial and, 214
 confidence intervals and, 295
 continuity correction and, 215–217
 examples, 212, 214–215
 normal approximation and, 211
Chi-square distribution
 critical values for, 465–466
 expectation, 89, 477
 graph, 90
 m.g.f., 89, 479
 p.d.f., 89, 477
 variance, 89, 477
Chi-square goodness-of-fit test. *See* Goodness-
 of-fit test
Combinations, 61
Completeness, 264

- Concept of probability and basic results.
See Probability and basic results, concept of
- Conditional and marginal probability density functions, conditional expectation and variance, 117–126
- examples, 118–123
- exercises, 123–126
- k random variables and, 138
- multinomial distribution and, 140
- Conditional expectations, 138
- Conditional probability and related results, 41–51
- Bayes formula, 46–48
- definition, 42
- examples, 42, 43–44, 45
- exercises, 48–51
- multiplicative theorem, 44
- total probability theorem, 45–46
- Conditional expectation, 120
- variance, 123, 138
- Confidence intervals, 228, 282–292
- with approximate confidence coefficient, 282, 294–298
- construction steps, 283, 407–408
- definition, 282
- examples, 283–285, 289–292, 295–296
- exercises, 285–289, 292, 296–298
- for quantiles of distribution function, 431–433
- in linear regression model, 374–375, 383–389
- nuisance parameters, presence of, 289–292
- random interval and, 230–231
- significance of, 282
- with given approximate confidence coefficient, 294–296, 429–431
- Confidence regions, 292–294, 360
- confidence interval and, 231
- examples, 292–294, 361–362
- testing hypotheses relationship to, 360–362
- theorem, 361
- Contingency tables, likelihood ratio tests in multinomial case and, 343, 345–348
- Continuity correction, 215–217
- Continuous case distribution, 86–95
- chi-square, 89, 90
- gamma, 86–87
- median and, 102, 103
- negative exponential, 88–89
- normal, 89, 90–93
- uniform (or rectangular), 94–95
- Continuous sample spaces, 13
- Convergence modes of random variables, applications, 202–226
- central limit theorem, 210–215
- continuity correction, 215–217
- in distribution or in probability, 202–208
- further limit theorems, 222–226
- weak law of large numbers, 208–210
- Correlation coefficient, 132
- Counting, basic concepts and results in, 59–67
- exercises, 64–67
- fundamental principle of counting, 59–60, 61–62
- problem of counting examples, 59, 62–64
- Covariance, 126, 129–135
- Cramér-Rao (C-R) inequality, 262–263, 265, 266
- examples, 264
- usage, 263
- Cramér-Wold device, 139
- Critical or rejection region, 233
- Critical values for chi-square distribution table, 465–466
- Critical values for F -distribution table, 467–476
- Critical values for student's t -distribution table, 463–464
- Cumulative binomial distribution table, 450–457
- Cumulative Poisson distribution table, 458–459
- Curve estimation, nonparametric, 442–449
- ## D
- Decision-theoretic approach to estimation, 229, 270–277
- Bayes estimate, 271, 274
- decision function, 272
- examples, 273, 274–275
- exercises, 275
- loss function, 270
- minimax, 270, 274, 275
- risk function, 270
- theorems, 271, 274
- Decision-theoretic approach to testing hypotheses, 234, 353–360
- Bayes decision function, 359
- examples, 356–359
- loss function, 354
- minimax, 354–355
- nonrandomized decision function and, 353–354
- risk function, 354
- theorem, 355
- Delta method, 225
- DeMorgan's laws, 15, 16
- Dependent events, 51
- r.v.'s, 151
- Discrete case distributions, 79–86
- binomial, 79–81

- geometric, 81–82, 83
- hypergeometric, 85–86
- median and, 102–103
- Poisson, 83–85
- Discrete sample spaces, 13
- Disjoint, 14
- Dispersion, 71
- Distribution(s)
 - 2-dimensional normal, 143, 147
 - bivariate normal, 143–146
 - characteristics table, 477–479
 - confidence intervals for quantiles of, 431–433
 - convergence, 202–222
 - function (d.f.), 34–35
 - graphs, 34–35, 203
 - joint, 110–117
 - k -variate normal, 146, 147
 - marginal and conditional, 117–126
 - multinomial, 140–142, 143
 - multivariate normal, 146–147
 - of random variable X properties, 34, 37
 - probability, 34, 69
 - reproductive property of certain distributions, 159–167
- Distribution of random variables, 33–41.
 - See also* Distribution and Probability density function
 - beta, 273
 - binomial, 79–81
 - Cauchy, 76, 325
 - chi-square, 89
 - continuous, 86–95
 - discrete, 79–86
 - double exponential, 247
 - exercises, 39–41
 - F , 179
 - function graphs, 34–35
 - gamma, 86–87
 - geometric, 81–82, 83
 - hypergeometric, 85–86
 - mode, 104–106
 - negative exponential, 88–89
 - normal, 89, 90–93
 - Poisson, 83–85
 - t , 177
 - uniform (or rectangular), 94–95
 - Weibull, 99
- E**
- Effects
 - column, 412, 418
 - row, 412, 418
- Error(s) sum of squares, 365, 402, 418
 - type I, 233
 - type II, 233
- Event(s), 8
 - certain, 8
 - complement of, 13
 - difference of, 14
 - disjoint, 14, 26
 - happens, 8
 - impossible, 8
 - intersection of, 13, 14
 - monotone, 16–17
 - occurs, 8
 - related results and independent events, 51–59
 - union of, 13
- Expectation of random variables, 68–77
 - definition, 69
 - examples, 70–74
 - exercises, 74–77
 - of selected discrete and continuous distributions, 477, 478
- Exponential type of families of probability density functions, 307–308
- F**
- F critical values for, 467–476
 - expectation, 181
 - graph, 180
 - p.d.f., 179
 - variance, 181
- Factorization theorem, 151–152, 154
- Failure rate, 99
- Fisher information, 259
- Fisher-Neyman factorization theorem, 256
- Fitted regression line, 366, 370, 372, 377, 378
- Fundamental concepts, 8–19
 - events, 8
 - exercises, 17–19
 - intersection of events, 13, 14
 - mutually or pairwise disjoint, 14
 - random experiment, 8
 - sample points, 8, 13
 - sample space, 8, 13
 - union of events, 13
 - Venn diagram, 8, 9, 13–15
- Fundamental principle of counting theorem, 60, 61
 - corollary to, 61
 - proof of, 61–62
 - proof of corollary, 62
- Further limit theorems, 222–226
- G**
- Gamma distribution
 - expectation, 87, 477
 - graph, 86–87
 - m.g.f., 87, 479

- p.d.f., 86, 477
 - variance, 87, 477
 - Gamma function, 86
 - recursive property, 87
 - Geometric distribution
 - expectation, 82, 83, 477
 - graphs, 82
 - m.g.f., 82, 83, 479
 - p.d.f., 82, 83, 477
 - variance, 82, 83, 477
 - Goodness-of-fit test, 234, 349–353
 - exercises, 351–353
 - multinomial distribution and, 349
 - numerical examples, 350–351
 - Graduate Management Aptitude Test (GMAT), 7, 364, 384
- H**
- Hypergeometric distribution
 - expectation, 85, 477
 - p.d.f., 85, 477
 - variance, 85, 477
- I**
- Independence of random variables and some
 - applications, 150–167
 - criteria, 150–159
 - definition, 151
 - examples, 152–153, 162–163
 - exercises, 156–159, 164–167
 - factorization theorem, 151–152, 154
 - reproductive property, 159–167
 - Independent events and related results, 51–55
 - definitions, 51, 52–53, 55
 - examples, 52, 54–55
 - exercises, 55–59
 - theorem, 53
 - Interval estimation, 228
 - basics, 230–231
 - confidence interval, 231
 - lower confidence limit, 231
 - random, 230–231, 282
 - statistic(s), 230
 - upper confidence limit, 231, 282
 - Inversion formula, 73, 129
- J**
- Joint moment generating function, 126–128, 146
 - Joint probability distribution functions, 110–117, 137
 - examples, 111–115
 - exercises, 115–117
 - k random variables and, 137–138
- K**
- k random variables
 - generalizations, 137–139
 - sample mean and sample variance of, 159
 - k -variate normal distribution, 146, 147, 478
 - kernel-estimation approach, 239, 442, 443
 - Kolmogorov, 25
- L**
- Lagrange multipliers, 409
 - Least squares (LS's), 230, 277
 - estimate (LSE), 230, 363, 366–374
 - examples, 369–374
 - fitted regression line, 366, 370, 372
 - minimize sum of squares error, 365–366
 - pairs relationship, 364–365
 - principle of, 363, 365–366
 - regression line, 366
 - theorems, 366, 371, 373
 - Lehmann-Scheffé theorem, 265
 - Level of significance of test employed, 233, 234
 - Likelihood equation(s), 242, 245
 - Likelihood function, 229, 234, 241, 247, 254, 375, 398, 413
 - maximum, 325
 - Likelihood ratio (LR) tests, 234, 299, 324–342
 - applications, 327–337, 345–347
 - examples, 325–327
 - exercises, 337–342, 348–349
 - in multinomial case and contingency tables, 343–349
 - linear regression model and, 384
 - motivation, 235, 324
 - normal case applications, 327–337
 - numerical examples, 328, 330, 331, 335, 344, 348
 - one-way layout and, 398
 - theorem, 343–344
 - two-way layout and, 415
 - Linear regression model, simple, 236, 363–396
 - concluding remarks, 395–396
 - confidence intervals, 374
 - confidence intervals and hypotheses testing problems, 383–389
 - errors, normally distributed, 374–383, 393–395
 - examples, 369–370, 376–379, 384–386
 - fitted regression line, 366, 370, 372, 377
 - general, 396, 421
 - least squares estimates of β_1 and β_2 , 366–374
 - least squares principle, 365–366
 - likelihood ratio tests and, 384
 - multiple, 396

- pairs relationship, 364–365
- prediction problems, 389–393
- regression line, 366
- setting up, 364–366
- theorems, 366, 371, 373, 375, 379, 383, 384, 390, 391
- Linear transformations, 185–191
 - definition, 186
 - exercises, 190–191
 - orthogonal, 186
 - theorems, 187, 189
- Lower confidence limit, 231, 282
- M**
- Marginal and conditional probability density functions, conditional expectation and variance, 117–126
 - examples, 118–123
 - exercises, 123–126, 139, 149
 - k random variables and, 138
 - multinomial distribution and, 140
- Marginal moment generating function, 128
- Markov inequality, 77
- Mathematical expectation. *See* Expectation of random variables
- Matrix
 - orthogonal, 186
 - transpose, 186
- Maximum likelihood estimates (MLE's), 229, 278
 - definition, 241, 242
 - Fisher-Neyman factorization theorem, 256
 - identification of, 242
 - invariance property, 254
 - linear regression model and, 374–383
 - motivation and examples, 240–253
 - one-way layout and parameters, 399
 - properties, 253–261
 - sufficient statistics, 257–258
 - theorems, 253, 254, 256, 258–259
 - two-way layout and parameters, 413–414
- Maximum likelihood function, 324–326
- Mean, mean value, *See* expectation
- Measure of
 - dispersion, 71
 - location, 69, 71
- Median and mode of random variables, 102–108
 - continuous case, 102, 103
 - definitions, 103, 104
 - discrete case, 102–103
 - examples, 102–106
 - exercises, 106–108
- Method of moments, 229, 277
 - exercises, 278–279
- Minimax
 - decision function, 354–357
 - estimate, 270, 274, 275, 278
- Minimum chi-square method, 277
- Mode of distribution, 104–106
- Moment estimates, 277
- Moment generating function (m.g.f.), 72–74, 93
 - inversion formula and, 73, 129
 - joint, 126–129, 138, 142, 146
 - marginal, 128, 130
 - of selected discrete and continuous distributions, 479
- Moments, 71
- Monotone events, 16–17
- Most powerful (MP) test, 234, 303
- Motivating examples, 1–8
- Multicomparison method of analysis of variance, 407–412
- Multinomial distribution
 - correlation coefficients, 143
 - examples, 140–143
 - expectations, 142, 478
 - goodness-of-fit test and, 349
 - likelihood ratio tests and, 343–349
 - marginal and conditional probability density functions and, 140
 - m.g.f., 142, 479
 - p.d.f., 140, 478
 - theorem, 141
 - variances, 142–143, 478
- Multiple random variables, transforming, 173–185
- Multiplicative theorem, 44
- Multivariate normal distribution, 146–147
- N**
- Negative exponential distribution, 88
 - expectation, 88, 477
 - graph, 88
 - m.g.f., 88, 479
 - p.d.f., 88, 477
 - variance, 88, 477
- Negatively correlated, 133
- Neyman-Pearson fundamental lemma, 299, 302–307
 - application examples, 305–307
 - most powerful test and, 303, 305–306
 - proof of theorem, 303–304
 - uniformly most powerful test and, 306
- Nonparametric curve estimation, 442–449
- Nonparametric inference, topics in, 428–449
 - basics, 238–239
 - confidence intervals with given confidence coefficient, 429–431

- confidence intervals for quantiles of
 - distribution function, 431–433
- curve estimation, 442–449
- definition, 227
- kernel method, 239, 442
- probability density function estimation, 442–444
- rank sum tests, 435–438, 440–442
- regression estimation, 444–447
- two-sample sign test, 433–435
- weak law of large numbers and, 239
- Wilcoxon-Mann-Whitney test, 438
- Nonrandomized decision function, 353
- Nonrandomized test, 233, 303
- Normal distribution, 89, 90–93, 95
 - central limit theorem, 90
 - expectation, 92, 477
 - graphs, 90, 91
 - importance of, 89, 90
 - likelihood ratio tests and, 327–337
 - m.g.f., 93, 479
 - p.d.f., 90, 477
 - standard, 91
 - table, 460–462
 - testing hypotheses about the mean, 317–319, 320–321
 - testing hypotheses about the variance, 319–320
 - variance, 92, 477
- Nuisance parameters, confidence intervals in presence of, 289–292
- Numerical characteristics of random variables, 68–108
 - expectation, variance, and moment generating function, 68–77
- O**
- One-way (classification) layout model analysis of variance, 237, 397, 398–407
- Order statistics, 193–201
 - definition, 193
 - examples, 196, 197–199
 - exercises, 199–201
 - theorems, 194, 196
- P**
- Parameter space, 227, 228
- Parametric statistical inference, 227
- Partition, 45
- Permutations, 61
- Point estimation, 227, 240–280
 - basics, 228–230
 - binomial distribution and, 228
 - decision-theoretic method, 229, 270–277
 - least squares, 230, 277, 364–374
 - maximum likelihood estimate, 229, 240–246
 - maximum likelihood estimation motivation and examples, 240–253
 - maximum likelihood estimation properties, 253–261
 - method of moments, 229, 277
 - other methods, 277–280
 - parameter(s), 228, 240
 - Poisson distribution and, 228
 - unbiasedness, 229
 - uniformly minimum variance unbiased, 229, 261–270
- Point of equilibrium, 69
- Poisson distribution, 83–85, 95, 215
 - application, 316–317
 - binomial distribution relationship to, 84–85
 - cumulative table for, 458–459
 - expectation, 84, 477
 - graph, 83
 - m.g.f., 84, 479
 - point estimation and, 228
 - p.d.f., 83, 477
 - uses for, 84
 - variance, 84, 477
- Principle of least squares. *See* Least squares (LS's)
- Probability
 - axiomatic definition, 25–26
 - classical definition, 24–25
 - conditional, 41
 - inequalities, 77–79
 - justification of basic properties, 28–29
 - relative frequency definition, 25
- Probability and basic results, concept of, 23–67
 - conditional probability and related results, 41–51
 - counting, basic concepts and results in, 59–67
 - definition, 24–26
 - examples in calculating probabilities, 26–31
 - independent events and related results, 51–59
 - random variable distribution, 33–41
 - theorems, 31, 44, 45, 46–47, 53, 60
- Probability density function (p.d.f.), 37
 - definition, 35, 36
 - graph examples, 80, 82, 83, 86–88, 90, 91, 94, 178, 180
 - nonparametric estimation of, 442–444
 - of selected discrete and continuous distributions, 477–479
 - probability inequalities, 77

Probability integral transform, 192–193
 Prior probabilities, 45

Q

Quantile(s), 103

R

Random experiment, 8
 examples, 2–8
 Random interval, 230–231, 282
 Random sample, 155
 Random variables (r.v.'s)
 continuous, 36, 86–95
 convergence modes of, 202–226
 definition, 20
 degrees of freedom (d.f.), 34
 denoting a, 20
 discrete, 36, 79–86
 distribution, 33–41
 exercises, 21–22
 expectation, variance, and moment
 generating function, 68–77
 independence of, 150–167
 introduction of, 19–21
 k, generalization to, 137–139
 median and mode of, 102–108
 numerical characteristics of, 68–108
 special, 79–101
 transformation of, 168–201
 types of, 21
 Randomized tests, 233, 303
 Rank sum test, 435–438, 440–441
 Rank test, 239, 429, 435–438, 440–442
 Rao-Blackwell theorem, 265
 Recursive relation for
 binomial p.d.f., 95, 96
 gamma function, 87
 hypergeometric p.d.f., 98
 Poisson p.d.f., 97
 Regression analysis
 basics, 235–236
 linear regression model, 236,
 363–396
 simplest form, 235–236
 Regression estimation, nonparametric,
 444–447
 Regression line, 366
 Regression model
 fixed design, 447
 linear. *See* Linear regression model, simple
 stochastic design, 447
 Relative frequency definition of probability,
 25
 Reproductive property of certain distribution,
 159–167

 examples, 162–163
 exercises, 164–167
 theorems, 160–161, 164
 Risk function, 270, 354

S

Sample mean, 159
 Sample points, 8, 13
 Sample range, 200
 Sample space, 8
 continuous, 13
 discrete, 13
 examples with countably infinite points,
 10–11
 examples with finitely many sample points,
 9–10
 examples with nondegenerate finite or
 infinite intervals in real line, 11
 random experiment and, 19, 20
 Sample variance, 159
 Sign test, 239, 433–435
 Single random variables, transforming, 168–173
 Standard deviation (s.d.), 71–72, 77
 Statistic, 229
 sufficient, 256
 Statistical analysis. *See* Analysis of variance
 Statistical hypothesis, 232
 alternative, 232
 null, 232
 Statistical inference overview, 227–239
 aim, 227
 analysis of variance basics, 236–238
 interval estimation basics, 228, 230–231
 nonparametric inference basics, 227,
 238–239
 parametric, 227
 point estimation basics, 227, 228–230
 regression analysis basics, 235–236
 testing hypotheses basics, 228, 231–235
 Stirling formula, 185

T

t distribution, 177, 178
 expectation, 172
 p.d.f., 177
 variance, 172
 Tchebichev inequality, 77, 208
 Test function, 232
 Testing hypotheses, 228, 299–342, 343–362
 acceptance region, 233, 361
 basics, 231–235
 binomial case application, 315–316
 concepts, general, 300–302
 confidence regions relationship to, 360–362
 critical or rejection region, 233

- decision-theoretic approach, 234, 353–360
- equality of means, 399–404
- exponential type families of probability
 - density functions, 307–308
- for parameters in single normal population, 327–332
- for parameters in two normal populations, 332–337
- formulating, 300–302
- goodness-of-fit test, 234, 349–353
- level of significance, 233, 234
- likelihood ratio tests, 234, 235, 299, 324–342
- linear regression model and problems in, 383–389, 389–392
- most powerful test, 234, 303, 305, 306
- Neyman-Pearson fundamental lemma, 299, 302–307
- nonrandomized test, 233, 303
- normal case application, 317–321
- p-value (probability value), 235
- Poisson case application, 316–317
- power, 234, 304, 309, 311, 312, 318, 319, 321, 328, 340
- randomized tests, 233, 303
- statistical hypotheses, 232
- two-way layout, 414–420
- type I error, 233
- type II error, 233–234
- uniformly most powerful tests, 234, 299, 301, 306
- uniformly most powerful tests for composite hypotheses, 308–312, 315–321
- Theorems, 31–32, 104, 105, 131, 134, 135
- Bayes formula, 46–48, 271, 359
- Cauchy-Schwarz inequality, 131
- central limit, 90, 208, 210–215
- confidence interval for contrasts, 408
 - LSE's (MLE's), 383
 - predictor, 390, 391
 - quantiles, 432
- confidence regions and testing hypotheses, 361
- continuity, 206
- convergence in distribution, 203, 206
- convergence in probability, 204, 206
- convergence of MLE, 258, 259
- correlation coefficient, 134
- Cramér-Rao inequality, 263
- decision-theoretic approach to estimation, 271, 274
- decision-theoretic approach to testing hypotheses, 355
- decomposition of total variability, 373–374
- distribution of LSE's (MLE's), 375
- distribution of sums of squares, 379, 393–395
- factorization, 151–152, 154
- Fisher-Neymann factorization, 256
- fundamental principle of counting, 60, 61–62
- further limit, 222–226
- independence of sample mean and sample variance in a normal distribution, 163, 189
- independent events, 53
- invariance property of MLE, 253, 254
- LSE's, 366
- likelihood ratio tests, 343–344
- linear regression model, 366, 370, 373, 375, 379, 383, 384, 390, 391
- linear transformations, 187, 189
- maximum likelihood estimates, 253, 254, 258–259
- minimax decision function, 355–356
- minimax estimate, 274
- mode, 104, 105
- multicomparison method in analysis of variance, 407
- multinomial distribution, 141
- multiplicative, 44
- Neyman-Pearson fundamental lemma, 302–303
- nonparametric curve estimation, 443–444
- nonparametric inference, 432
- nonparametric regression estimation, 445, 446
- one-way layout model, 399, 401
- order statistics, 194, 196
- probability inequalities, 77, 78
- probability integral transform, 192
- rank sum test, 437
- Rao-Blackwell and Lehmann-Scheffé, 265
- reproductive property of distributions, 160–161, 164
- sign test, 434
- Slutsky, 223
- testing hypotheses in linear regression model, 384–385
- total probability, 45–46
- transforming multiple random variables, 174, 182–183
- transforming single random variable, 169, 170, 171
- two-sample sign test, 434
- two-way layout, 414, 418
- uniformly most powerful test for composite hypotheses, 308–309, 311
- variance of LSE's, 371
- variance of sums of r.v.'s, 135, 139
- WLLN, 208
- Wilcoxon-Mann-Whitney test, 438

- Total probability theorem, 45–46
- Transformation of random variables,
168–201
examples, 168, 170–171, 173, 174–181,
192–193, 196–199
exercises, 171–173, 183–185, 190–191, 193,
199–201
linear, 185–191
order statistics, 193–201
orthogonal, 186
probability integral transform, 192–193
single, 168–173
two or more, 173–185
- Triangular probability density function,
176
- Two-sample sign test, 433–435
- Two-way (classification) layout model, 238,
412–420
examples, 412–413, 419, 420
exercises, 425–427
lemmas and proof, 413–414, 415–418,
420–425
maximum likelihood estimates parameters,
413–414
table, 418
testing hypotheses, 414–420
theorems, 414, 418
with one observation per cell, 412–427
- U**
- Unbiasedness, 229
- Uniform (or rectangular) distribution, 94–95,
477
expectation, 94, 477
m.g.f., 94, 479
p.d.f., 94, 477
variance, 94, 477
- Uniformly minimum variance unbiased
(UMVU) estimates, 229, 261–270, 278
- completeness, 264
- Cramér-Rao inequality, 262–264
- definition, 261, 262
- desirability of, 262
- examples, 261–262, 264, 265–266
- Rao-Blackwell and Lehmann-Scheffé
theorems, 265
- Uniformly most powerful (UMP) tests, 234, 299,
301
exercises, 313–314
for composite hypotheses, 308–312
Neyman-Pearson fundamental lemma and,
306
power for one-sided hypotheses, 310
power for two-sided hypotheses, 312
- Upper confidence limit, 231
- V**
- Variance analysis basics, 236–238
- Variance analysis models. *See* Analysis of
variance
- Variance of random variables, 71–72
of selected discrete and continuous
distributions, 477, 478
- Venn diagram, 9, 13–15
- W**
- Weak law of large numbers (WLLN), 208–210,
224, 225, 277
applications, 209–213
confidence intervals and, 295
example, 212–213
interpretation and most common use, 209
nonparametric inference and, 239
theorem, 208
- Weibull distribution, 99
- Wilcoxon-Mann-Whitney test, 429, 438
examples, 439–440

This Page Intentionally Left Blank

Table of Selected Discrete and Continuous Distributions and Some of their Characteristics

PROBABILITY DENSITY FUNCTIONS IN ONE VARIABLE			
Distribution	Probability Density Function	Mean	Variance
Binomial, $B(n, p)$	$f(x) = \binom{n}{x} p^x q^{n-x}, x = 0, 1, \dots, n;$ $0 < p < 1, q = 1 - p$	np	npq
(Bernoulli, $B(1, p)$)	$f(x) = p^x q^{1-x}, x = 0, 1$	p	pq
Geometric	$f(x) = pq^{x-1}, x = 1, 2, \dots;$ $0 < p < 1, q = 1 - p$	$\frac{1}{p}$	$\frac{q}{p^2}$
Poisson, $P(\lambda)$	$f(x) = e^{-\lambda} \frac{\lambda^x}{x!}, x = 0, 1, \dots; \lambda > 0$	λ	λ
Hypergeometric	$f(x) = \frac{\binom{m}{x} \binom{n}{r-x}}{\binom{m+n}{r}}, \text{ where}$ $x = 0, 1, \dots, r \left(\binom{m}{r} = 0, r > m \right)$	$\frac{mr}{m+n}$	$\frac{mnr(m+n-r)}{(m+n)^2(m+n-1)}$
Gamma	$f(x) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} \exp\left(-\frac{x}{\beta}\right), x > 0;$ $\alpha, \beta > 0$	$\alpha\beta$	$\alpha\beta^2$
Negative Exponential	$f(x) = \lambda \exp(-\lambda x), x > 0; \lambda > 0; \text{ or}$ $f(x) = \frac{1}{\mu} e^{-x/\mu}, x > 0; \mu > 0$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$
Chi-Square	$f(x) = \frac{1}{\Gamma\left(\frac{r}{2}\right) 2^{r/2}} x^{r/2-1} \exp\left(-\frac{x}{2}\right), x > 0;$ $r > 0 \text{ integer}$	r	$2r$
Normal, $N(\mu, \sigma^2)$	$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2}\right],$ $x \in \Re; \mu \in \Re, \sigma > 0$	μ	σ^2
(Standard Normal, $N(0, 1)$)	$f(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right), x \in \Re$	0	1
Uniform, $U(\alpha, \beta)$	$f(x) = \frac{1}{\beta - \alpha}, \alpha \leq x \leq \beta;$ $-\infty < \alpha < \beta < \infty$	$\frac{\alpha + \beta}{2}$	$\frac{(\alpha - \beta)^2}{12}$

(Continued)

PROBABILITY DENSITY FUNCTIONS IN MANY VARIABLES

Distribution	Probability Density Function	Means	Variances
Multinomial	$f(x_1, \dots, x_k) = \frac{n!}{x_1!x_2! \dots x_k!} \times$ $p_1^{x_1} p_2^{x_2} \dots p_k^{x_k}, x_i \geq 0 \text{ integers,}$ $x_1 + x_2 + \dots + x_k = n; p_j > 0, j = 1,$ $2, \dots, k, p_1 + p_2 + \dots + p_k = 1$	np_1, \dots, np_k	$np_1q_1, \dots, np_kq_k.$ $q_i = 1 - p_i, j = 1, \dots, k$
Bivariate Normal	$f(x_1, x_2) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \exp\left(-\frac{q}{2}\right),$ $q = \frac{1}{1-\rho^2} \left[\left(\frac{x_1-\mu_1}{\sigma_1}\right)^2 - 2\rho\left(\frac{x_1-\mu_1}{\sigma_1}\right) \right.$ $\left. \times \left(\frac{x_2-\mu_2}{\sigma_2}\right) + \left(\frac{x_2-\mu_2}{\sigma_2}\right)^2 \right],$ $x_1, x_2, \in \mathfrak{R}; \mu_1, \mu_2 \in \mathfrak{R}, \sigma_1, \sigma_2 > 0, -1 \leq \rho \leq 1, \rho = \text{correlation coefficient}$	μ_1, μ_2	σ_1^2, σ_2^2
k-Variate Normal, $N(\mu, \Sigma)$	$f(\mathbf{x}) = (2\pi)^{-k/2} \Sigma ^{-1/2} \times$ $\exp\left[-\frac{1}{2}(\mathbf{x} - \mu)' \Sigma^{-1}(\mathbf{x} - \mu),\right.$ $\mathbf{x} \in \mathfrak{R}^k; \mu \in \mathfrak{R}^k, \Sigma : k \times k$ <p style="margin-top: 10px;">nonsingular symmetric matrix</p>	μ_1, \dots, μ_k	Covariance matrix: Σ

Distribution	Moment Generating Function
Binomial , $B(n, p)$ (Bernoulli, $B(1, p)$)	$M(t) = (pe^t + q)^n, t \in \mathfrak{R}$ $M(t) = pe^t + q, t \in \mathfrak{R}$
Geometric	$M(t) = \frac{pe^t}{1 - qe^t}, t < -\log q$
Poisson , $P(\lambda)$	$M(t) = \exp(\lambda e^t - \lambda), t \in \mathfrak{R}$
Hypergeometric	—
Gamma	$M(t) = \frac{1}{(1 - \beta t)^\alpha}, t < \frac{1}{\beta}$
Negative Exponential	$M(t) = \frac{\lambda}{\lambda - t}, t < \lambda; \text{ or } M(t) = \frac{1}{1 - \mu t}, t < \frac{1}{\mu}$
Chi-Square	$M(t) = \frac{1}{(1 - 2t)^{r/2}}, t < \frac{1}{2}$
Normal , $N(\mu, \sigma^2)$	$M(t) = \exp\left(\mu t + \frac{\sigma^2 t^2}{2}\right), t \in \mathfrak{R}$
(Standard Normal, $N(0, 1)$)	$M(t) = \exp\left(\frac{t^2}{2}\right), t \in \mathfrak{R}$
Uniform , $U(\alpha, \beta)$	$M(t) = \frac{e^{t\beta} - e^{t\alpha}}{t(\beta - \alpha)}, t \in \mathfrak{R}$
Multinomial	$M(t_1, \dots, t_k) = (p_1 e^{t_1} + \dots + p_k e^{t_k})^n,$ $t_1, \dots, t_k \in \mathfrak{R}$
Bivariate Normal	$M(t_1, t_2) = \exp\left[\mu_1 t_1 + \mu_2 t_2 + \frac{1}{2}\left(\sigma_1^2 t_1^2 + 2\rho\sigma_1\sigma_2 t_1 t_2 + \sigma_2^2 t_2^2\right)\right],$ $t_1, t_2 \in \mathfrak{R}$
k-Variate Normal , $N(\mu, \Sigma)$	$M(\mathbf{t}) = \exp\left(\mathbf{t}'\mu + \frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}\right),$ $\mathbf{t} \in \mathfrak{R}^k$

This Page Intentionally Left Blank