Article

Land capability assessment by combining LESA and GIS in a calcareous watershed, Iran

March 2022 · Arabian Journal of Geosciences 15(5)
DOI: 10.1007/s12517-022-09729-5

Mohammad Akbari · Mohammad Tahmoures · Aliasghar Azma · Show all 5 authors · Farzad Tat Shahdost

Similar research

A new method is used to evaluate the strategic value of Fresno County farmland

Article Full-text available

July 2010 · California Agriculture

Evan E. Schmidt · James H. Thorne · Patrick R Huber · [...] · Michael Mccoy

Fresno County is a rich agricultural area that faces rapid urbanization and farmland conversion. The county is participating in a strategic, multi-county planning initiative aimed at making sustainable and regionally cohesive land-use decisions. To inform this effort, we conducted a farmland conservation assessment and identified strategic farmlands f…

98 Reads · 2 Citations

Download Recommend Follow Share

Land suitability assessment for maize farming using a GIS-AHP method for a semi- arid region, Iran

Article Full-text available

April 2020 · Journal of the Saudi Society of Agricultural Sciences

Behnam Tashayo · Afshin Honarbakhsh · Mohammad Akbari · Mobin Eftekhari

The aim of this study was to generate land suitability maps for maize farming in calcareous and saline-sodic soils in the Marvdasht plain, Iran. An analytic hierarchy process (AHP) multi-criteria method integrated with GIS and geostatictic was employed to estimate weighting of soil properties, climate and topography data. The results indicated that soil texture…

333 Reads · 21 Citations

Download Recommended Follow Share

Land Suitability Modeling For Ricecrop Based On An Integrated Multi-Criteria Decision Making In Quang Tri Province Of Vietnam

New Article Full-text available

October 2021 · GEOGRAPHY ENVIRONMENT SUSTAINABILITY

Thi Hanh Tong · Mai-Phuong Pham · Bui Thi Quyen · [...] · Altaf Hussain Lahori

The present study was aimed to determine the potential cultivated lands for rice crop production in Vietnam. Analytic Hierarchy Process (AHP) and the Technique for Order of Preference by Similarity to Ideal Solution (TOPSIS) were employed in order to analyze the structure of an ideal solution in agriculture that focuses mainly on environmental,…

104 Reads

Download Recommend Follow Share

Statistical Modeling for Spatial Groundwater Potential Map Based on GIS Technique

Article Full-text available

March 2021 · Sustainability

Aliasghar Azma · Esmaeil Narreie · Abouzar Shojaaddini · [...] · Afshin Davarpanah

In arid and semi-arid lands like Iran water is scarce, and not all the wastewater can be treated. Hence, groundwater

remains the primary and the principal source of water supply for human consumption. Therefore, this study attempted to spatially assess the groundwater potential in an aquifer in a semi-arid region of Iran using geographic information…

126 Reads · 7 Citations

Download

Recommended Follow Share

Soil erodibility prediction by Vis-NIR spectra and environmental covariates coupled with GIS, regression and PLSR in a watershed scale, Iran

New Article Full-text available

December 2021 · Geoderma Regional

Behnaz Salehi-Varnousfaderani · Afshin Honarbakhsh · Mohammad Tahmoures · Mohammad Akbari

Visible–Near-Infrared (Vis-NIR) spectroscopy as a rapid, cost-effective, and non-destructive technique has become an alternative approach to evaluate difficult-to-measured soil properties. Hence, this study aimed to evaluate RUSLE soil erodibility (so-called K-factor) in calcareous soils of a semi-humid watershed located in west of Iran using Vis-NIR…

23 Reads

Download

Recommended Follow Share

View more related research