

نقشه گسلهای فعال ایران

خاله حاسمی، فرشاد جمالی، هادی طیبی
گروه زلزله زمین ساخت - پژوهشکده زلزله شناسی

چاپ: ۱۳۸۲

Major Active Faults of Iran

Khaled Hessami, Farshad Jamali & Hadi Tabassi
Seismotectonic Department, Seismology Research Centre

Edition: 2003

The active faults of Iran are direct indicators of active crustal deformation due to the on-going continental collision between Arabia and Eurasia. Since Quaternary stratigraphy and geomorphic methods in the study of active faulting, except in very few cases have not been employed in Iran, the study of active faults is based on strong scrutiny of available earthquake data. However, this becomes difficult when studying structurally complex and inhomogeneous collision region such as Iran in which seismicity is not the result of the activity of a few faults but it is due to fault activity in zones hundreds kilometers wide. Due to the high density of active faults in Iran and the inaccuracy of the macroseismic data of the area, the source of some of the earthquakes have been related to more than one fault. Additionally lack of accurate fault and epicentral maps result in the inability to correlate seismic data with the active faulting in the area. Obviously even highly accurate seismic data will not solely suffice to locate all existing active faults. This is mainly due to: 1) the fact that many earthquakes are caused by blind faults such as earthquakes occurring in the Zagros region, and 2) the gradual movement along many active faults over a long period of time is the result of a creep regime which does not result in a large earthquake.

This map overviews the distribution of major active faults of Iran and demonstrates the relationship between the slip vectors and compressive axes, obtained from the solution of the focal mechanisms of the area's earthquakes, and GPS velocities in different areas of Iran.

Active faults in Zagros are blind and the focal mechanism solutions of the earthquakes of the region points to the presence of both thrust and transverse strike-slip faulting in its basement. Whereas in the rest of the country most active faults reach the surface. The earthquake mechanism solutions along active fault systems in eastern and central Iran imply dominance of strike-slip faulting in a transpression regime. Conversely, active faults of Azarbaijan (NW Iran) are transpressive. The Alborz and Kopeh-Dagh fault zones are relatively young fault zones in which location of individual active faults is difficult. Aside from raised terraces in the shores of the Oman Sea, information on active faults in the Makran region is scarce.

Although many of the active fault zones of Iran have been determined, many of the individual active faults are still to be studied. With the current level of information on the active faults of Iran, it is not possible to establish a dominant fault regime for the entire country. However, it can be generally stated that, in most cases a transpression regime governs the active faulting in Iran.

Several items included in this map are explained in the following.

ACTIVE FAULTS
An active fault is defined as a fault which has moved repeatedly in recent geological time and has the potential for reactivation in the future. Virtually all major faults in Iran are active and thus have great seismic potential. Since study of active faults has not been detailed enough in Iran, and since some damaging earthquakes with magnitude less than 6.5 may not leave ruptures on the ground surface we can not be certain that an active fault is completely free of earthquake risk. Active faults are classified into the following three types:

Earthquake Fault
During the last 500 years, surface ruptures associated with large earthquakes have appeared and documented in various places in Iran. Most of these ruptures occurred along the active faults which have moved repeatedly in the Quaternary, thus constituting evidence that these active faults have the potential of reactivating in the future.

Seismogenic Fault
An underground fault which generates an earthquake can be called a seismogenic fault in order to distinguish it from an earthquake fault. This type is seismically identified.

Documented Quaternary Fault
This type is recognized to be active based on observation of offset Quaternary landforms, however, they are not known to be seismically active. This could be due to very long recurrence interval along these faults or resulted from creeping, which continuously move slowly without necessarily causing earthquakes.

EARTHQUAKE DATA
The earthquake data are compiled primarily from the catalogue of centroid moment tensor solutions (CMT), which is nearly complete over the interval 1976-2002 for earthquakes greater than $M_w = 5.5$. All earthquakes with magnitude $m_b = 5.0$ or greater that occurred during the period 1948 to 1976 in the area are from Chandra (1984). The solutions denoted by numerals are from Ch. Shirokova (1962); ⊕, Jackson (1992); ⊙, Shirokova (1967) and ⊗, Jackson & McKenzie (1984).

Focal Mechanism Solutions
Focal mechanism solutions of the area's earthquakes have been displayed to reveal mechanisms of seismically active fault zones in Iran. These solutions indicate dominance of thrust and strike-slip faults in a compressive regime for vast majority of earthquakes of Iran.


Assignment of Fault Planes
Mechanism solutions for strike-slip faults indicate two steep nodal planes for all cases. The trend of the fault planes when earthquakes are associated with surface ruptures is known from the offset following recent earthquakes. However, it can be known from distribution of aftershocks. Thus there is no ambiguity between the fault plane and the auxiliary plane in these cases. There is, however, inherent ambiguity between the fault plane and the auxiliary plane when earthquakes have mechanism solutions with a predominance of thrusting and when both nodal planes have nearly the same azimuth, such as most of the mechanism solutions of the earthquakes associated with the Zagros basement faults. For these cases the fault plane is known from geological structures observed in the region. For instance, in the Zagros the nodal plane with shallower dip is assumed to be the fault plane. This is in agreement with the northeast dipping structures observed in the Zagros Mountains.

Slip Vectors
The slip vector of mechanism solutions for which the fault planes are identified, generally indicate a broad convergent zone in Iran. This convergence is in many cases resulted in oblique motion of crustal material with respect to the trend of the seismogenic faults. However, slip vectors for some other recent earthquakes indicate that the convergence is partitioned into pure strike-slip motion and pure thrust faulting. Such partitioning can be seen along the Rudbar, Golba-Sirch and Main Kopeh-Dagh fault zones. The slip vectors for strike-slip faults in central Iran and Azarbaijan (NW Iran) indicate lateral expulsion of crustal material due east-northeast. However, we could not assign slip vectors to the mechanism solutions of the earthquakes associated with the Zagros blind thrusts, as precise location of epicenters is not known with respect to the location of the seismogenic faults.

Compressive Axes
The direction of the horizontal projection of the maximum principal stress deduced from the focal mechanism solution of the area's earthquakes reveals a characteristic regional stress field along the colliding boundary. However, the distribution of horizontal compressive stress axes in different parts of Iran indicates that the local stress field is not everywhere consistent with the relative motion of the Iranian crust with respect to Eurasia. The compressive stress axes along the faults in central Iran are approximately N-S, parallel to relative motion of the two plates. However, they are NE-SW for the vast majority of earthquakes in Iran, which is not consistent with the direction of relative motion. For the earthquakes in the Zagros Mountains and the Kopeh-Dagh region, the compressive axes have roughly a N-N-E direction, and on the average, they are nearly perpendicular to the trend of the geological structures. In eastern Iran shortening is distributed over several active deforming zones. In eastern Iran shortening is distributed over the Makran subduction complex (up to 1.8 cm/yr) and the Kopeh-Dagh Mts. (about 0.5 cm/yr). To the west, shortening is distributed over the Zagros (about 0.8 cm/yr) and Alborz Mts. (about 0.1 cm/yr).

GPS VELOCITIES
GPS horizontal velocities are illustrated for the Iranian GPS stations in Eurasia-fixed reference frame for the period 1999-2001 (adopted from Nilforoushan et al., 2003). Preliminary motion estimates show that Arabia moves at 2.1±2.5 cm/yr due north relative to Eurasia. However, deformation is distributed differently over several active deforming zones. In eastern Iran shortening is distributed over the Makran subduction complex (up to 1.8 cm/yr) and the Kopeh-Dagh Mts. (about 0.5 cm/yr). To the west, shortening is distributed over the Zagros (about 0.8 cm/yr) and Alborz Mts. (about 0.1 cm/yr).

REFERENCES
This map is based on Ambraseys & Melville, 1982; Berberian, 1976, 1994, 1995; Chandra, 1984; Dziewonski et al., 1981; Hessami & Jamali, 1996; Huber, 1977; Mohajer-Ahaji et al., 1982; Nilforoushan et al., 2003 and other published sources and available data up to 2003.



گسلهای فعال ایران را می توان به عنوان شاخص مهم زمینشناسی کشور پسته در ایران دانست که حاصل برخورد قاره ای بین صفحات عربستان و اوراسیا می باشد. از آنجا که در مطالعه گسلهای فعال ایران به جز در معدود مواردی، از روشهای ژئومورفولوژیکی و چینه شناسی کواترنری استفاده نشده است. معرفی گسلهای فعال بر پایه تحقیق زلزله در داده های زمین لرزه ای صورت پذیرفته است. با وجود این، شناسایی گسلهای فعال براساس داده های لرزه ای در مناطق برخوردی قاره ای نظیر ایران، که از نظر ساختاری پیچیده و نامهمکن می باشد، بسیار دشوار است. این موضوع از آنجا ناشی می شود که لرزه خیزی در چنین مناطقی به تعداد معدودی گسل منفرد و مجزا محدود نمی باشد بلکه در زمینه فعالیت پهنه های گسلی به پهنای چند صد کیلومتری روی می دهند. به این ترتیب، به دلیل تراکم زیاد گسلهای فعال در ایران و وقت کم داده های مپارزه ای، گسلهای فعال بزرگ مربوط به گسلهای فعال و کانون یک گسل نسبت داده شده است. علاوه بر این، فلات نشنه های بزرگ میسای مربوط به گسلهای فعال و کانون میسای با دقت بالا به ناتیوی ما بر برقراری یک نشانه یک به یک بین زمین لرزه ها و گسلهای فعال برآیند انجامیده است. با این وصف روشن است که حتی با فرض داشتن نقشه های بسیار دقیق از کانون سلسله زمین لرزه ها، شناسایی تمام گسلهای فعال امکانپذیر نمی باشد. این امر از آنجا ناشی می شود که: 1- بسیاری از زمین لرزه ها در نتیجه فعالیت گسلهایی روی می دهند که در ریز رسوبات مدفون می باشند و مستقیماً به سطح زمین نمی رسند (مانند زمین لرزه های منطقه زاگرس). 2- حرکات تدریجی زمین در طیف گسلهای فعال در یک دوره زمانی طولانی (از ریز زمین لرزه ها) با ایجاد زمین لرزه بزرگ همراه نمی باشد.

این نقشه گسلهای فعال اصلی ایران، سرچشمه های لرزه خیزی شده با استفاده از GPS و بین بردار لرزه ای و محور فشاری که از حل مکانیک کانونی زمین لرزه های رده در مناطق مختلف ایران به دست آمده، نشان داده شده است. اگرچه گسلهای فعال در زاگرس مدفون می باشند ولی حل مکانیک کانونی زمین لرزه ها در این منطقه دلالت بر وجود گسلهای منگوس و امتدادده در پی سنگ آن دارد. در سایر مناطق ایران، بیشتر گسلهای فعال به سطح زمین می رسند. حل مکانیک کانونی زمین لرزه های رده در امتداد سیستم های گسلی شرق و مرکز ایران به نفع می رساند. گسلهای فعال در یک ریز زمین لرزه خیزی-برشی دلالت دارند. در صورتی که گسلهای فعال منطقه آذربایجان از نوع امتدادده بوده که در یک ریز زمین لرزه خیزی-برشی امتدادده شده اند. در مناطق لرزه خیز و گسلهای فعال گسلها وسیع و مترکام بوده، به طوری که شناسایی فعال منفرد و مجزا به دشواری صورت گرفته است. در نهایت، صرف نظر از روشهای ایش یافته در طول سواحل دریای مازندران با اطلاعات ما در مورد گسلهای فعال ناحیه مکران بسیار اندک است.

اگرچه بسیاری از گسلهای فعال ایران به خوبی شناسایی شده اند با وجود این، هنوز امکان شناسایی و مطالعه بسیاری از گسلهای فعال وجود دارد. به سطح اطلاعات فعلی از گسلهای فعال نمی توان زمین گسلی غالب در کل ایران را زمین گسلی اصلی می نامد. در زیر، مواردی که این نقشه را شامل می شود توضیح داده شده است.

گسلهای فعال
گسل فعال، گسلی است که بلوک های طریفان آن در طی زمانهای اخیر زمین شناسی (کواترنری-پلاستوسن) به حرکت درآمده باشد. براساس فعالیت مجدد در آینده را داشته باشد. تمام گسلهای اصلی-مترکام ایران فعال و از دستبند لرزه خیزی، با این معیار شناخته می شوند. از آنجا که از یک سو شناسایی گسلهای فعال ایران شناسایی شده اند و از سوی دیگر بسیاری از زمین لرزه های محرب با بزرگی کمتر از 5.5 با گسیختگی همراه نبوده اند، نباید منتظرمانی را که در روی این نقشه کاری از گسل فعال نشان داده شده، کاری از خطر زمین لرزه دانست. گسلهای فعال ایران به سه رده زیر تقسیم شده اند.

گسل زمین لرزه ای
گسیختگی های سطح زمین که در طی زمین لرزه های بزرگ 5.0-6.0 ساله اخیر در مناطق مختلف ایران ایجاد شده اند، به این گروه تعلق دارند. بیشتر گسیختگی های شناسایی شده در طول گسلهای فعال روی داده اند که کرات در طی زمان کواترنری دیده جا شده اند. این وقت گسلی است که این گسلهای زمین لرزه ای، با شناسایی فعالیت مجدد در آینده را دارا می باشد.

گسل لرزه زا
گسلهایی پهنایی که باعث ایجاد زمین لرزه می شوند، ولی با گسیختگی سطحی همراه نمی باشند. گسل لرزه زا نامیده می شوند تا بدینوسیله از گسل زمین لرزه ای متمایز شوند. در شناسایی این دسته از گسلها از لرزه خیزی بهره گرفته شده است.

گسل کواترنری
شناسایی فعالیت این گروه از گسلها برپایه مشاهده جلوه جایی مورخ سلسله کواترنری است. شناسایی سنی بر لرزه خیزی این گسلها در دست نیست این موضوع می تواند ناشی از دوره بازگشت بسیار طولانی زمین لرزه و با خزش بر روی این گسلها باشد.

داده های زمین لرزه ای
داده های زمین لرزه ای این نقشه عمدتاً از فعالیت زمین لرزه های بزرگتر از $M_w = 5.5$ منطبق از کاتالوگ CMT که در دوره زمانی 1976-2002 تقریباً کامل است، به دست آمده است. تمام زمین لرزه های بزرگتر از $M_w = 5.5$ که در سالهای 1976-1978 در ایران روی داده اند از چاندر (1984) گردیده است. مکانیک حالی که با داده شده اند در منابع زیر درج شده اند: (1) شیکووا (1962)، (2) جکسون (1992)، (3) شیکووا (1967) و (4) جکسون و مکینزی (1984).

حل مکانیک کانونی
به منظور نشان دادن مکانیک پهنه های فعال لرزه ای، حل مکانیک کانونی زمین لرزه های روی داده در ارتباط با گسلها نشان داده شده است. مکانیک کانونی زمین لرزه ها نشانه آن است که اکثر قریب به اتفاق گسلهای ایران از نوع تراشید و امتدادده می باشد که در یک ریز زمین لرزه خیزی-برشی فشاری-برشی فشاری-برشی فشاری است.

تجزیه ضحکات گسلی
حل مکانیک کانونی زمین لرزه های مربوط به گسلهای امتدادده در تمام موارد دو سطح خنثی برشید را نشان می دهد. به جز زمین لرزه های بزرگتر از $M_w = 5.5$ که در مناطق مختلف ایران شناسایی شده اند، زمین لرزه های کوچکتر از $M_w = 5.5$ را با زمین لرزه های بزرگتر از $M_w = 5.5$ در مناطق مختلف ایران شناسایی شده اند. زمین لرزه های کوچکتر از $M_w = 5.5$ در مناطق مختلف ایران شناسایی شده اند. زمین لرزه های کوچکتر از $M_w = 5.5$ در مناطق مختلف ایران شناسایی شده اند.

بردار لرزه ای
بردار لرزه ای زمین لرزه ای این نقشه عمدتاً از فعالیت زمین لرزه های بزرگتر از $M_w = 5.5$ منطبق از کاتالوگ CMT که در دوره زمانی 1976-2002 تقریباً کامل است، به دست آمده است. تمام زمین لرزه های بزرگتر از $M_w = 5.5$ که در سالهای 1976-1978 در ایران روی داده اند از چاندر (1984) گردیده است. مکانیک حالی که با داده شده اند در منابع زیر درج شده اند: (1) شیکووا (1962)، (2) جکسون (1992)، (3) شیکووا (1967) و (4) جکسون و مکینزی (1984).

محورهای فشاری
جهت تصویر فقی حرکت افقی استیکانهای GPS از سال 1999 تا 2001 (گرفته شده از Nilforoushan et al., 2003) در این نقشه سرعت حرکت افقی استیکانهای GPS ایران در یک سیستم مختصاتی که مرکز آن در روی صفحه ثابت شده، از طریق فرض گردیده برای یک دوره 10 ساله (1990-2000) به تصویر کشیده شده است. (نیل فرشان و همکاران، 2003). زمین لرزه ای هر حرکت مکانیک شده شده است. جهت که در منطقه عربستان با سرعتی بین 1.5 تا 2.5 سانتی متر در سال به طرف شمال شرق حرکت است. این با این فرض زمین لرزه ای در ایران، از طرف شمال غربی و به پهنای فعال زمینشناسی توزیع یافته است. در شرق ایران، کوه دشتی بر روی دو بخش مختلف طور متوسط حدود 2 سانتی متر در سال به طرف شمال شرقی حرکت می کند. در منطقه های جنوب شرقی زاگرس از جهت عمومی شمال-شمال شرقی الحرف حاصل کرده، به سمت شمال غرب امتداد یافته و ولی همباز بر راستای ساختارهای منطقه مدفون می باشد. جنوب غربی لرزه خیزی، هنوز تنها بخش از ایران است که در آن جهت محورهای تنش شرقی-غربی می باشد.

سرعتهای اندازه گیری شده با استفاده از GPS
در این نقشه سرعت حرکت افقی استیکانهای GPS از سال 1999 تا 2001 (گرفته شده از Nilforoushan et al., 2003) در این نقشه سرعت حرکت افقی استیکانهای GPS ایران در یک سیستم مختصاتی که مرکز آن در روی صفحه ثابت شده، از طریق فرض گردیده برای یک دوره 10 ساله (1990-2000) به تصویر کشیده شده است. (نیل فرشان و همکاران، 2003). زمین لرزه ای هر حرکت مکانیک شده شده است. جهت که در منطقه عربستان با سرعتی بین 1.5 تا 2.5 سانتی متر در سال به طرف شمال شرق حرکت است. این با این فرض زمین لرزه ای در ایران، از طرف شمال غربی و به پهنای فعال زمینشناسی توزیع یافته است. در شرق ایران، کوه دشتی بر روی دو بخش مختلف طور متوسط حدود 2 سانتی متر در سال به طرف شمال شرقی حرکت می کند. در منطقه های جنوب شرقی زاگرس از جهت عمومی شمال-شمال شرقی الحرف حاصل کرده، به سمت شمال غرب امتداد یافته و ولی همباز بر راستای ساختارهای منطقه مدفون می باشد. جنوب غربی لرزه خیزی، هنوز تنها بخش از ایران است که در آن جهت محورهای تنش شرقی-غربی می باشد.

منابع
برای تهیه این نقشه از تمام منابع منتشر شده در سال 2003-2004، بوزمانی از بهره بهره گرفته شده است. امبرسیز و ملویل (1982)، بربریان (1976، 1994، 1995) و چاندر (1984)، حاسمی و جمالی (1996)، دیوونسی و همکاران (1981)، مپاروشیمی و همکاران (1982)، نیل فرشان و همکاران (2003) و هوبر (1977) و هوبر (1977).